

## Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit [www.landfire.gov](http://www.landfire.gov). Please direct questions to [helpdesk@landfire.gov](mailto:helpdesk@landfire.gov).

### Potential Natural Vegetation Group (PNVG)

R#ABAMlw Pacific Silver Fir--Low Elevation

#### General Information

**Contributors** (additional contributors may be listed under "Model Evolution and Comments")

**Modelers**

Karen Kopper karen\_kopper@nps.gov  
Jane Kertis jkertis@fs.fed.us

**Reviewers**

Miles Hemstrom mhemstrom@fs.fed.us

**Vegetation Type**

Forested

**General Model Sources**

- Literature
- Local Data
- Expert Estimate

**Rapid Assessment Model Zones**

- California
- Pacific Northwest
- Great Basin
- South Central
- Great Lakes
- Southeast
- Northeast
- S. Appalachians
- Northern Plains
- Southwest
- N-Cent. Rockies

**Dominant Species\***

ABAM  
TSHE  
PSME  
ABPR

**LANDFIRE Mapping Zones**

1	8
2	9
7	

#### Geographic Range

The Pacific Silver fir PNVG occurs on the western slopes of the Cascades from British Columbia south to the Rogue and Umpqua River divide in the Southern Cascades. It can occur east of the Cascades crest south of Mt. Hood. This low elevation type (with a mixed-severity regime) is also found throughout the Ross Lake Drainage and on the eastern slopes of the North Cascades.

#### Biophysical Site Description

The Pacific Silver fir forests described in this PNVG occur at lower elevations within the Pacific Silver fir zone (450 - 800 meters in the north, 1600 - 1800 meters in the south). Pacific silver fir grows on soils developed from nearly every type of parent material found in the Northwest. Growth rates for Pacific silver fir are greatest at low elevations on fine-textured residual soils from sedimentary and basaltic rocks.

#### Vegetation Description

Pacific Silver fir is the dominant and climax tree species in the mature canopy, which it shares with a wide-variety of conifers depending upon locale. Douglas-fir and western hemlock are codominant throughout the range. Subalpine fir, Grand fir, White pine and Englemann spruce are common around Mount Adams and in parts of Oregon. Noble fir is commonly associated to the PNVG from Mount Rainier and south, and Lodgepole pine is common in the North Cascades. The understory is predominantly composed of a lush to moderate layer (depending upon the amount of moisture) of heath shrubs, forbs, ferns and mosses.

#### Disturbance Description

Pacific Silver fir forests in this PNVG are characterized by infrequent mixed severity fire regimes occurring at greater than 100 years. These fires produced variably sized patches throughout the landscape. Landscapes were reset at intervals greater than 200 years through stand-replacing events. Avalanches and wind events are also common disturbances in this PNVG.

#### Adjacency or Identification Concerns

The Pacific Silver fir PNVG occurs above the Western hemlock forests. This low elevation type is replaced

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

by moister and cooler plant associations (Silver fir - Mountain hemlock) at higher elevations. This PNVG is distinguished from the high elevation Pacific Silver Fir type (R#ABAMup) by elevation breaks: the low elevation type occurs below 800m in the north and 1800m in the south.

**Scale Description**

**Sources of Scale Data**  Literature  Local Data  Expert Estimate

Mixed-severity fire events occur on the scale of 1000's of acres, and are usually stand-replacing. Infrequent avalanches and wind disturbances also occur in this PNVG, but these disturbances more frequently occur at scales of 10's and 100's of acres.

**Issues/Problems**

Although windthrow and avalanches are known disturbances in this PNVG, the nature of these disturbances are based upon opinion only, and should be checked for validity. These disturbances were not modeled explicitly.

**Model Evolution and Comments**

Probability of mixed severity events was reduced (from 0.003 to 0.0025) during peer review and had negligible effect on the percentages in each class, but raised the fire return of 'All fire' from 111 to 146 years. (Component fire Interval values differed, too: Replacement fire from 300 to 350; and mixed fire from 175 to 250).

**Succession Classes\*\***  
*Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).*

**Class A 15%**

Early1 All Struct

**Description**

The early seral stand consists of low heath shrubs, seedlings and saplings. Sometimes, competition can keep the trees at no greater than 2" dbh. Silver fir is seral, however Douglas fir, western white pine and noble fir may also be seral, and, where they occur, they grow more quickly than silver fir. This stage can last for decades (up to 40 years). [A replacement fire sets the stand back forty years and occurs about every 300-350 years.]

**Dominant Species\* and Canopy Position**

PSME  
 ABPR  
 PIMO3  
 ABAM

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	0 %	60 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class B 20%**

Mid1 Closed

**Description**

Canopy closure occurs in the middle-aged stand. Trees grow to 20" dbh. The early seral species continue to dominate the stand, and the midstory fills in with increasingly larger amounts of Pacific Silver fir and a variety of more shade tolerant conifers

**Dominant Species\* and Canopy Position**

ABAM  
 PSME  
 PIMO3  
 ABPR

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	60 %	100 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

(Engelmann spruce, western hemlock, western red cedar).  
 [After 80 years in this class it would proceed to class E. Mixed fire has equal probability (~300-350 years for each type) to move this class to C (mid-open) as to making no change (stays in B). Replacement fire may also occur at the same probability.]

**Class C 3%**

Mid1 Open  
**Description**

Openings in the canopy are created by mixed severity fire [about every 300-350 years and maintains it in Class C]. Fire resistant Douglas-fir and Western white pine remain. They continue to grow to 20" dbh. Lodgepole pine (where it occurs) and Silver fir regenerate in the openings as the stand fills back in. [After 80 in this class it proceeds to Class E. Replacement fire about every 300-350 years.]

**Dominant Species\* and Canopy Position**

PSME  
 PIMO3  
 ABAM  
 PICO

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

	<i>Min</i>	<i>Max</i>
<i>Cover</i>	20 %	60 %
<i>Height</i>	no data	no data
<i>Tree Size Class</i>	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class D 10%**

Late1 Open  
**Description**

Douglas fir is resistant to mixed severity fire events. The overstory trees average 45" dbh. Silver fir and western hemlock regenerate in the understory. [After 50 years in this class it fills in to become class E. However, mixed fire can maintain it in D (300-350 years), and replacement fire may occur at the same frequency.]

**Dominant Species\* and Canopy Position**

PSME  
 ABAM  
 TSHE

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

	<i>Min</i>	<i>Max</i>
<i>Cover</i>	20 %	60 %
<i>Height</i>	no data	no data
<i>Tree Size Class</i>	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Class E 52%**

Late I Closed

**Description**

Silver fir is dominant in the late seral stand. The trees average 45" dbh, and range from 21" - 150" dbh. Douglas fir and Western hemlock are codominant. [Fire and Insect/disease occur in these old stands. Mixed fire could maintain the closed stand or move the stand to open-late (class D) - each occurs about every 300-350 years. Similarly, Insect/disease could either replace the stand (class A), or open it up to class D but at a low frequency (about every 1000 years.)]

**Dominant Species\* and Canopy Position**

ABAM  
PSME  
TSHE

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	60 %	100 %
Height	no data	no data
Tree Size Class	no data	

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Fuel Model** no data

**Disturbances**

**Disturbances Modeled**

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

**Fire Regime Group: 3**

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

**Fire Intervals (FI)**

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

**Historical Fire Size (acres)**

Avg: no data  
Min: no data  
Max: no data

**Sources of Fire Regime Data**

- Literature
- Local Data
- Expert Estimate

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	350	100	800	0.00286	46
Mixed	300	100	400	0.00333	54
Surface					
All Fires	162			0.00620	

**References**

Agee, James K. 1993. Fire Ecology of Pacific Northwest Forests. Island Press

Franklin, Jerry F. and C. T. Dyrness 1988. Natural Vegetation of Oregon and Washington. Oregon State University Press

Hemstrom, Miles Arthur 1979. A recent disturbance history of forest ecosystems at Mount Rainier National Park. Phd dissertation. Oregon State University.

Hemstrom, M.A., S.E. Logan, and W. Pavlat. 1987. Plant association and management guide, Willamette National Forest. Publication R6-Ecol-257b-1986. USDA Forest Service, Pacific Northwest Region, Portland,

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

OR. 312 p.

Lillybridge, T.R.; Kovalchik, B.L.; Williams, C.K.; Smith, B.G. 1995. Field guide for forested plant associations of the Wenatchee National Forest. PNW-GTR-359. Portland, OR: USDA Forest Service, Pacific Northwest Research Station. 337p.