

Supplemental Material

Rie Yanagisawa¹, Hirohisa Takano¹, Ken-ichiro Inoue¹, Eiko Koike¹, Kaori Sadakane²,
Takamichi Ichinose²

1: Environmental Health Sciences Division, National Institute for Environmental
Studies, Tsukuba, Japan.

2: Department of Health Sciences, Oita University of Nursing and Health Sciences, Oita,
Japan.

Address for correspondence:

Hirohisa Takano, MD, PhD

Environmental Health Sciences Division


National Institute for Environmental Studies

16-2 Onogawa, Tsukuba, 305-8506, Japan

Tel: +81-29-850-2336, Fax: +81-29-850-2334

E-mail: htakano@nies.go.jp, yanagi@nies.go.jp


Supplemental Material, Figure 1



Maternal exposure to DEHP during fetal periods does not affect the symptoms of atopic dermatitis-like skin lesions in offspring. We evaluated clinical scores 24 hours after each intradermal injection. Data are the means \pm SE of 7-12 animals per group.

*; $p < 0.01$; Dp-treated groups vs. nontreated group and saline(+vehicle) group

Supplemental Material, Figure 2



Maternal exposure to DEHP during neonatal periods enhances the symptoms of atopic dermatitis-like skin lesions in offspring. We evaluated clinical scores 24 hours after each intradermal injection.

Data are the means \pm SE of 12 animals per group.

*; $p < 0.01$; Dp-treated groups vs. nontreated group and saline(+vehicle) group.

#; $p < 0.01$; Dp(+DEHP 100 µg) group vs. Dp(+vehicle) group