

# Medical Checklist

**Mission Operations Directorate  
Operations Division**

**Generic, Rev K  
May 12, 2006**

**NOTE**

For STS-121 and subsequent flights

National Aeronautics and  
Space Administration

**Lyndon B. Johnson Space Center**  
Houston, Texas



**Verify this is the correct version for the pending operation (training, simulation or flight).**  
Electronic copies of FDF books are available. URL: <http://mod.jsc.nasa.gov/do3/FDF/index.html>



MEDICAL CHECKLIST

GENERIC, REV K (May 12, 2006)

## PCN-2 (Feb 2, 2007) Sheet 1 of 1

List of Implemented Change Requests (482s):

MED-320  
MED-321  
MED-322  
MED-323  
MED-324

**NOTE**

For STS-117 and subsequent flights

Incorporate the following:

1. Replace iii thru viii
2. Replace 1-1 and 1-2, 1-7 and 1-8, 1-11 thru 1-20, 1-25 and 1-26, 1-31 and 1-32, 1-35 and 1-36, 1-39 and 1-40, 1-43 and 1-44
3. Replace 4-3 and 4-4
4. Replace 7-7 and 7-8
5. Replace 8-1 thru 8-4, 8-9 thru 8-12, 8-19 and 8-20, 8-23 and 8-24
6. Replace section 9 (8 pages)

Prepared by:

  
Book Manager

Approved by:

  
Manager, Shuttle Procedures Management

Accepted by:

  
FDF Manager

Encl: 54 pages

**File this PCN immediately behind the front cover as a permanent record**



MEDICAL CHECKLIST

GENERIC, REV K (May 12, 2006)

## PCN-1 (Nov 10, 2006) Sheet 1 of 1

List of Implemented Change Requests (482s):


MED-317  
MED-318  
MED-319**NOTE**

For STS-116 and subsequent flights


Incorporate the following:

1. Replace iii thru vi
2. Replace 1-7 and 1-8
3. Replace 2-3 and 2-4
4. Replace 3-3 and 3-4, 3-11 and 3-12
5. Replace 5-5 and 5-6
6. Replace 6-5 and 6-6, 6-11 and 6-12
7. Replace 8-5 thru 8-12, 8-15 and 8-16, 8-19 and 8-20

Prepared by:

  
Book Manager

Approved by:

  
Manager, Shuttle Procedures Management

Accepted by:

  
FDF Manager

Encl: 30 pages

**File this PCN immediately behind the front cover as a permanent record**




MISSION OPERATIONS DIRECTORATE

**MEDICAL CHECKLIST**

GENERIC, REVISION K


May 12, 2006


PREPARED BY:


  
Jerry Woodard  
Book Manager

APPROVED BY:

  
Kimberly R. Johnson  
Manager, Shuttle  
Procedures Management

  
B. Scott Coleman  
Wyle Life Sciences

  
Rainer K. Effenhauser, M.D.  
Lead Shuttle Flight Surgeon  
Space Medicine –  
Medical Operations

  
James M. Polk, D.O.  
Manager  
Space Medicine –  
Medical Operations

  
Michael T. Hurt  
FDF Manager

This document is under the configuration control of the Crew Procedures Control Board (CPCB). All proposed changes must be submitted on FDF Workflow Crew Procedure Change Request (CR) to DO3/FDF Manager.

Additional distribution of this book, for official use only, may be requested in writing to DO3/FDF Manager. The request must include justification and requester's name, organization, position, and phone number. Contractor requests are made through the NASA or DOD organization supported. Deletions, reduction in quantity, or change of address may be submitted to DO3/FDF Management Office, 281-244-1184.

MED/ALL/GEN K

Incorporates the following:		
482#:	MED-0304	MED-0313
	MED-0306A	MED-0314
	MED-0310	MED-0315
	MED-0311	MED-0316
	MED-0312	

AREAS OF TECHNICAL RESPONSIBILITY

Book Manager	DO35/J. Woodard	281-483-9685
Backup Book Manager	DO352/M. Bruce	281-483-6083
Biomedical Engineer	SD24/B. Coleman	281-483-5285

This Medical Checklist (Med C/L, JSC-48031) now supports all missions including Extended Duration Orbiter (EDO). The EDO Medical Checklist (EDO Med C/L, JSC-48092) information has been incorporated and is no longer printed separately.



MEDICAL CHECKLIST  
**LIST OF EFFECTIVE PAGES**

GENERIC 01/25/88  
 REV K 05/12/06  
 PCN-1 11/10/06  
 PCN-2 02/02/07

Sign Off .....	* ALL/GEN K	1-20 .....	ALL/GEN K
ii .....	* ALL/GEN K	1-21 .....	ALL/GEN K
iii .....	* ALL/GEN K,2	1-22 .....	ALL/GEN K
iv .....	* ALL/GEN K,2	1-23 .....	ALL/GEN K
v .....	* ALL/GEN K,2	1-24 .....	ALL/GEN K
vi .....	* ALL/GEN K,2	1-25 .....	ALL/GEN K,2
vii .....	ALL/GEN K,2	1-26 .....	ALL/GEN K
viii .....	ALL/GEN K	1-27 .....	ALL/GEN K
ix .....	ALL/GEN K	1-28 .....	ALL/GEN K
x .....	ALL/GEN K	1-29 .....	ALL/GEN K
xi .....	ALL/GEN K	1-30 .....	ALL/GEN K
xii .....	ALL/GEN K	1-31 .....	ALL/GEN K
1-1 .....	ALL/GEN K,2	1-32 .....	ALL/GEN K,2
1-2 .....	ALL/GEN K	1-33 .....	ALL/GEN K
1-3 .....	ALL/GEN K	1-34 .....	ALL/GEN K
1-4 .....	ALL/GEN K	1-35 .....	ALL/GEN K,2
1-5 .....	ALL/GEN K	1-36 .....	ALL/GEN K,2
1-6 .....	ALL/GEN K	1-37 .....	ALL/GEN K
1-7 .....	ALL/GEN K,2	1-38 .....	ALL/GEN K
1-8 .....	ALL/GEN K	1-39 .....	ALL/GEN K,2
1-9 .....	ALL/GEN K	1-40 .....	ALL/GEN K
1-10 .....	ALL/GEN K	1-41 .....	ALL/GEN K
1-11 .....	ALL/GEN K	1-42 .....	ALL/GEN K
1-12 .....	ALL/GEN K,2	1-43 .....	ALL/GEN K
1-13 .....	ALL/GEN K,2	1-44 .....	ALL/GEN K,2
1-14 .....	ALL/GEN K	1-45 .....	ALL/GEN K
1-15 .....	ALL/GEN K,2	1-46 .....	ALL/GEN K
1-16 .....	ALL/GEN K,2	2-1 .....	ALL/GEN K
1-17 .....	ALL/GEN K,2	2-2 .....	ALL/GEN K
1-18 .....	ALL/GEN K	2-3 .....	ALL/GEN K,1
1-19 .....	ALL/GEN K,2	2-4 .....	ALL/GEN K

\* – Omit from flight book

2-5.....	ALL/GEN K	4-13.....	ALL/GEN K
2-6.....	ALL/GEN K	4-14.....	ALL/GEN K
2-7.....	ALL/GEN K	4-15.....	ALL/GEN K
2-8.....	ALL/GEN K	4-16.....	ALL/GEN K
3-1.....	ALL/GEN K	4-17.....	ALL/GEN K
3-2.....	ALL/GEN K	4-18.....	ALL/GEN K
3-3.....	ALL/GEN K,1	4-19.....	ALL/GEN K
3-4.....	ALL/GEN K	4-20.....	ALL/GEN K
3-5.....	ALL/GEN K	4-21.....	ALL/GEN K
3-6.....	ALL/GEN K	4-22.....	ALL/GEN K
3-7.....	ALL/GEN K	4-23.....	ALL/GEN K
3-8.....	ALL/GEN K	4-24.....	ALL/GEN K
3-9.....	ALL/GEN K	4-25.....	ALL/GEN K
3-10.....	ALL/GEN K	4-26.....	ALL/GEN K
3-11.....	ALL/GEN K	4-27.....	ALL/GEN K
3-12.....	ALL/GEN K,1	4-28.....	ALL/GEN K
3-13.....	ALL/GEN K	4-29.....	ALL/GEN K
3-14.....	ALL/GEN K	4-30.....	ALL/GEN K
3-15.....	ALL/GEN K	4-31.....	ALL/GEN K
3-16.....	ALL/GEN K	4-32.....	ALL/GEN K
3-17.....	ALL/GEN K	4-33.....	ALL/GEN K
3-18.....	ALL/GEN K	4-34.....	ALL/GEN K
3-19.....	ALL/GEN K	4-35.....	ALL/GEN K
3-20.....	ALL/GEN K	4-36.....	ALL/GEN K
3-21.....	ALL/GEN K	4-37.....	ALL/GEN K
3-22.....	ALL/GEN K	4-38.....	ALL/GEN K
3-23.....	ALL/GEN K	4-39.....	ALL/GEN K
3-24.....	ALL/GEN K	4-40.....	ALL/GEN K
4-1.....	ALL/GEN K	4-41.....	ALL/GEN K
4-2.....	ALL/GEN K	4-42.....	ALL/GEN K
4-3.....	ALL/GEN K,2	4-43.....	ALL/GEN K
4-4.....	ALL/GEN K	4-44.....	ALL/GEN K
4-5.....	ALL/GEN K	5-1.....	ALL/GEN K
4-6.....	ALL/GEN K	5-2.....	ALL/GEN K
4-7.....	ALL/GEN K	5-3.....	ALL/GEN K
4-8.....	ALL/GEN K	5-4.....	ALL/GEN K
4-9.....	ALL/GEN K	5-5.....	ALL/GEN K,1
4-10.....	ALL/GEN K	5-6.....	ALL/GEN K
4-11.....	ALL/GEN K	5-7.....	ALL/GEN K
4-12.....	ALL/GEN K	5-8.....	ALL/GEN K

5-9.....	ALL/GEN K	8-7.....	ALL/GEN K,1
5-10.....	ALL/GEN K	8-8.....	ALL/GEN K
5-11.....	ALL/GEN K	8-9.....	ALL/GEN K
5-12.....	ALL/GEN K	8-10.....	ALL/GEN K,2
5-13.....	ALL/GEN K	8-11.....	ALL/GEN K,2
5-14.....	ALL/GEN K	8-12.....	ALL/GEN K
5-15.....	ALL/GEN K	8-13.....	ALL/GEN K
5-16.....	ALL/GEN K	8-14.....	ALL/GEN K
5-17.....	ALL/GEN K	8-15.....	ALL/GEN K,1
5-18.....	ALL/GEN K	8-16.....	ALL/GEN K,1
5-19.....	ALL/GEN K	8-17.....	ALL/GEN K
5-20.....	ALL/GEN K	8-18.....	ALL/GEN K
6-1.....	ALL/GEN K	8-19.....	ALL/GEN K,2
6-2.....	ALL/GEN K	8-20.....	ALL/GEN K
6-3.....	ALL/GEN K	8-21.....	ALL/GEN K
6-4.....	ALL/GEN K	8-22.....	ALL/GEN K
6-5.....	ALL/GEN K	8-23.....	ALL/GEN K
6-6.....	ALL/GEN K,1	8-24.....	ALL/GEN K,2
6-7.....	ALL/GEN K	8-25.....	ALL/GEN K
6-8.....	ALL/GEN K	8-26.....	ALL/GEN K
6-9.....	ALL/GEN K	9-1.....	ALL/GEN K
6-10.....	ALL/GEN K	9-2.....	ALL/GEN K,2
6-11.....	ALL/GEN K,1	CC 9-3.....	ALL/GEN K,2
6-12.....	ALL/GEN K	CC 9-4.....	ALL/GEN K,2
6-13.....	ALL/GEN K	CC 9-5.....	ALL/GEN K,2
6-14.....	ALL/GEN K	CC 9-6.....	ALL/GEN K,2
7-1.....	ALL/GEN K	CC 9-7.....	ALL/GEN K,2
7-2.....	ALL/GEN K	CC 9-8.....	ALL/GEN K,2
7-3.....	ALL/GEN K		
7-4.....	ALL/GEN K		
7-5.....	ALL/GEN K		
7-6.....	ALL/GEN K		
7-7.....	ALL/GEN K		
7-8.....	ALL/GEN K,2		
8-1.....	ALL/GEN K		
8-2.....	ALL/GEN K,2		
8-3.....	ALL/GEN K,2		
8-4.....	ALL/GEN K		
8-5.....	ALL/GEN K,1		
8-6.....	ALL/GEN K		

MEDICAL CUE CARDS

<u>Title</u>	<u>Ref. Page</u>	<u>Card No.</u>
MEDICATION DATA LOG		
(Front).....	CC 9-3	MED-1a/O/C
(Back).....	CC 9-4	MED-1b/O/C
AIRWAY (ILMA)		
(Front).....	CC 9-5	MED-2a/O/D
(Back).....	CC 9-6	MED-2b/O/B
RESUSCITATOR		
(Front).....	CC 9-7	MED-4a/O/A
(Back).....	CC 9-8	MED-4b/O/A

<u>CONTENTS</u>	<u>PAGE</u>
<u>SECTION 1 – EMERGENCY</u>	
IMMEDIATE LIFE SUPPORT .....	1-3
BREATHING DIFFICULTY .....	1-3
CHOKING .....	1-4
OBSTRUCTED AIRWAY – CONSCIOUS ADULT .....	1-4
OBSTRUCTED AIRWAY – UNCONSCIOUS ADULT ..	1-5
CPR (CARDIOPULMONARY RESUSCITATION) .....	1-6
PATIENT RESTRAINT .....	1-6
CARDIAC MONITORING .....	1-8
ORAL AIRWAY .....	1-11
OXYGEN – SUPPLEMENTAL .....	1-12
INTUBATION (ILMA) .....	1-12
RESUSCITATOR (see RESUSCITATOR REFERENCE CARD, CC 9-7)	
INTUBATION, ENDOTRACHEAL .....	1-12
CRICOTHYROTOMY .....	1-19
OTHER EMERGENCIES .....	1-27
ABDOMINAL INJURY .....	1-27
ALLERGIC REACTION – SEVERE REACTION .....	1-29
BURNS – MAJOR .....	1-32
CHEST INJURY .....	1-35
CHEST PAIN .....	1-36
EYE – PENETRATING OBJECT .....	1-38
HEAD OR NECK INJURY .....	1-39
SEIZURES .....	1-42
SHOCK .....	1-44
<u>SECTION 2 – A THRU B</u>	
ABDOMINAL INJURY (see EMERGENCY, 1-27)	
ABDOMINAL PAIN .....	2-2
ALLERGIC REACTION .....	2-3
MILD-TO-MODERATE REACTION .....	2-3
SEVERE REACTION (see EMERGENCY, 1-29)	
ALTITUDE SICKNESS .....	2-4
BACK PAIN .....	2-4
BENDS (see DECOMPRESSION SICKNESS, 3-4)	
BLADDER INFECTION .....	2-5
BREATHING DIFFICULTY (see EMERGENCY, 1-3)	
BROKEN BONE (see FRACTURE/DISLOCATION, 4-3)	
BRONCHITIS (COUGH) .....	2-7
BURNS – MAJOR (see EMERGENCY, 1-32)	
BURNS – MINOR .....	2-8

SECTION 3 – C THRU E

CARDIAC MONITORING (see CPR  
(CARDIOPULMONARY RESUSCITATION), 1-8)  
CHEST INJURY (see EMERGENCY, 1-35)  
CHEST PAIN (see EMERGENCY, 1-36)  
CHOKING (see EMERGENCY, 1-4)  
CONGESTION, NASAL..... 3-3  
CONSTIPATION..... 3-3  
COUGH (see BRONCHITIS, 2-7)  
CPR (CARDIOPULMONARY RESUSCITATION) (see  
EMERGENCY, 1-6)  
CRICOTHYROTOMY (see EMERGENCY, 1-19)  
DCS – DECOMPRESSION SICKNESS (BENDS)..... 3-4  
    EVA DCS EXAMINATION SCORECARD ..... 3-5  
    IN-SUIT EXAM..... 3-5  
    OUT-OF-SUIT EXAM ..... 3-7  
    DCS TREATMENT (POST SUIT DOFFING)..... 3-8  
    AFTERCARE ..... 3-8  
    LATE DCS SYMPTOMS ..... 3-8  
    RECURRENT DCS..... 3-10  
DCS – OTHER DECOMPRESSION PROBLEMS  
(BAROTRAUMA)..... 3-10  
DIARRHEA..... 3-11  
EAR EXAMINATION..... 3-11  
EAR PROBLEMS ..... 3-12  
    CONGESTION (SEROUS OTITIS MEDIA) ..... 3-12  
    INTERNAL INFECTION (OTITIS MEDIA) ..... 3-12  
    EAR CANAL INFECTION (EXTERNAL OTITIS)..... 3-13  
    RUPTURED EARDRUM ..... 3-14  
    EAR/SINUS BLOCK ..... 3-14  
ELECTROCARDIOGRAPH (ECG)  
(see CPR (CARDIOPULMONARY RESUSCITATION),  
CARDIAC MONITORING, 1-8)  
EYE EXAMINATION..... 3-15  
EYE – PENETRATING OBJECT (see EMERGENCY, 1-38)  
EYE PROBLEMS..... 3-17  
    CHEMICAL BURN (SHUTTLE EMERGENCY  
    EYEWASH (SEE) GOGGLES) ..... 3-17  
    DECONTAMINATION PROCEDURES  
    (see CHEMICAL BURN, 3-17)  
    FOREIGN BODY/ABRASION ..... 3-20  
    CORNEAL ULCER ..... 3-22  
    HERPETIC VIRAL OCULAR INFECTION..... 3-23  
    IRRITATION/INFECTION..... 3-23

SECTION 4 – F THRU N

FRACTURE/DISLOCATION .....	4-3
HEADACHE (see PAIN RELIEF, 5-3)	
HEAD OR NECK INJURY (see EMERGENCY, 1-39)	
HEMORRHOID .....	4-5
HOT CABIN.....	4-5
INDIGESTION (UPSET STOMACH) .....	4-6
INJECTIONS.....	4-7
INTRAVENOUS (I.V.) .....	4-7
CATHETER DRESSING CHANGES (DAILY).....	4-22
REMOVAL OF Y-TYPE CATHETER EXTENSION.....	4-23
INTRAMUSCULAR.....	4-24
SUBCUTANEOUS.....	4-26
PREFILLED SYRINGE TECHNIQUE .....	4-27
TUBEX INJECTOR TECHNIQUE .....	4-28
INTUBATION (ILMA)	
(see AIRWAY(ILMA) CUE CARD, CC 9-4)	
INTUBATION, ENDOTRACHEAL (see CPR	
(CARDIOPULMONARY RESUSCITATION), 1-12)	
LACERATIONS.....	4-30
MAJOR LACERATIONS.....	4-30
MINOR LACERATIONS .....	4-37
MOTION SICKNESS (also see VOMITING, 5-20) .....	4-38
MILD MOTION SICKNESS .....	4-38
MODERATE-TO-SEVERE MOTION SICKNESS .....	4-39
MUSCLE SPASM (DRUG SIDE EFFECT).....	4-39
NASAL CONGESTION (see CONGESTION, NASAL, 3-3)	
NAUSEA/VOMITING .....	4-39
NOSEBLEED .....	4-40

SECTION 5 – O THRU Z

OBSTRUCTED AIRWAY – CONSCIOUS ADULT (see	
CHOKING, 1-4)	
OBSTRUCTED AIRWAY – UNCONSCIOUS ADULT (see	
CHOKING, 1-5)	
ORAL AIRWAY (see CPR (CARDIOPULMONARY	
RESUSCITATION), 1-11)	
OXYGEN – SUPPLEMENTAL (see CPR	
(CARDIOPULMONARY RESUSCITATION), 1-12)	
PAIN RELIEF (IN INCREASING ORDER OF	
STRENGTH).....	5-3
RADIATION.....	5-4
SEIZURES (see EMERGENCY, 1-42)	
SHOCK (see EMERGENCY, 1-44)	

<u>SECTION 5 – O THRU Z (Concluded)</u>	
SINUS PROBLEMS .....	5-5
SKIN INFECTION .....	5-6
SKIN – MINOR ABRASIONS .....	5-7
SKIN RASH.....	5-7
ALLERGIC RASH .....	5-7
FUNGAL RASH .....	5-8
SLEEPING PROBLEMS (INSOMNIA).....	5-8
SORE THROAT .....	5-9
SPACE MOTION SICKNESS (see MOTION SICKNESS, 4-38)	
STRAINS/SPRAINS .....	5-9
TOOTHACHE .....	5-10
DENTAL INJECTION TECHNIQUE .....	5-11
TEMPORARY FILLING .....	5-15
EXPOSED PULP .....	5-15
REPLACING CROWN w/TEMPORARY CEMENT .....	5-16
UPSET STOMACH (see INDIGESTION, 4-6)	
URINARY RETENTION.....	5-18
URINARY TRACT INFECTION (DYSURIA) (see BLADDER INFECTION, 2-5)	
VOMITING (also see MOTION SICKNESS – MODERATE-TO-SEVERE, 4-39) .....	5-20
YEAST INFECTION .....	5-20
 <u>SECTION 6</u>	
MEDICATION SIDE EFFECTS.....	6-2
 <u>SECTION 7 – TOXICOLOGY</u>	
HAZARDOUS SPILL LEVEL DEFINITIONS TABLE.....	7-2
CONTAMINANT CLEANUP PROCEDURES.....	7-5
EYE EXPOSURE TOXICOLOGY TREATMENT .....	7-6
SKIN EXPOSURE TOXICOLOGY TREATMENT .....	7-6
INHALATION EXPOSURE TOXICOLOGY TREATMENT (INCLUDES CARDIOVASCULAR AND CENTRAL NERVOUS SYSTEM) .....	7-7
 <u>SECTION 8 – SOMS STOWAGE AND USAGE</u>	
ALPHABETICAL LIST .....	8-2
USAGE LIST .....	8-15
 <u>SECTION 9 – CUE CARD CONFIGURATION .....</u>	
	CC 9-1



## FIGURES

Figure		Page
1-1	Restraints .....	1-6
1-2	Medical locker .....	1-8
1-3	Electrodes .....	1-9
1-4	Oral airway insertion .....	1-11
1-5	Patient restrained .....	1-13
1-6	Laryngoscope .....	1-14
1-7	Vocal cord anatomy .....	1-15
1-8	Upper airway anatomy .....	1-16
1-9	CO2 detector.....	1-17
1-10	Patient restrained .....	1-20
1-11	Tracheostomy tube .....	1-20
1-12	Airway anatomy .....	1-21
1-13	Trachea restraint.....	1-21
1-14	Tracheal hook .....	1-22
1-15	Tracheal hook insertion.....	1-22
1-16	Incision placement .....	1-23
1-17	Opening enlargement .....	1-23
1-18	Removal of dilator .....	1-24
1-19	Tracheostomy tube secure position .....	1-24
1-20	CO2 detector.....	1-25
3-1	Right eardrum .....	3-11
3-2	Eyelid eversion.....	3-16
3-3	Eye landmarks; 6 directions of gaze .....	3-16
3-4	SEE-WCS connection.....	3-18
4-1	Existing I.V. flow.....	4-7
4-2	I.V. assembly with Y-type catheter extension .....	4-8
4-3	Medication administration technique.....	4-9
4-4	Tourniquet placement .....	4-13
4-5	Removal of I.V. needle.....	4-14
4-6	Surface anatomy.....	4-25
4-7	Prefilled syringe .....	4-27
4-8	Tubex injector .....	4-29
4-9	Suture placement.....	4-32
4-10	Laceration closure using stapler .....	4-33
4-11	Skin staple remover .....	4-36
4-12	Nasal speculum .....	4-40
4-13	Nasal packing .....	4-42
4-14	Nasal catheter insertion .....	4-43

5-1	Dosimeter.....	5-4
5-2	Upper molar anesthesia.....	5-13
5-3a	Lower (mandibular) molar anesthesia #1.....	5-14
5-3b	Lower (mandibular) molar anesthesia #2.....	5-14
5-4	Catheter insertion.....	5-19

SECTION 1

EMERGENCY

EMERGENCY

<u>IMMEDIATE LIFE SUPPORT</u> .....	1-3
BREATHING DIFFICULTY .....	1-3
CHOKING .....	1-4
OBSTRUCTED AIRWAY – CONSCIOUS ADULT .....	1-4
OBSTRUCTED AIRWAY – UNCONSCIOUS ADULT ..	1-5
CPR (CARDIOPULMONARY RESUSCITATION) .....	1-6
PATIENT RESTRAINT .....	1-6
CARDIAC MONITORING .....	1-8
ORAL AIRWAY .....	1-11
OXYGEN – SUPPLEMENTAL .....	1-12
INTUBATION (ILMA) .....	1-12
RESUSCITATOR	
(see RESUSCITATOR REFERENCE CARD, CC 9-7)	
INTUBATION, ENDOTRACHEAL .....	1-12
CRICOTHYROTOMY .....	1-19
<u>OTHER EMERGENCIES</u> .....	1-27
ABDOMINAL INJURY .....	1-27
ALLERGIC REACTION – SEVERE REACTION .....	1-29
BURNS – MAJOR .....	1-32
CHEST INJURY .....	1-35
CHEST PAIN .....	1-36
EYE – PENETRATING OBJECT .....	1-38
HEAD OR NECK INJURY .....	1-39
SEIZURES .....	1-42
SHOCK .....	1-44

FIGURES

1-1	Restraints.....	1-6
1-2	Medical locker.....	1-8
1-3	Electrodes.....	1-9
1-4	Oral airway insertion.....	1-11
1-5	Patient restrained.....	1-13
1-6	Laryngoscope.....	1-14
1-7	Vocal cord anatomy.....	1-15
1-8	Upper airway anatomy.....	1-16
1-9	CO2 detector.....	1-17
1-10	Patient restrained.....	1-20
1-11	Tracheostomy tube.....	1-20
1-12	Airway anatomy.....	1-21
1-13	Trachea restraint.....	1-21
1-14	Tracheal hook.....	1-22
1-15	Tracheal hook insertion.....	1-22
1-16	Incision placement.....	1-23
1-17	Opening enlargement.....	1-23
1-18	Removal of dilator.....	1-24
1-19	Tracheostomy tube secure position.....	1-24
1-20	CO2 detector.....	1-25

## **IMMEDIATE LIFE SUPPORT**

### **BREATHING DIFFICULTY**

If difficulty develops during eating and person can talk/cough/whisper:

1. No action but continue to monitor

If difficulty develops during eating and person unable to breathe:

2. Go to CHOKING, 1-4 >>

If difficulty develops after exposure to dust/chemicals and person has wheezing/shortness of breath:

3. Remove from exposure to chemical/irritant
4. Protect others from exposure
5. Contact Surgeon

Med  
Locker

6. Using stethoscope (Airway Subpack-9), listen to breath sounds for wheezing, crackles, or decreased breath sounds

If wheezing, crackles, or decreased breath sounds present:



7. \*Proventil Inhaler (Drug Subpack-11)

Dose: 2 puffs initially. May repeat in 1-2 hr if necessary, then 2 puffs every 4 to 6 hr as needed for continued wheezing




8. Don QDM for 100% O2 until MCC Surgeon advises otherwise

# CHOKING






## OBSTRUCTED AIRWAY – CONSCIOUS ADULT

	Step	Activity
	1.	“Are you choking?”  Victim may be using “Universal Distress Signal” of choking: clutching neck between thumb, index finger
 <p data-bbox="462 777 673 808">Abdominal Thrust</p>	2. 3.	Perform Heimlich Maneuver: 2. Deliver 6-10 abdominal thrusts 3. Repeat thrusts until either foreign body expelled or victim becomes unconscious (see below)

## Adult W/Obstructed Airway Becomes Unconscious


	4.	Perform finger sweep
	5.	Open airway – head-tilt/chin lift  Attempt to ventilate by mouth-to-mouth resuscitation – Two Breaths
	6. 7.	Perform 6-10 abdominal thrusts If unsuccessful: 7. Repeat steps 4-6 <u>Two Times</u>  If unable to ventilate: 8. Go to CPR (CARDIOPULMONARY RESUSCITATION), 1-6; prepare for CRICOTHYROTOMY, 1-19

## OBSTRUCTED AIRWAY – UNCONSCIOUS ADULT

	Step                      Activity
	1. Open airway – head-tilt/chin lift
	2. Attempt to ventilate/mouth-to-mouth  3. Airway remains obstructed, <u>reattempt</u> ventilation
	4. Perform 6-10 abdominal thrusts
	5. Perform finger sweep
	6. Attempt to ventilate If unsuccessful: 7. Repeat steps 4-6 <u>Two Times</u>    If unable to ventilate: 8. Go to CPR (CARDIOPULMONARY RESUSCITATION), 1-6. For several failed attempts to intubate and unable to mask ventilate, contact Surgeon for possible cricothyrotomy (CRICOTHYROTOMY, 1-19)

# CPR (CARDIOPULMONARY RESUSCITATION)

## PATIENT RESTRAINT







### NOTE

A second rescuer restraint belt may be used to position an additional rescuer to assist w/CPR

Figure 1-1.- Restraints.



**CPR (CARDIOPULMONARY RESUSCITATION) (Cont)**

	Step	Activity
	1.	Open airway Look, feel, listen  2. If blockage suspected, refer to <b>CHOKING; OBSTRUCTED AIRWAY, 1-4</b>  3. If not breathing, insert oral airway (see 1-11)
	4.	Unstow <b>RESUSCITATOR</b> and follow Cue Card procedures  5. Ventilate twice per Cue Card procedure, step 7, or perform mouth-to-mouth resuscitation  6. Observe chest rise
	7.	Check for circulation by feeling for carotid pulse
	8.	Begin compression/ventilation cycles: 30 compressions/2 ventilations for one (1) person CPR  30 compressions/2 ventilations for two (2) person CPR  Rate of compression is 80-100 per min  Feel for pulse after 1 min, then every 2 min  Contact Surgeon



2. Swab electrode sites w/Alcohol Pads; let dry
3. Connect: IVA Cable to side of Signal Conditioner, silver end of Sternal Harness to top of Signal Conditioner (see fig 1-3)
4. Don OBS Belt w/Signal Conditioner
5. Apply small amount of electrode gel to electrodes
6. Remove stomaseal protective coverings from electrodes
7. Attach electrodes to chest; secure w/tape




Figure 1-3.- Electrodes.


- R10      Operation  
BIOMED CH 1(2):
1. PS (for CDR, MS2 seats; connect Biomed Cable to A15)
  2. MS (for PLT, MS1 seats; connect Biomed Cable to A11)
  3. MIDDECK L (for middeck port seat S7; connect Biomed Cable to MO62M; LEFT)
  4. MIDDECK CTR (for middeck center seat S5; connect Biomed Cable to MO62M; CENTER)
  5. MIDDECK R (for middeck stbd seat S6; connect Biomed Cable to MO62M; RIGHT)
6. Connect IVA Cable to Biomed Cable

Doffing and Stowage

1. Remove electrodes, doff belt, discard stomaseals (3)
2. Disconnect IVA Cable from Signal Conditioner, Biomed Cable
3. Using Alcohol Pads, clean electrodes, air dry, or wipe dry
4. Apply new stomaseals to electrodes (3)
5. Stow OBS:

Med Locker      OBS Belt w/Signal Conditioner  
                         Sternal Harness  
                         IVA Cable  
                         Biomed Cable  
                         Electrode Attachment Kit

ORAL AIRWAY



Oral Airway (Airway Subpack-3)

1. Insert oral airway while pointing it toward head (as shown)



2. Rotate oral airway 180 deg so it points toward feet



3. Final position



Figure 1-4.- Oral airway insertion  
(for unconscious, breathing patient).

OXYGEN – SUPPLEMENTAL

- \* Use Quick Don Mask (QDM) for oxygen \*
- \* delivery to crewmember who is awake, \*
- \* breathing but needs protection of airway \*
- \* from toxins, smoke, or DCS (bends) \*

NOTE

Use RESUSCITATOR for unconscious patient

INTUBATION (ILMA)

See cover of Airway Subpack or Cue Card section

RESUSCITATOR

See inside cover of Medical Accessory Kit or Cue Card section

INTUBATION, ENDOTRACHEAL

Indications: Inability of rescuer to ventilate patient who is not breathing w/conventional methods (mouth-to-mouth, mouth-to-mask); inability of patient to protect her(his) own airway (coma, loss of gag reflex, or cardiac arrest); need for prolonged artificial ventilation

1. Unstow:

Med	Tracheal Tube	(Airway Subpack-20)
Locker	Stylet	(Airway Subpack-6)
	Syringe (10 cc)	<b>(Airway Subpack-15,</b> Trauma Subpack-22, EENT Subpack-20)
	Lubricant (water-soluble)	<b>(Airway Subpack-13,</b> Trauma Subpack-9)
	Laryngoscope Handle w/med Blade	(Airway Subpack-5)
	End-Tidal CO2 Detector	(Airway Subpack-7)
	Toomey Syringe suction device	(Airway Subpack-4)
	Ziplock Bag (12 in X 12 in)	<b>(Airway Subpack-4,</b> IV Admin-1, CCK)
	Stethoscope	(Airway Subpack-12)
	RESUSCITATOR	
	Restraints:	
	Patient	
Misc	Gray Tape	
Stowage	Towels	

Contact Surgeon while proceeding w/following:

2. Restrain crewmember w/CPR patient restraints (see fig 1-5, Restraints)

**CAUTION**

If any suspicion of neck trauma, crewmember's head will require continuous immobilization. One crewmember should use both hands to stabilize patient's head. Continue with step 5

3. If no neck trauma suspected, place rolled towel(s) under crewmember's hyperextended neck




Figure 1-5.- Patient restrained.

4. Place Gray Tape over crewmember's forehead to immobilize head against middeck locker
5. Continue to ventilate/oxygenate patient (see RESUSCITATOR. See inside cover of Medical Accessory Kit or Cue Card section) while preparing for intubation
6. Remove Tracheal Tube from pkg; leave stylet in place and unfold tube; lubricate cuffed end of Tracheal Tube w/water soluble lubricant

7. Firmly insert syringe, filled w/6-cc air, into one-way valve/pilot balloon on Tracheal Tube; leave connected
8. Do not inflate balloon

Assemble, if necessary:




Figure 1-6.- Laryngoscope.

9. Extend Laryngoscope blade to 90 deg position  
√Laryngoscope light
10. Open mouth w/fingers of right hand

**WARNING**  
Avoid pressure on lips, teeth

11. Suction fluid from mouth,throat w/Toomey Syringe suction device; discharge syringe into Ziplock Bag or towel
12. Hold Laryngoscope in left hand  
Insert blade in right side of mouth displacing tongue to left



13. Advance blade into space between base of tongue, epiglottis (vallecula)




Figure 1-7.- Vocal cord anatomy.

14. Lift tongue w/Laryngoscope blade tip to expose vocal cords

- \* If unable to see vocal cords due to fluid, \*
- \* remove laryngoscope then use Toomey \*
- \* Syringe suction device (Airway \*
- \* Subpack-4) to remove fluid. Ventilate \*
- \* patient w/RESUSCITATOR for another \*
- \* 1-2 minutes (about once every 3-4 sec) \*
- \* before again attempting intubation. \*
- \* Re-insert Laryngoscope and again \*
- \* attempt intubation \*

Whenever needed:

15. Suction fluid from back of throat w/Toomey Syringe suction device; discharge syringe into Ziplock Bag or towel




Figure 1-8.- Upper airway anatomy.

16. While watching, advance cuffed end of Tracheal Tube along right side of mouth into trachea until entire cuff is about 1 cm below vocal cords

**WARNING**  
 Hold Tracheal Tube **FIRMLY** in place, until proper placement confirmed, tube secured

17. Inflate Tracheal Tube cuff w/air (8-10 cc) from syringe. Remove syringe, remove stylet from tube
18. Remove mask from RESUSCITATOR
19. Connect RESUSCITATOR to Tracheal Tube

Ventilate/oxygenate lungs:

20. Squeeze RESUSCITATOR until chest rises (no more than 1-2 sec)
21. Release RESUSCITATOR to allow passive exhalation
22. Repeat every 4-5 sec

**WARNING**  
Do not interrupt ventilation for > 15 sec




Figure 1-9.- CO2 detector.

23. Connect End-Tidal CO2 Detector between Tracheal Tube and RESUSCITATOR
  24. ✓ End-Tidal CO2 Detector color changes from purple to yellow w/each breath
    - \* If no color change, deflate cuff w/syringe; \*
    - \* remove tube. Ventilate/oxygenate \*
    - \* w/RESUSCITATOR/mask for 1-2 \*
    - \* minutes. Reattempt intubation from step 5 \*
  25. If reqd, continue ventilation
  26. W/stethoscope, listen to chest for equal breath sounds on both sides
- If breath sounds not equal:
27. Withdraw 0.5 in; recheck breath sounds
28. If reqd, continue ventilation
  29. Secure tube to face using Tape

30. Unstow:  
Blood Pressure Cuff (IV Admin-1)  
Thermometer (EENT Subpack-3)

31. Obtain vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

32. Apply ECG electrodes (see CPR  
(CARDIOPULMONARY RESUSCITATION),  
CARDIAC MONITORING, 1-8)
33. Continue resuscitation as indicated

## CRICOTHYROTOMY

(for victim w/obstructed airway or failed intubation attempts)

Indications: Obstruction of upper airway that cannot be cleared by CPR techniques; inability to obtain/maintain open airway w/tracheal intubation

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

- Unstow:
  - Med Locker
    - End-Tidal CO2 Detector (Airway Subpack-7)
    - Alcohol Pads (**Airway Subpack-10**, IV Admin-18, Drug Subpack-16, Drug Subpack-Spine, Trauma Subpack-6)
    - ^Povidone-Iodine Swab (Airway Subpack-10)
    - Gauze Pads (sterile) (Airway Subpack-11)
    - Toomey Syringe suction device (Airway Subpack-4)
    - Tape (**Airway Subpack-9**, IV Admin-13, Trauma Subpack-34, EENT Subpack-9)
    - Gloves (**Airway Subpack-8**, IV Admin-14, CCK, EENT Subpack-7)
    - Lubricant (water-soluble) (**Airway Subpack-13**, Trauma Subpack-9)
    - Tracheostomy Items:
      - Alcohol Pads (Airway Subpack-10)
      - Scissors, Straight (Airway Subpack-2)
      - Forceps, Curved (Airway Subpack-19)
      - Tracheal Hook (Airway Subpack-2)
      - Silk Sutures (2-3) (Airway Subpack-11)
      - Tracheostomy Tube (Airway Subpack-1)
      - Tracheostomy Tube holder (Airway Subpack-1)
    - Scalpel (Airway Subpack-2)
    - Stethoscope (Airway Subpack-12)
  - RESUSCITATOR
  - Restraints:
    - Patient (2 belts)
    - Rescuer (2 belts, if needed)
  - Misc Stowage
    - Gray Tape
    - Towels

Contact Surgeon while proceeding w/following:

2. Restrain crewmember w/CPR patient restraints (fig 1-10)

**CAUTION**

If any suspicion of neck trauma, crew-member's head will require continuous immobilization. One crewmember should use both hands to stabilize patient's head. Continue with step 5

3. If no neck trauma suspected, position the patient:  
Flex the neck and extend the chin
4. Place Gray Tape over crewmember's forehead to immobilize head against middeck locker




Figure 1-10.- Patient restrained.




Figure 1-11.- Tracheostomy tube.

5. Lubricate cuffed Tracheostomy Tube w/water-soluble lubricant

Locate Cricothyroid membrane (fig 1-12):

6. Don Gloves. Use index finger to locate Adam's Apple (thyroid cartilage)
7. Slide down, feel for next ridge (cricoid cartilage)




Figure 1-12.- Airway anatomy.

8. Feel for small triangular soft spot above this ridge. This is the Cricothyroid membrane
9. Cleanse neck w/ Povidone-Iodine Swab. Cleanse neck w/ Alcohol Pads
10. If right-handed, grasp trachea w/ left hand and hold between middle finger and thumb until procedure is finished (see figure 1-13). Use Scalpel to make 1-cm long midline vertical incision in skin over Cricothyroid membrane



Figure 1-13.- Trachea restraint.

11. Insert left index finger and feel cricoid cartilage and larynx
12. Insert sharp end of Tracheal Hook into Cricothyroid membrane (fig 1-15). Use blunt end to elevate and retract trachea. The trachea should become superficial



Figure 1-14.- Tracheal hook.




Figure 1-15.- Tracheal hook insertion.



13. Place blade of Scalpel next to Tracheal Hook. Make 1-cm long horizontal incision into membrane. Air should bubble out of trachea (see figure 1-16)

**WARNING**  
Briskly bleeding blood vessels may be clamped w/forceps




Figure 1-16.- Incision placement.

14. Enlarge incision opening by spreading the Straight Scissors or Curved Forceps




Figure 1-17.- Opening enlargement.

15. Insert Tracheostomy Tube sideways until tip is in trachea
16. Turn tube so that it moves down the trachea toward the feet
17. To advance tube into trachea, gently push until faceplate rests against skin

18. Leave tube in position in trachea  
Remove dilator (fig 1-18)




Figure 1-18.- Removal of dilator.

19. Do not let go of Tracheostomy Tube until it is  
securely tied around patient's neck (fig 1-19)




Figure 1-19.- Tracheostomy tube secure position.

20. Fill syringe w/3-cc air and connect to one-way  
valve/pilot balloon on Tracheostomy Tube. Inflate  
the cuff balloon

**WARNING**  
Do not remove hook until cricothyrotomy  
airway is functioning well

21. Remove mask from RESUSCITATOR. Connect RESUSCITATOR directly to end of the Tracheostomy Tube
22. Check for proper tube placement by ventilating the lungs. Squeeze RESUSCITATOR for 1-2 sec, and release. Look for the symmetric rise and fall of the chest
23. Once proper tube placement has been verified, connect End-Tidal CO2 Detector between Tracheostomy Tube and RESUSCITATOR (fig 1-20)




Figure 1-20.- CO2 detector.

**NOTE**

End-Tidal CO2 Detector color changes from purple to yellow w/each breath. No color change may indicate improper placement of the tube

**WARNING**  
If no color change, verify chest movement w/respirations. If no chest movement, contact surgeon IMMEDIATELY (if surgeon directs removal of tube, deflate cuff )

24. Manually ventilate crewmember with RESUSCITATOR for 1-2 sec and then releasing to allow passive exhalation. Repeat every 4-5 sec

NOTE

If vomiting occurs, clear mouth w/Toomey  
Syringe suction device. Expel vomitus  
into Ziplock Bag

25. W/stethoscope, listen to chest for equal breath  
sounds on both sides

26. Continue ventilation; ensure tube securely in place

27. Unstow:

Med	Blood Pressure Cuff	(IV Admin-1)
Locker	Thermometer	(EENT Subpack-3)

28. Obtain vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

29. Continue resuscitation as indicated. Apply ECG  
electrodes (see fig 1-2, Electrodes, 1-8)

**OTHER EMERGENCIES**

**ABDOMINAL INJURY**

- \* If no pulse or respiration, perform \*
- \* CPR using patient/rescuer \*
- \* restraints (see CPR \*
- \* (CARDIOPULMONARY \*
- \* RESUSCITATION), 1-6) \*
- \* \*
- \* If inadequate respiration, provide \*
- \* supplemental O2 (see CPR \*
- \* (CARDIOPULMONARY \*
- \* RESUSCITATION), OXYGEN – \*
- \* SUPPLEMENTAL, 1-12) \*

1. Control bleeding w/direct pressure, Gauze Pads  
(Trauma Subpack-14, IV Admin-14, Airway  
Subpack-11, EENT Subpack-2)

2. Unstow:  
Med Blood Pressure Cuff (IV Admin-1)  
Locker Stethoscope (Airway Subpack-12)  
Thermometer (EENT Subpack-3)

3. Evaluate vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

4. Bandage with:  
Kling (Trauma Subpack-12)  
Kerlix (Trauma Subpack-2)  
Tape (**Trauma Subpack-34**,  
IV Admin-13,  
Airway Subpack-9,  
EENT Subpack-9)

**WARNING**  
Do not replace protruding intestines

If intestines protruding:

5. Cover w/Kerlix, tape in place, moisten w/Saline (IV Admin-1, Saline Supply Bag, Middeck Locker, EENT Subpack-1), bandage

**WARNING**  
Any abdominal trauma may cause internal bleeding which can produce abdominal rigidity. Continue to monitor vital signs, assess for developing shock (refer to SHOCK, 1-44) or abdominal rigidity

6. Perform abdominal exam

Visual inspection  
(bruises, swelling): \_\_\_\_\_

Bowel sounds (type,  
frequency): \_\_\_\_\_

Palpation (rigid, pain): \_\_\_\_\_

If vomiting:

7. Inspect for blood (black/red coffee grounds); keep airway clear
8. Contact Surgeon
9. Start I.V. line (see INJECTIONS, INTRAVENOUS (I.V.), Intravenous Fluid Infusion w/Y-Type Catheter Extension and Administration Set, 4-11)

## ALLERGIC REACTION

### SEVERE REACTION

#### NOTE

Institute rapid treatment if pallor, sweating, or breathing difficulty occur

Symptoms: Difficulty breathing

Signs: Pallor, sweating, difficulty breathing, low blood pressure, rapid pulse, wheezing

#### WARNING

Symptoms may follow exposure to any substance to which an individual has become allergic, including substances to which he has never had previous reaction. Severe reaction w/development of shock possible (see SHOCK, 1-44)

1. Provide supplemental O2 (see CPR (CARDIOPULMONARY RESUSCITATION), OXYGEN – SUPPLEMENTAL, 1-12)

2. Unstow:

Med	Blood Pressure Cuff	(IV Admin-1)
Locker	Stethoscope	(Airway Subpack-12)
	Epi Pen Injector	(Drug Subpack-9)

3. Remove Epi Pen from plastic yellow vial. Remove gray safety cap and discard cap in trash. Simultaneously provide O2
4. Place black tip on outer thigh

#### NOTE

Do not put thumb over end of unit. Epi Pen may NOT be used through clothing

5. Using quick motion, press firmly and hold in place for 10 sec

6. Remove Epi Pen injector from patient and restow used unit, needle end first, into plastic yellow vial. Place white cap back on yellow vial and mark as "USED"

7. Restow yellow vial in Drug Subpack-9

8. Monitor pulse, respiration, blood pressure

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

9. Contact Surgeon

10. While waiting for Epinephrine to take effect, unstow Tubex Injector (Drug Subpack-19) and Injectable Epinephrine (Drug Subpack-5) and temp stow. If time allows, review INJECTIONS, TUBEX INJECTOR TECHNIQUE (4-28) and SUBCUTANEOUS (4-26)

11. If no improvement in breathing or blood pressure has occurred within 5 min from first injection, give additional Epinephrine as follows: Load Tubex Injector w/injectable medication (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28)

\*Epinephrine (Drug Subpack-5)

Dose: Inject 0.3 to 0.5-ml subcutaneously (see INJECTIONS, SUBCUTANEOUS, 4-26)

12. Reload Tubex Injector w/injectable medication:

\*Benadryl (Injectable) (Drug Subpack-13)

(see general WARNING, 6-2)

Dose: Inject 1 ampule intramuscularly (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28 and INTRAMUSCULAR, 4-24)



13. Reload Tubex Injector w/injectable medication:  
\*Dexamethasone (Drug Subpack-28)  
Dose: Inject 1 cc (10 mg/cc) intramuscularly  
(see INJECTIONS, TUBEX  
INJECTOR TECHNIQUE, 4-28 and  
INTRAMUSCULAR, 4-24)
14. If blood pressure and pulse are stable and wheezing  
persists, use \*Proventil Inhaler (Drug Subpack-11).  
Inhaler should be discharged (puffed) during  
inhalation

NOTE

Shake inhaler gently back and forth a  
few times

Before putting inhaler into mouth, take one deep  
breath and exhale normally. Next, slowly take  
another deep breath and, about halfway into this,  
seal mouth around Inhaler Tube and squeeze to  
discharge while inhaling. At this point, hold the  
breath for several seconds, then exhale normally.  
Repeat for the second puff

Dose: 2 puffs initially. May repeat in 1 hr, then  
2 puffs every 4 hr

Possible side effects: Tremor, palpitations,  
fast heart rate,  
nausea, increased  
blood pressure

## BURNS – MAJOR

### NOTE

Electrical burns may cause severe internal injury in spite of minimal skin damage. May also produce irregular heart rhythm including cardiac arrest

- \* If unable to detect pulse and/or \*  
 \* respiration, perform CPR using \*  
 \* patient/rescuer restraints (see CPR \*  
 \* (CARDIOPULMONARY \*  
 \* RESUSCITATION), 1-6) \*  
 \* \*  
 \* If unconscious and patient cannot \*  
 \* maintain airway, insert oral airway \*  
 \* (see CPR (CARDIOPULMONARY \*  
 \* RESUSCITATION), ORAL AIRWAY, \*  
 \* 1-11, or INTUBATION, \*  
 \* ENDOTRACHEAL, 1-12) and provide \*  
 \* manual ventilation using \*  
 \* RESUSCITATOR (see inside cover of \*  
 \* Medical Accessory Kit or Cue Card \*  
 \* section) \*  
 \* \*  
 \* If conscious, provide supplemental O2 \*  
 \* with QDM \*

1. Unstow:

Med Locker	Blood Pressure Cuff Stethoscope Thermometer	(IV Admin-1) (Airway Subpack-12) (EENT Subpack-3)
---------------	---	---

2. Evaluate vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

3. Estimate degree of burn:
  - First degree – superficial reddening of skin, pain (like a sunburn)
  - Second degree – blisters, red skin, swelling, pain
  - Third degree – pain often minimal, skin may appear white, charred, may resemble second degree burn

NOTE

Area of palm of hand equals approx  
1% (for estimation)

4. Estimate area of burn:
  - Head – 9%
  - Chest, Abdomen – 18%
  - Back – 18%
  - Arms (each) – 9%
  - Legs (each) – 18%
5. Contact Surgeon for further treatment options. Those options may include the following steps

WARNING  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

6. Apply thin layer of ^Silvadene Cream (Trauma Subpack-20) to burn using tongue depressor (EENT Subpack-34)

7. Unstow:
 

Med Locker	Sterile Drapes Saline	(Trauma Subpack-24) <b>(Saline Supply Bag,</b> EENT Subpack-1, IV Admin-1, Middeck Locker)
	Kling Tape	(Trauma Subpack-12) <b>(Trauma Subpack-34,</b> IV Admin-13, Airway Subpack-9, EENT Subpack-9)

Misc Stowage      Towel

8. Cover burn w/clean Sterile Drapes moistened w/Saline

9. Wrap area loosely w/Kling bandage and tape

NOTE

Will need to give a volume of I.V. fluids equal to 2ml/kg/% surface area of burn (if second or third degree) over 8-hr period

10. Increase fluid intake. Consult Surgeon whether to establish I.V. or not (see INJECTIONS, INTRAVENOUS (I.V.), 4-11)

11. For pain relief, see PAIN RELIEF, 5-3

## CHEST INJURY

- \* If no pulse or respiration, perform CPR \*
- \* using patient/rescuer restraints (see \*
- \* CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), 1-6) \*
- \* \*
- \* If unconscious and patient cannot \*
- \* maintain airway, insert oral airway \*
- \* (see CPR (CARDIOPULMONARY \*
- \* RESUSCITATION) ORAL AIRWAY, \*
- \* 1-11, or INTUBATION, \*
- \* ENDOTRACHEAL, 1-12); provide \*
- \* manual ventilation using \*
- \* RESUSCITATOR (see inside cover of \*
- \* Medical Accessory Kit or Cue Card \*
- \* section) \*
- \* \*
- \* If conscious, provide supplemental O2 \*
- \* with QDM \*

1. Unstow:

Med Locker	Blood Pressure Cuff Stethoscope Thermometer	(IV Admin-1) (Airway Subpack-12) (EENT Subpack-3)
---------------	---	---

### 2. Evaluate vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

**WARNING**

Do not remove an impaled object from chest. Bandage around object using Xeroform gauze (Trauma Subpack-11)

3. Cover any puncture wound to chest w/Xeroform Gauze (Trauma Subpack-11)
4. Control bleeding, provide airtight bandage w/following:
  - Gauze Pads (Trauma Subpack-14, IV Admin-14, Airway Subpack-11, EENT Subpack-2)
  - Kling (Trauma Subpack-12)
  - Kerlix (Trauma Subpack-2)
  - Ace Bandages (Trauma Subpack-3,13)
  - Telfa Pads (EENT Subpack-22)
5. Use stethoscope (Airway Subpack-12) to listen to lungs. Note absence or decrease in breath sounds (may indicate lung collapse)
6. Contact Surgeon

**CHEST PAIN**

- \* If no pulse or respiration, perform CPR \*
- \* using patient/rescuer restraints (see \*
- \* CPR (CARDIOPULMONARY \*
- \* RESUSCITATION)), 1-6) \*
- \* \*
- \* If unconscious and patient cannot \*
- \* maintain airway, insert oral airway \*
- \* (see CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), ORAL AIRWAY, \*
- \* 1-11, or INTUBATION, \*
- \* ENDOTRACHEAL, 1-12); provide \*
- \* manual ventilation using \*
- \* RESUSCITATOR (see inside cover of \*
- \* Medical Accessory Kit or Cue Card \*
- \* section) \*
- \* \*
- \* If conscious, provide supplemental O2 \*
- \* with QDM \*

Med Locker	1. Unstow:			
	Blood Pressure Cuff			(IV Admin-1)
	Stethoscope			(Airway Subpack-12)
	Thermometer			(EENT Subpack-3)
	2. Evaluate vital signs			
	MET:	_____	_____	_____
	Blood Pressure:	_____	_____	_____
	Pulse:	_____	_____	_____
	Respiratory Rate:	_____	_____	_____
	Temperature:	_____	_____	_____
	3. Chest Examination			
	Heart Sounds:			
	Regular/Irregular	_____	_____	_____
	Breath Sounds:			
	Increased	_____	_____	_____
	Decreased	_____	_____	_____
	Absent	_____	_____	_____
	Crackles	_____	_____	_____
	Wheezes	_____	_____	_____
	Chest Wall Motion			
	symmetrical/ asymmetrical	_____	_____	_____
	Skin Color			
	(bluish, grayish, pale)	_____	_____	_____

4. Attach ECG leads (see CPR (CARDIOPULMONARY RESUSCITATION), CARDIAC MONITORING, 1-8)
5. Ascriptin (Drug Subpack-Spine)  
(see WARNING, 6-3)  
Dose: 1 tab
6. Contact Surgeon

## EYE – PENETRATING OBJECT

WARNING  
Do not remove object penetrating eyeball. Do not remove any tissue extruding from eyeball

1. Unstow:
 

Med Locker	Eye Pads (2) Eye Shield Tape	(EENT Subpack-4) (EENT Subpack-4) <b>(Trauma Subpack-34,</b> IV Admin-13, Airway Subpack-9, EENT Subpack-9)
---------------	------------------------------------	--
2. Place single eye patch lightly over injured eye w/eyelid closed, if possible. Tape metal Eye Shield over eye patch
3. Close unaffected eye w/eye patch also. Tape in place firmly enough so that patient cannot blink
4. Contact Surgeon



## HEAD OR NECK INJURY

### WARNING

W/trauma to neck, possibility of fracture and/or spinal cord damage must be considered. Never move victim's neck

- \* If no pulse or respiration, perform CPR \*
- \* using patient/rescuer restraints (see \*
- \* CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), 1-6) \*
- \* \*
- \* If unconscious and patient cannot \*
- \* maintain airway, insert oral airway \*
- \* (see CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), ORAL AIRWAY, \*
- \* 1-11, or INTUBATION, \*
- \* ENDOTRACHEAL, 1-12); provide \*
- \* manual ventilation using \*
- \* RESUSCITATOR (see inside cover of \*
- \* Medical Accessory Kit or Cue Card \*
- \* section) \*
- \* \*
- \* If conscious, provide supplemental O2 \*
- \* with QDM \*

1. Unstow:

Med	Blood Pressure Cuff	(IV Admin-1)
Locker	Stethoscope	(Airway Subpack-12)
	Thermometer	(EENT Subpack-3)

### 2. Evaluate vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

3. If bleeding, control w/pressure, Gauze Pads (IV Admin-14, Airway Subpack-11, Trauma Subpack-14, EENT Subpack-2) or Telfa Pads (EENT Subpack-22)

4. Bandage with:  
 Kling (Trauma Subpack-12)  
 Kerlix (Trauma Subpack-2)  
 Ace Bandages (Trauma Subpack-3,13)

If neck injury suspected:

5. Immobilize head, neck using:  
 Med Kling (Trauma Subpack-12)  
 Locker Kerlix (Trauma Subpack-2)  
 Ace Bandages (Trauma Subpack-3,13)  
 Misc Gray Tape  
 Stowage Towels

6. Quantify level of consciousness:

Eye Opening (E)	Verbal Response (V)	Motor Response (M)
4 = Spontaneous	5 = Normal conversation	6 = Normal
3 = To voice	4 = Disoriented conversation	5 = Localizes to pain
2 = To pain	3 = Words, but not coherent	4 = Withdraws to pain
1 = None	2 = No words, only sounds	3 = Elbows flexed (Decorticate posture)
	1 = None	2 = Elbows extended (Decerebrate posture)
		1 = None
		Total = E + V + M

Record Total (E + V + M) every 15 min if total less than 15:

MET: \_\_\_\_\_  
 Total: \_\_\_\_\_

7. Examine eyes

Pupil size:

Pinpoint \_\_\_\_\_ Normal \_\_\_\_\_ Dilated \_\_\_\_\_

Difference in pupil size:

R > L \_\_\_\_\_ R = L \_\_\_\_\_ R < L \_\_\_\_\_

Med  
Locker

Reaction to light (Otoscope, head removed,  
EENT Subpack-32):

Present \_\_\_\_\_ Absent \_\_\_\_\_

8. Examine ears w/Otoscope, nose for blood or clear fluid
9. Contact Surgeon
10. Watch for vomiting; keep airway clear

## SEIZURES

### NOTE

Do not use Valium Autoinjector through clothing

- Med Locker 1. \*Valium Autoinjector (Drug Subpack-22)  
(see WARNING, 6-13)  
Dose: 2-cc intramuscularly ASAP

\* If patient conscious, provide supplemental \*  
\* O2 (see CPR (CARDIOPULMONARY \*  
\* RESUSCITATION), OXYGEN – \*  
\* SUPPLEMENTAL, 1-12) \*

2. Tear open protective plastic packet by pulling apart the split end of the Autoinjector

### WARNING

If patient has a thin lateral thigh muscle, injection must be administered in the upper outer quadrant of the buttocks

3. Determine whether to give the injection in the thigh or buttocks. The outer middle of the thigh is the preferred injection site

### WARNING

DO NOT cover or hold the black needle end with your hand or fingers

4. Use your dominant hand and grasp the Autoinjector with the black needle end extending beyond the thumb and index finger  
5. Use your other hand to pull the safety cap off the Autoinjector base. When safety cap is off, Autoinjector can be fired  
6. Position the black end of the Autoinjector perpendicular against the injection site

**WARNING**

DO NOT inject into areas close to the hip, knee, or thigh bone. For buttocks injections, inject the Valium only into the upper outer portion of the buttocks

**WARNING**

DO NOT use a jabbing motion when injecting. It may result in injury to the crewmember

7. Apply firm even pressure to the Autoinjector until it activates and pushes the needle into the injection site. Hold the Autoinjector in place for at least 10 sec. You should be able to feel the Autoinjector activate
8. Carefully pull the Autoinjector straight out from the injection site. Massage the injection site if time permits
- Med Locker 9. If seizure is continuing 5 min after first Valium injection, give additional Valium as follows:  
\*Valium (Injectable) (Drug Subpack-22)  
Dose: SLOWLY 2 ml intravenously (over 1-2 min) (see INJECTIONS, PREFILLED SYRINGE TECHNIQUE, 4-27, and Medication Administration thru Y-Type Catheter Extension Set, 4-18)
10. Unstow patient/rescuer restraints (Med Locker)
11. Secure patient to middeck lockers w/restraints (see CPR (CARDIOPULMONARY RESUSCITATION), 1-6)
12. Start I.V. (see INJECTIONS, INTRAVENOUS (I.V.), 4-11)
13. Contact Surgeon
14. If seizure stops, it can recur. Have \*Valium Autoinjector (Drug Subpack-22) ready
15. Attach ECG leads (see CPR (CARDIOPULMONARY RESUSCITATION), CARDIAC MONITORING, 1-8)

# SHOCK

## NOTE

Most critical step is identifying/treating underlying cause.

Basic causes of shock are:

- Heart damage
- Loss of circulating blood volume (bleeding, burns, dehydration)
- Blood pooling (allergy, pain, drugs, heat stroke, infection)

Signs:

- Pulse – rapid, weak, thready
- Respiration – shallow, irregular, labored
- Blood Pressure – low, falling
- Mental State – confused, sluggish, anxious
- Eyes – pupils dilated
- Skin – cold, clammy, sweating

- \* If no pulse or respiration, perform CPR \*
- \* using patient/rescuer restraints (see \*
- \* CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), 1-6) \*
- \* \*
- \* If unconscious and patient cannot \*
- \* maintain airway, insert oral airway \*
- \* (see CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), ORAL AIRWAY, \*
- \* 1-11, or INTUBATION, \*
- \* ENDOTRACHEAL, 1-12); provide \*
- \* manual ventilation using \*
- \* RESUSCITATOR (see inside cover of \*
- \* Medical Accessory Kit or Cue Card \*
- \* section) \*
- \* \*
- \* If conscious, provide supplemental O2 \*
- \* with QDM \*

### 1. Unstow:

- |        |                     |                     |
|--------|---------------------|---------------------|
| Med    | Blood Pressure Cuff | (IV Admin-1)        |
| Locker | Stethoscope         | (Airway Subpack-12) |
|        | Thermometer         | (EENT Subpack-3)    |
- 1-44 MED/ALL/GEN K,2

2. Evaluate vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_


Temperature: \_\_\_\_\_

If bleeding:

3. Apply pressure w/Gauze Pads (Trauma Subpack-14, IV Admin-14, Airway Subpack-11, EENT Subpack-2)
4. Prevent loss of body heat
5. Start I.V. (see INJECTIONS, INTRAVENOUS (I.V.), 4-11). Begin infusing 1 L of normal Saline (IV Admin-1, Saline Supply Bag, EENT Subpack-1)
6. Attach ECG leads (see CPR (CARDIOPULMONARY RESUSCITATION), CARDIAC MONITORING, 1-8)
7. Contact Surgeon



This Page Intentionally Blank





SECTION 2

A THRU B

ABDOMINAL INJURY (see EMERGENCY, 1-27)	
ABDOMINAL PAIN .....	2-2
ALLERGIC REACTION .....	2-3
MILD-TO-MODERATE REACTION .....	2-3
SEVERE REACTION (see EMERGENCY, 1-29)	
ALTITUDE SICKNESS .....	2-4
BACK PAIN .....	2-4
BENDS (see DECOMPRESSION SICKNESS, 3-4)	
BLADDER INFECTION .....	2-5
BREATHING DIFFICULTY (see EMERGENCY, 1-3)	
BROKEN BONE (see FRACTURE/DISLOCATION, 4-3)	
BRONCHITIS (COUGH).....	2-7
BURNS – MAJOR (see EMERGENCY, 1-32)	
BURNS – MINOR .....	2-8

A THRU B

### ABDOMINAL PAIN

Med Locker

1. Unstow:

Blood Pressure Cuff	(IV Admin-1)
Stethoscope	(Airway Subpack-12)
Thermometer	(EENT Subpack-3)

2. Evaluate vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

3. Perform abdominal exam, evaluate all 4 quadrants

Visual inspection  
(bruises, swelling): \_\_\_\_\_

Bowel sounds (presence  
or absence, type,  
frequency): \_\_\_\_\_

Palpation (rigid, pain,  
rebound tenderness): \_\_\_\_\_

Prior to treatment:

4. Contact Surgeon

If urinary symptoms (burning, frequency, pain,  
discomfort):

5. Go to BLADDER INFECTION, 2-5

If unable to urinate:

6. Go to URINARY RETENTION, 5-18

## ALLERGIC REACTION

### MILD-TO-MODERATE REACTION

Symptoms: Eyes may itch, burn, water. Nose may be congested.  
Throat, lungs may feel "tight"

Signs: Red, watery eyes. Clear discharge from nose. Rash/hives  
may be present on chest, back, abdomen, arms, or legs

Med Locker 1. Refresh Plus (artificial tears, eye drops)  
(EENT Subpack-16)  
as needed for dry, itchy eyes

Antihistamines:

2. Benadryl (oral) (Drug Subpack-42) for relief of  
itching (see general WARNING, 6-2)  
Dose: 1-2 cap every 4-6 hr as needed

-OR-

\*Claritin (Drug Subpack-43) antihistamine,  
anti-allergy (see general WARNING, 6-2)  
Dose: 10 mg once daily

Decongestants:

3. Sudafed (Drug Subpack-Spine) for relief of  
congestion (see WARNING, 6-12)  
Dose: 2 tabs every 6 hr as needed

-OR-

Guaifenes PSE (Drug Subpack-44)  
Dose: 1 tab every 12 hr as needed

-OR-

Afrin (EENT Subpack-19)  
Dose: 1-2 sprays every 12 hr as needed

## ALTITUDE SICKNESS

### NOTE

Symptoms of altitude sickness may be result of planned or unplanned cabin depressurization or PPO2 reduction that results in pressure altitude equivalent  $\geq 8000$  ft

Symptoms: Headache, nausea, vomiting, shortness of breath, cough, difficulty sleeping, lethargy, weakness

1. \*Diamox (Drug Subpack-43)  
(see WARNING, 6-5)  
Dose: 1/2-1 tab every 24 hr for total of 3 doses
2. Contact Surgeon

## BACK PAIN

1. Knee to chest position

Pain relief:

- Med Locker
2. Ascriptin (Drug Subpack-Spine)  
(see WARNING, 6-3)  
Dose: 1-2 tabs every 4-6 hr as needed  
-OR-  
Tylenol (Drug Subpack-Spine)  
Dose: 1-2 tabs every 4-6 hr as needed  
-OR-  
Motrin (Drug Subpack-Spine)  
(see WARNING, 6-9)  
Dose: 1 tab every 4-6 hr as needed w/food and drink

\* If unsuccessful, see \*  
\* PAIN RELIEF, 5-3 \*

Muscle relaxant:

3. \*Valium (oral) (Drug Subpack-20)  
(see WARNING, 6-2, 6-13)  
Dose: 1/2-1 tab every 6-8 hr as needed for severe spasm/pain

## BLADDER INFECTION

Symptoms: Pain, spasms, feeling of “heaviness” in groin; fever may or may not be present

Signs: Frequent, small volume, painful urination, cloudy or bloody urine

- Med Locker
1. Don non-sterile Gloves (IV Admin-12, Airway Subpack-8, EENT Subpack-7, CCK)
  2. Check urine w/Chemstrip 10 SG (Trauma Subpack-15):
  3. Hold two Cotton Swabs (EENT Subpack-34) together under urine stream to saturate
  4. Pass two Cotton Swabs over Chemstrip, compare w/color chart
- Wet Trash
5. Dispose of Cotton Swabs, Chemstrip

6. Record Chemstrip results in following table:

Chemstrip #	#1	#2	#3	#4	#5	#6	#7
	Name	Name	Name	Name	Name	Name	Name
	Date	Date	Date	Date	Date	Date	Date
	Flight Day	Flight Day	Flight Day	Flight Day	Flight Day	Flight Day	Flight Day
Specific Gravity							
pH							
Leukocytes (neg, trace, pos)							
Nitrate (neg, pos)							
Protein (mg/dL)							
Glucose (mg/dL)							
Ketones (neg, +, ++, +++)							
Urobilinogen (mg/dL)							
Bilirubin (neg, pos)							
Hgb (ery/ $\mu$ L)							

Table cont next page

Chemstrip #	#8	#9	#10	#11	#12	#13
	Name	Name	Name	Name	Name	Name
	Date	Date	Date	Date	Date	Date
	Flight Day	Flight Day	Flight Day	Flight Day	Flight Day	Flight Day
Specific Gravity						
pH						
Leukocytes (neg, trace, pos)						
Nitrate (neg, pos)						
Protein (mg/dL)						
Glucose (mg/dL)						
Ketones (neg, +, ++, +++)						
Urobilinogen (mg/dL)						
Bilirubin (neg, pos)						
Hgb (ery/ $\mu$ L)						

7. Increase daily fluid intake, take antibiotic

Antibiotics:

- Med Locker 8. \*Bactrim DS (Drug Subpack-38)  
 (see WARNING, 6-3)  
 Dose: 1 tab 2X/day for 3 days  
 -OR-  
\*Cipro (Drug Subpack-40)  
 Dose: 1 tab 2X/day for 3 days  
 -OR-  
\*Duricef (Drug Subpack-40)  
 (see WARNING, 6-6)  
 Dose: 1 cap 2X/day for 3 days  
 -OR-  
\*Augmentin (Drug Subpack-37)  
 Dose: 1 tab 2X/day for 7 days

Pain/spasm relief:

9. Pyridium (Drug Subpack-17)

Dose: 1 tab 3X/day as needed for pain  
(may color urine orange)

\* If no improvement in 24 hr, \*  
\* contact Surgeon \*

## **BRONCHITIS (COUGH)**

Symptoms: Lung congestion, "tightness" in chest, fatigue, fever,  
productive cough (w/mucus)

Med 1. Cough Lozenges (Drug Subpack-45)

Locker Dose: As needed

Antibiotics:

2. \*Zithromax (Drug Subpack-36)

Dose: 2 tabs initial dose, then 1 tab/day  
for 5 days

-OR-

\*Cipro (Drug Subpack-40)

Dose: 1 tab 2X/day for 10 days

-OR-

\*Duricef (Drug Subpack-40)

(see WARNING, 6-6)

Dose: 1 cap 2X/day for 10 days

-OR-

\*Augmentin (Drug Subpack-37)

Dose: 1 tab 3X/day for 10 days

## BURNS – MINOR

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

- Med Locker
1. Cool first, second degree burns w/Gauze Pads (IV Admin-14, Airway Subpack-11, Trauma Subpack-14, EENT Subpack-2) moistened w/CHILLED water
  2. Clean skin around burn w/^Povidone-Iodine Swabs (IV Admin-19, Airway Subpack-10, Trauma Subpack-7)
  3. Apply ^Silvadene Cream (Trauma Subpack-20) to burn
  4. Bandage with:
    - Adaptic Bandages or Gauze Pads (Trauma Subpack-14, IV Admin-14, Airway Subpack-11, Trauma Subpack-5, EENT Subpack-2)
    - Kling (Trauma Subpack-12)
    - Kerlix (Trauma Subpack-2)
    - Tape (Trauma Subpack-34, IV Admin-13, Airway Subpack-9, EENT Subpack-9)
    - Telfa Pads (EENT Subpack-22)
  5. For pain relief, see PAIN RELIEF, 5-3
  6. Contact Surgeon



SECTION 3

C THRU E

CARDIAC MONITORING (see CPR  
(CARDIOPULMONARY RESUSCITATION), 1-8)

CHEST INJURY (see EMERGENCY, 1-35)

CHEST PAIN (see EMERGENCY, 1-36)

CHOKING (see EMERGENCY, 1-4)

CONGESTION, NASAL..... 3-3

CONSTIPATION..... 3-3

COUGH (see BRONCHITIS, 2-7)

CPR (CARDIOPULMONARY RESUSCITATION) (see  
EMERGENCY, 1-6)

CRICOTHYROTOMY (see EMERGENCY, 1-19)

DCS – DECOMPRESSION SICKNESS (BENDS)..... 3-4

    EVA DCS EXAMINATION SCORECARD ..... 3-5

    IN-SUIT EXAM ..... 3-5

    OUT-OF-SUIT EXAM ..... 3-7

    DCS TREATMENT (POST SUIT DOFFING)..... 3-8

    AFTERCARE ..... 3-8

    LATE DCS SYMPTOMS ..... 3-8

    RECURRENT DCS..... 3-10

DCS – OTHER DECOMPRESSION PROBLEMS  
(BAROTRAUMA)..... 3-10

DIARRHEA..... 3-11

EAR EXAMINATION..... 3-11

EAR PROBLEMS ..... 3-12

    CONGESTION (SEROUS OTITIS MEDIA) ..... 3-12

    INTERNAL INFECTION (OTITIS MEDIA) ..... 3-12

    EAR CANAL INFECTION (EXTERNAL OTITIS)..... 3-13

    RUPTURED EARDRUM ..... 3-14

    EAR/SINUS BLOCK ..... 3-14

ELECTROCARDIOGRAPH (ECG)  
(see CPR (CARDIOPULMONARY RESUSCITATION),  
CARDIAC MONITORING, 1-8)

EYE EXAMINATION..... 3-15

EYE – PENETRATING OBJECT (see EMERGENCY, 1-38)

EYE PROBLEMS..... 3-17

    CHEMICAL BURN (SHUTTLE EMERGENCY EYEWASH  
    (SEE) GOGGLES)..... 3-17

    DECONTAMINATION PROCEDURES  
    (see CHEMICAL BURN, 3-17)

    FOREIGN BODY/ABRASION ..... 3-20

    CORNEAL ULCER ..... 3-22

C THRU E

HERPETIC VIRAL OCULAR INFECTION.....	3-23
IRRITATION/INFECTION.....	3-23

FIGURES

3-1	Right eardrum .....	3-11
3-2	Eyelid eversion.....	3-16
3-3	Eye landmarks; 6 directions of gaze .....	3-16
3-4	SEE-WCS connection.....	3-18

C T H R U E

## CONGESTION, NASAL

Decongestants:

Med            Sudafed (Drug Subpack-Spine)  
Locker        (see WARNING, 6-12)  
                  Dose: 2 tabs every 6 hr as needed

-OR-

Guaifenes PSE (Drug Subpack-44)  
Dose: 1 tab every 12 hr as needed

-OR-

Afrin (EENT Subpack-19)  
Dose: 1-2 sprays every 12 hr as needed

## CONSTIPATION

1. Increase fluid intake; exercise

Med            2. Dulcolax (oral) (Drug Subpack-30)  
Locker        Dose: 1-2 tabs as needed for constipation  
                  -OR-

Dulcolax (suppository) (Drug Subpack-39)  
Dose: One suppository as needed for constipation

## DCS – DECOMPRESSION SICKNESS (BENDS)

### NOTE

Symptoms of DCS may be result of:  
Inadequate or interrupted prebreathe time  
Inadequate nitrogen purge from EMU  
Strenuous, prolonged EVA  
Severe dehydration during EVA  
Loss of cabin or suit pressure.

Primary treatment principles consist of repressurization and 100% O2 over time

1. Determine if following symptoms present:

### CUFF CLASS

- 1: Mild pain, at single or multiple sites and/or single extremity paresthesia. Difficult to distinguish from suit pressure points.  
**Symptoms do not interfere with performance**
  - 2: Moderate CUFF 1 symptoms that **interfere with performance or symptoms that resolve upon repress**
  - 3: **Severe CUFF 1 symptoms** or migratory, truncal or multiple site paresthesia, unusual headache
  - 4: Serious symptoms – Central neurological (spotted vision, slurred speech, coordination difficulty, loss of sensation, headache, seizures, unconsciousness), cardiopulmonary (chest pain, cough, shortness of breath)
2. If symptoms present, go to 19.1 DCS TREATMENT (EVA, EMER PROCS)

## EVA DCS EXAMINATION SCORECARD

IN-SUIT EXAM (Questions 1-20 only); OUT-OF-SUIT EXAM (Questions 1-27)

**Instructions:**

- 1) To be performed at periodic intervals throughout DCS treatment procedure
- 2) Mark the MET time of examination in the top row
- 3) Circle any abnormal finding in the appropriate item row for the correct exam time column  
(AB = abnormal, NL = normal, Y = yes, N = no)
- 4) Report any abnormal findings during PMC
- 5) Questions 21 through 27 included only in DCS Out-of-Suit examinations
- 6) Questions in bold should be asked verbatim

**IN-SUIT EXAM**

Item #	Examination Question/ Challenge	MET time of examination											
<b>Level of Consciousness</b>													
<b>1</b>	<b>Level of Consciousness:</b> Crewmember is awake, alert and responsive	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
<b>2</b>	<b>Orientation: "What Flight Day or day of the week is it? What are the names and positions of your fellow Crewmembers?"</b> (Crewmember answers all correctly)	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
<b>Verbal Function</b>													
<b>3</b>	<b>Object Naming:</b> Crewmember correctly names three readily identifiable objects (Can be chosen at random from available objects)	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
<b>4</b>	<b>Reading, Speech:</b> Crewmember correctly reads from EVA checklist without difficulty and without speech changes or difficulties	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
<b>Emotional Status</b>													
<b>5</b>	The Crewmember's emotional display is normal and appropriate	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL

3-5

MED/ALL/GEN K

IN-SUIT EXAM (Cont)

Item #	Examination Question/ Challenge	MET time of examination											
<b>Symptoms</b> (Crewmember answers the following questions)													
6	“Do you have any chest pain?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
7	“Do you have any difficulty breathing?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
8	“Do you have any coughing, or pain with deep breathing?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
9	“Are you experiencing fatigue beyond what you would expect from the EVA?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
10	“Do you feel lightheaded or dizzy?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
11	“Do you have moderate to severe headache?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
12	“Do you have any pain beyond what you would expect from post-EVA suit discomfort?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
13	“Do you feel any tingling or numbness?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
14	“Do you have any trouble moving or weakness in your arms or legs?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
15	“Do you have any double vision, loss of vision, or distortion of vision?”	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N	Y N
<b>Facial Nerves</b>													
16	<b>Facial Muscles:</b> Crewmember raises eyebrows, squeezes eyes shut, and puffs up cheeks without difficulty	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
<b>Strength</b>													
17	<b>Arm Bending Strength:</b> Crewmember bends elbow, with palm toward face and holds for two seconds against examiner resistance. Repeat both sides	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
18	<b>Leg Bending Strength:</b> Crewmember bends knee and holds for two seconds against examiner resistance. Repeat both sides	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL

3-6

MED/ALL/GEN K

**IN-SUIT EXAM (Cont)**

Item #	Examination Question/ Challenge	MET time of examination											
<b>Coordination Functions</b>													
19	<b>Finger-to-Finger:</b> Starting with hands wide apart, Crewmember easily and accurately touches fingertips together with eyes closed		AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
<b>Sensory Function</b>													
20	<b>Gross Sensation:</b> Examiner squeezes Crewmember's forearms, feet, and knees through suit. Crewmember should feel squeezing of the forearms, feet, and knees		AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL

**OUT-OF-SUIT EXAM (Includes questions 21-27)**

21	<b>Blood Pressure:</b> Record systolic and diastolic blood pressure using blood pressure cuff, stethoscope												
22	<b>Pulse:</b> Record pulse in beats/minute												
23	<b>Temperature:</b> Record temperature												
24	<b>Breath Sounds:</b> Breath sounds should be equal and symmetric on both sides, without crackles or wheezes		AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
25	<b>Skin examination:</b> Skin should be normal in appearance without areas of abnormal paleness, redness, or marbling		AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
26	<b>Fine Sensation:</b> Crewmember uses own hands to identify any areas of altered sensation. No abnormal areas should be identified (If any, mark boundaries with pen)		AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL
27	<b>Hearing:</b> Crewmember identifies and touches snapping fingers without difficulty with eyes closed		AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL	AB NL

## DCS TREATMENT (POST SUIT DOFFING)

### NOTE

If patient experiencing difficulty breathing, chest discomfort, or is otherwise in distress, go to CPR (CARDIOPULMONARY RESUSCITATION), 1-6

1. Perform DCS OUT-OF-SUIT EXAM, 3-7

## AFTERCARE

### NOTE

Expect possible 'S66 CABIN PPO2' msg

1. Apply QDM and deliver oxygen for a total of 2 hours
2. ✓LEH O2 SPLY 1,2 (two) – OP
3. ✓LEH O2 vlv – OP
4. Minimize physical activity
5. Hydrate w/ 1 liter/hour for 2 hours orally or by Saline I.V. if unable to drink. (Breaking QDM seal to hydrate is allowed) (Refer to INJECTIONS, INTRAVENOUS (I.V.), 4-11)
6. Contact Surgeon (consider medications)
7. Perform BENDS TREATMENT ADAPTER (BTA) INSTALLATION; BTA PREP (EVA, EMU CONT PROCS) for possible RECURRENT DCS symptoms

## LATE DCS SYMPTOMS

Definition: First presentation of symptoms occurs Post-EVA after recompression and EMU doffing

### A. CUFF CLASS 1,2,3

### NOTE

If symptoms worsen or do not improve after 20 minutes of treatment, increase cabin pressure to 14.7 psi.

Expect possible 'S66 CABIN PPO2' msg

1. Apply QDM and deliver oxygen for a total of 2 hours
2. ✓LEH O2 SPLY 1,2 (two) – OP
3. ✓LEH O2 vlv – OP



4. Perform DCS OUT-OF-SUIT EXAM (3-7) every 15-30 minutes and report changes to Surgeon
5. Minimize physical activity
6. Hydrate w/ 1 liter/hour for 2 hours orally or by Saline I.V. if unable to drink. (Breaking QDM seal to hydrate is allowed) (Refer to INJECTIONS, INTRAVENOUS (I.V.), 4-11)
7. Perform BENDS TREATMENT ADAPTER (BTA) INSTALLATION; BTA PREP (EVA, EMU CONT PROCS) for DCS symptoms refractory to oxygen/hydration therapy
8. Contact Surgeon (consider medications and BTA Ops)

B. CUFF 4 (Cardiopulmonary and CNS symptoms)

NOTE

If patient experiencing difficulty breathing, chest discomfort, or is otherwise in distress, go to CPR (CARDIOPULMONARY RESUSCITATION), 1-6.

Expect possible 'S66 CABIN PPO2' msg

- C7  
C6,  
MO32M,  
MO69M
1. Apply QDM and deliver oxygen
  2. √LEH O2 SPLY 1,2 (two) – OP
  3. √LEH O2 vlv – OP

If docked to ISS and hatches open:

4. Remove ODS air duct extension from vestibule, stow
5. Close ODS Hatch per decal
6. √Equal vlv (two) – OFF, capped

NOTE

Expect possible 'S66 CABIN P', 'S66 CABIN PPO2' msgs

- C5
7. Increase cabin pressure to 15.56 psi  
DIRECT O2 vlv – OP  
When CABIN P = 15.56 psia,  
DIRECT O2 vlv – CL
  8. Perform DCS OUT-OF-SUIT EXAM (3-7) every 15-30 minutes and report changes to Surgeon
  9. Perform BENDS TREATMENT ADAPTER (BTA) INSTALLATION; BTA PREP (EVA, EMU CONT PROCS). Continue w/subsequent steps concurrently until ready to initiate BTA treatment in EMU

10. Minimize physical activity
11. Hydrate w/ 1 liter/hour for 2 hours orally or by Saline I.V. if unable to drink. (Breaking QDM seal to hydrate is allowed) (Refer to INJECTIONS, INTRAVENOUS (I.V.), 4-11)
12. Contact Surgeon (consider medications and BTA Ops)
13. Check MCC for DEORBIT PREP

### RECURRENT DCS

Definition: Reappearance of symptoms after initial treatment completed

FOR ALL CUFF SYMPTOMS: Go to LATE DCS SYMPTOMS CUFF 4 (3-9) treatment procedure

### **DCS – OTHER DECOMPRESSION PROBLEMS (BAROTRAUMA)**

#### NOTE

Symptoms result from reduction in barometric pressure and expansion of trapped gas. Symptoms may occur during decompression preceding EVA or following loss of cabin pressure

Symptoms: Abdominal distention, ear pain, inability to clear ear, loss of hearing acuity, sinus pain, toothache, jaw pain

If progressive pain:

1. REPRESS (DEPRESS/REPRESS Cue Card) thru AIRLOCK REPRESS
2. Go to PAIN RELIEF, 5-3

## DIARRHEA

1. Increase fluid intake
2. Avoid solid foods, milk products for 24 hr

If fever present:

3. Contact Surgeon

Antidiarrheal medication:

- Med Locker 4. Pepto-Bismol (Drug Subpack-31)  
Dose: 2 tabs every 30-60 min as needed to maximum of 16 tabs/day

-OR-

### NOTE

Use only if symptoms severe  
w/very frequent stools

Imodium (Drug Subpack-30)

Dose: 2 caps initially, then 1 cap after each unformed stool to maximum of 8 caps/day

5. Contact Surgeon

## EAR EXAMINATION

- Med Locker 1. Unstow Otoscope (EENT Subpack-32)
2. Evaluate eardrum (see fig 3-1). Determine presence of Congestion, Internal Infection, Ruptured Eardrum, or Ear Canal Infection and see appropriate procedure for treatment




Figure 3-1.- Right eardrum.

## **EAR PROBLEMS**

### **CONGESTION (SEROUS OTITIS MEDIA)**

Symptoms: Decreased hearing, feeling of fullness in ear, possible pain, inability to clear ear

Signs: Dull/retracted eardrum, bubbles behind eardrum

Decongestants:

- Med Locker 1. Sudafed (Drug Subpack-Spine)  
(see WARNING, 6-12)  
Dose: 2 tabs every 6 hr as needed  
-OR-  
Guaifenes PSE (Drug Subpack-44)  
Dose: 1 tab every 12 hr as needed  
-OR-  
Afrin (EENT Subpack-19)  
Dose: 1-2 sprays in each nostril every 12 hr as needed

### **INTERNAL INFECTION (OTITIS MEDIA)**

Symptoms: Pain, sensation of warmth, decreased hearing in affected ear

Signs: Red, bulging eardrum, no normal landmarks on exam (see illustration)

1. Contact Surgeon

Antibiotics:

- Med Locker 2. \*Augmentin (Drug Subpack-37)  
Dose: 1 cap every 8 hr for 10 days  
-OR-  
\*Bactrim DS (Drug Subpack-38)  
(see WARNING, 6-3)  
Dose: 1 tab every 12 hr for 10 days

Decongestants:

3. Sudafed (Drug Subpack-Spine)  
(see WARNING, 6-12)  
Dose: 2 tabs every 6 hr as needed  
-OR-  
Guaifenes PSE (Drug Subpack-44)  
Dose: 1 tab every 12 hr as needed

Pain medications:

4. Ascriptin (Drug Subpack-Spine)  
(see WARNING, 6-3)  
Dose: 1-2 tabs every 4 hr as needed

-OR-

- Tylenol (Drug Subpack-Spine)  
Dose: 1-2 tabs every 4 hr as needed

### EAR CANAL INFECTION (EXTERNAL OTITIS)

Symptoms: Pain, decreased hearing in affected ear

Signs: Redness, debris, pus in external ear canal, pain w/movement of ear

#### WARNING

Do not proceed w/steps 1 or 2 if ruptured eardrum suspected. Refer to RUPTURED EARDRUM, 3-14

- Med Locker
1. Pope Otowick (EENT Subpack-27):  
Insert Pope Otowick in affected ear canal w/tweezers (EENT Subpack-34)

Eardrops:

2. Tobradex (EENT Subpack-28)  
Note: Tobradex is a dual-purpose ophthalmic and ear medication  
Dose: Saturate Pope Otowick w/drops until fully expanded in ear canal. Repeat saturation 4X/day

Pain medications:

3. Ascriptin (Drug Subpack-Spine)  
(see WARNING, 6-3)  
Dose: 1-2 tabs every 4 hr as needed

-OR-

- Tylenol (Drug Subpack-Spine)  
Dose: 1-2 tabs every 4 hr as needed

-OR-

- \*Vicodin (Drug Subpack-21)  
(see WARNING, 6-2, 6-13)  
Dose: 1-2 tabs every 4-6 hr as needed

## RUPTURED EARDRUM

Symptoms: Pain, difficulty hearing

Signs: Perforated eardrum, blood on eardrum

1. Contact Surgeon

Antibiotics:

- Med Locker
2. \*Augmentin (Drug Subpack-37)  
(see WARNING, 6-3)  
Dose: 1 cap every 8 hr for 10 days  
-OR-  
\*Bactrim DS (Drug Subpack-38)  
(see WARNING, 6-3)  
Dose: 1 tab every 12 hr for 10 days

Pain medications:

3. Ascriptin (Drug Subpack-Spine)  
(see WARNING, 6-3)  
Dose: 1-2 tabs every 4 hr as needed  
-OR-  
Tylenol (Drug Subpack-Spine)  
Dose: 1-2 tabs every 4 hr as needed

## EAR/SINUS BLOCK

### NOTE

Symptoms may occur during recompression from EVA

Symptoms: Ear pain, inability to clear middle ear, loss of hearing acuity, sinus pain, facial pain

1. Stop recompression; gently attempt equalization of pressure in middle ear

If unsuccessful or sinus pain present:

2. If possible, depressurize airlock until symptoms relieved
3. Slowly resume recompression
4. Contact Surgeon

## EYE EXAMINATION

### WARNING

For chemical burn or penetrating object, immediately refer to appropriate procedure (EYE – PENETRATING OBJECT, 1-38, or CHEMICAL BURN, 3-17)

1. Unstow:
- |        |  |  |
|--------|--|--|
| Med    | Otoscope   | (EENT Subpack-32)  |
| Locker | Ophthalmoscope Head                              | (EENT Subpack-5)   |
|        | Fluorescein Strips                               | (EENT Subpack-6)   |
|        | Cotton Swabs                                     | (EENT Subpack-34)  |
|        | Gauze Pads                                       | (EENT Subpack-2,<br>IV Admin-14,<br>Airway Subpack-11,<br>Trauma Subpack-14) |
|        | Alcaine  | (EENT Subpack-12)  |
|        | Refresh Plus (artificial tears,<br>eye drops) or | (EENT Subpack-16)  |
|        | Magnifying Glasses                               | (EENT Subpack-17)  |
|        | Magnifying Glasses Strap                         | (EENT Subpack-1)   |

2. Remove Otoscope head; remove light source; attach Ophthalmoscope head

#### Appearance:

3. Observe lids, lashes, conjunctiva, sclera; note any matted material, redness, swelling of lids, excess tearing, etc
4. Note size of pupils, particularly any size difference of two pupils. Shine light into each eye, one at a time; note change in pupil size

#### Check for redness:

5. Observe white of eye for redness; note what quadrant(s) of eye is(are) red
6. Determine if one or both eyes affected

If in addition to redness there is blurred vision, pain, abnormal sensitivity to light, or matted material (pus) in eye:

7. Contact Surgeon

√For foreign body:

8. Observe upper and lower lids, cornea for foreign body
9. √Lower lid. Have patient look up, pull down on lid
10. √Upper lid. Have patient look down; place cotton swab at top of upper lid; grasp upper lashes. Pull out and up on lashes; fold lid over swab, which can then be removed (fig 3-2)
11. √Eyeball. Have patient look in all six directions of gaze (fig 3-3)
12. Return eyelid to its normal position by having patient blink or look up




Figure 3-2.- Eyelid eversion.




Figure 3-3.- Eye landmarks; 6 directions of gaze.



Fluorescein stain (check for corneal abrasion):

**WARNING**  
Do not use more than two drops  
Alcaine per treatment as corneal  
damage may occur. Do not rub  
eyes for 30 min following anesthetic  
administration

13. \*Alcaine (EENT Subpack-12)  
Dose: 1-2 drops in affected eye

**WARNING**  
Do NOT repeat Alcaine dose w/o  
contacting Surgeon

14. Place 1-2 drops of Refresh Plus artificial tears on  
Fluorescein strip. Touch strip to lower, inner  
eyelid until yellow-green film covers eye
15. Use blue light setting on Ophthalmoscope; shine  
light on cornea, sclera
16. Area of more intense staining, which does not  
clear w/blinking, may be abrasion
17. Rinse twice w/Refresh Plus artificial tears

## **EYE PROBLEMS**

### CHEMICAL BURN (SHUTTLE EMERGENCY EYEWASH (SEE) GOGGLES)

1. Unstow:  
Food                      Drinking water container, straw  
Lockers  
CCK                        SEE (goggles, tubing, connectors), pH strip  
Misc                        Towels  
Stowage
2. Prior to SEE configuration, use drinking water  
container, straw for eye irrigation. Use towels to  
prevent water from escaping
3. Remove PHS hose QD from Galley auxiliary port,  
making sure orange stopcock is perpendicular to  
direction of flow; replace w/SEE QD
4. Turn Galley knob MV2 cw until it stops (ambient)

5. Twist lock SEE-WCS connector to Waste Collection System (WCS) urinal hose




Figure 3-4.- SEE-WCS connection.

6. Don SEE goggles (place outflow side of goggles on affected eye)
7. Configure WCS for urine collection
8. Activate water flow by turning orange stopcock parallel to direction of flow
9. Flush eyes – keep eyes open

If unable to keep affected eye open due to pain, apply:

- Med  
Locker
10. \*Alcaine (EENT Subpack-12)  
Dose: 1-2 drops in affected eye

**WARNING**  
Do NOT repeat Alcaine dose w/o  
contacting Surgeon

11. Continue flushing for at least 10 min. Contact Surgeon
12. During (after) SEE use, absorb excess water w/towels

If obstruction exists in WCS urine hose:

13. Hose filter may need to be cleaned/replaced
14. Touch sclera w/pH strip to wet  
If pH neutral, go to step 15  
If pH not neutral, continue flushing w/SEE until pH neutral
15. Perform EYE EXAMINATION, Fluorescein stain, steps 13 thru 17, 3-17, for affected eye(s)

**WARNING**

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

If corneal staining is seen w/Fluorescein stain:

- Med Locker
16. If available, use Video Camera for photo-documentation (closeup and macroviews)
  17. \*Cyclogyl (EENT Subpack-11)  
Dose: 1 drop in affected eye
  18. \*Ciloxan Ointment (EENT Subpack-15)  
Dose: Apply 1/2-inch ribbon of ointment in lower lid 3X/day
  19. Repeat Fluorescein stain every 12 hr until staining no longer seen

For pain:

- Med Locker
20. \*Vicodin (Drug Subpack-21)  
(see WARNING, 6-2, 6-13)  
Dose: 1-2 tabs every 4-6 hr as needed

21. Restain eye in approx 12 hr  
Contact Surgeon w/results

If no corneal staining seen w/Fluorescein stain:

- Med Locker
22. \*Ciloxan Ointment (EENT Subpack-15)  
Dose: Apply 1/2-inch ribbon of ointment in lower lid 3X/day
  23. Restain eye in 6-12 hr  
Contact Surgeon w/results

FOREIGN BODY/ABRASION

Symptoms: "Grain of sand" feeling under eyelid, pain, sensitivity to light, blurred vision

Signs: Red eye, excess tears, may see foreign body; area of intense staining on Fluorescein examination, usually irregular in shape (see EYE EXAMINATION, 3-11)

1. Contact Surgeon

WARNING  
Do not use more than two drops Alcaine per treatment as corneal damage may occur. Do not rub eyes for 30 min following anesthetic administration

Pain relief:

- Med Locker
2. \*Alcaine (EENT Subpack-12)  
Dose: 1-2 drops in affected eye

WARNING  
Do NOT repeat Alcaine dose w/o contacting Surgeon

Mechanical removal:

If foreign body seen:

3. Try to remove by flushing eye w/Refresh Plus artificial tears (EENT Subpack-16). Blot excess fluid

If unsuccessful:

- Med Locker
4. Moisten Cotton Swab (EENT Subpack-34); try to gently dislodge foreign body by rotating swab
  5. If everted, return eyelid to its normal position by having patient blink
  6. Stain eye w/Fluorescein strip (EENT Subpack-6), using blue light examination (see EYE EXAMINATION, steps 14-15, 3-17)

If unsuccessful in removing foreign body:

7. Contact Surgeon

**WARNING**

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

- If cornea shows scratch highlighted by flourescein:
- Med Locker
8. \*Ciloxan Ointment (EENT Subpack-15)  
Dose: Apply 1/2-inch ribbon of ointment in lower lid 3X/day  
-OR-  
\*Genoptic Ophthalmic Ointment (EENT Subpack-14)  
Dose: Apply small amount to eye prior to patching
  9. \*Cyclogyl (EENT Subpack-11)  
Dose: 1 drop in affected eye

**WARNING**

Do not patch if signs of infection present, patient wears contact lens, or corneal ulcer suspected

- Patch eye if needed for relief of discomfort:
- Med Locker
10. Unstow:  
Eye Pads (2) (EENT Subpack-4)  
Tape (EENT Subpack-9,  
IV Admin-13,  
Airway Subpack-9,  
Trauma Subpack-34)
  11. Fold one Eye Pad in half; place over closed eyelid; cover w/second unfolded pad
  12. Tape in place firmly enough so that patient cannot blink eyelid
  13. Remove patch after 24 hr; repeat Fluorescein eye exam

## CORNEAL ULCER

Symptoms: Marked eye discomfort, pain, moderate-to-significant sensitivity to light, blurred vision

### NOTE

Overnight wear of contact lenses can result in corneal ulcer

Signs: Marked redness of eye, eye examination shows white lesion on cornea, and Fluorescein examination shows intensely staining well-circumscribed round lesion on cornea (see EYE EXAMINATION, Fluorescein stain, steps 13 thru 17, 3-17)

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

1. Contact Surgeon

### WARNING

Do not patch eye

Antibiotics:

Med Locker 2. \*Ciloxan Ointment (EENT Subpack-15)  
Dose: Apply 1/2-inch ribbon of ointment in lower lid 3X/day

Pain relief:

3. \*Vicodin (EENT Subpack-21)  
(see WARNING, 6-2, 6-13)  
Dose: 1-2 tabs every 4-6 hr as needed for pain

4. Do not use contact lenses if ulcer suspected

HERPETIC VIRAL OCULAR INFECTION

Symptoms: Tearing, photophobia, pain

Signs: Red eye, corneal ulcers showing branch-like pattern when Fluorescein-stained

1. Contact Surgeon
2. Perform EYE EXAMINATION, Fluorescein stain, steps 13 thru 17, 3-17

<p><u>WARNING</u> ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use</p>
---

If Fluorescein stain suggestive of herpetic keratitis:

- Med Locker
3. \*Viroptic (Drug Subpack-1)  
Dose: 1 drop every 3 hr; max 9 drops/day
  4. \*^Ciloxan Ointment (EENT Subpack-15)  
Dose: Apply 1/2-inch ribbon of ointment in lower lid 3X/day

IRRITATION/INFECTION

Symptoms: Mild discomfort (if any), burning, itching, blurred vision

Signs: Red eye, "matted" material in lids and at corners of eye, swelling around eye


1. Contact Surgeon

Antibiotics:

- Med Locker
2. \*Genoptic Ophthalmic Ointment (EENT Subpack-14)  
Dose: Apply small amount to eye 2X/day



This Page Intentionally Blank





SECTION 4

F THRU N

FRACTURE/DISLOCATION .....	4-3
HEADACHE (see PAIN RELIEF, 5-3)	
HEAD OR NECK INJURY (see EMERGENCY, 1-39)	
HEMORRHOID .....	4-5
HOT CABIN.....	4-5
INDIGESTION (UPSET STOMACH) .....	4-6
INJECTIONS.....	4-7
INTRAVENOUS (I.V.) .....	4-7
CATHETER DRESSING CHANGES (DAILY).....	4-22
REMOVAL OF Y-TYPE CATHETER EXTENSION.....	4-23
INTRAMUSCULAR.....	4-24
SUBCUTANEOUS.....	4-26
PREFILLED SYRINGE TECHNIQUE .....	4-27
TUBEX INJECTOR TECHNIQUE .....	4-28
INTUBATION (ILMA)	
(see AIRWAY(ILMA) CUE CARD, CC 9-4)	
INTUBATION, ENDOTRACHEAL (see CPR	
(CARDIOPULMONARY RESUSCITATION), 1-12)	
LACERATIONS.....	4-30
MAJOR LACERATIONS.....	4-30
MINOR LACERATIONS .....	4-37
MOTION SICKNESS (also see VOMITING, 5-20) .....	4-38
MILD MOTION SICKNESS .....	4-38
MODERATE-TO-SEVERE MOTION SICKNESS .....	4-39
MUSCLE SPASM (DRUG SIDE EFFECT).....	4-39
NASAL CONGESTION (see CONGESTION, NASAL, 3-3)	
NAUSEA/VOMITING .....	4-39
NOSEBLEED .....	4-40

F THRU N

FIGURES

4-1	Existing I.V. flow.....	4-7
4-2	I.V. assembly with Y-type Catheter Extension .....	4-8
4-3	Medication administration technique.....	4-9
4-4	Tourniquet placement .....	4-13
4-5	Removal of I.V. needle.....	4-14
4-6	Surface anatomy .....	4-25
4-7	Prefilled syringe .....	4-27
4-8	Tubex injector .....	4-29
4-9	Suture placement.....	4-32
4-10	Laceration closure using skin stapler.....	4-33
4-11	Skin staple remover .....	4-36
4-12	Nasal speculum .....	4-40
4-13	Nasal packing .....	4-42
4-14	Nasal catheter insertion .....	4-43

F T H R U N

## FRACTURE/DISLOCATION

- \* If no pulse or respiration, perform CPR \*
- \* using patient/rescuer restraints (see \*
- \* CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), 1-6) \*
- \* \*
- \* If unconscious and patient cannot \*
- \* maintain airway, insert oral airway (see \*
- \* CPR (CARDIOPULMONARY \*
- \* RESUSCITATION), ORAL AIRWAY, \*
- \* 1-11), or INTUBATION, \*
- \* ENDOTRACHEAL, 1-12); provide \*
- \* manual ventilation using \*
- \* RESUSCITATOR (see inside cover of \*
- \* Medical Accessory Kit or Cue Card \*
- \* section) \*
- \* \*
- \* If conscious, provide supplemental O2 \*
- \* with QDM \*

1. Control bleeding w/direct pressure, Gauze Pads  
(Trauma Subpack-14)

2. Unstow:

Med Locker	Blood Pressure Cuff	(IV Admin-1)
	Stethoscope	(Airway Subpack-12)
	Thermometer	(EENT Subpack-3)
	Gauze Pads	(Trauma Subpack-14, IV Admin-14, Airway Subpack-11, EENT Subpack-2)

3. Evaluate vital signs

MET: \_\_\_\_\_

Blood Pressure: \_\_\_\_\_

Pulse: \_\_\_\_\_

Respiratory Rate: \_\_\_\_\_

Temperature: \_\_\_\_\_

- |               |                  |              |                       |
|---------------|------------------|--------------|-----------------------|
| Med<br>Locker | 4. Bandage with: |              |                       |
|               |                  | Kling        | (Trauma Subpack-12)   |
|               |                  | Kerlix       | (Trauma Subpack-2)    |
|               |                  | Ace Bandages | (Trauma Subpack-3,13) |
|               |                  | Telfa Pads   | EENT Subpack-22)      |

**WARNING**  
 Fractures of pelvis, upper leg can result in large blood loss and shock. Dislocations treated as fractures

If fracture open (bone protruding thru skin or caused by projectile):

- |               |                  |              |  |
|---------------|------------------|--------------|--|
| Med<br>Locker | 5. Bandage with: |              |  |
|               |                  | Gauze Pads   | (Trauma Subpack-14,<br>IV Admin-14,<br>Airway Subpack-11,<br>EENT Subpack-2) |
|               |                  | Kerlix       | (Trauma Subpack-2)   |
|               |                  | Kling        | (Trauma Subpack-12)  |
|               |                  | Ace Bandages | (Trauma Subpack-3,13)  |

6. ✓Pulse distal to fracture location

If no pulse or if joint dislocated:

7. Contact Surgeon immediately for instructions to set fracture

If pulse intact:

8. Splint fracture

9. Splint w/Kerlix, Kling, Ace, any available rigid materials such as FDF Cover, rolled checklist, IFM tools, etc.

10. If finger fracture suspected, splint finger w/Finger Splint (Trauma Subpack-23)

11. Contact Surgeon

## HEMORRHOID

Signs: Anal pain or itching

- Med Locker
1. Anusol HC Suppositories (Drug Subpack-41)  
Dose: 1 suppository every 12 hr as needed
  2. Drink plenty of fluids to prevent constipation

## HOT CABIN (Total loss thermal cntl – TIG > 2.5 hr)

1. Wear minimal clothing; limit physical activity
- Med Locker
2. Unstow Skin Temp Monitors (EENT Subpack-23).  
Temp which appears in green is actual temp to be recorded
  3. After drying skin, apply Skin Temp Monitors to forehead or trunk
  4. Record pulse rate and skin temp every hr
  5. All crewmembers immediately begin drinking 8-oz chilled water every 15 min. Consume one Salt Tab (Drug Subpack-Spine) w/every other container of water

## INDIGESTION (UPSET STOMACH)

### WARNING

Refer to CHEST PAIN, 1-36, for any associated chest pain, pressure  
or sweating  
or difficulty breathing  
or arm pain

Med Locker      1. Pepto-Bismol (Drug Subpack-31)  
Dose: 2 tabs every 30-60 min as needed to maximum of 16 tabs/day  
-OR-

Antacid:

2. Mylanta (Drug Subpack-35)  
Dose: 2-4 tabs well chewed between meals and at bedtime. May take 2 tabs between this schedule as needed to maximum of 12 tabs/day

## INJECTIONS

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

### INTRAVENOUS (I.V.)

- Butterfly Needle (one-time dose only)
- Intravenous Fluid Infusion w/Y-Type Catheter Extension and Administration Set
- Medication Administration w/Existing I.V. Flow
- Removal of IV Administration Set/Intracatheter w/Y-Type Catheter Extension Set left in place
- Medication Administration thru Y-Type Catheter Extension Set
- Y-Type Catheter Extension Maintenance Flush

If necessary:

1. Perform CPR (CARDIOPULMONARY RESUSCITATION), CARDIAC MONITORING, Donning, 1-8, then:
2. Contact Flight Surgeon before giving any I.V. medication




Figure 4-1.- Existing I.V. flow.




Figure 4-2.- I.V. assembly with Y-type Catheter Extension.

Butterfly Needle Procedure (one-time dose only)  
DO NOT leave in vein

- |            |   |   |
|------------|---|---|
| Misc       | 1. Prepare Gray Tape to secure supplies                               |   |
| Stowage    | 2. Unstow:  |   |
| Med Locker | Tubex Injector  | (Drug Subpack-19, Drug Subpack-Spine, Trauma Subpack-31)                                |
|            | Injectable Medication<br>(Contact Surgeon for appropriate medication) | (Drug Subpack, Trauma Subpack, EENT Subpack)  |
|            | Gloves, non-sterile   | (IV Admin-12, Airway Subpack-8, EENT Subpack-7, CCK)                                    |
|            | ^Povidone-Iodine Swab   | (IV Admin-19, Airway Subpack-10, Trauma Subpack-7)                                      |
|            | Alcohol Pads  | (IV Admin-18, Airway Subpack-10, Drug Subpack-16, Drug Subpack-Spine, Trauma Subpack-6) |
|            | Butterfly Needle, 21ga  | (IV Admin-17)   |
|            | Penrose Tubing (tourniquet)   | (IV Admin-16)   |



Gauze Pads

(IV Admin-14,  
Airway Subpack-11,  
Trauma Subpack-14,  
EENT Subpack-2)

Bandaid

(IV Admin-20,  
Drug Subpack-Spine,  
Drug Subpack-12,  
Trauma Subpack-11)

Sharps Container

3. Open Butterfly Needle, temp stow package
4. Load Tubex Injector w/desired medication and unsheathe needle (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28)

Unstow Butterfly Needle (IV Admin-17)




Figure 4-3.- Medication administration technique.

5. Insert needle of Medication Tubex Injector into rubber port of Butterfly Needle
6. Remove air bubbles from tubing by expelling 1-2 drops of medication. Temp stow assembly
7. Tighten tourniquet around upper arm. Don Gloves
8. Select site. Cleanse site w/ Povidone-Iodine Swab; fan to dry. Repeat w/Alcohol Pad; fan to dry
9. Unsheathe Butterfly Needle. Capture vein w/thumb and index finger of non-dominant hand. W/needle bevel up, while holding plastic wings, puncture skin over vein
10. Pull back Medication Tubex Injector plunger, check for blood return

NOTE

If no blood visible, gentle traction may be used on needle to establish flow.  
Procedure may need to be repeated

11. REMOVE TOURNIQUET. Inject medication. Retrieve Gauze Pads, remove Butterfly Needle quickly, apply pressure w/Gauze Pads
12. Withdraw Medication Tubex Injector from Butterfly port. Carefully remove medication cartridge and place in Sharps container. Return Tubex Injector to Drug Subpack
13. Return Butterfly Needle to package, wrap securely w/Gray Tape and ensure it is returned to IV Admin Kit. Do not put needles in Wet Trash

Intravenous Fluid Infusion w/Y-Type Catheter Extension and Administration Set

Equipment setup:

Misc 1. Prepare Gray Tape to secure supplies

Stowage

2. Consult Flight Surgeon for appropriate saline quantity

3. Unstow:

Med Locker	Saline, 500 ml Bag OR 250 ml Bag	(IV Admin-1, Saline Supply Bag) (Middeck Locker)
	I.V. Administration Set (Transfer Set)	(IV Admin-17)
	Blood Pressure Cuff	(IV Admin-1)
	Gauze Pads	(IV Admin-14, Airway Subpack-11, Trauma Subpack-14, EENT Subpack-2)
	Lever Lock Cannula	(IV Admin-17)
	Y-Type Catheter Extension Set	(IV Admin-17)
	Gloves, non-sterile	(IV Admin-12, Airway Subpack-8, CCK, EENT Subpack-7)
	^Povidone-Iodine Swab	(IV Admin-19, Airway Subpack-10, Trauma Subpack-7)
	Alcohol Pads	(IV Admin-18, Airway Subpack-10, Drug Subpack-Spine, Drug Subpack-16, Trauma Subpack-6)
	I.V. Intracatheter:	
	18ga OR	(IV Admin-8,9)
	20ga	(IV Admin-10,11)
	Penrose Tubing (tourniquet)	(IV Admin-16)
	Op Site (transparent dressing)	(IV Admin-14,
	Tape (Four 4-inch strips)	(IV Admin-13, Airway Subpack-9, Trauma Subpack-34, EENT Subpack-9)

4. Remove from packages:
  - Saline Bag
  - I.V. Administration Set (Transfer Set)
  - Y-Type Catheter Extension Set

NOTE

Make sure roller clamp on I.V.  
Administration Set is CLOSED  
(roller is snug at narrow end)

5. Remove saline bag cap (blue) from Saline bag.  
Remove cap from spike on I.V. Administration Set  
(Transfer Set). Plug spike into Saline port (see  
fig 4-1)
6. Relocate and open Lever Lock Cannula package.  
ONLY TOUCH EDGES OF CANNULA
7. Remove cap on free end of I.V. Administration Set  
and connect to uncovered end of Lever Lock  
Cannula (tighten w/clockwise rotation)
8. Clean one access port on Y-Type Catheter Extension  
Set w/Alcohol Pad
9. Remove cap on Lever Lock Cannula and insert into  
access port on Y-Type Catheter Extension Set

NOTE

Do NOT remove cap on  
Y-Type Catheter Extension

10. Open Gauze Pad package. Open roller clamp;  
squeeze Saline bag to prime I.V. line and purge air;  
absorb Saline w/ Gauze Pad. Close roller clamp
11. Wrap Blood Pressure Cuff tightly around bag; inflate  
to 100 mmHg

NOTE

Blood Pressure Cuff can be removed later  
for patient use by clamping I.V. line shut.  
Roller clamp can be left CLOSED while  
patient's blood pressure being taken.  
When roller clamp is OPEN, bag can be  
manually squeezed to infuse Saline

Patient preparation:

12. Place tourniquet around upper arm (see fig 4-4). Don Gloves. Relocate open Gauze. Relocate Sharps Container




Figure 4-4.- Tourniquet placement.

13. Select site. Cleanse site w/ Povidone-Iodine Swab; fan to dry. Repeat w/ Alcohol Pad; fan to dry
14. Remove I.V. Intracatheter from package.
15. Rotate Teflon catheter 360 deg in its sheath to loosen catheter
16. Capture vein w/ thumb and index finger of non-dominant hand. W/ needle bevel up, puncture skin over vein. Observe for blood return in casing

NOTE

If no blood visible, gentle traction may be used on needle to establish flow.  
Procedure may need to be repeated

17. Drop angle of needle and advance 1-2 mm more to ensure catheter properly placed in vein
18. Advance/slide Teflon catheter off needle into vein while holding needle stationary

NOTE

If resistance met, **DO NOT FORCE** catheter into vein. If unable to establish flow, dispose of catheter and repeat procedure at a new site w/new equipment

19. While holding needle in place, REMOVE TOURNIQUET FROM UPPER ARM
20. Remove cap from free end of Y-Type Catheter Extension Set
21. Occlude vein by applying pressure above catheter tip w/finger
22. Remove I.V. needle leaving catheter in vein and quickly attach Y-Type Catheter Extension Set. Stop Occlusion (see fig 4-5)



Figure 4-5.- Removal of I.V. needle.

23. Discard I.V. Needle into Sharps container. Clean excess blood at site w/Gauze
24. Secure Catheter Assembly to skin w/Opsite (transparent dressing) or a tape "chevron"
25. Secure excess I.V. tubing by looping from site to wrist area and back to site w/tape "chevrons"
26. OPEN roller clamp. Adjust rate per Surgeon consultation
27. Refer to INJECTIONS, CATHETER DRESSING CHANGES (DAILY), 4-22

28. For removal procedures, refer to INJECTIONS, REMOVAL OF Y-TYPE CATHETER EXTENSION, 4-23

NOTE

If NO flow established, gently manipulate catheter to re-establish flow. If swelling around site is visible or if unable to re-establish flow, remove catheter, dispose of set and repeat procedure at a new site

Medication Administration with Existing I.V. Flow

- |  |   |                     |  |              |   |                |  |  |  |                  |  |
|--|---|---------------------|--|--------------|---|----------------|--|--|--|------------------|--|
| Misc<br>Stowage  | <ol style="list-style-type: none"> <li>1. Prepare Gray Tape to secure supplies</li> <li>2. Contact Flight Surgeon before giving any I.V. medication</li> </ol>  |                     |  |              |   |                |  |  |  |                  |  |
| Med<br>Locker  | <ol style="list-style-type: none"> <li>3. Unstow:           <table border="0" style="margin-left: 20px;"> <tr> <td style="padding-right: 20px;">Gloves, non-sterile</td> <td>(IV Admin-12,<br/>Airway Subpack-8,<br/>CCK, EENT Subpack-7)</td> </tr> <tr> <td>Alcohol Pads</td> <td>(IV Admin-18,<br/>Airway Subpack-10,<br/>Drug Subpack-Spine,<br/>Drug Subpack-16,<br/>Trauma Subpack-6)</td> </tr> <tr> <td>Tubex Injector</td> <td>(Drug Subpack-19,<br/>Drug Subpack-Spine,<br/>Trauma Subpack-31)</td> </tr> <tr> <td>Injectable Medication or<br/>Prefilled Syringe<br/>(Contact Surgeon for<br/>appropriate medication)</td> <td>(Drug Subpack,<br/>Trauma Subpack,<br/>EENT Subpack)</td> </tr> <tr> <td>Sharps Container</td> <td></td> </tr> </table> </li> <li>4. Don Gloves</li> <li>5. Load Tubex Injector w/desired medication cartridge or attach plunger to prefilled syringe (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28, or PREFILLED SYRINGE TECHNIQUE, 4-27)</li> </ol> | Gloves, non-sterile | (IV Admin-12,<br>Airway Subpack-8,<br>CCK, EENT Subpack-7) | Alcohol Pads | (IV Admin-18,<br>Airway Subpack-10,<br>Drug Subpack-Spine,<br>Drug Subpack-16,<br>Trauma Subpack-6) | Tubex Injector | (Drug Subpack-19,<br>Drug Subpack-Spine,<br>Trauma Subpack-31) | Injectable Medication or<br>Prefilled Syringe<br>(Contact Surgeon for<br>appropriate medication) | (Drug Subpack,<br>Trauma Subpack,<br>EENT Subpack) | Sharps Container |  |
| Gloves, non-sterile  | (IV Admin-12,<br>Airway Subpack-8,<br>CCK, EENT Subpack-7)  |                     |  |              |   |                |  |  |  |                  |  |
| Alcohol Pads   | (IV Admin-18,<br>Airway Subpack-10,<br>Drug Subpack-Spine,<br>Drug Subpack-16,<br>Trauma Subpack-6)   |                     |  |              |   |                |  |  |  |                  |  |
| Tubex Injector   | (Drug Subpack-19,<br>Drug Subpack-Spine,<br>Trauma Subpack-31)  |                     |  |              |   |                |  |  |  |                  |  |
| Injectable Medication or<br>Prefilled Syringe<br>(Contact Surgeon for<br>appropriate medication) | (Drug Subpack,<br>Trauma Subpack,<br>EENT Subpack)  |                     |  |              |   |                |  |  |  |                  |  |
| Sharps Container   |   |                     |  |              |   |                |  |  |  |                  |  |

6. Stop flow to I.V. Assembly, CLOSE roller clamp on I.V. tubing (see fig 4-1)
7. Clean free access port on Y-type Catheter Extension OR I.V. Administration Set w/Alcohol Pad
8. Unsheathe Medication Tubex Injector Needle and insert into I.V. port
9. Inject medication. Withdraw Medication Tubex Injector from I.V. port. Open roller clamp to resume flow. Carefully remove medication cartridge or prefilled syringe and place in Sharps container
10. Return Tubex Injector to Drug Subpack



Removal of I.V. Administration Set/Intracatheter w/Y-Type Catheter Extension Set left in place

- Misc 1. Prepare Gray Tape to secure supplies  
Stowage
2. Contact Flight Surgeon before removing I.V. Administration Set
3. Unstow:
- |            |                     |   |
|------------|---------------------|---|
| Med Locker | Gloves, non-sterile | (IV Admin-12, Airway Subpack-8, CCK, EENT Subpack-7)                |
|            | Gauze Pads          | (IV Admin-14, Airway Subpack 11, Trauma Subpack 14, EENT Subpack-2) |
4. Stop flow to I.V. Assembly, CLOSE roller clamp on I.V. tubing (see fig 4-1)
5. Close clamps on Y-Type Extension Set by sliding clamp over tubing to lock
6. Untape I.V. Administration tubing from arm, leaving Intracatheter w/Y-Type Catheter Extension Set secure
7. Don Gloves
8. Remove Lever Lock Cannula and I.V. Administration Set from Y-Type Catheter Extension Set port by disconnecting Lever Lock Cannula and Y-Type Catheter Extension Set
9. Wrap empty Saline bag and I.V. Administration Set w/Gray Tape and discard

NOTE

Y-Type Catheter Extension Set must be flushed w/Saline every 8 hr. Medication administration can be substituted as a flushing procedure

Medication Administration thru Y-Type Catheter Extension Set

- Misc Stowage
1. Prepare Gray Tape to secure supplies
  2. Contact Flight Surgeon before giving any I.V. medication
  3. Unstow:

Med Locker	Syringe (3 cc)	(IV Admin-2,3,4,5,6,7)
	Saline, 500 ml Bag OR	(IV Admin-1, Saline Supply Bag)
	250 ml Bag	(Middeck Locker)
	Tubex Injector	(Drug Subpack-19, Drug Subpack-Spine, Trauma Subpack-31)
	Injectable Medication or Prefilled Syringe (Contact Surgeon for appropriate medication)	(Drug Subpack, Trauma Subpack, EENT Subpack)
	Alcohol Pads	(IV Admin-18, Airway Subpack-10, Drug Subpack-16, Drug Subpack-Spine, Trauma Subpack-6)
	Sharps Container	
  4. Load Tubex Injector w/desired medication cartridge or attach plunger to prefilled syringe (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28, or PREFILLED SYRINGE TECHNIQUE, 4-27)
  5. Clean rubber medication port of Saline bag w/Alcohol Pad. Uncap 3-cc syringe, temp stow cap on Gray Tape
  6. Fill 3-cc syringe w/saline from Saline bag and recap syringe needle by spearing cap; temp stow
  7. Clean free access port on Y-Type Catheter Extension Set w/Alcohol Pad
  8. Unsheathe Medication Tubex Injector Needle and insert into Y-Type Catheter Extension Set port
  9. Unclamp tubing w/sliding lock; inject medication from Medication Tubex Injector

NOTE

If NO flow established, gently  
manipulate catheter to re-establish flow.  
If swelling around site visible or if unable  
to re-establish flow, remove catheter,  
dispose of set and repeat procedure at  
new site

10. Reclamp tubing and withdraw needle. Re-sheath needle by spearing cap on Gray Tape; temp stow
11. Uncap 3-cc syringe; insert into Y-Type Catheter Extension Set port and unclamp tubing w/sliding lock
12. OPEN roller clamp. Inject entire amount saline from 3-cc syringe. Reclamp tubing and remove 3-cc syringe. Discard syringe with needle in Sharps container
13. Carefully disassemble Medication Tubex Injector. Remove medication cartridge w/needle or prefilled syringe and place in Sharps container
14. Return Tubex Injector to Drug Subpack

Y-Type Catheter Extension Maintenance Flush

NOTE

Y-Type Catheter Extension device must be flushed w/Saline every 8 hr. Medication administration can be substituted as a flushing procedure

Misc 1. Prepare Gray Tape to secure supplies  
Stowage

2. Unstow:  
Med Syringe (3 cc) (IV Admin-2,3,4,5,6,7)  
Locker Saline, 500 ml Bag OR (IV Admin-1,  
Saline Supply Bag)  
250 ml Bag (Middeck Locker)  
Alcohol Pads (IV Admin-18,  
Airway Subpack-10,  
Drug Subpack-16,  
Drug Subpack-Spine,  
Trauma Subpack-6)  
Sharps Container

3. Clean rubber medication port of Saline bag w/Alcohol Pad. Uncap 3-cc syringe, temp stow cap on Gray Tape
4. Fill 3-cc syringe w/saline from Saline bag and recap syringe needle by spearing cap; temp stow
5. Clean access port on Y-Type Catheter Extension Set w/Alcohol Pad
6. Uncap 3-cc syringe; insert into Y-Type Catheter Extension Set port and unclamp tubing w/sliding lock
7. Inject saline from 3-cc syringe until NO BLOOD is visible in tubing

NOTE

If NO flow established, gently manipulate catheter to re-establish flow. If swelling around site visible or if unable to re-establish flow, remove catheter, dispose of set and repeat procedure at new site

8. Reclamp tubing w/sliding lock and remove 3-cc syringe
9. If blood visible in second access port, repeat 3-cc syringe flush on second port
10. Discard syringe(s) in Sharps container

NOTE

A small amount of blood may remain in tubing close to insertion site

CATHETER DRESSING CHANGES (DAILY)

1. Unstow:
- |            |                     |  |
|------------|---------------------|--|
| Med Locker | Gauze Pads          | (IV Admin-14,<br>Airway Subpack-11,<br>Trauma Subpack-14,<br>EENT Subpack-2) |
|            | Tape                | (IV Admin-13,<br>Airway Subpack-9,<br>Trauma Subpack-34,<br>EENT Subpack-9)  |
|            | Gloves, non-sterile | (IV Admin-12,<br>Airway Subpack-8, CCK,<br>EENT Subpack-7)                   |

NOTE

W/routine flushing, saline lock (Y-type Catheter Extension) is normally kept in place for 3 days, or until evidence of infection



2. Don Gloves
3. Remove existing gauze dressing
4. Examine site for evidence of infection. Consult Flight Surgeon
5. Tape new folded Gauze Pad in place
6. Use Coban (IV Admin-1) over dressing for stability, if desired



## INTRAMUSCULAR

1. Unstow:
  - Med Locker Tubex Injector (Drug Subpack-19, Drug Subpack-Spine, Trauma Subpack-31)
  - Injectable Medication (Drug Subpack, Trauma Subpack, EENT Subpack)
  - Alcohol Pads (IV Admin-18, Airway Subpack-10, Drug Subpack-16, Drug Subpack-Spine, Trauma Subpack-6)
  - Gauze Pads (IV Admin-14, Airway Subpack 11, Trauma Subpack 14, EENT Subpack-2)
2. Recommended site is upper outer quadrant of buttocks (see fig 4-6, Surface anatomy, 4-25); shoulder may also be used w/nonirritating medication. Consult w/Surgeon for nonirritating medications
3. Load Tubex Injector w/desired medication (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28)
4. Clean wide area w/Alcohol Pad
5. Insert needle w/quick thrust (perpendicular to muscle) to 3/4 of needle length
6. ✓For blood return
  - If blood return:
    7. Withdraw needle; expel blood onto Gauze Pad; reinsert needle
  - If no blood return:
    8. Inject medication over several sec
9. Quickly remove needle
10. Apply pressure to injection site (2-3 min) w/Alcohol Pad or Gauze Pad
11. Carefully remove medication cartridge from Tubex Injector
12. Dispose of medication cartridge in Sharps container (Med Locker). Restow Tubex Injector to original location





N-45 = Shoulder (deltoid) injection site  
 K-54 = Buttocks injection site

Figure 4-6.- Surface anatomy.

## SUBCUTANEOUS

1. Unstow:
    - Med Locker Tubex Injector (Drug Subpack-19, Drug Subpack-Spine, Trauma Subpack-31)
    - Injectable Medication (Drug Subpack, Trauma Subpack, EENT Subpack)
    - Alcohol Pads (IV Admin-18, Airway Subpack-10, Drug Subpack-16, Drug Subpack-Spine, Trauma Subpack-6)
    - Gauze Pads (IV Admin-14, Airway Subpack 11, Trauma Subpack 14, EENT Subpack-2)
  2. Load Tubex Injector w/desired medication (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28)
  3. Clean shoulder area w/Alcohol Pad (see fig 4-6, Surface anatomy, 4-25)
  4. Remove cap from needle. Pinch skin between thumb, fingers of free hand
  5. Insert needle w/quick thrust to 1/2 needle length at shallow (no more than 30 deg) angle to skin
  6. ✓For blood return
- If blood return:
7. Withdraw needle; expel blood onto Gauze Pad; insert needle again
- If no blood return:
8. Inject medication over several sec. Should produce bump under skin
  9. Carefully remove medication cartridge from Tubex Injector
  10. Dispose of medication cartridge in Sharps container (Med Locker). Restow Tubex Injector to original location

**PREFILLED SYRINGE TECHNIQUE**

- Med Locker
1. Unstow:  
Syringe, Plunger (Drug Subpack)
  2. Screw plunger on syringe
  3. Express air from syringe





Figure 4-7.- Prefilled syringe.


TUBEX INJECTOR TECHNIQUE

- |               |   |  |
|---------------|---|--|
| Med<br>Locker | 1. Unstow:<br>Tubex Injector<br><br>Injectable Medication | (Drug Subpack-Spine,<br>Drug Subpack-19,<br>Trauma Subpack-31)<br>(Drug Subpack,<br>Trauma Subpack,<br>EENT Subpack) |
|---------------|---|--|
2. Screw Tubex Injector plunger rod into plunger of medication until resistance is felt
  3. Place fingers on wings and with other hand loosen ribbed collar to OPEN position (labeled on collar)
  4. Slide ribbed collar + wings over medication cartridge
  5. Holding wings, tighten ribbed collar onto medication cartridge
  6. To inject medication, see appropriate drug administration technique (INTRAVENOUS (I.V.), 4-7, INTRAMUSCULAR, 4-24, or SUBCUTANEOUS, 4-26)
  7. Once used, hold wings with one hand and unscrew ribbed collar with other hand. Slide wings + ribbed collar back toward plunger. Unscrew entire Tubex Injector from medication cartridge
  8. Dispose of medication cartridge w/needle in Sharps container (Med Locker)
  9. Restow Tubex Injector in original location

Tubex Injector  
(Drug Subpack-Spine,  
Drug subpack-19,  
Trauma Subpack-31)



**Tubex Injector (Stowage Configuration)**



**Tubex Injector (Operational Configuration)**

Figure 4-8.- Tubex injector.

## LACERATIONS

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

### MAJOR LACERATIONS

1. Don non-sterile Gloves (IV Admin-12, Airway Subpack-8, EENT Subpack-7)
- Med Locker 2. Control bleeding by applying pressure over wound w/Gauze Pads (Trauma Subpack-14, IV Admin-14, Airway Subpack-11, EENT Subpack-2)
3. Unstow:
  - ^Povidone-Iodine Swabs (Trauma Subpack-7, IV Admin-19, Airway Subpack-10)
  - Saline, 100 ml (EENT Subpack-1)
  - Gloves, Sterile (Trauma Subpack-24)
  - Drape, Sterile (Trauma Subpack-24)
  - Tubex Injector (Trauma Subpack-31, Drug Subpack-19, Drug Subpack-Spine)
  - Surgical Instrument Assembly (Trauma Subpack-33)
  - Needle, 18ga (Trauma Subpack-29, EENT Subpack-18)
  - Syringe (10 cc) (Airway Subpack-15, EENT Subpack-20)
  - Alcohol Pads (Trauma Subpack-6, IV Admin-18, Airway Subpack-10, Drug Subpack-16, Drug Subpack-Spine)
  - Suture (Surgeon will advise) (Trauma Subpack-17,18)
  - \*Xylocaine, 2% (Trauma Subpack-30)
  - Gauze Pads (Trauma Subpack-14, IV Admin-14, Airway Subpack-11, EENT Subpack-2)
  - Kling (Trauma Subpack-12)
  - Tegaderm transparent dressing (Trauma Subpack-4)
  - ^Bactroban Ointment (Trauma Subpack-19)

4. Contact Surgeon and photodocument the wound.  
Continue to photodocument the wound every 24 hrs
5. Gently cleanse wound edge w/ Povidone-Iodine Swab
6. Insert Xylocaine, 2% (see WARNING, 6-14) into Tubex Injector (see INJECTIONS, TUBEX INJECTOR TECHNIQUE, 4-28)
7. Insert needle into wound edge
8. For blood return

If blood return:

9. Withdraw needle
10. Expel blood onto Gauze Pad
11. Repeat steps 7, 8

If no blood return:

12. Inject enough Xylocaine to raise small bump under skin. Advance needle parallel to wound edge
13. Repeat process to anesthetize wound edge
14. Repeat process on opposite side

Wound cleansing/preparation; clean all wounds before repairing:

15. Twist and lock 18ga Needle onto 10 cc Syringe
16. Clean rubber port on Saline bag w/Alcohol Pad
17. Remove cap and insert needle into rubber port on Saline bag; fill syringe w/saline
18. Carefully recap needle and remove from syringe
19. Flush laceration and adjacent skin w/saline; use Gauze Pads to assist cleaning as needed
20. Repeat until wound is thoroughly flushed (10 times is recommended)
21. Clean w/ Povidone-Iodine Swabs. Clean skin several inches beyond wound edge
22. Irrigate wound w/remaining saline. Blot dry w/sterile Gauze Pads

Cont next page

23. Don sterile Gloves w/o touching outer Glove surface
24. Obtain sterile drape and remove protective covering over adhesive. Place center hole in drape over wound; tape drape edges to patient
25. Inspect wound closely for foreign bodies. Remove any foreign bodies found and irrigate again
26. Consult Surgeon for appropriate suture technique (sutures, tissue adhesive, staples)

#### Laceration Closure Using Sutures

- Med Locker
1. Unstow suture material (see list, 4-30)
  2. Have assistant secure Surgical Instrument Assembly on lockers w/Velcro
  3. Set needle in needle holder. Place small point blunt forceps in nondominant hand; gently pick up skin edge; place first suture at midpoint of laceration, 2-3 mm from wound edge
  4. Reset Needle w/needle holders. Pick up opposite skin edge w/Forceps, repeat process from "inside out" (i.e., from wound edge to skin). (Needle should enter, exit skin at points equally distant from both wound edges)
  5. Pull suture together until skin edges are approximated (do not pull tight); secure suture w/instrument tie. Trim sutures, leaving 1 cm for suture removal
  6. Place additional sutures at midpoint of each wound section; sutures should be spaced 4-5 mm apart (fig 4-9)




Figure 4-9.- Suture placement.



Wound dressing:

7. Clean suture line gently w/Saline solution on Gauze Pads; apply ^Bactroban Ointment
8. Photodocument the laceration every 24 hrs
9. Apply Tegaderm transparent dressing to wound; cover w/Gauze Pads; secure w/Tape or Kling roll bandage
10. Reapply ointment once per day for 4-5 days; keep wound clean, dry, covered
11. Consult Surgeon for suture removal schedule (see Suture and Staple Removal, 4-35)

Laceration Closure Using Skin Stapler

- Med Locker
1. Don sterile Gloves (Trauma Subpack-24) without touching outer Glove surface
  2. Remove Skin Stapler from packaging (Trauma Subpack-21) and handle as sterile item
  3. Approximate and evert skin surfaces w/fingers or Forceps (Trauma Subpack-25)
  4. Position Skin Stapler on midpoint of wound
  5. Squeeze Skin Stapler actuating lever while maintaining pressure on wound




Figure 4-10.- Laceration closure using skin stapler.

6. Place additional staples at midpoint of each wound section; should be 1 cm apart when complete
7. If staple placed abnormally across wound, complete laceration repair if possible, then refer to Suture and Staple Removal, 4-35
8. When repair complete, restow Skin Stapler in pack
9. Consult Surgeon on staple removal schedule (see Suture and Staple Removal, 4-35)

## Laceration Closure Using Tissue Adhesive

Use for small wounds of face, trunk, and proximal extremities

Do not use on the following:

- Wounds crossing a joint
- Dirty/contaminated wounds
- Wounds w/jagged edges
- Wounds that require deep sutures
- Non-superficial wounds on hands/feet

- Med Locker 1. Unstow: ^Dermabond (Trauma Subpack-27) |

<p><u>WARNING</u> Wear Goggles (CCK) and Gloves (CCK) while applying ^Dermabond</p>
---

2. Dry wound edges and skin
3. Approximate wound edges with gloved fingers or Forceps (Trauma Subpack-25), maintaining proper eversion of skin
4. Remove applicator. Apply pressure at applicator midpoint, crushing internal glass ampule
5. Gently squeeze applicator to express the liquid

### NOTE

^Dermabond must be used immediately before polymerization |

6. Use gentle brushing strokes to apply a thin film of liquid to the wound edge

### NOTE

Avoid seepage into wound as it may delay healing

7. Adhesive should extend at least 1/2 cm on each side of apposed wound edges
8. Build up 3-4 layers of adhesive

NOTE

Wound can open if it gets wet

9. Maintain approximation of the wound edges until adhesive sets and forms a flexible film (should occur 1 min after last application)

NOTE

DO NOT apply ointment or medication on top of adhesive.

Avoid getting adhesive wet

10. Apply a dry adhesive using Tegaderm (Trauma Subpack-4) or Bandaid (Trauma Subpack-11) over wound, but NOT Steri-Strips
11. Adhesive will slough off in 5-10 days

Suture and Staple Removal

NOTE

Remove suture on face in 4 days. Remove sutures from other parts of the body in 5-7 days. Sutures across a joint or sutures in the scalp may be left in place for 8-10 days

Suture Removal:

- |                                |   |
|--------------------------------|---|
| Med<br>Locker                  | 1. Unstow:  |
|                                | Surgical Instrument Assembly<br>(Trauma Subpack-33)   |
|                                | Ziplock Bag<br>(IV Admin-1,<br>Airway Subpack-4, CCK) |
| Forceps<br>(Trauma Subpack-25) |   |
2. Carefully remove wound dressing

WARNING

DO NOT cut both sides of suture; a small portion of suture will be left under skin

3. Using Forceps, gently lift suture knot away from skin and cut one side of suture

4. Slowly pull to remove entire suture, discard in Ziplock Bag
5. Continue until all sutures are removed
6. Contact Surgeon w/description of laceration following suture removal and for advice on wound dressing

Staple Removal:

Med  
Locker

1. Unstow:  
Skin Staple Remover (Trauma Subpack-21)  
Ziplock Bag (IV Admin-1,  
Airway Subpack-4, CCK)
2. Remove Skin Staple Remover from packaging
3. Carefully remove dressing from stapled wound
4. Insert teeth of Skin Staple Remover between staple and skin
5. Squeeze levers of Skin Staple Remover together (see fig 4-11)




Figure 4-11.- Skin staple remover.

6. Place removed staple in Ziplock Bag
7. Continue until all staples removed

NOTE

If Skin Staple Remover required to remove abnormally placed staple during repair, wipe thoroughly w/Alcohol Pad (IV Admin-18, Airway Subpack-10, Drug Subpack-16, Drug Subpack-Spine, Trauma Subpack-6)

- Med Locker 8. Place new dressing using ^Bactroban Ointment (Trauma Subpack-19), Gauze Pads (IV Admin-14, Airway Subpack-11, Trauma Subpack-14, EENT Subpack-2) and Kling (Trauma Subpack-12)

MINOR LACERATIONS

- Med Locker 1. Stop bleeding by applying pressure over wound w/Gauze Pads (IV Admin-14, Airway Subpack-11, Trauma Subpack-14, EENT Subpack-2)

If bleeding uncontrolled:

2. Apply pressure bandage over cut
3. Go to MAJOR LACERATIONS, 4-30
  
4. After bleeding controlled, cleanse skin adjacent to wound w/^Povidone-Iodine Swabs (IV Admin-19, Airway Subpack-10, Trauma Subpack-7) or Alcohol Pads (IV Admin-18, Airway Subpack-10, Drug Subpack-16, Drug Subpack-Spine, Trauma Subpack-6)
5. Photodocument the laceration now and every 24 hrs
6. Use Steri-Strips (Trauma Subpack-16), ^Benzoin swabs (Trauma Subpack-8) to close cut
  
7. Contact Surgeon for healing timeline

8. Dress wound with:
- |                     |                       |
|---------------------|-----------------------|
| ^Bactroban Ointment | (Trauma Subpack-19)   |
| Adaptic Bandages    | (Trauma Subpack-5)    |
| or Gauze Pads       | (Trauma Subpack-14,   |
|                     | IV Admin-14,          |
|                     | Airway Subpack-11,    |
|                     | EENT Subpack-2)       |
| or Band-aids        | (Trauma Subpack-11,   |
|                     | IV Admin-20,          |
|                     | Drug Subpack-8,       |
|                     | Drug Subpack-12,      |
|                     | Drug Subpack-Spine)   |
| Kling               | (Trauma Subpack-12)   |
| Kerlix              | (Trauma Subpack-2)    |
| Ace Bandage         | (Trauma Subpack-3,13) |
| Telfa Pads          | (EENT Subpack-22)     |

9. Contact Surgeon

**MOTION SICKNESS (also see VOMITING, 5-20)**

Symptoms: Headache, sleepiness, lethargy, stomach awareness, decreased appetite, flushed feeling, "tumbling gyros" w/head movements, excess salivation, nausea, vomiting

Countermeasures:

1. Rest
2. Extra fluids, bland diet as able
3. Move slowly; avoid head movements

MILD MOTION SICKNESS

No vomiting, mild nausea; give antinausea, antiemetic drugs:

- |        |   |
|--------|---|
| Med    | <u>*Phenergan (suppository) (Drug Subpack-10)</u> |
| Locker | (see WARNING, 6-2, 6-10)                          |
|        | Dose: 1 every 6 hr as needed                      |
|        | -OR-  |
|        | <u>*Phenergan (oral) (Drug Subpack-15)</u>        |
|        | (see WARNING, 6-2, 6-10)                          |
|        | Dose: 1-2 tabs every 4-6 hr as needed             |

## MODERATE-TO-SEVERE MOTION SICKNESS

Nausea, vomiting, fatigue, inability to eat; give  
antinausea, antiemetic drugs:

- Med Locker      \*Phenergan (Injectable) (Drug Subpack-Spine)  
(see WARNING, 6-2, 6-10)  
Dose: Inject 1/2 to 1 cc intramuscularly every 6  
hr (see INJECTIONS, INTRAMUSCULAR,  
4-24)  
-OR-  
\*Phenergan (suppository) (Drug Subpack-10)  
(see WARNING, 6-2, 6-10)  
Dose: 1 every 6 hr as needed

## **MUSCLE SPASM (DRUG SIDE EFFECT)**

Signs: Muscles tight, contracted; eyes may be “rolled up” into  
head (rare reaction to medication such as Phenergan)

1. Contact Surgeon

- Med Locker      2. \*Benadryl (Injectable) (Drug Subpack-13):  
(see general WARNING, 6-2)  
Dose: 1 ml IM (see INJECTIONS,  
INTRAMUSCULAR, 4-24, and TUBEX  
INJECTOR TECHNIQUE, 4-28)

## **NAUSEA/VOMITING**

- Med Locker      \*Phenergan (oral) (Drug Subpack-15)  
(see WARNING, 6-2, 6-10)  
Dose: 1-2 tabs every 4-6 hr as needed  
-OR-  
\*Phenergan (suppository) (Drug Subpack-10)  
(see WARNING, 6-2, 6-10)  
Dose: 1 every 6 hr as needed  
-OR-  
\*Phenergan (Injectable) (SMS Kit, Drug Subpack-  
Spine)  
(see WARNING, 6-2, 6-10)  
Dose: Inject 1/2 ml intramuscularly every 6 hr  
as needed (see INJECTIONS,  
INTRAMUSCULAR, 4-24)

## NOSEBLEED

### WARNING

Do not administer Ascriptin or Motrin. Use Tylenol for pain. If both nasal cavities are packed, patient may require supplemental oxygen

1. Pinch nose just below nasal bone applying direct, uninterrupted pressure for 10 min

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

If bleeding continues:

Med  
Locker

2. Unstow:

Cotton Balls	(EENT Subpack-22)
Nasal speculum	(EENT Subpack-26)
Forceps, Blunt	(Trauma Subpack-25)
Afrin	(EENT Subpack-19)
Silver Nitrate Stick	(EENT Subpack-34)
^Bactroban Ointment	(Trauma Subpack-19)

3. Spray Afrin on Cotton Balls. Using Forceps, pack nose, apply pressure for 10-15 min
4. Remove Cotton Balls; check nose for bleeding site w/nasal speculum (fig 4-12)




Figure 4-12.- Nasal speculum.



If bleeding site located:

5. Touch area 3-4 times w/Silver Nitrate stick
6. Apply ^Bactroban Ointment to affected area using clean cotton ball

If bleeding site not located or bleeding continues:

7. Contact Surgeon
8. Unstow:

Med Locker	Saline, 100 ml Needle, 18ga  Syringe (10 cc)  Tape  ^Neosporin Plus Nasal Packing	(EENT Subpack-1) (EENT Subpack-18, Trauma Subpack-29) (EENT Subpack-20, Airway Subpack-15, Trauma Subpack-22) (EENT Subpack-9, IV Admin-13, Airway Subpack-9, Trauma Subpack-34) (EENT Subpack-25)   (EENT Subpack-21)
---------------	---	---

**CAUTION**  
Do not insert cotton balls beyond point of  
visibility, where they can be easily removed

9. To anesthetize nasal cavity, apply small amount of ^Neosporin Plus to cotton balls (do not exceed 5-g ointment). Using forceps, pack nose
10. Apply pressure for 5 min; remove cotton balls
11. Grasp drawstring end of packing; quickly insert along floor of nasal cavity until drawstring reaches anterior nasal opening (see fig 4-13)

If packing not fully expanded 30 sec after placement:

12. Place 18ga Needle on end of Syringe
13. Insert into port on bag of Saline
14. Fill Syringe w/Saline
15. Remove Needle from Syringe
16. Irrigate Nasal Packing w/Saline

17. Tape drawstring over end of nose, trim excess
18. Nasal Packing may be left in place up to 48 hr

Med 19. \*Augmentin (Drug Subpack-37)  
 Locker Dose: 1 tab 3X/day

20. Contact Surgeon before removing packing




Figure 4-13.- Nasal packing.

If bleeding continues:

21. Contact Surgeon
22. Unstow:

Med Nasal Catheter (EENT Subpack-21)  
 Locker

Test Nasal Catheter prior to insertion:

23. Inject 10 cc saline into valve on Nasal Catheter
24. Inspect balloon for integrity
25. Withdraw saline from catheter w/syringe
26. Apply ^Neosporin Plus to Nasal Catheter device to lubricate for insertion
27. Remove nasal packing from nose and replace w/Nasal Catheter; quickly insert along floor of nasal cavity until sponge fully inserted (fig 4-14)
28. Gently inject 10 cc saline into valve on Nasal Catheter





Figure 4-14.- Nasal catheter insertion.



If sponge on Nasal Catheter device not fully expanded  
30 sec after placement:

29. Irrigate sponge w/saline using syringe
30. Adjust retaining ring on Nasal Catheter until it gently rests against the nose
31. Nasal Catheter may be left in place up to 48 hrs
32. \*Augmentin (Drug Subpack-37)  
Dose: 1 tab 3X/day
33. Contact Surgeon before removing catheter

Med  
Locker



This Page Intentionally Blank



SECTION 5

O THRU Z

O THRU Z

OBSTRUCTED AIRWAY – CONSCIOUS ADULT (see CHOKING, 1-4)

OBSTRUCTED AIRWAY – UNCONSCIOUS ADULT (see CHOKING, 1-5)

ORAL AIRWAY (see CPR (CARDIOPULMONARY RESUSCITATION), 1-11)

OXYGEN – SUPPLEMENTAL (see CPR (CARDIOPULMONARY RESUSCITATION), 1-12)

PAIN RELIEF (IN INCREASING ORDER OF STRENGTH).. 5-3

RADIATION..... 5-4

SEIZURES (see EMERGENCY, 1-42)

SHOCK (see EMERGENCY, 1-44)

SINUS PROBLEMS..... 5-5

SKIN INFECTION ..... 5-6

SKIN – MINOR ABRASIONS ..... 5-7

SKIN RASH..... 5-7

    ALLERGIC RASH ..... 5-7

    FUNGAL RASH ..... 5-8

SLEEPING PROBLEMS (INSOMNIA)..... 5-8

SORE THROAT..... 5-9

SPACE MOTION SICKNESS (see MOTION SICKNESS, 4-38)

STRAINS/SPRAINS ..... 5-9

TOOTHACHE ..... 5-10

    DENTAL INJECTION TECHNIQUE ..... 5-11

    TEMPORARY FILLING ..... 5-15

    EXPOSED PULP ..... 5-15

    REPLACING CROWN w/TEMPORARY CEMENT..... 5-16

UPSET STOMACH (see INDIGESTION, 4-6)

URINARY RETENTION..... 5-18

URINARY TRACT INFECTION (DYSURIA) (see BLADDER INFECTION, 2-5)

VOMITING (also see MOTION SICKNESS – MODERATE-TO-SEVERE, 4-39)..... 5-20

YEAST INFECTION..... 5-20

FIGURES

**O THRU Z**

5-1	Dosimeter.....	5-4
5-2	Upper molar anesthesia.....	5-13
5-3a	Lower (mandibular) molar anesthesia #1.....	5-14
5-3b	Lower (mandibular) molar anesthesia #2.....	5-14
5-4	Catheter insertion.....	5-19

**PAIN RELIEF (IN INCREASING ORDER OF STRENGTH)**

Med  
Locker

Ascriptin (Drug Subpack-Spine)

(see WARNING, 6-3)

Dose: 1-2 tabs every 4-6 hr as needed

-OR-

Tylenol (Drug Subpack-Spine)

Dose: 1-2 tabs every 4-6 hr as needed

-OR-

Motrin (Drug Subpack-Spine)

(see WARNING, 6-9)

Dose: 1 tab every 4-6 hr as needed w/food and  
drink

-OR-

\*Vicodin (Drug Subpack-21)

(see WARNING, 6-2, 6-13)

Dose: 1-2 tabs every 4-6 hr

-OR-

NOTE

For Demerol and Morphine Sulfate, see  
INJECTIONS, INTRAMUSCULAR (4-24)  
and TUBEX INJECTOR TECHNIQUE (4-28)

\*Demerol (Drug Subpack-25)

(see WARNING, 6-2, 6-5)

Dose: Inject 1 cc intramuscularly every 3-4 hr as  
needed

-OR-

\*Morphine Sulfate (Drug Subpack-23,24)

(see general WARNING, 6-2)

Dose: Inject 1/2 cc intramuscularly every 3-4 hr  
as needed

-OR-

NOTE

Handle Duragesic Patch using non-sterile Gloves (IV Admin-12, Airway Subpack-8, EENT Subpack-7, CCK). Pain relief w/patch does not occur rapidly

WARNING

If contact w/gel is made by other crew-members, wash contaminated area w/generous amounts of water

\*Duragesic Patch (Drug Subpack-8)

(see general WARNING, 6-2, 6-6)

Dose: 1 patch. Apply to shoulder, upper arm, abdomen, flank, chest, or back. Remove after 72 hr

**RADIATION**

Med  
Locker




Figure 5-1.- Dosimeter.



### RADIATION INSTRUMENTATION LOG

Pouch #	Serial #	MET/Rdng	MET/Rdng	MET/Rdng	MET/Rdng	Initials

### SINUS PROBLEMS

Symptoms: Facial pain, ear pain, tooth pain, congestion

Med Locker 1. Unstow: Thermometer (EENT Subpack-3)

Temperature: \_\_\_\_\_

Decongestants:

2. Sudafed (Drug Subpack-Spine)  
 (see WARNING, 6-12)  
 Dose: 2 tabs every 6 hr as needed

-OR-

Guaifenes PSE (Drug Subpack-44)  
 Dose: 1 tab every 12 hr as needed

-OR-

Afrin (EENT Subpack-19)  
 Dose: 1-2 sprays every 12 hr as needed

If fever:

Antibiotics:

3. \*Augmentin (Drug Subpack-37)  
 Dose: 1 cap 2X/day for 10-14 days  
 -OR-

\*Bactrim DS (Drug Subpack-38)

(see WARNING, 6-3)

Dose: 1 tab 2X/day for 10-14 days

-OR-

\*Duricef (Drug Subpack-40)

(see WARNING, 6-6)

Dose: 1 cap every 12 hr for 10-14 days

Pain medication:

4. Ascriptin (Drug Subpack-Spine)

Dose: 1-2 tabs every 4-6 hr as needed

-OR-

Tylenol (Drug Subpack-Spine)

Dose: 1-2 tabs every 4-6 hr as needed

5. Contact Surgeon

## SKIN INFECTION

Signs: Skin is red, warm, tender; pus may be seen coming out of localized point

1. Contact Surgeon. Place Ruler (Trauma Subpack-26) in proximity of infection and photodocument the infection every 24 hrs

Antibiotics:

Med  
Locker

2. \*Augmentin (Drug Subpack-37)

Dose: 1 tab 2X/day for 5 days

-OR-

\*Cipro (Drug Subpack-40)

Dose: 1 tab 2X/day for 10 days

-OR-

\*Duricef (Drug Subpack-40)

(see WARNING, 6-6)

Dose: 1 cap every 12 hr for 10 days

-OR-

\*Zithromax (Drug Subpack-36)

Dose: 2 tabs initial dose, then 1 tab/day for 5 days

## SKIN – MINOR ABRASIONS

1. Clean affected area thoroughly w/Saline (IV Admin-1, EENT Subpack-1, Saline Supply Bag, Middeck Locker); blot dry gently w/Gauze Pads (IV Admin-14, Airway Subpack-11, Trauma Subpack-14, EENT Subpack-2)
2. Place Ruler (Trauma Subpack-26) in proximity of the wound and photodocument the wound every 24 hrs

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

3. Dress wound w/^Bactroban Ointment (Trauma Subpack-19). Cover w/Bandaid (IV Admin-20, Drug Subpack-12, Drug Subpack-Spine, Trauma Subpack-11) or w/Gauze Pads for large abrasions. Wrap Gauze Pads w/Kerlix, Kling (Trauma Subpack-2,12) to hold in place
4. Repeat above procedure and inspect abrasion once daily or at any time wound becomes wet or contaminated
5. Notify Surgeon at next routine Private Medical Conference (PMC)

## SKIN RASH

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

## ALLERGIC RASH

Signs: Skin is red, warm; may have hives or blisters

Med Locker 1. ^Topicort (EENT Subpack-28)  
Dose: Apply thin layer 2X/day until rash resolved (rub in well, avoid contact w/eyes)

Antihistamines for relief of itching:

2. Benadryl (oral) (Drug Subpack-42)  
(see general WARNING, 6-2)  
Dose: 1 cap every 6 hr as needed

-OR-

\*Claritin (Drug Subpack-43)  
(see general WARNING, 6-2)  
Dose: 10 mg once daily

3. Contact Surgeon for further dosing. Place Ruler (Trauma Subpack-26) in proximity of rash and photodocument the rash every 24 hrs

### FUNGAL RASH

Signs: Skin is red, scaly; oval/round patches w/central scales may be present

Med Locker 1. Lotrimin AF Cream (Trauma Subpack-26)  
Dose: Apply thin layer 2X/day for duration of flight (rub in well; avoid contact w/eyes)

2. Contact Surgeon for further dosing. Place Ruler (Trauma Subpack-26) in proximity of rash and photodocument the rash every 24 hrs

### **SLEEPING PROBLEMS (INSOMNIA)**

<u>WARNING</u> Both PLT and CDR -OR- Both CMOs <u>may not</u> use sedatives/hypnotics simo
--

#### NOTE

Prepare for bed before taking Restoril or Ambien. DO NOT continue work after taking medication

Med Locker \*Ambien (Drug Subpack-Spine)  
(see general WARNING, 6-2)  
Dose: 1 tab 30 min prior to sleep period  
-OR-  
\*Restoril (Drug Subpack-Spine)  
(see general WARNING, 6-2)  
Dose: 1 or 2 caps 30 min prior to sleep period



## TOOTHACHE

Infection, signs/symptoms: Swelling, fever, tender/raised lymph nodes under lower jaw, redness around affected tooth, dull ache when biting, usually no pain w/hot or cold, but may be present w/multi-rooted teeth

Treatment: Antibiotics, avoidance of tooth, and pain medication

Cracked Tooth, signs/symptoms: Sharp stabbing pain when biting down or upon release, unusual cold sensitivity. May or may not be visible signs of a crack

Treatment: Avoidance of tooth, and pain medication

Pulpitis, signs/symptoms: Painful stimulus to either heat and/or cold, may have pain to bite pressure, spontaneous pain w/o provocation

Treatment: Avoidance of tooth, and pain medication

1. Contact Surgeon

Antibiotics:

Med 2. \*Augmentin (Drug Subpack-37)  
Locker Dose: 1 tab 2X/day for 10 days w/food and drink

Pain relief, mild to moderate:

Med 3. Eugenol (contained in 3 needleless Tubex  
Locker syringes - 1 ml) (Toothache Kit, EENT  
Subpack-35)  
Unstow Tubex Injector (Trauma Subpack-31,  
Drug Subpack-Spine,  
Drug Subpack-19)

(see Tubex Injector Technique, 4-28)

Apply drop of Eugenol directly on affected tooth or  
apply using cotton pellet and tweezers

4. Ascriptin (Drug Subpack-Spine)  
(see WARNING, 6-3)  
Dose: 2 tabs every 4-6 hr as needed

-OR-

Tylenol (Drug Subpack-Spine)  
Dose: 2 tabs every 4-6 hr as needed

-OR-

Motrin (Drug Subpack-Spine)  
(see WARNING, 6-9)  
Dose: 1 tab every 4-6 hr as needed w/food and  
drink

Pain relief, severe:

5. \*Vicodin (Drug Subpack-21)

(see WARNING, 6-2, 6-13)

Dose: 1-2 tabs every 4-6 hr as needed

-OR-

\*Xylocaine w/Epinephrine (Injectable)

(EENT Subpack-30) (see WARNING, 6-14)

(see DENTAL INJECTION TECHNIQUE, step 6)

DENTAL INJECTION TECHNIQUE

1. Unstow:

Dental Syringe (EENT Subpack-33)


\*Xylocaine w/Epinephrine  
(anesthetic) (EENT Subpack-30)

Needle, Short (upper injection)  
(EENT Subpack-36)


Needle, Long (lower injection)  
(EENT Subpack-29)

2. Load dental syringe:

Withdraw plunger; insert metal end of dental  
carpule into syringe body by sliding toward  
threaded end; snap into place




3. Push plunger arrow into dental carpule; ensure head  
of arrow firmly seated



4. Open needle by twisting off clear cap

**NOTE**

For upper injections use short needle;  
For lower injections use long needle



5. Screw needle cartridge into syringe by turning cw until firmly seated
6. To inject, remove yellow needle cap. For upper injections, see figure 5-2. For lower injections, see figures 5-3a and 5-3b

**WARNING**

Draw back gently on syringe before injection. Reposition needle if blood is seen

7. Upper Dental Anesthesia:  
Place needle at height of moveable membrane (mucobuccal fold) above fixed gum tissue. Insert needle; direct needle to root apex of tooth to be anesthetized. Draw back gently on syringe to ensure no blood
8. Inject 1/2 carpule of dental anesthetic directly over root apex. Wait 5 min; pain should subside
9. Lower (mandibular) Anesthesia:  
Place thumb in deepest portion of coronoid notch of mandibular ramus and use center of thumb as guide for needle height (see figs 5-3a and 5-3b)






Figure 5-2.- Upper molar anesthesia.

10. Insert needle at point where seam between the cheek and throat turns upward (fig 5-3a)
11. Barrel of syringe is kept over the third and fourth teeth on opposite side of the injection (fig 5-3b). Insert needle until bone is contacted and only 5 to 10 mm of needle remains exposed
  - If bone not contacted:
    12. Withdraw needle 5-10 mm; reinsert needle w/syringe barrel over teeth 4 and 5
  - If bone contacted and more than 5-10 mm of needle remains exposed:
    13. Withdraw needle 5-10 mm; reinsert needle w/syringe barrel over teeth 2 and 3. Draw back gently on syringe to ensure no blood
14. Inject slowly, over 1 min, the entire contents of syringe; remove needle
15. Wait 5 min; pain should subside
16. Contact Surgeon and re-inject if necessary




Figure 5-3a.- Lower (mandibular) molar anesthesia #1.




Figure 5-3b.- Lower (mandibular) molar anesthesia #2.

## TEMPORARY FILLING

### WARNING

Use only if missing portion of filling or tooth does not include an exposed pulp. If dental pulp exposed, see EXPOSED PULP

1. Unstow:

Med	Gauze Pads	(EENT Subpack-2)
Locker	Temporary filling	(EENT Subpack-24)
	Dental carver	(EENT Subpack-26)
2. Squeeze filling material onto finger
3. Place in tooth, wipe off excess w/Gauze Pad (EENT Subpack-2)
4. Bite and rub teeth together, then open mouth and adjust filling slightly out of bite using Gauze Pad (EENT Subpack-2) and/or dental carver (EENT Subpack-26)
5. Allow 5 min for initial set; be careful with filling for at least 24 hr

## EXPOSED PULP

### NOTE

Don non-sterile Gloves (CCK) and Goggles (CCK) to apply Eugenol. Consider use of Dental Injection if painful prior to exposure of pulp

1. Unstow:

Med	Tubex Injector	(Trauma Subpack-31, Drug Subpack-Spine, Drug Subpack-19)
Locker	Toothache Kit (tweezers, cotton pellets, Tubex Syringes)	(EENT Subpack-35)
	Gauze Pads	(EENT Subpack-2)
	Dental Explorer Probe	(EENT Subpack-26)
2. Assemble Tubex Injector (see Tubex Injector Technique, 4-28)

3. Remove 1 cotton pellet from box with supplied tweezers (EENT Subpack-35), resizing cotton pellet as necessary to fit opening of pulp
4. Remove red cap from Tubex Injector and apply enough Eugenol to cotton pellet to lightly moisten it. Remove excess Eugenol by squeezing out excess w/Gauze Pad (EENT Subpack-2)
5. Place lightly moistened cotton pellet into exposed pulp area of tooth, using tweezers and Dental Explorer Probe (EENT Subpack-26) to push pellet into the 'hole'
6. Keep pellet in place for remainder of mission. Check daily and replace if covered w/food debris. Replace w/new pellet and Eugenol if persistent pain

**WARNING**  
Do not place filling in tooth w/  
exposed pulp

REPLACING CROWN w/TEMPORARY CEMENT

NOTE

If no pain, especially when eating or drinking, stow crown in secure location; can safely be replaced upon return to Earth

1. Unstow:
 

Cotton Swabs	(EENT Subpack-34)
Dental carver	(EENT Subpack-26)
Dycal Base	(EENT Subpack-1)
Dycal Catalyst	(EENT Subpack-1)
Dental Explorer Probe	(EENT Subpack-26)
Dental floss	(Personal Hygiene Kits)
Tongue depressor	(EENT Subpack-34)
Gauze Pads (4)	(EENT Subpack-2)
2. Remove residual cement from crown and tooth, utilizing dental carver (EENT Subpack-26)
3. Carefully check fit of crown by replacing on tooth and biting down. Remove and dry crown. Dry and isolate tooth as well as possible w/rolled Gauze Pad (EENT Subpack-2)

4. Place a 1/2-inch line of both Dycal Base and Dycal Catalyst (EENT Subpack-1) on one end of Tongue Depressor (EENT Subpack-34) and mix well using stick end of cotton swab (EENT Subpack-34)
5. Place small portion of Dycal mixture around inside walls of crown; seat crown on tooth using a positive rocking force. Place cotton tip end of swab over crown and have patient bite down for 3 sec. Remove cotton swab and have patient bite down to determine if crown is fully seated
  - If crown not fully seated:
    6. Carefully remove crown by prying up at different locations on crown margin w/dental carver (EENT Subpack-26) until crown is loose. Remove crown; return to step 2
  - If crown fully seated:
    7. Replace cotton swab over crown and have patient continue biting on swab w/moderate pressure for 5 min
    8. Gently clean remaining cement from around gum w/Dental Explorer Probe (EENT Subpack-26) and dental floss (Personal Hygiene Kits). (Place a knot in center of 18-inch piece of dental floss; glide it back and forth gently between the crown and the adjacent teeth to clean cement from between the teeth)

## URINARY RETENTION

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

1. Unstow:

Med Locker	Gloves, sterile	(Trauma Subpack-24)
	Urine Chemstrips 10 SG	(Trauma Subpack-15)
	^Povidone-Iodine Swabs	(Trauma Subpack-7, IV Admin-19, Airway Subpack-10)
	Cotton Swabs	(EENT Subpack-34)
	Catheter, Foley OR Catheter, Straight	(Trauma Subpack-10)
	Lubricant (water soluble)	(Trauma Subpack-9, Airway Subpack-13)
	Ambulatory Leg-Bag	(Trauma Subpack-1)
	Syringes (10 cc)	(Airway Subpack-15, Trauma Subpack-22, EENT Subpack-20)
	Saline	(IV Admin-1, Saline Supply Bag, Middeck Locker, EENT Subpack-1)
	Needle, 22ga	(Trauma Subpack-22)
  2. Consult Surgeon for catheter type. Surgeon will provide use and resterilization techniques for Straight Catheter
  3. Expose patient
  4. Don sterile Gloves
  5. Clean catheter insertion site on patient w/^Povidone-Iodine Swabs
- For males:
6. Cleanse tip of penis w/^Povidone-Iodine Swab from urethra outward
- For females:
7. Cleanse area of urethra w/^Povidone-Iodine Swab, cleansing from urethra towards vagina only (front to back)

8. Connect catheter to Ambulatory Leg-Bag
9. Cover tip of catheter w/Water Soluble Lubricant
10. Insert catheter w/firm pressure until urine flows.  
Continue inserting to approx 2 cm depth (fig 5-4)




Figure 5-4.- Catheter insertion.

11. Maintain catheter in place until urine flow ceases

To determine if catheter should be removed:

12. Contact Surgeon

**WARNING**  
Do not pull on catheter while balloon is inflated

If catheter to remain inserted:

13. Inflate balloon w/syringe using 5-cc saline
14. Release small amount of urine from Leg Bag
15. Perform urine Chemstrip 10 SG testing
16. Go to BLADDER INFECTION, 2-5

## VOMITING (also see MOTION SICKNESS, 4-38)

1. Avoid solid foods, milk products for 24 hr
2. Maintain fluid intake

After vomiting:

3. Wait 20 min prior to drinking frequent small amounts of fluid

If abdominal pain present, or blood noted in vomitus:

4. Contact Surgeon, see ABDOMINAL PAIN, 2-2

Anti-Vomiting Medication:

- Med Locker
5. \*Phenergan (oral) (Drug Subpack-15)  
Dose: 1-2 tabs every 4-6 hr as needed  
-OR-  
\*Phenergan (suppository) (Drug Subpack-10)  
Dose: 1 every 6 hr as needed  
-OR-  
\*Phenergan (Injectable) (Drug Subpack-Spine)  
Dose: Inject 1/2 to 1 cc intramuscularly every 6 hr (see INJECTIONS, INTRAMUSCULAR, 4-24)

## YEAST INFECTION

Symptoms: Vaginal itching/burning  
White vaginal discharge

Signs: White vaginal discharge  
Vaginal redness

Treatment:

- Med Locker
1. \*Diflucan (Drug Subpack-18)  
Dose: 1 tab by mouth

Possible side effects: Headache, nausea, abdominal pain, diarrhea. Symptoms are usually transient

2. Contact Surgeon if not resolved in 2-3 days



SECTION 6

MEDICATION SIDE EFFECTS

MEDICATION SIDE EFFECTS..... 6-2

**SIDE EFFECTS**

## MEDICATION SIDE EFFECTS

\*Contact Surgeon before giving any medication marked w/asterisk. In emergency or when orbiter LOS, begin appropriate treatment; call Surgeon as soon as possible

### WARNING

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

### WARNING

Following drugs should not be used together as they may cause excessive drowsiness in combination: Ambien, Benadryl, Claritin, Demerol, Duragesic Patch (Fentanyl), Morphine Sulfate, Phenergan, Restoril, Valium, Vicodin

## SIDE EFFECTS

acetaminophen – See TYLENOL

acetaminophen w/codeine – See TYLENOL #3

\*acetazolamide – See DIAMOX

AFRIN – Long-acting topical nasal decongestant

Possible side effects: Burning, stinging, sneezing, or increased nasal discharge. Usage more frequently than directed can produce chronic nasal congestion

\*albuterol – See PROVENTIL INHALER

\*ALCAINE (proparacaine) – Anesthetic eye drops for eye exams

- \* Do NOT repeat Alcaine dose \*
- \* w/o contacting Surgeon \*

ALEVE (naproxen) – Pain reliever

Possible side effects: Abdominal pain, nausea, heartburn

\*AMBIEN (zolpidem) – Used to induce sleep

Possible side effects: Anxiety, fatigue, headache, irritability, nausea, vomiting, abdominal pain, tremors, double vision (see general WARNING, 6-2)

\*amikacin – See AMIKIN

\*AMIKIN (amikacin) – I.V. antibiotic

\*amoxicillin/clavulanate – See AUGMENTIN

ANUSOL HC – Suppository for relief of pain/itching from hemorrhoids or other rectal problems

Possible side effects: Discontinue if anal irritation develops/worsens

Artificial Tears – See REFRESH PLUS

**WARNING**

Do not use Ascriptin in patients w/asthma, history of aspirin allergy, or GI or other bleeding problems

ASCRIPTIN (aspirin) – Pain reliever, anti-inflammatory

Possible side effects: Upset stomach, ringing in ears

aspirin – See ASCRRIPTIN

\*ATROPINE – Given I.V. only

Use only in cardiopulmonary arrest. Contact Surgeon

\*AUGMENTIN (amoxicillin/clavulanate) – Antibiotic

Possible side effects: Nausea, diarrhea, rash

AYR SALINE MIST – Used to treat congestion and nasal dryness

Possible side effects: Nasal irritation

\*azithromycin - See ZITHROMAX

**WARNING**

Do not use Bactrim DS if allergy to sulfa or trimethoprim exists. Give w/caution to patients w/severe allergy or asthma

\*BACTRIM DS (trimethoprim/sulfamethoxazole) – Antibiotic

Possible side effects: Skin rash, headache, stomach upset

^BACTROBAN OINTMENT (mupirocin) – Topical antibiotic

beclamethazone – See BECONASE INHALER

BECONASE INHALER (beclamethazone) – Bronchodilator used for treatment of allergy and congestion

Possible side effects: Nasal irritation, nose bleed, headache

BENADRYL (diphenhydramine) – (oral and \*injectable) –  
Antihistamine; used for allergic reactions or for drug-induced  
muscle spasms

Possible side effects: Drowsiness, inability to concentrate,  
dry mouth, blurred vision, rash, sensitivity to light, headache,  
rapid heart rate, dizziness, fatigue (see general WARNING,  
6-2)

bisacodyl – See DULCOLAX

\*bupivacaine plain – See MARCAINE PLAIN

\*cefadroxil – See DURICEF

ceftriaxone – See ROCEPHIN

\*CILASTIN/PRIMAXIN (imipenem IV) – Antibiotic

Possible side effects: Nausea, diarrhea, vomiting, rash,  
possible pain at injection site

Note: Mix powder w/100 cc normal Saline and shake to  
obtain a clear solution. Administer intravenously. Give  
slowly over 20-30 min period

\*^CILOXAN OINTMENT (ciprofloxacin) – Antimicrobial eye  
ointment

Possible side effects: Local burning or discomfort; use may  
result in overgrowth of nonsusceptible organisms

\*CIPRO (ciprofloxacin) – Antibiotic

Possible side effects: Nausea, diarrhea, vomiting, abdominal  
pain/discomfort, headache, restlessness

\*^ciprofloxacin – See ^CILOXAN OINTMENT, CIPRO

\*CLARITIN (loratadine) – Antihistamine, anti-allergy

Possible side effects: Headache, sleepiness/drowsiness,  
fatigue, dry mouth (see general WARNING, 6-2)

^clotrimazole cream – See ^LOTRIMIN AF CREAM

COUGH LOZENGES – For cough suppression

\*CYCLOGYL (cyclopentolate) – Eye drops for pupil dilation  
Possible side effects: Transient burning, blurred vision due to pupil dilation, sensitivity to light

\*cyclopentolate – See CYCLOGYL

deltasone – See PREDNISONONE

**WARNING**

Sedation w/Demerol may be additive if given w/ other drugs that depress central nervous system (see general WARNING, 6-2); may cause low blood pressure, reduced breathing rate

\*DEMEROL (meperidine) – Narcotic pain reliever  
Possible side effects: Dizziness, lightheadedness, sedation, nausea, vomiting, sweating, dry mouth, blurred vision, constipation

^desoximetasone – See ^TOPICORT

\*dexamethasone – Injectable steroid for treatment of increased intracranial pressure, severe allergic reactions

**WARNING**

Do not use Dexedrine in patient w/high blood pressure; may impair ability to concentrate

\*DEXEDRINE (dextroamphetamine) – For motion sickness  
Possible side effects: Dizziness, rapid heart rate, restlessness, tremor, headache, loss of appetite

\*dextroamphetamine – See DEXEDRINE

\*DIAMOX (acetazolamide) – For altitude sickness  
Possible side effects: Minimal w/short-term therapy – tingling sensation in extremities, loss of appetite, increased urine output, occasional drowsiness, confusion

\*diazepam – See VALIUM

\*DIFLUCAN (fluconazole) – Oral antifungal medication for vaginal yeast infections

Possible side effects: Headache, nausea, abdominal pain, diarrhea. Symptoms are usually transient

\*DILANTIN (phenytoin) – Anticonvulsant

Possible side effects: Confusion, dizziness, decreased coordination

diphenhydramine – See BENADRYL

DULCOLAX (bisacodyl) – Tablets or suppositories; for constipation

Possible side effects: Abdominal cramps

\*DURAGESIC PATCH (fentanyl) – Use only for severe pain. (see general WARNING, 6-2)

Possible side effects: Respiratory depression, low blood pressure. Other crewmembers should avoid contact w/skin

GUAIFENEX PSE (guaifenesin/pseudoephedrine) – Decongestant

Possible side effects: Headache, nervousness, nausea, vomiting

**WARNING**

Do not use Duricef if there is history of Penicillin or Cephalosporin allergy

\*DURICEF (cefadroxil) – Antibiotic

Possible side effects: Diarrhea, nausea, vomiting, abdominal pain, rash

DYCAL (base and catalyst) – Used for crown replacement

Possible side effects: Skin and eye irritant

\*EPINEPHRINE (Injectable) – Use only in cardiopulmonary arrest or severe allergic reaction. Contact Surgeon

EUGENOL – Local antiseptic and analgesic

Possible side effects: Skin and eye irritant

\*fentanyl – See DURAGESIC PATCH

- \*FLAGYL (metronidazole) – Antibiotic  
Possible side effects: Nausea, headache, vomiting, diarrhea, loss of appetite, metallic taste, rash
- \*fluconazole – See DIFLUCAN
- flumazenil – See ROMAZICON
- \*furosemide – See LASIX
- \*GENOPTIC OPHTHALMIC OINTMENT (gentamicin) – Antibiotic eye ointment  
Possible side effects: Occasional burning or stinging of eye
- \*gentamicin – See GENOPTIC OPHTHALMIC OINTMENT
- \*HALDOL (haloperidol) – Injectable major tranquilizer. Use only after contact w/Surgeon
- \*haloperidol – See HALDOL
- \*hydrocodone/acetaminophen – See VICODIN
- ibuprofen – See MOTRIN

<p style="text-align: center;"><u>WARNING</u> Do not use Imipenem if there is history of Penicillin or Cephalosporin allergy</p>
--

- \*imipenem IV – See CILASTIN/PRIMAXIN
- IMODIUM (loperamide HCl) – Used to stop diarrhea  
Possible side effects: Abdominal discomfort, nausea, vomiting, constipation, drowsiness, dizziness, dry mouth
- \*INDERAL (propranolol hydrochloride) – Oral medication that decreases heart rate and blood pressure, decreases heart workload; used for high blood pressure, chest pain
- \*ISOPTIN (verapamil) – Given I.V. only; use only for very rapid heart rates that do not resolve w/other therapy. Contact Surgeon

KENALOG IN ORABASE (triamcinolone acetonide) – Topical steroid for treatment of mouth ulcers and fever blisters

ketorolac tromethamine – See TORADOL

LASIX (furosemide) – Potent intravenous diuretic used in severe edema, hypertension

Possible side effects: Orthostatic hypotension, dizziness, nausea, headache

LEVAQUIN (levofloxacin) – Antibiotic

Possible side effects: Nausea, headache, diarrhea

levofloxacin – See LEVAQUIN

\*lidocaine plain – See XYLOCAINE PLAIN

\*lidocaine w/epinephrine – See XYLOCAINE W/EPINEPHRINE

\*lidocaine/cardiac – See XYLOCAINE/CARDIAC

loperamide HCl – See IMODIUM

\*loratadine – See CLARITIN

^LOTRIMIN AF CREAM (clotrimazole cream) – Antifungal skin medication

magnesium hydroxide – See MILK OF MAGNESIA

WARNING

Low blood pressure, heart block, seizures may occur w/overdose of Marcaine Plain. Do not inject intravenously; give smallest amount possible subcutaneously

WARNING

Do not use Marcaine Plain for anesthesia on fingers, toes, earlobes, or nose where constriction of blood vessels may lead to tissue death

\*MARCINE PLAIN (bupivacaine plain) – Local anesthetic used prior to laceration repair



\*meperidine – See DEMEROL

\*metronidazole – See FLAGYL

MILK OF MAGNESIA (magnesium hydroxide) – Antacid used to treat constipation

Possible side effects: Diarrhea

\*MORPHINE SULFATE – Use only for severe pain. Contact Surgeon (see general WARNING, 6-2)

Possible side effects: Respiratory depression, low blood pressure

**WARNING**

Do not take Motrin if allergic to aspirin

MOTRIN (ibuprofen) – Intermediate-strength pain reliever, anti-inflammatory agent

Possible side effects: Upset stomach, diarrhea; if possible, take w/food and drink

^mupirocin – See ^BACTROBAN OINTMENT

MYLANTA – Antacid/antiflatulent

\*naloxone – See NARCAN

naproxen – See ALEVE

\*NARCAN (naloxone) – Intravenous narcotic antagonist. Use only under Surgeon's direction

^NEOSPORIN PLUS – Aids in prevention of infection and provides temporary pain relief in minor cuts, scrapes, burns

NITROGLYCERIN – patch and tablets – Used in treatment of angina

Possible side effects: Headache, postural hypotension, dizziness

\*NITROSTAT – nitroglycerin tabs – Used only for treatment of heart attacks, other cardiac emergencies. Contact Surgeon

\*norgestrel/ethinyl estradiol – See OVRAL-21

\*OVRAL-21 (norgestrel/ethinyl estradiol) – Female hormone mixture for controlling abnormal menstrual bleeding  
Possible side effects: Nausea, increased blood pressure, fluid retention

omeprazole – See PRILOSEC

PEPTO-BISMOL – Bismuth compound used to treat nausea, indigestion, diarrhea  
Possible side effects: May turn tongue, stool black

phenazopyridine – See PYRIDIUM

**WARNING**

Phenergan may have an additive sedative effect w/other central nervous system depressants (see general WARNING, 6-2). Spasms of head, neck muscles may occur w/intramuscular injection. Do not exceed 100 mg in a 24-hr period

\*PHENERGAN (promethazine) (oral, suppository, and injectable) – Antinausea, antihistamine  
Possible side effects: Sedation, inability to concentrate, drowsiness, dizziness, blurred or double vision, nausea, rash

phenytoin – See DILANTIN

PREDNISONE (deltasone) – Steroid for treatment of severe allergic reactions  
Possible side effects: High blood pressure, salt and water retention, mood changes, increased sweating, headache, vertigo

PRILOSEC (omeprazole) – Used in treatment of heartburn or gastric hyperacidity  
Possible side effects: Headache, diarrhea, dizziness, nausea

\*promethazine – See PHENERGAN

**WARNING**

Do not use more than two drops of Proparacaine eye drops per treatment as corneal damage may occur. Do not rub eyes for 30 min following anesthetic administration

\*proparacaine eye drops – See ALCaine

\*propranolol hydrochloride – See INDERAL

\*PROVENTIL INHALER (albuterol) – (Bronchodilator) Given for shortness of breath due to wheezing

Possible side effects: Palpitations, tachycardia, increase blood pressure, tremor, nausea

pseudoephedrine – See SUDAFED

guaifenesin/pseudoephedrine – See GUAIFENEX PSE

PYRIDIUM (phenazopyridine) – Relieves pain from bladder infection; compatible w/antibacterial treatment

Possible side effects: Colors urine orange, occasional stomach or intestinal upset

REFRESH PLUS (Artificial Tears) – Saline eye drops for dryness, irritation

\*RESTORIL (temazepam) – Benzodiazepine used to treat insomnia

Possible side effects: Residual drowsiness, dizziness, lethargy (see general WARNING, 6-2)

ROCEPHIN (ceftriaxone) – Antibiotic

Possible side effects: Rash, diarrhea, headache, dizziness

ROMAZICON (flumazenil) – Intravenous med used in benzodiazepine overdose (such as Valium)

Possible side effects: Seizure, agitation, emotional liability, cardiac dysrhythmia

^SILVADENE CREAM (silver sulfadiazine) – Topical burn therapy

Possible side effects: Burning on application, potential allergic reaction. Avoid if allergic to sulfa drugs

^silver sulfadiazine – See ^SILVADENE CREAM

<p style="text-align: center;"><u>WARNING</u> Do not use Sudafed in a patient w/high blood pressure. Can cause high blood pressure, rapid heart rate</p>
--

SUDAFED (pseudoephedrine) – Decongestant

\*temazepam – See RESTORIL

TOBRADEX (tobramycin and dexamethasone) – Used in external eye infection with inflammatory conditions

Possible side effects: Delayed wound healing, lacrimation, itching eyes, edema of the eyelid, allergic contact dermatitis, increased intraocular pressure, glaucoma and cataract formation

tobramycin and dexamethasone – See TOBRADEX

^TOPICORT (desoximetasone) – Cream for skin rashes

TORADOL (ketorolac tromethamine) – Nonsteroidal anti-inflammatory used for pain

Possible side effects: Stomach ulcers/gastric bleeding, allergic reaction, dizziness, nausea, headache, hallucinations

triamcinolone acetonide – See KENALOG IN ORABASE

\*trifluridine – See VIROPTIC

\*trimethoprim/sulfamethoxazole – See BACTRIM DS

TYLENOL (acetaminophen) – Aspirin-free pain reliever

Possible side effects: Overdose could cause sedation

TYLENOL #3 (acetaminophen w/codeine) – Pain reliever

Possible side effects: Sedation, dizziness, nausea

\*valacyclovir – See VALTREX

WARNING

Valium may cause low blood pressure, reduced breathing rate, sedation when given I.V. Additive sedative effects may occur w/other central nervous system depressants (see general WARNING, 6-2)

\*VALIUM (diazepam) – (Injectable, oral, Autoinjectable) sedative, anticonvulsant (antiseizure drug)

Possible side effects: Drowsiness, fatigue, nausea, slow heart rate, blurred vision

\*VALTREX (valacyclovir) – (Oral) Antiviral tablets

Possible side effects: Headache, dizziness, depression, nausea, diarrhea, constipation, abdominal pain, anorexia, vomiting, neuromuscular weakness

\*verapamil – See ISOPTIN

WARNING

Vicodin may seriously impair concentration and cause drowsiness/sedation (see general WARNING, 6-2)

\*VICODIN (hydrocodone/acetaminophen) – Narcotic pain reliever

Possible side effects: Lightheadedness, dizziness, nausea, vomiting, constipation

\*VIROPTIC (trifluridine) – Antiviral eyedrops

Possible side effects: Mild transient burning or stinging upon instillation, eyelid swelling, swelling of the eye, irritation, dry eyes, redness of the eyes, increased intraocular pressure

WARNING

Rapid heart rate, seizures may occur w/overdose Xylocaine Plain. Do not inject intravenously; give smallest amount possible subcutaneously

\*XYLOCAINE PLAIN (lidocaine plain) – Injectable used for laceration repair and as dental anesthetic

**WARNING**

Do not use Xylocaine w/Epinephrine for anesthesia on fingers, toes, earlobes, or nose where constriction of blood vessels may lead to tissue death. Other warnings apply as for Xylocaine above

- \*XYLOCAINE W/EPINEPHRINE (lidocaine w/Epinephrine) –  
Injectable local anesthetic used prior to selected laceration repair
- \*XYLOCAINE/CARDIAC (lidocaine/cardiac) – Use only in cardiopulmonary arrest or w/serious heart rhythm abnormalities. Contact Surgeon
- \*ZITHROMAX (azithromycin) – Antibiotic  
Possible side effects: Diarrhea, nausea, abdominal pain
- \*zolpidem – See AMBIEN

SECTION 7

TOXICOLOGY

HAZARDOUS SPILL LEVEL DEFINITIONS TABLE ..... 7-2  
CONTAMINANT CLEANUP PROCEDURES ..... 7-5  
EYE EXPOSURE TOXICOLOGY TREATMENT ..... 7-6  
SKIN EXPOSURE TOXICOLOGY TREATMENT ..... 7-6  
INHALATION EXPOSURE TOXICOLOGY TREATMENT  
(INCLUDES CARDIOVASCULAR AND CENTRAL  
NERVOUS SYSTEM) ..... 7-7

TOXICOLOGY

## TOXICOLOGY

**HAZARDOUS SPILL LEVEL DEFINITIONS TABLE**

COLOR/ LEVEL	STATE	SYSTEMIC/ INTERNAL OR DAMAGE	FLAMMABILITY	IRRITANCY	SUMM OF HAZARD LEVEL	PROTECTIVE EQUIPMENT
RED 4	Gas, volatile liquid, fumes  Not containable  ARS used to decontaminate	Appreciable effects: Coord, perception, memory, etc  Potential for long-term or delayed serious injury (e.g., cancer) or may result in internal tissue damage	Can produce flammable vapors/fine mist  Sufficient qty to produce hazard	Mod-to-severe  Potential for long-term performance decrement  Eye hazards: Risk of permanent eye damage  Therapy reqd	Catastrophic:  Can cause disabling injury  Potential for systemic tox  Mod-to-severe tissue damage  e.g., Metal vapor like Mercuric Iodide	All crew-members don, activate: QDMs or SEBS
ORANGE 3	Solid or non-volatile liquid  Containable	Appreciable effects: Coord, perception, memory, etc  Potential for long-term or delayed serious injury (e.g., cancer) or may result in internal tissue damage	Can produce flammable vapors/fine mist  Sufficient qty to produce hazard	Possible irritation accompanies systemic tox concerns  Irritancy alone would not drive to Level 3  Therapy reqd	Catastrophic:  Can cause disabling injury or systemic tox, internal tissue damage  e.g., Acetonitrile	All crew-members don, activate: QDMs or SEBS, Silver Shield gloves



**HAZARDOUS SPILL LEVEL DEFINITIONS TABLE (Cont)**

COLOR/ LEVEL	STATE	SYSTEMIC/ INTERNAL DAMAGE	OR FLAMMABILITY	IRRITANCY	SUMM OF HAZARD LEVEL	PROTECTIVE EQUIPMENT
YELLOW 2	Solid or non- volatile liquid  Containable	None	Can produce flammable solids/liquids, but not vapors  Sufficient qty to produce hazard	Mod-to-severe  Lasts > 30 min  Potential for long-term performance decrements  Eye hazards: May risk permanent damage, can affect visual acuity > 30 min  Therapy reqd	Catastrophic:  Can cause disabling injury  No systemic tox concerns  e.g., Sodium Hydroxide w/very high pH (>12)	All crew- members in vicinity*: Don goggles, surgical masks, Silver Shield gloves

\*In vicinity: Crewmembers in cabin or module w/spill

**HAZARDOUS SPILL LEVEL DEFINITIONS TABLE (Concluded)**

COLOR/ LEVEL	STATE	SYSTEMIC/ INTERNAL OR DAMAGE	FLAMMABILITY	IRRITANCY	SUMM OF HAZARD LEVEL	PROTECTIVE EQUIPMENT
BLUE 1	Gas, solid, liquid  May or may not be containable	Minimal effects  No potential for lasting internal tissue damage	Low flammability  High flash point  Low vapor pressure	Slight-to-mod  Lasts > 30 min  Eye only: can affect visual acuity > 30 min  May/May not req therapy	Critical:  Can cause non disabling injury  e.g., 15% Sodium Chloride	All crew- members in vicinity*: Don goggles, surgical masks  Cleanup crew also don surgical gloves
GREEN 0	Gas, solid, liquid  May or may not be containable	None	None	Slight  Lasts < 30 min  All effects resolved w/in 30 min  No therapy reqd	Nonhazard:  Neither health nor fire. If not containable, MCC  e.g., Silicone oil or weak hypertonic solutions	None

\*In vicinity: Crewmembers in cabin or module w/spill

## CONTAMINANT CLEANUP PROCEDURES

1. Remove individual from further exposure
2. Don reqd protective equipment:
  - Tox Level 4: Not containable by cleanup crew.  
Don QDM or SEBS
  - CCK Tox Level 3: QDM or SEBS, Silver Shield Gloves
  - Tox Level 2: Surgical masks, goggles, Silver Shield Gloves
  - Tox Level 1: Surgical masks, goggles, Surgical gloves
  - Tox Level 0: No protective equipment reqd
3. Follow Toxicology Treatment Protocols, section 7
4. Perform cleanup utilizing CCK contents

### If Chemical Spill:

1. Don Silver Shield Gloves
2. Don Yellow Mess-Up Mitt over gloves
3. Clean up spill
4. Turn Mitt inside out and seal w/adhesive strip
5. Place Mitt and gloves in Chemical Resistant Bag(s)
6. Seal Bag(s) w/double stick tape on bag
7. Identify hazard level w/appropriate Toxicology Identification decal
8. Once closed, dispose of Chemical Resistant Bag(s) in Wet Trash

### If Biological/Medical Event:

- CCK 1. Don non-sterile Surgical gloves
2. Don Red Bio-Wipe Bag over gloves
3. Clean up biological waste
4. Turn Red Bio-Wipe Bag inside out and seal with adhesive strip
5. Place gloves and Red Bio-Wipe Bag in large Ziplock Containment Bag(s)
6. Identify hazard level w/appropriate Biohazard Identification decal
7. Once closed, dispose of Ziplock Containment Bag(s) in Wet Trash

## EYE EXPOSURE TOXICOLOGY TREATMENT

1. Remove individual from further exposure

If reqd:

2. Don Quick Don Mask (QDM) to protect against further exposure
3. Perform EYE PROBLEMS, CHEMICAL BURN, 3-17
- Galley 4. Immediately activate Shuttle Emergency Eyewash (SEE)
5. Perform EYE EXAMINATION, 3-15; evaluate for EYE PROBLEMS, FOREIGN BODY/ABRASION, 3-20
6. Contact Surgeon

## SKIN EXPOSURE TOXICOLOGY TREATMENT

1. Remove individual from further exposure

If reqd:

2. Don Quick Don Mask (QDM) to protect against further exposure
3. Use Contaminant Cleanup Kit (CCK)
4. Remove contaminated clothing
5. Remove remaining toxic material from skin
6. Irrigate affected area w/copious amounts of water

If unable to flush:

- Med Locker 7. Use Gauze Pads (IV Admin-14, Airway Subpack-11, Trauma Subpack-14, EENT Subpack-2) wetted w/cold water

As reqd:

8. Perform MAJOR BURNS (1-32) or MINOR BURNS (2-8)
9. Contact Surgeon

**INHALATION EXPOSURE TOXICOLOGY TREATMENT  
(INCLUDES CARDIOVASCULAR AND CENTRAL  
NERVOUS SYSTEM)**

1. Remove individual from further exposure
2. Don Quick Don Mask (QDM) to protect against further exposure

**WARNING**  
Signs of toxicity may be delayed.  
If in doubt about situation, maintain  
crewmember on QDM until Surgeon  
contacted

If respiratory distress:

3. ✓Respiratory rate, record \_\_\_\_\_

Symptoms	✓	Signs	✓
Shortness of breath		Pale	
Difficulty breathing		Sweating	
Anxious		Retractions	
Fatigue		Wheezing	
		Coarse breath sounds	
		Changes in mental status	

If conscious, breathing on own:

4. Continue to use QDM

**WARNING**  
Do not administer Epinephrine if exposed  
to Halogenated Hydrocarbons (i.e., Freon  
Halon 1301)

5. As reqd, perform ALLERGIC REACTION – SEVERE REACTION, 1-29, then:
6. Contact Surgeon for drug administration

~  
If unconscious, not breathing:

7. As reqd, perform CPR (CARDIOPULMONARY RESUSCITATION), ORAL AIRWAY, 1-11, or INTUBATION, ENDOTRACHEAL, 1-12, then:
8. As reqd, perform RESUSCITATOR procedures (see inside cover of Medical Accessory Kit or Cue Card section), then:
9. Contact Surgeon for drug administration

10. ✓Pulse rate, record \_\_\_\_\_

If pulse present:

11. Continue w/oxygen
12. As reqd, perform CPR (CARDIOPULMONARY RESUSCITATION), CARDIAC MONITORING, 1-8, then:
13. Contact Surgeon for drug administration

If no pulse:

14. Continue w/oxygen
15. As reqd, perform CPR (CARDIOPULMONARY RESUSCITATION), 1-6, steps 8-11 (compression/ventilation cycles), then:
16. As reqd, perform INJECTIONS, INTRAVENOUS (I.V.), INTRAVENOUS FLUID INFUSION, 4-11, then:
17. As reqd, perform CPR (CARDIOPULMONARY RESUSCITATION), CARDIAC MONITORING, 1-8, then:
18. Contact Surgeon for drug administration

NOTE

Blood pressure will be present only when crewmember has pulse

19. ✓Blood Pressure (BP); record \_\_\_\_\_

20. Contact Surgeon for drug administration

21. As reqd, perform SEIZURES, 1-42, then:

22. Contact Surgeon for drug administration

SECTION 8

SOMS STOWAGE AND USAGE

ALPHABETICAL LIST ..... 8-2  
USAGE LIST ..... 8-15

**STOWAGE/  
USAGE**

## ALPHABETICAL LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Absorbant Wipes	CCK		72
Ace Bandage	Trauma Subpack-3 Trauma Subpack-13	4 in wide 6 in wide	1 1
acetaminophen (Tylenol)	Drug Subpack-Spine	325 mg	30 tabs
*acetazolamide (Diamox)	Drug Subpack-43	250 mg	30 tabs
Adaptic Bandages	Trauma Subpack-5	3-in X 3-in	3
Afrin (nasal spray)	EENT Subpack-19	3-ml bottles	6
Air temp monitors	EENT Subpack-10	90-120 degF 58-88 degF	2 2
Airway	Airway Subpack-3	oral	1
*albuterol (Proventil Inhaler)	Drug Subpack-11	17-g container	1
*Alcaine (proparacaine) (eye drops)	EENT Subpack-12	15-ml bottle	1
Alcohol Pads	Electrode Attachment Kit IV Admin-18 Airway Subpack-10 Drug Subpack-16 Drug Subpack-Spine Trauma Subpack-6		8 6 10 15 10 10
*Ambien (zolpidem)	Drug Subpack-Spine	10 mg	75 tabs
Ambulatory leg-bag	Trauma Subpack-1	600-ml bag	1
*amikacin (Amikin)	Drug Subpack-29	250 mg/cc, 2-cc unit	1
*Amikin (amikacin)	Drug Subpack-29	250 mg/cc, 2-cc unit	1
*amoxicillin/clavulanate (Augmentin)	Drug Subpack-37	875/125 mg	20 tabs
Anusol HC suppositories	Drug Subpack-41		6
artificial tears (eye drops) Refresh Plus	EENT Subpack-16	0.3 cc	12
Ascriptin (aspirin)	Drug Subpack-Spine	325-mg aspirin w/Maalox	25 tabs
aspirin (Ascriptin)	Drug Subpack-Spine	325-mg aspirin w/Maalox	25 tabs
*Atropine	Drug Subpack-4	1 mg/cc, 2-cc unit	2
*Augmentin (amoxicillin/clavulanate)	Drug Subpack-37	875/125 mg	20 tabs
*azithromycin (Zithromax)	Drug Subpack-36	250 mg	18 tabs

**STOWAGE/  
USAGE**



## ALPHABETICAL LIST

<b>WARNING</b>
* Indicates item to be used only after Surgeon approval or as directed in C/L
^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
*Bactrim DS (trimethoprim/sulfamethoxazole)	Drug Subpack-38		28 tabs
^Bactroban Ointment (mupirocin)	Trauma Subpack-19		1
Bags			
Chemical Resistant	CCK	16 in X 12 in Double Stick Tape closure	8
Containment, Ziplock	CCK	12 in X 12 in Ziplock closure	7
Mess-Up Mitts, Yellow	CCK	Chemical Absorbent Mitt Tape closure	2
Red Bio-Wipe	CCK	Biological Absorbent Mitt Tape closure	2
Ziplock	IV Admin-1 Airway Subpack-4	12 in X 12 in	1 1
Band-aids	Trauma Subpack-11 Drug Subpack-12 Drug Subpack-8 Drug Subpack-Spine (SMS Kit) IV Admin-20	1 in X 3 in Sheer Spot 1 in X 3 in Sheer Spot 1 in X 3 in Sheer Spot	10 10 10 10 6
Batteries	Airway Subpack-5 EENT Subpack-32	AA size AA size	2 2
*Benadryl, injectable (diphenhydramine)	Drug Subpack-13	50 mg/cc, 1-cc unit	2
Benadryl, oral (diphenhydramine)	Drug Subpack-42	25 mg	20 caps
^Benzoin swabs	Trauma Subpack-8	Sepp applicators	5
Biohazard Identification Labels	CCK		20
bisacodyl (Dulcolax)	Drug Subpack-30 Drug Subpack-39	oral, 5 mg suppository, 10 mg	30 tabs 6
Blistex lip balm	EENT Subpack-24		1
Blood pressure cuff	IV Admin-1	w/aneroid sphyg	1
*bupivacaine (Marcaine)	Trauma Subpack-30	0.5% plain, 2 cc	1
Butterfly Needle	IV Admin-18	21 ga	2

## ALPHABETICAL LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Catheter, Foley	Trauma Subpack-10	16 Fr, 30-cc balloon, silastic	1
Straight	Trauma Subpack-10	16 French, Red Rubber	1
	Trauma Subpack-24	14 French, Female	5
*cefadroxil (Duricef)	Drug Subpack-40	500 mg	20 caps
Chemstrip 10 SG	Trauma Subpack-15		13
*cilastin/primaxin (Imipenem IV)	Drug Subpack-34	500 mg, powder form	2 vials
*^Ciloxan (ciprofloxacin) ointment	EENT Subpack-15	0.3%, 3.5-g tube	3
*Cipro (ciprofloxacin)	Drug Subpack-40	500 mg	25 tabs
	Drug Subpack-22		25 tabs
*ciprofloxacin (Cipro)	Drug Subpack-40	500 mg	25 tabs
	Drug Subpack-22		25 tabs
*^ciprofloxacin ointment (Ciloxan)	EENT Subpack-15	0.3%, 3.5-g tube	3
*Claritin (loratadine)	Drug Subpack-43	10 mg	20 tabs
^clotrimazole cream (Lotrimin AF)	Trauma Subpack-26	24-g tube	1
Coban	IV Admin-1	5 in X 5 yard self-adherent wrap	1
Cotton balls	EENT Subpack-22		10
Cotton swabs	EENT Subpack-34		6
Cough lozenges	Drug Subpack-45	5 mg dextromethorphan	15 tabs
*Cyclogyl (cyclopentolate)	EENT Subpack-11	1%, 15-ml bottle	1
*cyclopentolate (Cyclogyl)	EENT Subpack-11	1%, 15-ml bottle	1
deltasone (Prednisone)	Drug Subpack-33	20 mg	30 tabs
*Demerol (meperidine)	Drug Subpack-25	50 mg/cc, 1-cc unit	2
Dental Items	EENT Subpack		
Carver/File	EENT Subpack-26		1
Explorer Probe	EENT Subpack-26		1
Kenalog in Orabase (triamcinolone acetonide)	EENT Subpack-24	5-g tube	1
Mirror	EENT Subpack-26		1
Needles	EENT Subpack-29	long: 27 ga, 1.25 in	6
	EENT Subpack-36	short: 27 ga, 0.75 in	6
Orangewood Sticks	EENT Subpack-26		2
Syringe	EENT Subpack-33		1

## ALPHABETICAL LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Dental Items (Cont)			
Temporary Filling	EENT Subpack-24		1
Toothache Kit	EENT Subpack-35		1
Eugenol dental anesthetic drops		1 ml tubex	3
Tweezers			
Cotton Pellets			
*Xylocaine (lidocaine)	EENT Subpack-30	2% w/Epinephrine 1:100,000, 2% Plain	3 dental carpules 3 dental carpules
Dycal (Base)	EENT Subpack-1	13-g tube	1
Dycal (Catalyst)	EENT Subpack-1	11-g tube	1
^Dermabond	Trauma Subpack-27	skin adhesive	1
^desoximetasone (Topicort)	Trauma Subpack-28	0.25% cream 15-g tube	1
*dexamethasone	Drug Subpack-28	10 mg/cc, 1-cc unit	2
*Dexedrine (dextroamphetamine)	Drug Subpack-14	5 mg	30 tabs
*dextroamphetamine (Dexedrine)	Drug Subpack-14	5 mg	30 tabs
*Diamox (acetazolamide)	Drug Subpack-43	250 mg	30 tabs
*diazepam, injectable (Valium)	Drug Subpack-22	5 mg/cc, 2-cc unit	1
*diazepam, oral (Valium)	Drug Subpack-20	5 mg	30 tabs
*diazepam Autoinjector (Valium)	Drug Subpack-22	10 mg in 2 ml	1
*Diflucan (fluconazole)	Drug Subpack-18	150 mg	6 tabs
*diphenhydramine, injectable (Benadryl)	Drug Subpack-13	50 mg/cc, 1-cc unit	1
diphenhydramine, oral (Benadryl)	Drug Subpack-42	25 mg	20 caps
Drape, sterile	Trauma Subpack-24		1
Dulcolax (bisacodyl)	Drug Subpack-30 Drug Subpack-39	oral, 5 mg suppository, 10 mg	30 tabs 6
*Duragesic Patch (fentanyl)	Drug Subpack-8	25 mcg	1
*Duricef (cefadroxil)	Drug Subpack-40	500 mg	20 caps
Dycal (Base)	EENT Subpack-1	13-g tube	1
Dycal (Catalyst)	EENT Subpack-1	11-g tube	1

## ALPHABETICAL LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
*Duricef (cefadroxil)	Drug Subpack-40	500 mg	20 caps
Dycal (Base)	EENT Subpack-1	13-g tube	1
Dycal (Catalyst)	EENT Subpack-1	11-g tube	1
Ear viewer (Otoscope) + Specula	EENT Subpack-31,32		1 each
Electrode Attachment Kit	Med Locker		
End-Tidal CO2 Detector	Airway Subpack-7		1
*Epinephrine	Drug Subpack-5	1:1000, 1-cc unit	3
Epi Pen w/Injectable Epinephrine	Drug Subpack-9	2 ml, 1:1000	2
Eugenol dental anesthetic drops	EENT Subpack-35	1 ml tubex	3
Eye drops (artificial tears) Refresh Plus	EENT Subpack-16	0.3 cc	12
Eye pads	EENT Subpack-4		2
Eye Shield	EENT Subpack-4	metallic eyepatch	1
Eye viewer (ophthalmoscope head)	EENT Subpack-5,32		1
*fentanyl (Duragesic Patch)	Drug Subpack-8	25 mcg patch	1
Finger splint	Trauma Subpack-23		1
*Flagyl (metronidazole)	Drug Subpack-36	250 mg	28 tabs
*fluconazole (Diflucan)	Drug Subpack-18	150 mg	6 tabs
Fluorescein strips	EENT Subpack-6	0.6 mg	8
Foley catheter	Trauma Subpack-10	16 Fr, 30-cc balloon, silastic	1
Forceps Magill	Trauma Subpack-25 Airway Subpack-19	Blunt curved	1 1
Gauze Pads	Trauma Subpack-14 IV Admin-14 Airway Subpack-11 EENT Subpack-2	4 in X 4 in 2 in X 2 in 4 in X 4 in 4 in X 4 in	5 6 5 5
*Genoptic Ophthalmic Ointment (gentamicin)	EENT Subpack-14	3.5-g tube	1
*gentamicin (Genoptic Ophthalmic Ointment)	EENT Subpack-14	3.5-g tube	1
Gloves, Silver Shield	CCK	chemical resistant	7 pair

## ALPHABETICAL LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Gloves	IV Admin-12 Airway Subpack-8 EENT Subpack-7 CCK	nonsterile, surgical	4 pair 2 pair 2 pair 7 pair
Gloves	Trauma Subpack-24	sterile, surgical	2 pair
Goggles	CCK	eye protection	7
Guaifenesin/pseudophedrine (Guaifenex PSE)	Drug Subpack-44	600-mg guaifenesin, 120-mg pseudophedrine	40 tabs
Guaifenex PSE (guaifenesin/pseudophedrine)	Drug Subpack -44	600-mg guaifenesin, 120-mg pseudophedrine	40 tabs
*Haldol (haloperidol)	Drug Subpack-26	5mg/cc, 1-cc unit	2
*haloperidol (Haldol)	Drug Subpack-26	5mg/cc, 1-cc unit	2
Hazard Identification Labels	CCK	decals yellow,green,blue, orange,red Biohazard IDs	6 each  20
Hemostat	Trauma Subpack-32 Trauma Subpack-33 Airway Subpack-2	small small curved	1 1 1
*hydrocodone/acetaminophen (Vicodin)	Drug Subpack-21	10 mg/660 mg	20
ibuprofen (Motrin)	Drug Subpack-Spine	400 mg	30 tabs
ILMA (Intubating Laryngeal Mask Airway)	Airway Subpack-17 Airway Subpack-18	small large	1 1
*Imipenem IV (cilastin/primaxin)	Drug Subpack-34	500 mg, powder form	2 vials
Imodium (loperamide HCl)	Drug Subpack-30	2 mg	32 caps
*Inderal (propranolol hydrochloride)	Drug Subpack-7	40 mg	24 tabs
Injector (Tubex)	Drug Subpack-Spine Drug Subpack-19 Trauma Subpack-31		1 1 1
Intubating Laryngeal Mask Airway (ILMA)	Airway Subpack-17 Airway Subpack-18	small large	1 1
*Isoptin (verapamil)	Drug Subpack-3,19	2.5 mg/cc, 2-cc unit	2
I.V. Administration Set	IV Admin-17		2
I.V. Intracatheter	Airway Subpack-11 Trauma Subpack-29 IV Admin-8,9 IV Admin-10,11	14 ga 18 ga 18 ga 20 ga	2 1 2 2
Kenalog in Orabase (triamcinolone acetate)	EENT Subpack-24	5-g tube	1
Kerlix	Trauma Subpack-2	4.5 in wide	1 roll

## ALPHABETICAL LIST

WARNING  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Kling	Trauma Subpack-12	3 in wide gauze dressing	2 rolls
Laryngoscope	Airway Subpack-5	Handle w/ Macintosh 3 blade	1
Lever Lock Cannula	IV Admin-17		4
*Lidocaine (Xylocaine)	EENT Subpack-30	2% w/Epinephrine 1:100,000, 1.8-cc unit	3
*Lidocaine (Xylocaine) Plain	EENT Subpack-30	2% w/o Epinephrine, 1.8-cc unit	3
	Trauma Subpack-30	2% w/o Epinephrine, 2-cc unit	2
*Lidocaine/cardiac	Drug Subpack-1,2	20 mg/cc, 5-cc unit	2
Lidocaine/cardiac plunger	Drug Subpack-1,2		2
loperamide HCl (Imodium)	Drug Subpack-30	2 mg	32 caps
*loratadine (Claritin)	Drug Subpack-43	10 mg	20 tabs
^Lotrimin AF Cream (clotrimazole)	Trauma Subpack-26	24-g tube	1
Lubricant (water-soluble)	Trauma Subpack-9	3 g use w/Foley catheter	10
	Airway Subpack-13		5
Magill Forceps	Airway Subpack-19	curved	1
Magnifying glasses and strap	EENT Subpack-1,17	10X magnification	1
*Marcaine (bupivacaine) Plain	Trauma Subpack-30	0.5% plain, 2 cc	1
Masks, surgical	CCK		7
Medical Data Logs	Drug Subpack-Cover	crew size + generic	variable
*meperidine (Demerol)	Drug Subpack-25	50 mg/cc, 1-cc unit	2
*metronidazole (Flagyl)	Drug Subpack-36	250 mg	28 tabs
*Morphine Sulfate	Drug Subpack-23,24	10 mg/cc, 1-cc unit	5
Motrin (ibuprofen)	Drug Subpack-Spine	400 mg	30 tabs
^mupirocin (Bactroban Ointment)	Trauma Subpack-19		1
Mylanta	Drug Subpack-35		24 tabs
*naloxone (Narcan)	Drug Subpack-27	0.4 mg/cc, 1-cc unit	2
*Narcan (naloxone)	Drug Subpack-27	0.4 mg/cc, 1-cc unit	2
nasal catheter	EENT Subpack-21		1
nasal packing, posterior	EENT Subpack-21	10 cm	1
		5.5 cm	1
nasal speculum	EENT Subpack-26		1

## ALPHABETICAL LIST

WARNING  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
nasal spray (Afrin)	EENT Subpack-19	3-ml bottles	6
Needles  Butterfly	Trauma Subpack-22	22 ga, 1.5 in	2
	EENT Subpack-18	18 ga, 1.5 in	1
	Trauma Subpack-29	18 ga, 1.5 in	1
	IV Admin-18	21 ga, 0.75 in	1
^Neosporin Plus	EENT Subpack-25	0.5-oz tube	1
*nitroglycerin patch	Drug Subpack-8	15 mg	1
*nitroglycerin tabs (Nitrostat)	Drug Subpack-6	0.4 mg (1/150)	25
*Nitrostat (nitroglycerin tabs)	Drug Subpack-6	0.4 mg (1/150)	25
*norgestrel/ethinyl estradiol (Ovral-21)	Drug Subpack-17		21 tabs
Op Site (transparent dressing)	IV Admin-14	transparent dressing	6
Operational Bioinstrumentation System (OBS)	Med Locker		
Electrode Attachment Kit (EAK)			1
OBS Belt w/Signal Conditioner			1 ea
Sternal Harness			1
IVA Cable			1
Biomed Cable		2	
Ophthalmoscope head	EENT Subpack-5		1
Ophthalmoscope, spare bulb	EENT Subpack-8		1
Oral airway	Airway Subpack-3		1
Otoscope	EENT Subpack-32		1
Otoscope, spare bulb	EENT Subpack-8		1
*Ovral-21 (norgestrel/ethinyl estradiol)	Drug Subpack-17		21 tabs
Patient/Rescuer Restraints	Med Locker		2 sets
Penrose Tubing (tourniquet)	IV Admin-16	1 in X 18 in	1
Pepto-Bismol	Drug Subpack-31		24 tabs
pH Strips	CCK		10 strips
phenazopyridine (Pyridium)	Drug Subpack-17	200 mg	20 tabs
*Phenergan (promethazine) injectable	Drug Subpack-Spine	50 mg/cc, 1-cc unit	10
*Phenergan (promethazine)	Drug Subpack-15	oral, 25 mg	30 tabs
	Drug Subpack-10	suppository, 25 mg	14

## ALPHABETICAL LIST

<u>WARNING</u>
* Indicates item to be used only after Surgeon approval or as directed in C/L
^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Pope Otowicks	EENT Subpack-27		6
Posterior Nasal Packing	EENT Subpack-21	10 cm, 5.5 cm	2
^Povidone-Iodine swabs	Trauma Subpack-7 Airway Subpack-10 IV Admin-19		10 4 6
Prednisone (deltasone)	Drug Subpack-33	20 mg	30 tabs
*promethazine (Phenergan) injectable	Drug Subpack-Spine	50 mg/cc, 1-cc unit	10
*promethazine (Phenergan)	Drug Subpack-15 Drug Subpack-10	oral, 25 mg suppository, 25 mg	30 tabs 14
*proparacaine (Alcaine) (eye drops)	EENT Subpack-12	15-ml bottle	1
*propranolol hydrochloride (Inderal)	Drug Subpack-7	40 mg	24 tabs
*Proventil Inhaler (albuterol)	Drug Subpack-11	17-g container	1
pseudoephedrine (Sudafed)	Drug Subpack-Spine	30 mg	100 tabs
Pyridium (phenazopyridine)	Drug Subpack-17	200 mg	20 tabs
Radiation dosimeters	Med Locker		
Refresh Plus (artificial tears, eye drops)	EENT Subpack-16	0.3 cc	12
*Restoril (temazepam)	Drug Subpack-Spine	15 mg	40 caps
Restraints Patient Rescuer	Med Locker		2 belts 2 belts
Resuscitator	Med Locker		1
Ruler	Trauma Subpack-26		1
Saline	EENT Subpack-1 Middeck Locker IV Admin-1 Saline Supply Bag	100 ml 250 ml 500 ml 500 ml  All are 0.9% sodium chloride (NaCl) Purpose: I.V. and irrigation, including eyes	1 2 1 4
Salt tablets	Drug Subpack-Spine	1-g NaCl	128 tabs
Scalpels No. 10, 11 No. 10	Trauma Subpack-25 Airway Subpack-2		1 ea 1



## ALPHABETICAL LIST

<b>WARNING</b>
* Indicates item to be used only after Surgeon approval or as directed in C/L
^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Scissors, curved	Trauma Subpack-33	Surgical instrument assembly	1 pair
straight	Airway Subpack-2		1 pair
Sharps Container	Med Locker		1
Shuttle Emergency Eyewash (SEE) System	CCK	Irrigation goggles	1
Silk Sutures	Airway Subpack-11	4-0 Ethicon, Silk	1
^Silvadene Cream (silver sulfadiazine)	Trauma Subpack-20	20-g tube	1
Silver Nitrate Sticks	EENT Subpack-34		5
Silver Shield, Gloves	CCK	chemical resistant	7 pair
^silver sulfadiazine (Silvadene Cream)	Trauma Subpack-20	20-g tube	1
Skin Stapler	Trauma Subpack-21	15-shot stapler	1
Skin Staple Remover	Trauma Subpack-21		1
Skin temp monitors	EENT Subpack-23	84-106 degF	15
Space Motion Sickness (SMS) Kit Alcohol Pads (10) Band-aids (10) *Phenergan Injectables (10) Tubex injector (1)	Drug Subpack-Spine	for SMS treatment  Sheer Spot	1
Splint	Trauma Subpack-23	finger	1
Stabilizer Rod	Airway Subpack-15		1
Steri-Strip skin closures	Trauma Subpack-16		3
Sterile Drapes	Trauma Subpack-24		1
Sterile Gloves	Trauma Subpack-24		2 pair
Stethoscope	Airway Subpack-12		1
Straight Catheter	Trauma Subpack-10 Trauma Subpack-24	16 French, Red Rubber 14 French, Female	1 5
Stylet	Airway Subpack-6		1
Suction device	Airway Subpack-4	Toomey Syringe	1
Sudafed (pseudoephedrine)	Drug Subpack-Spine	30 mg	100 tabs

## ALPHABETICAL LIST

WARNING  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Surgical Instrument Assembly Forceps, small point Hemostat, small Needle Holder Scissors, curved Tweezers, fine point	Trauma Subpack-33		1 ea
Suture	Trauma Subpack-18 Trauma Subpack-17 Trauma Subpack-17 Trauma Subpack-17 Trauma Subpack-18 Airway Subpack-11	4-0 Dexon, w/needle 5-0 Ethilon, w/needle 4-0 Ethilon, w/needle 3-0 Ethilon, w/needle 2-0 Vicryl w/CT-1 needle 4-0 Ethilon, silk	1 1 2 2 1 1
Syringes	Airway Subpack-16 Airway Subpack-15 Trauma Subpack-22 EENT Subpack-20 IV Admin-2,3,4,5,6,7	35 cc 10 cc  3 cc	1 1 1 1 1 in ea
Tape	IV Admin-13 Airway Subpack-9 Trauma Subpack-34 EENT Subpack-9	0.5 in wide 1 in wide 1 in wide 0.5 in wide	1 roll 1 roll 1 roll 1 roll
Tegaderm (transparent dressing)	Trauma Subpack-4	transparent dressing	5
Telfa pads	EENT Subpack-22	3-in X 4-in nonstick bandages	2
*temazepam (Restoril)	Drug Subpack-Spine	15 mg	40 caps
Thermometers, air temp	EENT Subpack-10 EENT Subpack-10	90-120 degF 58-88 degF	2 2
Thermometers, oral	EENT Subpack-3	96-104 degF	18
Thermometers, skin temp	EENT Subpack-23	84-106 degF	15
Tobradex (tobramycin and dexamethasone)	EENT Subpack-28	ophthalmic suspension 7.5 ml	1
tobramycin and dexamethasone (Tobradex)	EENT Subpack-28	ophthalmic suspension 7.5 ml	1
Tongue depressors	EENT Subpack-34		5
Toomey Syringe suction device	Airway Subpack-4		1

## ALPHABETICAL LIST

<b>WARNING</b>
* Indicates item to be used only after Surgeon approval or as directed in C/L
^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
Toothache Kit	EENT Subpack-35		1
Eugenol dental anesthetic drops		1 ml tubex	3
Tweezers			
Cotton pellets			
^Topicort (dexamethasone)	Trauma Subpack-28	0.25% cream 15-g tube	1
Tourniquet (Penrose Tubing)	IV Admin-16	1 in x 18 in	1
Towels	Misc Stowage		
Tracheal Tube (ET Tube)	Airway Subpack-20	w/stylet	1
Tracheostomy Items	Airway Subpack		1
Alcohol Pads	Airway Subpack-10		
Straight scissors	Airway Subpack-2		
Curved forceps	Airway Subpack-19		
Tracheal hook	Airway Subpack-2		
Tracheostomy tube	Airway Subpack-1		
Tracheostomy tube holder	Airway Subpack-1		
Scalpel	Airway Subpack-2	No. 10	
Povidone-Iodine swabs	Airway Subpack-10		4
Silk Sutures	Airway Subpack-11	4-0 Ethicon, Silk	1
transparent dressing (Tegaderm)	Trauma Subpack-4	Tegaderm	5
	IV Admin-14	Op Site	6
triamcinolone acetonide (Kenalog in Orabase)	EENT Subpack-24	5-g tube	1
*trifluridine (Viroptic)	Drug Subpack-1	7.5 ml	1
*trimethoprim/sulfamethoxazole (Bactrim DS)	Drug Subpack-38		28 tabs
Tubex Injector	Drug Subpack-Spine		1
	Drug Subpack-19		1
	Trauma Subpack-31		1
tweezers	EENT Subpack-34	4.5-in blunt, use w/Otowicks	1
Tylenol (acetaminophen)	Drug Subpack-Spine	325 mg	30 tabs
Urine Test Package	Trauma Subpack-15		1
Chemstrip 10 SG			13 strips
Color Chart			

## ALPHABETICAL LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description	Amount
*valacyclovir (Valtrex)	Drug Subpack-32	1 g	21 tabs
*Valium, injectable (diazepam)	Drug Subpack-22	5 mg/cc, 2-cc unit	1
*Valium, oral (diazepam)	Drug Subpack-20	5 mg	30 tabs
*Valium (diazepam) Autoinjector	Drug Subpack-22	10 mg in 2 ml	1
*Valtrex (valacyclovir)	Drug Subpack-32	1 g	21 tabs
*verapamil (Isoptin)	Drug Subpack-3,19	2.5 mg/cc, 2-cc unit	2
*Vicodin (hydrocodone/acetaminophen)	Drug Subpack-21	10 mg/660 mg	20
*Viroptic (trifluridine)	Drug Subpack-1	7.5 ml	1
Xeroform Gauze	Trauma Subpack-11	5 in X 9 in	1
*Xylocaine (Lidocaine)	EENT Subpack-30	2% w/Epinephrine, 1:100,000, 1.8-cc unit	3
*Xylocaine (Lidocaine) Plain	EENT Subpack-30	2% w/o Epinephrine, 1.8-cc unit	3
	Trauma Subpack-30	2% w/o Epinephrine, 2-cc unit	2
Y-Type Catheter Extension	IV Admin-17		3
Ziplock Bags	IV Admin-1	12 in X 12 in	1
	Airway Subpack-4		1
	CCK		7
*Zithromax (azithromycin)	Drug Subpack-36	250 mg	18 tabs
*zolpidem (Ambien)	Drug Subpack-Spine	10 mg	75 tabs

## USAGE LIST

WARNING

\* Indicates item to be used only after Surgeon approval or as directed in C/L

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Abdominal Injury	Go to 1-27	
Abdominal Pain	Go to 2-2	
Airway	Go to 1-11	
Oral airway	Airway Subpack-3	
ILMA (Intubating Laryngeal Mask Airway)	Airway Subpack-17 Airway Subpack-18	small large
Stabilizer Rod	Airway Subpack-15	
Syringe	Airway Subpack-16	35 cc
Tracheal tube	Airway Subpack-20	
Stylet	Airway Subpack-6	
Laryngoscope	Airway Subpack-5	
Tracheostomy Items	Airway Subpack	
Alcohol Pads	Airway Subpack-10	
Straight scissors	Airway Subpack-2	
Curved forceps	Airway Subpack-19	
Tracheal hook	Airway Subpack-2	
Tracheostomy tube	Airway Subpack-1	
Tracheostomy tube holder	Airway Subpack-1	
Scalpel	Airway Subpack-2	No. 10
^Povidone-Iodine swabs	Airway Subpack-10	
Silk Sutures	Airway Subpack-11	4-0 Ethicon, Silk
Toomey Syringe	Airway Subpack-4	suction device
Tubing	Airway Subpack-6	use w/Toomey Syringe
Allergic Reaction	Go to 1-29,2-3	
*Benadryl, injectable (diphenhydramine)	Drug Subpack-13	50 mg/cc
Benadryl, oral (diphenhydramine)	Drug Subpack-42	25 mg caps
*Claritin (loratadine)	Drug Subpack-43	10 mg
*dexamethasone	Drug Subpack-28	10 mg/cc
*Epinephrine	Drug Subpack-5	
Epi Pen Injector	Drug Subpack-9	
Guaifenesin PSE (guaifenesin/pseudoephedrine)	Drug Subpack-44	40 tabs
Prednisone (deltasone)	Drug Subpack-33	20-mg tabs
*Proventil Inhaler (albuterol)	Drug Subpack-11	17-g container
Sudafed (pseudoephedrine)	Drug Subpack-Spine	30 mg pseudoephedrine
Altitude Sickness	Go to 2-4	
*Diamox tablets (acetazolamide)	Drug Subpack-43	250 mg

## USAGE LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Anesthetic – local injection *Marcaine (bupivacaine) Plain *Xylocaine (Lidocaine) *Xylocaine (Lidocaine) Plain	Trauma Subpack-30 EENT Subpack-30 EENT Subpack-30 Trauma Subpack-30	0.5% plain 2% w/Epinephrine 2% w/o Epinephrine 1.8-cc unit 2% w/o Epinephrine 2-cc unit
Antacids Mylanta	Drug Subpack-35	chewable tab
Antibiotics *Amikin (amikacin) *Augmentin (amoxicillin/clavulanate) *Bactrim DS (trimethoprim/ sulfamethoxazole) ^Bactroban Ointment (mupirocin) *^Ciloxan Ointment (ciprofloxacin) *Cipro (ciprofloxacin) *Duricef (cefadroxil) *Flagyl (metronidazole) *Genoptic Ophthalmic Ointment (gentamicin) *Imipenem IV (Cilastin/Primaxin) ^Neosporin Plus *Viroptic (trifluridine), Ophthalmic Drops *Zithromax (azithromycin)	Drug Subpack-29 Drug Subpack-37 Drug Subpack-38 Trauma Subpack-19 EENT Subpack-15 Drug Subpack-40 Drug Subpack-22 Drug Subpack-40 Drug Subpack-36 EENT Subpack-14 Drug Subpack-34 EENT Subpack-25 Drug Subpack-1 Drug Subpack-36	250 mg/cc 875/125 mg 250 mg 500 mg 500 mg 500 mg 250 mg 3.5-g tube 500 mg, powder form 0.5 oz tube 7.5 ml 250 mg
Antiseptics Alcohol Pads  ^Povidone-Iodine swabs	Electrode Attachment Kit IV Admin-18 Airway Subpack-10 Drug Subpack-16 Drug Subpack-Spine Trauma Subpack-6 Trauma Subpack-7 Airway Subpack-10 IV Admin-19	

## USAGE LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Back Pain	Go to 2-4	
Bandages		
Ace Bandage	Trauma Subpack-3 Trauma Subpack-13	4 in 6 in
Adaptic Bandages	Trauma Subpack-5	3 in X 3 in
Band-aids	Drug Subpack-Spine Drug Subpack-8 Trauma Subpack-11 Drug Subpack-Spine Drug Subpack-12 IV Admin-20	1 in X 3 in 1 in X 3 in 1 in X 3 in Sheer Spot Sheer Spot Sheer Spot
Gauze Pads	Airway Subpack-11 EENT Subpack-2 IV Admin-14 Trauma Subpack-14	4 in X 4 in 4 in X 4 in 2 in X 2 in 4 in X 4 in
Kerlix	Trauma Subpack-2	4.5 in
Kling	Trauma Subpack-12	3 in
Op Site (transparent dressing)	IV Admin-14	
Tegaderm, transparent dressing	Trauma Subpack-4	
Telfa pads	EENT Subpack-22	3 in X 4 in
Bends (DCS)	Go to 3-4	
Bladder Infection	Go to 2-5	
*Augmentin (amoxicillin/clavulanate)	Drug Subpack-37	875/125 mg
*Bactrim DS (trimethoprim/sulfamethoxazole)	Drug Subpack-38	
Chemstrip 10 SG	Trauma Subpack-15	
*Cipro (ciprofloxacin)	Drug Subpack-22 Drug Subpack-40	500 mg 500 mg
Color chart	Trauma Subpack-15	
*Duricef (cefadroxil)	Drug Subpack-40	500 mg
Pyridium (phenazopyridine)	Drug Subpack-17	200 mg
Breathing Difficulty	Go to 1-3	
Stethoscope	Airway Subpack-12	
*Proventil Inhaler	Drug Subpack-11	
Bronchitis	Go to 2-7	
Burns	Go to 1-32,2-8	
^Silvadene Cream (silver sulfadiazine)	Trauma Subpack-20	20-g tube

## USAGE LIST

**WARNING**

\* Indicates item to be used only after Surgeon approval or as directed in C/L

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Cardiac Medications *Atropine *Epinephrine *Isoptin (verapamil) *Lidocaine Lidocaine plunger *Nitroglycerin patch *Nitrostat tabs (nitroglycerin)	Drug Subpack-4 Drug Subpack-5 Drug Subpack-3,19 Drug Subpack-1,2 Drug Subpack-1,2 Drug Subpack-8 Drug Subpack-6	1 mg/cc, 2-cc unit 1:1000, 1-cc unit 2.5 mg/cc, 2-cc unit 20 mg/cc, 5-cc unit 15 mg/24 hr (0.6 mg/hr) 0.4 mg (1/150)
Cardiac Monitoring Alcohol Pads Electrode Attachment Kit (EAK) OBS Belt w/Signal Conditioner Sternal Harness IVA Cable Biomed Cable	Go to 1-8 Electrode Attachment Kit Med Locker	for ECG Downlink
Chest Injury	Go to 1-35	
Chest Pain	Go to 1-36	
Choking	Go to 1-4	
Cleanup Supplies SEE, masks, Red Bio-Wipe Bags, Yellow Mess-Up Mitts, Hazard Identification labels, chemical resistant gloves, surgical gloves, goggles, chemical resistant bags (16 in H X 12 in W), Ziplock Bags (12 in X 12 in) pH Strips Absorbent Wipes	CCK	for contaminant cleanup
Congestion Afrin (nasal spray) Sudafed (pseudoephedrine)	Go to 3-3 EENT Subpack-19 Drug Subpack-Spine	3-ml bottles 30 mg
Constipation Dulcolax (bisacodyl), oral Dulcolax (bisacodyl), suppositories	Go to 3-3 Drug Subpack-30 Drug Subpack-39	5 mg tabs 10 mg
Cough cough lozenges	Go to 2-7 Drug Subpack-45	5-mg dextromethorphan



## USAGE LIST

WARNING

\* Indicates item to be used only after Surgeon approval or as directed in C/L

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
CPR (Cardiopulmonary Resuscitation) Resuscitator & Patient/Rescuer Restraints	Go to 1-6 Med Locker MAK	
Cricothyrotomy Alcohol Pads Curved forceps Scalpel Silk Sutures Straight scissors Tracheal hook Tracheostomy tube Tracheostomy tube holder	Go to 1-19 Airway Subpack-10 Airway Subpack-19 Airway Subpack-2 Airway Subpack-11 Airway Subpack-2 Airway Subpack-2 Airway Subpack-1 Airway Subpack-1	
Decompression Sickness (DCS)	Go to 3-4	
Decongestants Afrin (nasal spray) Guaifenesin PSE (guaifenesin/ pseudophedrine)  Sudafed (pseudoephedrine)	EENT Subpack-19 Drug Subpack-44  Drug Subpack-Spine	3-ml bottles 600-mg guaifenesin, 120-mg pseudophedrine 30 mg
Dental Equipment Carver/File Explorer Probe Kenalog in Orabase (triamcinolone acetonide) Mirror Needles  Orangewood sticks Syringe Temporary Filling Toothache Kit Eugenol dental anesthetic drops Tweezers Cotton Pellets *Xylocaine (lidocaine)	EENT Subpack EENT Subpack-26 EENT Subpack-26 EENT Subpack-24  EENT Subpack-26 EENT Subpack-29 EENT Subpack-36 EENT Subpack-26 EENT Subpack-33 EENT Subpack-24 EENT Subpack-35  EENT Subpack-30	5-g tube  long: 27 ga, 1.25 in short: 27 ga, 0.75 in  1 ml tubex  2% w/Epinephrine 1:100,000 2% Plain

## USAGE LIST

WARNING

\* Indicates item to be used only after Surgeon approval or as directed in C/L

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Dental Equipment (Cont)	EENT Subpack	
Dycal (Base)	EENT Subpack-1	13-g tube
Dycal (Catalyst)	EENT Subpack-1	11-g tube
Diagnostic Equipment		
Blood pressure cuff	IV Admin-1	
Magnifying glasses and strap	EENT Subpack-1,17	
Ophthalmoscope head	EENT Subpack-5	
Otoscope	EENT Subpack-32	
Stethoscope	Airway Subpack-12	
Thermometers, oral	EENT Subpack-3	Tempadot
Tongue depressors	EENT Subpack-34	
Urine Test Package	Trauma Subpack-15	
Chemstrip 10 SG		
Color Chart		
Diarrhea	Go to 3-11	
Imodium (loperamide HCl)	Drug Subpack-30	2 mg
Pepto-Bismol	Drug Subpack-31	chewable tablets
Difficulty Breathing	Go to 1-3	
*Proventil Inhaler	Drug Subpack-11	17-g containers, 90-mcg albuterol per actuation
Dysbarism	Go to 3-4	see DCS
Ear Drops		
Tobradex Ophthalmic Drops (tobramycin and dexamethasone)	EENT Subpack-28	7.5 ml
Ear Problems	Go to 3-12	
Otoscope	EENT Subpack-32	
Pope Otowicks	EENT Subpack-27	
Tweezers	EENT Subpack-34	4.5-in blunt
Eye Treatment	Go to 1-38,3-17	
*^Ciloxan (ciprofloxacin), Ophthalmic Ointment	EENT Subpack-15	0.3%, 5-ml bottle
*Cyclogyl (cyclopentolate)	EENT Subpack-11	1%, 15-ml bottle
Eye pads	EENT Subpack-4	
Eye Shield	EENT Subpack-4	
Fluorescein strips	EENT Subpack-6	

## USAGE LIST

**WARNING**  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Eye Treatment (Cont)	Go to 1-38,3-17	
*Genoptic Ophthalmic Ointment (gentamicin)	EENT Subpack-14	3.5-g tube
Ophthalmoscope head	EENT Subpack-5	
Otoscope	EENT Subpack-32	
pH Strips	CCK	
*proparacaine (Alcaine) eye drops	EENT Subpack-12	15-ml bottle
Refresh Plus (artificial tears, eye drops)	EENT Subpack-16	0.3 cc
Saline, irrigation solution	EENT Subpack-1	100 ml, 0.9% sodium chloride (NaCl)
Irrigation – SEE goggles	CCK	Irrigation goggles
*Viroptic (trifluridine), ophthalmic drops	Drug Subpack-1	7.5 ml
Fever		
Thermometers, oral	EENT Subpack-3	Tempadot
Thermometers, temp	EENT Subpack-23	
Fever Blister		
*Valtrex (valacyclovir)	Drug Subpack-32	1 g
Fluid Loading		
Salt Tablets (Sodium chloride)	Drug Subpack-Spine	1 g
Fracture/Dislocation	Go to 4-3	
Head Injury	Go to 1-39	
Headache	Go to 5-3	see PAIN RELIEF
Heart Medications		
*Atropine	Drug Subpack-4	1 mg/cc, 2-cc unit
*Epinephrine	Drug Subpack-5	1:1000, 1-cc unit
*Isoptin (verapamil)	Drug Subpack-3,19	2.5 mg/cc, 2-cc unit
*Lidocaine/cardiac	Drug Subpack-1,2	20 mg/cc, 5-cc unit
Lidocaine/cardiac plunger	Drug Subpack-1,2	
*Nitroglycerin patch	Drug Subpack-8	15 mg/24 hr (0.6 mg/hr)
*Nitrostat tabs (Nitroglycerin)	Drug Subpack-6	0.4 mg (1/150)
Hemorrhoids		
Anusol HC suppositories	Drug Subpack-41	
Hot Cabin	Go to 4-5	
Air temp monitors	EENT Subpack-10	90-120 degF
Skin temp monitors	EENT Subpack-23	84-106 degF

## USAGE LIST

WARNING

\* Indicates item to be used only after Surgeon approval or as directed in C/L

^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Injection Equipment		
22 Gauge Hypodermic Needle	Trauma Subpack-22	
18 Gauge Hypodermic Needle	EENT Subpack-18 Trauma Subpack-29	
21 Gauge Butterfly Needle	IV Admin-18	
Coban	IV Admin-1	5 in X 5 yard self-adherent wrap
Cotton Balls	EENT Subpack-22	
Gloves, non-sterile	IV Admin-12 Airway Subpack-8 EENT Subpack-7 CCK	
I.V. Intracatheter	IV Admin-10,11 IV Admin-8,9 Trauma Subpack-29 Airway Subpack-11	20 ga 18 ga 18 ga 14 ga
Y-type catheter extension	IV Admin-17	
I.V. Administration Set	IV Admin-17	
Lever Lock cannula	IV Admin-17	
Penrose Tubing (Tourniquet)	IV Admin-16	
Syringes	Airway Subpack-16 Airway Subpack-15 EENT Subpack-20 Trauma Subpack-22 IV Admin-2,3,4,5,6,7	35 cc 10 cc  3 cc
Tubex Injector	Drug Subpack-Spine Drug Subpack-19 Trauma Subpack-31	
Irrigation Equipment		
I.V. Intracatheter	IV Admin-10,11 IV Admin-8,9	20 ga 18 ga
Needle	Trauma Subpack-22	22 ga

## USAGE LIST

<b>WARNING</b>
* Indicates item to be used only after Surgeon approval or as directed in C/L
^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Irrigation Equipment (Cont) Saline	IV Admin-1 EENT Subpack-1 Middeck Locker Saline Supply Bag	500 ml, 0.9% sodium chloride (NaCl) 100 ml, 0.9% sodium chloride (NaCl) 250 ml, 0.9% sodium chloride (NaCl) 500 ml, 0.9% sodium chloride (NaCl)
Syringes	Airway Subpack-16 Airway Subpack-15 EENT Subpack-20 Trauma Subpack-22	35 cc 10 cc
Lacerations	Go to 4-30	
Lips – Cracked, Chapped, Dry Blistex lip balm	EENT Subpack-24	
Motion Sickness *Dexedrine (dextroamphetamine) *Phenergan, injectable (promethazine) *Phenergan, oral (promethazine) *Phenergan, suppository (promethazine) Space Motion Sickness (SMS) Kit Alcohol Pads (10) Band-aids (10) *Phenergan Injectables (10) Tubex Injector (1)	Go to 4-38 Drug Subpack-14 Drug Subpack-Spine Drug Subpack-15 Drug Subpack-10 Drug Subpack-Spine	5 mg 1-cc unit, 50 mg/cc 25 mg 25 mg for SMS treatment  Sheer Spot
Muscle Relaxants *Valium, injectable (diazepam) *Valium, oral (diazepam) *Valium (diazepam) Autoinjector	Drug Subpack-22 Drug Subpack-20 Drug Subpack-22	5 mg/cc, 2-cc unit 5 mg 10 mg in 2 ml
Neck Injury	Go to 1-39	
Nosebleed Afrin, nasal spray Foley Catheter Nasal Catheter Nasal speculum Posterior Nasal Packing Silver Nitrate Stick	Go to 4-40 EENT Subpack-19 Trauma Subpack-10 EENT Subpack-21 EENT Subpack-26 EENT Subpack-21 EENT Subpack-34	3-ml bottles    10 cm, 5.5 cm

## USAGE LIST

WARNING  
 \* Indicates item to be used only after Surgeon approval or as directed in C/L  
 ^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
<b>Pain Medications</b>		
Ascriptin (aspirin)	Drug Subpack-Spine	325-mg aspirin w/Maalox
*Demerol (meperidine)	Drug Subpack-25	50 mg/cc, 1-cc unit
*Duragesic Patch (fentanyl)	Drug Subpack-8	25 mcg
*Morphine Sulfate	Drug Subpack-23,24	10 mg/cc, 1-cc unit
Motrin (ibuprofen)	Drug Subpack-Spine	400 mg
Tylenol (acetaminophen)	Drug Subpack-Spine	325 mg
*Vicodin (hydrocodone/acetaminophen)	Drug Subpack-21	10 mg/660 mg
<b>Radiation</b>		
Dosimeters	Go to 5-4 Med Locker	
<b>Resuscitation</b>		
Resuscitator	Med Locker	
<b>Seizures</b>		
*Valium, injectable (diazepam)	Go to 1-42 Drug Subpack-22	5 mg/ml, 2-cc unit
*Valium (diazepam) Autoinjector	Drug Subpack-22	10 mg in 2 ml
<b>Shock</b>		
	Go to 1-44	
<b>Skin Rash and Itching</b> (for allergic rash, see ALLERGIC REACTION)		
Benadryl, oral (diphenhydramine)	Go to 5-7 Drug Subpack-42	25 mg
*Claritin (loratadine)	Drug Subpack-43	10 mg
^Lotrimin AF Cream (clotrimazole)	Drug Subpack-26	24-g tube
Ruler	Drug Subpack-26	photodocumentation
^Topicort (desoximetasone)	Drug Subpack-28	0.25% cream, 15-g tube
<b>Sleeping Pills</b>		
*Ambien (zolpidem)	Drug Subpack-Spine	10 mg
*Restoril (temazepam)	Drug Subpack-Spine	15 mg
<b>Space Motion Sickness (SMS) Kit</b>		
Alcohol Pads (10)	Drug Subpack-Spine	for SMS treatment
Band-aids (10)		Sheer Spot
*Phenergan Injectables (10)		
Tubex Injector (1)		
<b>Splints</b>		
Finger	Trauma Subpack-23	



## USAGE LIST

<b>WARNING</b>
* Indicates item to be used only after Surgeon approval or as directed in C/L
^ Do not allow this item to come in contact with EMU. EV and IV crewmembers must contact Surgeon prior to use

Name	Location	Description
Toothache Toothache Kit Eugenol dental anesthetic drops Tweezers Cotton pellets *Xylocaine (Lidocaine) Dycal (Base + Catalyst)	Go to 5-10 EENT Subpack-35  EENT Subpack-30  EENT Subpack-1	1 ml tubex  2% w/Epinephrine 2% Plain
Upset Stomach Pepto-Bismol Mylanta	Go to 4-6 Drug Subpack-31 Drug Subpack-35	chewable tablets chewable tablets
Urinary Retention Catheter, Foley  Catheter, Straight  Lubricant, water-soluble  Ambulatory leg-bag Chemstrip 10 SG Color Chart	Go to 5-18 Trauma Subpack-10  Trauma Subpack-10  Trauma Subpack-24 Trauma Subpack-9 Airway Subpack-13 Trauma Subpack-1 Trauma Subpack-15 Trauma Subpack-15	16 Fr, 30-cc balloon, silastic  16 French, Red Rubber 14 French, Female  600-ml bag
Vomiting *Phenergan, injectable (promethazine) *Phenergan, oral (promethazine) *Phenergan, suppository (promethazine)	Go to 5-20 Drug Subpack-Spine Drug Subpack-15 Drug Subpack-10	50 mg/cc, 1-cc unit 25 mg 25 mg
Yeast Infection (Vaginitis) *Diflucan (fluconazole)	Drug Subpack-18	oral, 150 mg tablets



SECTION 9

CUE CARD CONFIGURATION

**CUE CARDS**

This Page Intentionally Blank





TOP

### AIRWAY (ILMA)


1. Unstow from Med Locker:  
Airway Subpack-17,18



2. Check integrity of ILMA cuff  
Inject 8 cc air;  
confirm inflation;  
deflate cuff  
Cuff tip must be in shape above.  
Apply Surgilube to bottom of cuff
3. Insert cuff in mouth and rub Surgilube on roof of mouth



4. While maintaining contact with roof of mouth, hug chin while inserting ILMA




5. Inflate mask w/o holding tube or handle:  
sm = 30 cc; lg = 40 cc



6. Provide ventilation using AMBU Bag
7. Contact Surgeon prior to Endotracheal ET tube insert

8. Lubricate ET tube w/ Surgilube. Hold tube with black line facing ILMA handle
9. Holding ILMA handle, gently insert ET tube into metal shaft



10. Advance ET tube (24 cm for small; 26 cm for large)
11. Inflate ET tube cuff (8-10 cc); confirm placement w/ Stethoscope



12. Remove all air from ILMA cuff. Remove tracheal tube connector and ease ILMA out by gently rotating towards neck
13. Use stabilizing rod to keep ET tube in place; remove ILMA until ET tube can be grasped above the teeth



14. Remove stabilizing rod and gently unthread cuff valve of ET tube
15. Replace ET tube connector



16. Attach AMBU Bag to ET tube and give patient two breaths
17. Check for breath sounds in both lungs; reposition ET tube until breath sounds confirmed
18. If successful, tape ET tube securely in place

MED-2a/O/D

(reduced copy)

CC 9-5

MED/ALL/GEN K,2



TOP, BACK OF 'AIRWAY (ILMA)'



Intentionally Blank

MED-2b/O/B



(reduced copy)  
CC 9-6


MED/ALL/GEN K,2



HOOK  
VELCRO

TOP

**RESUSCITATOR**



1. Deploy SOMS Resuscitator if not already instructed by the Medical Checklist

2. Verify SMART Valve is in ENABLE position (see Figure 1)  
Do not disable unless given direction from the flight surgeon

- \* If SMART Valve is not in ENABLE position, \*
- \* rotate the bar so that it points forward towards \*
- \* the patient valve \*




Figure 1 - SMART Valve  
ENABLE position

3. Verify Relief Valve is in UNLOCK position (see Figure 2)  
Do not lock unless given direction from the flight surgeon

- \* If Relief Valve is not in UNLOCK position, \*
- \* rotate the top Relief Valve knob so that the \*
- \* Lock arrow points at the patient valve \*
- \* DO NOT unscrew knurled knob \*




Figure 2.- Relief Valve  
UNLOCK position

4. Remove clean bag covering the Quick Disconnect

(Continue on BACK)

MED-4a/O/A

HOOK  
VELCRO

(reduced copy)

CC 9-7

MED/ALL/GEN K,2

TOP, BACK OF 'RESUSCITATOR'

HOOK  
VELCRO

5. If SOMS Resuscitator is to be used with oxygen, connect Quick Disconnect with Shuttle oxygen port LEH O<sub>2</sub> 8 on Middeck panel MO69M. Turn oxygen switch to OPEN at LEH O<sub>2</sub> 8

NOTE

LEH O<sub>2</sub> ports located on panels MO32M and C6 may be used if panel MO69M is not available

6. Interface SOMS Resuscitator with patient based on if Face Mask (see Figure 3), Intubating Laryngeal Mask Airway (ILMA) (see Figure 4), or endotracheal tube (ETT) (see Figure 5) is used



Figure 3.- Face Mask



Figure 4.- ILMA



Figure 5.- ETT

7. Squeeze SOMS Resuscitator until chest rises. Allow patient to passively exhale
  - \* If the AMBU bag becomes difficult to squeeze \*
  - \* or the chest fails to rise, check position of \*
  - \* ILMA or endotracheal tube using CO<sub>2</sub> \*
  - \* detector and listening for breath sounds \*
  - \* Refer to INTUBATION, ENDOTRACHEAL \*
  - \* procedure in SOMS Medical Checklist \*
8. Release SOMS Resuscitator to allow AMBU Bag to refill
9. Repeat every 5 seconds (approximately 12 breaths per minute) until spontaneous breathing resumes

NOTE

If oxygen is used with the SOMS Resuscitator, as patient care is stabilized, the Oxygen Hose should be tethered to the Middeck with Velcro straps

MED-4b/O/A

HOOK  
VELCRO


CC 9-8

MED/ALL/GEN K,2



Space Shuttle Program  
FLIGHT DATA FILE

JSC-48031  
GENERIC, REV K



**MEDICAL  
CHECKLIST**

**STS  
ALL**

Flight Cover (trim bottom to expose tabs)