

# Aqua-AIRS and MetOP-IASI Temperature and Water Vapor Retrievals Evaluation with Global RAOBs and Model Forecasts



By

Murty Divakarla, Chris Barnet, Mitch Goldberg,  
Tom King, Eric Maddy, Xingpin Liu, Zhaohui Cheng,  
Antonia Gambakorta, Fengying Sun, and Lihang Zhou

NOAA/NESDIS  
Camp Springs, MD 20746

**An evaluation of the IASI and AIRS Retrievals using Matched RAOB Measurements, ECMWF and NCEP-GFS Forecasts is presented. Differences due to Algorithm Differences (Channels, Cloud-Clearing), Sounding Geometry, and Spatial Sampling of the Matched Data Sets are discussed.**

**Comments/Suggestions are welcome**

E-Mail Contact: [Murty.Divakarla@noaa.gov](mailto:Murty.Divakarla@noaa.gov)

Acknowledgements:

- OSDPD (Frank Tilley and Tony Reale for RAOB Match Files)
- ECMWF and NCEP for Model Forecasts/Analysis

# Validation Data - Algorithms



## MetOp-IASI/Aqua-AIRS Match-up Database

(Adapted to MetOP-IASI from Aqua-AIRS Validation System)

- RAOB Measurements Matched to Aqua (1:30 AM/PM) and MetOP(9:30 AM/PM) Satellite Observations
- MetOp-IASI/AMSU-A/AMSU-B Level1B Radiances
- IASI Level-2 Retrievals
- Aqua-AIRS/AMSU-A Level1B Radiances
- AIRS Level-2 Retrievals
- NCEP-GFS (AVN) Level2-Forecast/Analysis
- ECMWF Level-2 Forecast/Analysis
- MetOp-ATOVS/NOAA-18 Level-2 Retrievals

Collocated Within  $\pm 3$  Hrs. & 100 Km Radius  
Data Used In this Study : November, 2007-2009

- Emulate IASI/AIRS Retrieval Algorithms
- Reprocessing Options with Algorithm Upgrades, New Data, Versions
- Test New Ideas in the Retrieval Algorithm

# What We Did

## Data Used : 01/20/2008 – 08/02/2008



- IASI/AIRS Temperature and Water Vapor Profile Retrieval Statistics with
  - » Collocated RAOBs and other Data Sets (ECMWF, NCEP-GFS, MetOp-ATOVS) for the Same Period (Regions Sampled by Aqua and MetoP Orbits Differ)
    - » 12,000 Accepted Matches for IASI
    - » ~ 6,000 Accepted Matches for AIRS
- Looked into Some Cloud-Clearing Aspects
  - » Noise Amplification in Cloud-Cleared Radiances
    - with IASI-4 FOVs *and* AIRS-9 FOVs
- Stats for A subset of Matches Where IASI & AIRS are looking at the same Ground Location but at different times ~ 1000 Collocations
  - » IASI Retrievals with RAOB Matches
  - » AIRS Retrievals with RAOB Matches
  - » AIRS Like IASI (Use 4 FOVs of AIRS) Retrievals with RAOB Matches
- MetOp-IASI/AIRS Validation Web-site




# Questions Addressed


- **What are the expectations from IASI Level-2 Temperature and Water Vapor Products ?**
  1. **Whether IASI Retrievals are Comparable to AIRS Retrievals (Similar Sounding Instruments)**
    - Yield
    - RMS Difference and Bias wrt RAOBs
  2. **Improvement seen with the IASI/AIRS Retrievals Compared to the Currently Operating Baseline Systems**
    - M2-ATOVS (9:30 AM/PM) Retrievals from the Same MetOp Platform (HIRS + AMSU-A/B)
    - NOAA-18 ATOVS (1:30 PM/AM) Retrievals are Time Coincident with Aqua-AIRS Retrievals
    - How IASI/AIRS Retrievals, ECMWF and AVN Models Compare with RAOBs ?
  3. **Cloud Clearing Aspects Aqua-AIRS MetOp IASI**
    - » **Differences and Impacts of 9 FOVs vs. 4 FOVs**



# Instruments, Algorithms and Products



## Channels Used in Physical Retrieval



### Pondering Questions That Need Careful Attention While Evaluating Retrievals



- Y Instrument Differences: **AIRS vs. IASI**, Geometry, Channels, S/N, Spectral Resolution, and other Trade-Offs
- Y Algorithm Differences: Three Pivotal Points: (a) Cloud Clearing IASI (4 FOVs) vs. AIRS (9 FOVs), (b) Channels Used in the Physical Retrieval (and the FG Reg), (c) Retrieval Conv. Criteria, Quality Indicators, Rejection Thresholds, etc.
- Y Consequences of Satellite Orbit Time (IASI: 9:30 AM/PM; AIRS: 1:30 PM/AM) (Diurnal Differences, Regional Sampling, and Collocations with in-situ Measurements with a Stipulated Time and Distance criteria (+/- 3 HRs, 100 Km Radius)

	AIRS	IASI
CC	58	69
T	103	152
Q	41	87
O3	41	53
CO	36	33
CH4	59	59
CO2	70	79
HNO3	14	14
N2O	58	58
SO2	60	60



# IASI and AIRS RAOB Matches


## Data Used : 01/20/2008 – 08/02/2008




**IASI- Accepted Matches (After RAOB Selection):12566**  
 (NH:77%, SH:23%); (LAND:78%, Sea:22%); (Day:60%,Night:40%)  
**Tropics:28%; Mid-Lat:64% Polar:8%**

**AIRS- Accepted Matches (After RAOB Selection):5993**  
 (NH:79%,SH:21%); (LAND:91%, Sea:9%); (Day:46% Night:54%)  
**Tropics:10%; Mid-Lat:68% Polar:22%**

**Long-term AIRS Yield (2002-2007) was a little higher than 50%. Recently we started seeing lower yield in the year 2008**



**Land D N ALL Sea D N ALL AD AN AA**  
 Legend: TROP\_ACCEP (purple), MIDLAT\_ACCEP (orange), POLAR\_ACCEP (blue)





**Land D N ALL Sea D N ALL AD AN AA**  
 Legend: TROP\_ACCEP (purple), MIDLAT\_ACCEP (orange), POLAR\_ACCEP (blue)




# IASI- T(p), q(p) RMS Difference

Global (L+S+Coast) NSAMP=12,666 Yield: 55%  
Acceptance Criteria: Mid-Troposphere Temp Flag = 0





**RAOB vs. IASI**   **AVN**   **ATOVS**   **ECMWF**   **FG**



# AIRS- T(p), q(p) RMS Difference

Global (L+S+Coast) NSAMP=5,993, Yield: 36%  
Acceptance Criteria: Mid-Troposphere Temp Flag = 0




—□— AIRS\_PR    —○— NCEP\_GFS    - -◇- N18\_ATOVS  
—×— AIRS\_FG    —△— ECMWF

—□— AIRS\_PR    —○— NCEP\_GFS    - -◇- N18\_ATOVS  
—×— AIRS\_FG    —△— ECMWF




**RAOB vs. AIRS**    **AVN**    **ATOVS**    **ECMWF**    **FG**  
**Dotted Lines : AIRS**



# IASI & AIRS Physical Ret vs. First Guess T(p), q(p) RMS Difference (Global)



•AIRS has difficulty close to the surface with –rej3 QA. The V5 QA improves the Statistics for the lowest layers, but the yield reduces.





**Solid lines : IASI, Dotted Lines : AIRS**

**RAOB vs. IASI\_PR AIRS\_PR IASI\_FG AIRS\_FG**

# IASI & AIRS T(p), q(p) RMS Difference Physical Retrievals: *LAND SEA ALL*



MetOp IASI T(p),  
q(p) as of Today



Aqua-AIRS V5: T(p),  
q(p) with -Rej3 Option



N-IASI(ALL)=12,666, N-AIRS(ALL)=5,993;  
N-IASI(Sea) 2,848, N-AIRS (Sea):540

# Experiments: Noise Amplification in Cloud Cleared Radiance




(<http://eol.jsc.nasa.gov>). STS104-724-50 on right (July 20, 2001).

**Cloud Amounts (Day time)**  
AIRS (1:30 pm) and IASI (9:30 am)



**MODIS Aqua-Land**

**MODIS Terra-Land**




	<b>AIRS</b>	<b>IASI</b>
<b>FOR (50Km)</b>	<b>9 FOVs</b>	<b>4 FOVs</b>
<b>Channels Used for CC</b>	<b>58 (LW, Window)</b>	<b>69 (LW,SW Window)</b>
<b>Ampl Factor</b>	<b>1/3 = A = 10</b>	<b>1/2 = A = 10</b>
<b>Error in <math>\hat{\eta}</math></b>	<b>Probably Lower</b>	<b>Probably Higher</b>
<b>Cloud Contrast</b>	<b>Probably Higher</b>	<b>Probably Lower</b>


**MODIS Derived Time Series of Cloud Fraction during the Daytime (M. D. King, S. Platnick et al. – NASA GSFC)**



**IASI Retrievals (4 FOVs)**



**AIRS Retrievals (9 FOVs)**



**AIRS Like IASI 4 FOVs 2,4,6,8**

**THREE Experiments**

$$R_1(n) = (1 - \alpha_1) \cdot R_{clr}(n) + \alpha_1 \cdot R_{cld}(n)$$


$$R_2(n) = (1 - \alpha_2) \cdot R_{clr}(n) + \alpha_2 \cdot R_{cld}(n)$$

$$\eta = \alpha_1 / (\alpha_2 - \alpha_1)$$


$$R_{CCR}(n) = \overline{R_j(n)} + \sum_{j=1} \eta_j \cdot (\overline{R_j(n)} - R_j)$$


$$A = \sqrt{1/N_f(1 + \sum \hat{\eta}^2) + \sum \hat{\eta}^2}$$

# Noise Amplification in Cloud Clearing Procedure Global Samples AIRS (9 FOVs)




# Noise Amplification in Cloud Clearing Procedure Global Samples IASI (4 FOVs)








# AIRS and IASI Accepted Samples Global




— □ AIRS\_G-ALL-ALL\_AC — ■ IASI\_G-ALL-ALL\_AC — □ AIRS\_G-ALL-ALL\_AC — ■ IASI\_G-ALL-ALL\_AC




# AIRS (FOVs 2, 4, 6, 8) like IASI and IASI Polar Samples






# AIRS (FOVs 2, 4, 6, 8) like IASI and IASI Polar Samples




■ AIRS\_P-ALL-NIGHT\_AC 
 ■ IASI\_P-ALL-NIGHT\_AC 
 ■ AIRS\_P-ALL-NIGHT\_AC 
 ■ IASI\_P-ALL-NIGHT\_AC



# IASI and AIRS Retrieval Stats Using A Common Data Set



Matched Locations of MetOp-IASI, Aqua-AIRS with RAOBs




## Experiments and Retrieval Statistics with IASI and AIRS Matches at the Same Locations

(1) IASI Retrievals



(2) AIRS Retrievals

(3) AIRS Like IASI Retrievals (AIRS FOVs:2,4,6,8)




# IASI & AIRS and AIRS Like IASI with -Rej3 T(p), q(p) RMS Difference

Common Data Set Yield : 55%, 36%, 32%



IASI\_PR AIRS\_PR\_4FOV\_WGT AIRS\_PR IASI\_PR AIRS\_PR\_4FOV\_WGT AIRS\_PR


IASI, AIRS with -Rej3 Option, AIRS with 4 FOVs -Rej3  
IASI & AIRS Collocated to the same Ground Location




# IASI & AIRS with -Rej3 and V50 QA

## T(p), q(p) RMS Difference

Common Data Set Yield: 55%, 36%, 20%



IASI\_PR AIRS\_PR AIRS\_PR50



IASI\_PR AIRS\_PR50 AIRS\_PR

IASI, AIRS with -Rej3 Option, AIRS with V5 QA  
IASI & AIRS Collocated to the same Ground Location

# Questions Addressed

## IASI and AIRS - With Global RAOBs



- Whether IASI Retrievals are Comparable with the AIRS Retrievals In terms of
  - Accuracy – **YES**
  - Yield – **YES - Higher than AIRS Yield (Currently)**
  - Global Temperature Retrieval: IASI and AIRS are comparable.
    - Over Oceans: AIRS Mid-Troposphere Temperatures are slightly better
  - Global Water Vapor Retrieval: IASI is slightly better
    - Over Oceans: AIRS and IASI are comparable, with AIRS showing better upper level water vapor RMS
  - **IASI Retrievals Need to Improve Over the Polar Regions both in terms of yield and RMS**
- How much improvement is seen with the IASI (and AIRS) retrievals compared to the ATOVS
  - At Least half a Degree RMS improvement over M2-ATOVS Retrievals
  - Substantial Improvement over M2-ATOVS water vapor retrievals
- In the presence of a Hyper-spectral Sounder like the AIRS, the AMSU-B adds little or NO Improvement to the Retrieval of Water Vapor.

# Summary- Cloud-Clearing



- AIRS retrievals appear to enjoy some of the advantages with 9FOV cloud-clearing compared to IASI 4 FOV cloud clearing. Noise amplification plots reveal better distribution of total and accepted samples that eventually improves cloud-cleared radiances despite the fact that there could be more clouds in the afternoon over land.
- AIRS like IASI experiment (4 FOV AIRS) with real data shows a slight advantage in cloud-clearing compared to IASI retrievals. This could be due to the geometry of the AIRS FOVs (Overlapping and spread-out) vs. IASI FOVs (Circular and Closer) where AIRS might be having a better S/N compared IASI.

# Summary-Future Work



- **IASI Retrieval Algorithm - Optimization**
  - » **Cloud-Clearing**
    - 4 FOVs – Show Inability to identify uniform scenes and tendency of Confusion with Overcast (Especially over Polar Regions) – Better QA, and Residual Tests
    - Cloud Clearing /Channel Optimization (Use of shortwave channels in cloud-clearing (YES/NO ?)
    - Clear Flag Detection
  - » **Retrieval Convergence Criteria, Residual Tests, Rejection Criteria**
  - » **FG improvement with more Training Data**
- **IASI global retrievals for the region 60S – 60 N are quite comparable to AIRS with a relatively larger yield than the yield currently seen with the AIRS.**
- **With improvements expected in the FG IASI training samples, and expected optimization in cloud-clearing, IASI Retrievals are bound to improve.**
- **However even with the current Algorithm, the results are quite comparable. In next 3 months with all the expected changes we are planning, IASI Instrument and the Algorithm shows lot of potential.**

# Web Interface for IASI/AIRS Validations

## You Name it – You Have it



NOAA/NESDIS/ORA/SPB/IOSSPDT Home Page --- MAR 27, 2008 - 9:33:43 AM - Mozilla Firefox

http://www.orbit2.nesdis.noaa.gov/smcd/spb/iOSSPDT/iOSSPDT.php#1

NOAA/NESDIS/ORA/SPB/IOSSPDT Home Page

HOME AIRS MODIS IASI CHS AMSU

IASI HOME  
Global Radiance  
PCS  
Channel Monitoring  
Near Real Time (demo)  
NOAA Unique Products  
Validation  
System Updates  
Spectral Ranges

Yearly Monthly Weekly Daily WholePeriod

ReadMe RAOB Locations Map

RAOB Validation ● Data Exist. ● No Data.

Period	CCR ALL		CCR OC DAY&NGT		CCR OC NGT		CLR ALL		CLR OC		
	Bias	RMS	Bias	RMS	Bias	RMS	Bias	RMS	Bias	RMS	
Region: *GLOB NH SH TROP MLAT POLAR											
11202007_03032008	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	<span style="color: green;">●</span>	


Please refer comments or questions to the [webmaster](#). This page was last modified on: 03/27/2008 09:33:13

Welcome to the NOAA/NESDIS/ORA/SPB/IOSSPDT Home Page - Mozilla Firefox

http://www.orbit2.nesdis.noaa.gov/smcd/spb/iOSSPDT/valid\_all.php?imggh=/(asi)/ifs/validation/Monthly/2008/&dfpr=Jan2008

Available Files and Descriptions

File#	RMS	Cloud_Cleared	Ocean_Day	RAOB vs SET_Final	RAOB vs ATOVS	RAOB vs NCEP_GFS	RAOB vs BCMWF	RAOB vs RET_F0
File#1	RMS	Cloud_Cleared	Ocean_Day	RAOB vs SET_Final	RAOB vs ATOVS	RAOB vs NCEP_GFS	RAOB vs BCMWF	RAOB vs RET_F0
File#2	Bias	Cloud_Cleared	Ocean_Day	RAOB vs SET_Final	RAOB vs ATOVS	RAOB vs NCEP_GFS	RAOB vs BCMWF	RAOB vs RET_F0
File#3	RMS	Cloud_Cleared	Ocean_Day	RAOB vs SET_Final	ECMWF vs RET_Final	ECMWF vs SET_Final	RAOB vs BCMWF	RAOB vs RET_F0
File#4	Bias	Cloud_Cleared	Ocean_Day	RAOB vs SET_Final	ECMWF vs RET_Final	ECMWF vs SET_Final	RAOB vs BCMWF	RAOB vs RET_F0
File#5	RMS	Clear_Case	Ocean_Day	RAOB vs SET_Final	RAOB vs BCMWF	RAOB vs ATOVS	ECMWF vs RET_Final	ECMWF vs RET_Final
File#6	Bias	Clear_Case	Ocean_Day	RAOB vs SET_Final	RAOB vs BCMWF	RAOB vs ATOVS	ECMWF vs RET_Final	ECMWF vs RET_Final
File#7	RMS	Clear_Case	Ocean_Day	RAOB vs SET_Final	RAOB vs RET_F0	RAOB vs RET_F0	ECMWF vs RET_Final	ECMWF vs RET_Final
File#8	Bias	Clear_Case	Ocean_Day	RAOB vs SET_Final	RAOB vs RET_F0	RAOB vs RET_F0	ECMWF vs RET_Final	ECMWF vs RET_Final
File#9	RMS	Cloud_Cleared	Land_Day	RAOB vs SET_Final	RAOB vs ATOVS	RAOB vs NCEP_GFS	RAOB vs BCMWF	RAOB vs RET_F0
File#10	Bias	Cloud_Cleared	Land_Day	RAOB vs SET_Final	RAOB vs ATOVS	RAOB vs NCEP_GFS	RAOB vs BCMWF	RAOB vs RET_F0
File#11	RMS	Cloud_Cleared	Land_Day	RAOB vs SET_Final	ECMWF vs RET_Final	ECMWF vs SET_Final	RAOB vs BCMWF	RAOB vs RET_F0
File#12	Bias	Cloud_Cleared	Land_Day	RAOB vs SET_Final	ECMWF vs RET_Final	ECMWF vs SET_Final	RAOB vs BCMWF	RAOB vs RET_F0
File#13	RMS	Clear_Case	Land_Day	RAOB vs SET_Final	RAOB vs BCMWF	RAOB vs ATOVS	ECMWF vs RET_Final	ECMWF vs RET_Final



<http://www.orbit2.nesdis.noaa.gov/smcd/spb/iOSSPDT/iOSSPDT.php#1>



**Thank You for your Attention**

**Backup Slides**



# Earlier Papers Using the Results from the NOAA/NESDIS IASI/AIRS Validation System



## • AIRS-V4 T(p), q(p) Validations, JGR

Validation of Atmospheric Infrared Sounder temperature and water vapor retrievals with matched radiosonde measurements and forecasts

Murty G. Divakarla,<sup>1</sup> Chris D. Barnet,<sup>2</sup> Mitchell D. Goldberg,<sup>2</sup> Larry M. McMillin,<sup>2</sup> Eric Maddy,<sup>3</sup> Walter Wolf,<sup>3</sup> Lihang Zhou,<sup>3</sup> and Xingpin Liu<sup>3</sup>

Received 21 April 2005; revised 3 November 2005; accepted 23 November 2005; published 6 April 2006.

<http://www.agu.org/journals/jd/jd0607/2005JD006116/2005JD006116.pdf>

## • AIRS-V4, V5 O3(p), Total Ozone Validations, JGR

Evaluation of Atmospheric Infrared Sounder ozone profiles and total ozone retrievals with matched ozonesonde measurements, ECMWF ozone data, and Ozone Monitoring Instrument retrievals

Murty Divakarla,<sup>1</sup> Christopher Barnet,<sup>2</sup> Mitchell Goldberg,<sup>2</sup> Eric Maddy,<sup>3</sup> Fredrick Irion,<sup>4</sup> Mike Newchurch,<sup>5</sup> Xingpin Liu,<sup>3</sup> Walter Wolf,<sup>2</sup> Lawrence Flynn,<sup>2</sup> Gordon Labow,<sup>6</sup> Xiaozhen Xiong,<sup>3</sup> Jennifer Wei,<sup>3</sup> and Lihang Zhou<sup>3</sup>


Received 23 August 2007; revised 12 May 2008; accepted 29 May 2008; published 13 August 2008.

<http://www.agu.org/journals/jd/jd0815/2007JD009317/>

## • AIRS



Improving Weather Forecasting and Providing New Data on Greenhouse Gases

BY MOUSTAFA T. CHAHINE, THOMAS S. PAGANO, HARTMUT H. AUMANN, ROBERT ATLAS,\* CHRISTOPHER BARNET, JOHN BLAISDELL, LUKE CHEN, MURTY DIVAKARLA, ERIC J. FETZER, MITCH GOLDBERG, CATHERINE GAUTIER, STEPHANIE GRANGER, SCOTT HANNON, FREDRICK W. IRION, RAMESH KAKAR, EUGENIA KALNAY, BJORN H. LAMBRIGTSEN, SUNG-YUNG LEE, JOHN LE MARSHALL, W. WALLACE MCMILLAN, LARRY MCMILLIN, EDWARD T. OLSEN, HENRY REVERCOMB, PHILIP ROSENKRANZ, WILLIAM L. SMITH, DAVID STAELIN, L. LARRABEE STROW, JOEL SUSSKIND, DAVID TOBIN, WALTER WOLF, AND LIHANG ZHOU, BAMS, July 2006








# IASI- T(p), q(p) RMS Difference Sea Samples NSAMP=2,848




**RAOB vs. IASI** **AVN** **ATOVS** **ECMWF** **FG**



# AIRS - T(p), q(p) RMS Difference Sea Samples NSAMP=540




**RAOB vs. AIRS**    **AVN**    **ATOVS**    **ECMWF**    **FG**





# AIRS (9FOVs), AIRS (4FOVs) Like IASI and IASI




**TOTSAMP AIRS, IASI**



**TOTSAMP  
AIRS Like IASI, IASI**



**Accepted Sample  
AIRS, IASI**



**Accepted Sample  
AIRS Like IASI, IASI**



# Datasets and Evaluation

- **Bias and RMS difference Statistics with Reference to RAOBs.**
  - » **Statistics for MetOp-IASI (9:30AM/PM)**
    - **Acceptance Criteria: Mid-Troposphere Temp Flag = 0**
    - **Solid Lines for IASI (9:30 AM/PM)**
      - **RAOB vs. IASI Final - Physical Retrievals**
      - **RAOB vs. IASI Fast Regression (FG)**
      - **RAOB vs. MetOp-ATOVS Retrievals**
      - **RAOB vs. ECMWF Forecast**
      - **RAOB vs. NCEP-GFS Analysis/Forecast Fields**
  - » **Similar Statistics for Aqua-AIRS (1:30 PM/AM)**
    - **Dotted Lines**
      - **Similar Color Convention**
    - **Acceptance Criteria: Mid-Troposphere Temp Flag = 0**
    - **(I do have comparison with V5 QA also)**
  - » **These Figures are only a few selected ones. Go to our website to see RMS and Bias Plots for**
    - **Any region (tropics, midlat, polar, global)**
    - **Any category: land/sea; day/night etc.**

<http://www.orbit2.nesdis.noaa.gov/smcd/spb/iosspdt/iosspdt.php#1>




# Datasets and Evaluation

- **Bias and RMS difference Statistics with Reference to RAOBs.**
  - » **Statistics for MetOp-IASI (9:30AM/PM)**
    - **Acceptance Criteria: Mid-Troposphere Temp Flag = 0**
    - **Solid Lines for IASI (9:30 AM/PM)**
      - **RAOB vs. IASI Final - Physical Retrievals**
      - **RAOB vs. IASI Fast Regression (FG)**
      - **RAOB vs. MetOp-ATOVS Retrievals**
      - **RAOB vs. ECMWF Forecast**
      - **RAOB vs. NCEP-GFS Analysis/Forecast Fields**
  - » **Similar Statistics for Aqua-AIRS (1:30 PM/AM)**
    - **Dotted Lines**
      - **Similar Color Convention**
    - **Acceptance Criteria: Mid-Troposphere Temp Flag = 0**
    - **(I do have comparison with V5 QA also)**
  - » **These Figures are only a few selected ones. Go to our website to see RMS and Bias Plots for**
    - **Any region (tropics, midlat, polar, global)**
    - **Any category: land/sea; day/night etc.**

<http://www.orbit2.nesdis.noaa.gov/smcd/spb/iosspdt/iosspdt.php#1>

# Spectral Coverage of Thermal Sounders (Example Radiances: AIRS, IASI, & CrIS)



CO<sub>2</sub>

O<sub>3</sub>

CH<sub>4</sub>



CO CO<sub>2</sub>

# IASI & AIRS T(p), q(p) RMS Difference

## LAND SEA ALL




MetOp IASI T(p),  
q(p) as of Today




Aqua-AIRS V5: T(p),  
q(p) with -Rej3 Option






# IASI & AIRS with -Rej3 and V50 QA Flags T(p), q(p) Bias (Common Data Set)




- - AIRS\_PR50    - IASI\_PR    - AIRS\_PR






- IASI\_PR    - - AIRS\_PR50    - AIRS\_PR

IASI, AIRS with -Rej3 Option, AIRS with V5 QA  
IASI & AIRS Collocated to the same Ground Location




# IASI & AIRS and AIRS Like IASI with -Rej3 T(p), q(p) Bias Common Data Set




IASI\_PR AIRS\_PR\_4FOV AIRS\_PR PR IASI\_PR AIRS\_PR\_4FOV AIRS\_PR PR

IASI, AIRS with -Rej3 Option, AIRS with 4 FOVs -Rej3  
IASI & AIRS Collocated to the same Ground Location

# Yield.




# Yield.




# Cloud-Clearing Aspects IASI 4 FOVs and AIRS 9 FOVs





- Noise Amplification in the Cloud Cleared Radiance





~ -0.15 K fringes in the shortwave end:  
RTA problems?  
Instrument problems?  
**(Antonia)**



# Statistics AIRS QA -rej3 option (Whole Profile is Accepted/Rejected)



Temperature			Water Vapor			Ozone		
Layer	Upper Prs (hPa)	Lower Prs (hPa)	Layer	Upper Prs (hPa)	Lower Prs (hPa)	Layer	Upper Prs (hPa)	Lower Prs (hPa)
15	103	126	1	0	201	1	0	2
16	126	142	2	201	314	2	2	4
17	142	160	3	314	407	3	4	8
18	160	190	4	407	516	4	8	16
19	190	223	5	516	618	5	16	32
20	223	273	6	618	707	6	32	66
21	273	314	7	707	853	7	66	126
22	314	344	8	853	1100	8	126	223
23	344	407				9	223	359
24	407	478				10	359	535
25	478	535				11	535	754
26	535	618				12	754	1100
27	618	684						
28	684	778						
29	778	879						
30	879	1100						

- RAOB/O3SND is taken as Reference.
- Temperature Stats for 1 Km Layers 1000-0.1 hPa
- Water Vapor Stats 2 km layers 1000-200 hPa
- Ozone – Umkehr Layers 1000-10 hPa
- %RMS is computed by Weighting RMS difference with Truth in the layer.
- %Bias is computed by (AIRS-Truth)/Truth)

# IASI and AIRS RAOB Matches p16



**AIRS- Accepted Matches (After RAOB Selection):5993**  
 (NH:79%,SH:21%); (LAND:91%, Sea:9%); (Day:46% Night:54%)  
 Tropics:10%; Mid-Lat:68% Polar:22%

**IASI- Accepted Matches (After RAOB Selection):12566**  
 (NH:77%, SH:23%); (LAND:78%, Sea:22%); (Day:60%,Night:40%)  
 Tropics:28%; Mid-Lat:64% Polar:8%



Accepted % WRT Total Accepted (N)

Accepted % WRT Total Sample

AIRS - Global Data Accepted Samples (NSAMP = 5993)  
 Percentages are With respect to the Total Number of Accepted Samples 5993

## AIRS



AIRS - Global Data Accepted Samples (NSAMP = 5993)  
 Percentages are With respect to the Total Sample Size of :15903



IASI - Global Accepted Samples (NSAMP = 12566)  
 Percentages are With respect to the Total Number of Accepted Samples 12566

## IASI

IASI - Global Data Accepted Samples (NSAMP = 12566)  
 Percentages are With respect to the Total Sample Size of :22760







# Time Series of Cloud Fraction during the Daytime (M. D. King, S. Platnick et al. - NASA GSFC)




July 2002 - January 2007



# Noise Amplification in Cloud Clearing Procedure AIRS (9 FOVs)




# Noise Amplification in Cloud Clearing Procedure IASI (4 FOVs)







# IASI Yield Compared to AIRS Long-term Yield




IASI Yield: Total, Accepted, Clear Cases




Aqua-AIRS V4 Yield: Total, Accepted, Clear Cases








IASI Yield: %Accepted, %Clear Cases



Aqua-AIRS V4 Yield: %Accepted, %Clear Cases




# Noise Amplification in Cloud Clearing Procedure AIRS (9 FOVs)




# Noise Amplification in Cloud Clearing Procedure IASI (4 FOVs)




**Tropical**




**Mid-lat**








**Polar**




**Global**



# AIRS (9 FOVs) and IASI 4 FOV Percentage of Accepted Samples for each bin wrt to Total Number of Accepted Samples






# AIRS like IASI


## Choose FOVs 2, 4, 6, 8 to Form IASI




### Tropical




### Mid-lat



### Polar



### Global









# Noise Amplification in Cloud Clearing Procedure IASI (4 FOVs)




## Tropical




## Mid-lat



## Polar



## Global





# Validation Data

## MetOp-IASI/Aqua-AIRS Match-up Database

(Adapted to MetOP-IASI from Aqua-AIRS Validation System)

- RAOB Measurements Matched to Aqua (1:30 AM/PM) and MetOP(9:30 AM/PM) Satellite Observations
- MetOp-IASI/AMSU-A/AMSU-B Level1B Radiances
- IASI Level-2 Retrievals
- Aqua-AIRS/AMSU-A Level1B Radiances
- AIRS Level-2 Retrievals
- NCEP-GFS (AVN) Level2-Forecast/Analysis
- ECMWF Level-2 Forecast/Analysis
- MetOp-ATOVS/NOAA-18 Level-2 Retrievals


Collocated Within  $\pm 3$  Hrs. & 100 Km Radius

Data Used In this Study : November, 2007-2008

Source of RAOB Data : OSDPD MDB



# Validation Data - Algorithms



## MetOp-IASI/Aqua-AIRS Match-up Database




(Adapted to MetOP-IASI from Aqua-AIRS Validation System)

- RAOB Measurements Matched to Aqua (1:30 AM/PM) and MetOP(9:30 AM/PM) Satellite Observations
- MetOp-IASI/AMSU-A/AMSU-B Level1B Radiances
- IASI Level-2 Retrievals
- Aqua-AIRS/AMSU-A Level1B Radiances
- AIRS Level-2 Retrievals
- NCEP-GFS (AVN) Level2-Forecast/Analysis
- ECMWF Level-2 Forecast/Analysis
- MetOp-ATOVS/NOAA-18 Level-2 Retrievals

Collocated Within  $\pm 3$  Hrs. & 100 Km Radius  
Data Used In this Study : November, 2007-2008

- Emulate IASI/AIRS Retrieval Algorithms/Versions
- Reprocessing Options with Algorithm Upgrades, New Data
- Testing New Ideas in the Retrieval Algorithm

# IASI & AIRS Physical Ret vs. First Guess T(p), q(p) RMS Difference (Global)



**Solid lines : IASI, Dotted Lines : AIRS**

**RAOB vs. IASI\_PR   AIRS\_PR   IASI\_FG   AIRS\_FG**