




Exploiting high-spectral resolution satellite data for studies of volcanic emissions: AIRS and IASI retrievals of SO₂ and fine ash particles

Dr Fred Prata
Department of Atmosphere and Climate
Norsk Institutt for Luftforskning (NILU)
Kjeller, Norway







15 October 2008

AIRS Science Team Meeting, Greenbelt

4







15 October 2008

AIRS Science Team Meeting, Greenbelt


7





Aura/OMI - 04/23/2006 02:20-05:38 UT

SO₂ mass: 21.646 kt; Area: 740221 km²; SO₂ max: 7.36 DU at lon: 145.21 lat: -3.41





Frequency of Eruptions

Mt St Helens (5) 1 in 10 years


Krakatau (6) 1 in 100 years

Tambora (7) 1 in 1000 years


VEI ~ 2 1 every 2 weeks

VEI > 7 (~100,000 years - Toba)

Volcanoes and Climate




Cooling Effects $\sim 18 \text{ Tg}$



Aviation Hazards




- Plugged cooling holes
- Erosion of edges
- Ash build-up





Eruptions Affecting North Pacific Air Routes During the 1990's





Growth in aviation traffic


Region	Annual average growth rate of passengers (percentage)	
	2001-2005	
Europe	2.3	4.5
Africa	3.3	5.0
Middle East	6.4	4.0
Asia and Pacific	4.1	7.5
North America	1.8	5.0
Latin America and Caribbean	2.9	4.5
World	2.7	6.0

Freight & mail
(% change in ton-km)



Source: International Civil Aviation Organisation

Global aviation threat





SO₂ measurements from satellites

INSTRUMENT	From
• TOMS	1979
• TOVS	1979
• MLS	1991
• MODIS	1999
• ASTER	1999
• <i>ACE</i>	<i>2003</i>
• <i>TES</i>	<i>2004</i>
• SCIAMACHY/GOME	1998
• AIRS	2002
• OMI	2004
• SEVIRI	2005
• IASI	2005

Measuring SO₂ from satellites

JOURNAL OF GEOPHYSICAL RESEARCH, VOL. 112, D20204, doi:10.1029/2006JD007955, 2007

Click Here for Full Article


Retrieval of volcanic SO₂ column abundance from Atmospheric Infrared Sounder data

A. J. Prata¹ and C. Bernardo²

Received 23 August 2006; revised 25 June 2007; accepted 16 July 2007; published 19 October 2007.


¹Norwegian Institute for Air Research, Kjeller, Norw

²Auspace Ltd., Mitchell, ACT, Australia.




Spectral matching

- Many molecules have distinctive absorption bands



- Using theoretically-derived spectral shapes, molecules can be identified by comparing (or matching) the observed shape with theory.





Spectral matching (1)

Radiative transfer equation

$$I_\nu = I_{\nu,s} + \int_0^\infty B_\nu[T(z)] \left(\frac{\partial \tau_\nu[z, q_1(z), q_2(z) \dots q_n(z)]}{\partial z} \right) dz,$$


Choose absorption region and split integral into 3 parts:

$$\begin{aligned} I_\nu &\approx \int_0^{z_1} B_\nu[T(z)] \left(\frac{\partial \tau_\nu[z, q_2(z) \dots q_n(z)]}{\partial z} \right) dz && \text{Below gas layer} \\ &+ \int_{z_1}^{z_2} B_\nu[T(z)] \left(\frac{\partial \tau_\nu[z, q_1(z), q_2(z) \dots q_n(z)]}{\partial z} \right) dz && \text{In gas layer} \\ &+ \int_{z_2}^\infty B_\nu[T(z)] \left(\frac{\partial \tau_\nu[z, q_2(z) \dots q_n(z)]}{\partial z} \right) dz. && \text{Above gas layer} \end{aligned}$$

Spectral matching (2)

$$I'_\nu = I_{\nu,0} \exp \left\{ - \int_0^\infty k_\nu(z) q(z) dz \right\} \quad \text{Beer-Bougier-Lambert law}$$

$$A_\nu = - \ln \left\{ \frac{I'_\nu}{I_{o,\nu}} \right\} = \int_0^\infty k_\nu(z) q(z) dz. \quad \text{Absorbance spectrum}$$





Spectral matching (3)


Problem: How do you find the “best” reference pixel?

Solution: Search all pixels for best spectral match

$$R = \frac{\frac{1}{n-1} \sum_{i=0}^{n-1} \tilde{A}_i \tilde{S}_i}{\sqrt{\frac{1}{n-1} \sum_{i=0}^{n-1} \tilde{A}_i^2} \sqrt{\frac{1}{n-1} \sum_{i=0}^{n-1} \tilde{S}_i^2}}$$

$$\tilde{A}_i = \frac{A_i - \min(A_i)}{\max(A_i) - \min(A_i)},$$

$$\tilde{S}_i = \frac{S_i - \min(S_i)}{\max(S_i) - \min(S_i)}$$





Single Lorentz line

Retrieval

$$A\Delta\nu = \int_{\Delta\nu} \left\{ 1 - \exp\left[-\frac{1}{\pi} \int_{z_1}^{z_2} \frac{\alpha q(z)}{(\nu - \nu_0)^2 + \alpha^2} dz\right] \right\} d\nu,$$

$$A\Delta\nu = 2\pi\alpha\psi [L_0(\psi) + L_1(\psi)] \exp\{-\psi\},$$

$$\psi = \frac{Su}{2\pi\alpha},$$


For a band with temperature dependent line strengths and line broadening, an “off-line” radiative transfer model is needed

$$\vec{x} = (\mathbf{K}^T \mathbf{K})^{-1} \mathbf{K}^T \vec{y}. \quad \text{Least squares solution}$$



\vec{x} : layer absorber amounts (N layers)

\vec{y} : observed absorbances (M wavenumbers)

\mathbf{K} : is an MxN matrix of absorption cross sections






Validation




Carn et al., GRL, 32, L02301, doi:10.1029/2004GL021034, 2005




Manam Eruption Jan 28 2005









$$SNR = \frac{\sqrt{\frac{1}{\Delta\nu} \int_{\Delta\nu} \Delta S_{\nu}^2 d\nu}}{NE\Delta I}$$


$$\Delta S_{\nu} = |I_{\nu,z'} - I_{\nu,0}|,$$










Kasatochi 08 August



Other spectral regions





H₂S: 1332–1344 cm⁻¹




Al Mishraq Sulphur fire

NOAA HYSPLIT MODEL
Forward trajectories starting at 00 UTC 29 Jun 03
FNL Meteorological Data



AIRS 4 μm


AIRS 4 μm Retrieval: Date: 2007.04.06 04:23:25.000UTC Granule:2007.04.06.044



AIRS Lower Tropospheric SO₂

Produced by: Dr Fred Prata, NILU, [fred.prata@nilu.no]

METOP GOME2 SO₂: 2007/04/06



Anthropogenic SO₂ emissions



IASI

Atmos. Chem. Phys. Discuss., 8, 16917–16949, 2008
www.atmos-chem-phys-discuss.net/8/16917/2008/
© Author(s) 2008. This work is distributed under
the Creative Commons Attribution 3.0 License.



This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

Tracking and quantifying volcanic SO₂ with IASI, the September 2007 eruption at Jebel at Tair

L. Clarisse¹, P. F. Coheur¹, A. J. Prata², D. Hurtmans¹, A. Razavi¹, T. Phulpin³, J. Hadji-Lazaro⁴, and C. Clerbaux^{4,1}








Fig. 1. Top panel. Two spectra (in brightness temperature) from 30 September 2007. The spectrum in blue is taken from inside the initial plume of the Jebel at Tair eruption, the spectrum in red is from a nearby pixel, just outside the plume. Bottom panel. Ratio of the two spectra from the top panel.



IASI uses 1407.25, 1408.75, 1371.50 and 131.75 cm^{-1}



Volcanic ash – Detection and discrimination







Prata, A. J., 1989, Infrared radiative transfer calculations for volcanic ash, Geophys. Res. Lett., 16(11), 1293-1296

$$T_4 - T_5 < 0$$

$$\Delta T = \Delta T_c [X - X^\beta],$$

$$X = 1 - \frac{\Delta T_4}{\Delta T_c},$$


$$\Delta T = T_4 - T_5, \quad \Delta T_c = T_s - T_c, \quad \Delta T_4 = T_s - T_4.$$






Operational use of the “reverse absorption” algorithm


Kasatochi eruption



AVHRR 4m5 – Left: NOAA-15 2008/08/08 16:41 UTC
AVHRR 4m5 – Right: NOAA-16 2008/08/08 17:09 UTC




Spectral signature of volcanic ash



Prata, A.J., Gangale, G., and L. Clarisse, 2008, On the spectral signature of volcanic ash, Submitted to *JGR Atmos.*

Chaitén, Chile


First rhyolitic ash eruption in the satellite era (~30 years)



S.A. Carn, J.S. Pallister, L. Lara, J. Ewert, G. Villarosa, M. Fromm, S. Watt, A.J. Prata, D.M. Pyle, T.A. Mather, N. Matthews, R.S. Martin, A. Pavez, R. Aguilera, R. Thomas, W. Rison, P. Krehbiel, J. Johnson, A. Folch, D. Basualto, T.J. Casadevall, M. Guffanti, C. Benitez, J.G. Viramonte, 2008, **The awakening of Chaitén volcano, Chile**, EOS Trans. (to appear).






Concavity index increases detection time and distance





Sensitivity to particle size and composition






2008.08.08.137 08 August



1070-1130 cm^{-1}




820-960 cm^{-1}



2008.08.09.128 09 August 2008




Mass= 0.1619 Tg Area= 345762 km² Max SO₂= 49.9DU Date: 2008.08.10 21:23:25.000UTC




AVHRR 4m5 - NOAA-18 2008/08/10 22:31 UTC

2008.08.10.214 10 August



2008.08.10.230 10 August


4 days later



2008.08.12.083 12 August



IASI – Dust detection (based on the concavity index)



August 2008 pm.

(Courtesy Levien Clarisse)



Further information

AIRS

Transport

OMI

SEVIRI

Aviation

UV imager

SO₂ Pinatubo

TOVS

- **Prata, A. J.**, and C. Bernardo, 2007, Retrieval of volcanic SO₂ column abundance from Atmospheric Infrared Sounder data, *J. Geophys. Res.*, 112, D20204, doi: 10.1029/2006JD007955.
- **Prata, A. J.**, S. A. Carn, A. Stohl, and J. Kerkmann, 2007, Long range transport and fate of a stratospheric volcanic cloud from Soufrière Hills volcano, Montserrat, *Atmos. Chem. Phys.*, 7, 5093–5103.
- Carn, S. A., Krotkov, N. A., Yang, K., Hoff, R. M., **Prata, A. J.**, Krueger, A. J., Loughlin, S. C., and P. F. Levelt, 2007, Extended observations of volcanic SO₂ and sulphate aerosol in the stratosphere, *Atmos. Chem. Phys. Discuss.*, 7, 2857–2871.
- **Prata, A. J.**, and J. Kerkmann, 2007, Simultaneous retrieval of volcanic ash and SO₂ using MSG-SEVIRI measurements, *Geophys. Res. Lett.* 34, L05813, doi:10.1029/2006GL028691.
- **Prata, A. J.**, 2008, Satellite detection of hazardous volcanic clouds and the risk to global air traffic, *Nat. Hazards*, DOI 10.1007/s11069-008-9273-z.
- Tupper, A., I. Itikarai, M. S. Richards, **A. J. Prata**, S. Carn, and D. Rosenfeld, 2007: Facing the challenges of the International Airways Volcano Watch: the 2004/05 eruptions of Manam, Papua New Guinea. *Weather and Forecasting*, Vol. 22, No. 1, 175-191.
- Bluth, G. J. S., J.M. Shannon, I.M. Watson, **A.J. Prata** and V.J. Realmuto, 2007, Development of an ultra-violet digital camera for volcanic SO₂ imaging, *J. Volcanol. Geothermal Res.*, 161, 47–56.
- Guo, S., Bluth, G. J. S., Rose, W. I., Watson, I. M., and **Prata, A. J.**, Re-evaluation of SO₂ release of the 15 June 1991 Pinatubo eruption using ultraviolet and infrared satellite sensors, *Geochem, Geophys, Geosys*, 5(4), Q04001, doi:10.1029/2003GC000654, 2004.
- **Prata, A. J.**, Rose, W. I., Self, S., and D. M. O'Brien, Global, long-term sulphur dioxide measurements from TOVS data: A new tool for studying explosive volcanism and climate, In *Volcanism and the Earth's Atmosphere* (ed. Robock and Oppenheimer), *Geophysical Monograph*. 139, 75-92, 2003.