

## Is Bone Mineral Composition Disrupted by Organochlorines in East Greenland Polar Bears (*Ursus maritimus*)?

Christian Sonne,<sup>1,2</sup> Rune Dietz,<sup>1</sup> Erik W. Born,<sup>3</sup> Frank F. Riget,<sup>1</sup> Maja Kirkegaard,<sup>1</sup> Lars Hyldstrup,<sup>4</sup> Robert J. Letcher,<sup>5</sup> and Derek C. G. Muir<sup>6</sup>

<sup>1</sup>National Environmental Research Institute, Department of Arctic Environment, Roskilde, Denmark; <sup>2</sup>Department of Basic Animal and Veterinary Sciences, Royal Veterinary and Agricultural University, Frederiksberg, Denmark; <sup>3</sup>Greenland Institute of Natural Resources, Nuuk, Greenland, Denmark; <sup>4</sup>University Hospital of Hvidovre, Hvidovre, Denmark; <sup>5</sup>Great Lakes Institute for Environmental Research, University of Windsor, Windsor, Ontario, Canada; <sup>6</sup>National Water Research Institute, Environment Canada, Burlington, Ontario, Canada

We analyzed bone mineral density (BMD) in skulls of polar bears (*Ursus maritimus*) ( $n = 139$ ) from East Greenland sampled during 1892–2002. Our primary goal was to detect possible changes in bone mineral content (osteopenia) due to elevated exposure to organochlorine [polychlorinated biphenyls (PCBs), dichlorodiphenyl trichloroethane (DDT) and its metabolites, chlordanes (CHLs), dieldrin, hexacyclohexanes, hexachlorobenzene] and polybrominated diphenyl ether (PBDE) compounds. To ensure that the BMD value in skull represented the mineral status of the skeletal system in general, we compared BMD values in femur and three lumbar vertebrae with skull in a subsample. We detected highly significant correlations between BMD in skull and femur ( $r = 0.99$ ;  $p < 0.001$ ;  $n = 13$ ) and skull and vertebrae ( $r = 0.97$ ;  $p < 0.001$ ;  $n = 8$ ). BMD in skulls sampled in the supposed pre-organochlorine/PBDE period (1892–1932) was significantly higher than that in skulls sampled in the supposed pollution period (1966–2002) for subadult females, subadult males, and adult males (all,  $p < 0.05$ ) but not adult females ( $p = 0.94$ ). We found a negative correlation between organochlorines and skull BMD for the sum of PCBs ( $\Sigma$ PCB;  $p < 0.04$ ) and  $\Sigma$ CHL ( $p < 0.03$ ) in subadults and for dieldrin ( $p < 0.002$ ) and  $\Sigma$ DDT ( $p < 0.02$ ) in adult males; indications for  $\Sigma$ PBDE in subadults were also found ( $p = 0.06$ ). In conclusion, the strong correlative relationships suggest that disruption of the bone mineral composition in East Greenland polar bears may have been caused by organochlorine exposure. **Key words:** BMD, bone mineral density, chlordanes, DDT, dieldrin, endocrine disruption, osteoporosis, PCBs, polar bear, polychlorinated biphenyls, *Ursus maritimus*. *Environ Health Perspect* 112:1711–1716 (2004). doi:10.1289/ehp.7293 available via <http://dx.doi.org/> [Online 13 September 2004]

Bone mineral composition in mammals is based on a complex set of interrelated mechanisms and is influenced by various nutritional and environmental factors (e.g., Ganong 1991; Johansson and Melhus 2001; Johansson et al. 2002; Leder et al. 2001; Michaelsson et al. 2003; Promislow et al. 2002; Sarazin et al. 2000). Furthermore, environmental stressors such as exposure to harmful chemicals, starvation, temperature extremes, and noise have been shown to disrupt bone mineral composition in laboratory mammals (Brandt and Siegel 1978; Doyle et al. 1977; Mooney et al. 1985; Nilsson 1994; Siegel and Doyle 1975a, 1975b; Siegel et al. 1977, 1992; Siegel and Mooney 1987). The pathogenesis of stress-induced bone mineral changes is an activation of the hypophyseal–adrenal/thyroid axis, leading to enhanced parathyroid and cortisol hormone secretion, increased bone resorption, and decreased bone formation (Colborn et al. 1993; Damstra et al. 2002; Feldman 1995; Ganong 1991; Selye 1973). Other hypotheses on disruption of bone mineral status include altered mitotic rates, changes in local subcellular calcium transport, and decreased protein synthesis (Siegel and Mooney 1987).

Organochlorines such as polychlorinated biphenyls (PCBs), dichlorodiphenyl trichloro-

ethane (DDT), chlordanes (CHLs), hexacyclohexanes (HCHs), dieldrin, hexachlorobenzene (HCB), polybrominated diphenyl ethers (PBDEs), and aryl hydrocarbon receptor (AhR)–active organochlorines (e.g., polychlorinated dibenzo-*p*-dioxins, dibenzofurans, and non-*ortho*-chlorine-substituted PCBs) are all lipophilic (low degradable) chemicals, pesticides, or unwanted chemical by-products (e.g., de March et al. 1998). In general, the presence of such compounds in the arctic marine environment is the result of long-range atmospheric transport from lower-latitude sources in more industrial areas of the world, where outputs and use of, for example, PCB peaked in the 1960s (de March et al. 1998). Because of their lipophilicity, organochlorines and PBDEs persist in the environment [Arctic Monitoring and Assessment Programme (AMAP) 2004; Colborn et al. 1993; Damstra et al. 2002; de March et al. 1998]. In polar bears, organochlorines are consequently transferred transplacentally from mother to fetus and via lactation, resulting in fetal and neonatal exposures that have the potential for adverse health effects, for example, on growth and development (Bernhoft et al. 1997; Birnbaum 1994; Polischuk et al. 1995, 2002).

In humans, PCB and DDT and its metabolites have been associated with low

bone mineral density (BMD) (Alveblom et al. 2003; Beard and et al. 2000; Glynn et al. 2000) through their action as exogenous agonists and antagonists to naturally endogenous hormones (Damstra et al. 2002). Various organochlorines have also been linked to periodontitis and osteoporosis in marine fish and mammal wildlife (Bengtsson et al. 1985; Bergman et al. 1992; de Guise et al. 1995; Lind et al. 2003, 2004; Mortensen et al. 1992; Schandorff 1997; Zakharov and Yablokov 1990) and in the laboratory (Fernie et al. 2003; Jamsa et al. 2001; Lind et al. 1999, 2000a, 2000b; Render et al. 2000a, 2000b, 2001; Singh et al. 2000; Valentine and Soulé 1973). In various mammalian wildlife [e.g., gray seal (*Halichoerus grypus*), ringed seal (*Phoca hispida*), harbor seal (*Phoca vitulina*), and alligator (*Alligator mississippiensis*)], osteopenia and macroscopic pathology have been examined in bone during distinct periods of exposure to anthropogenic pollutants (Bergman et al. 1992; Lind et al. 2003, 2004; Mortensen et al. 1992, Schandorff 1997; Sonne-Hansen et al. 2002; Zakharov and Yablokov 1990). The studies showed relationships between organochlorines and exostosis, periodontitis, loss of alveolar bone structures, osteoporosis, widening of the canine opening, and enlargement of the foramen mentalia.

Polar bears from East Greenland, Svalbard, and the Kara Sea carry higher loads of organochlorines than do polar bears elsewhere in the Arctic due to the different atmospheric transport routes (AMAP 2004; de March et al. 1998; Lie et al. 2003; Norstrom et al. 1998). Subsequently, the organochlorines

Address correspondence to C. Sonne, National Environmental Research Institute, Department of Arctic Environment, Frederiksborgvej 399, DK4000 Roskilde, Denmark. Telephone: 45-46-30-19-54. Fax: 45-46-30-19-14. E-mail: csh@dmu.dk

We thank J. Brønlund and local hunters for organizing sampling in East Greenland, the Zoological Museum of Copenhagen for skull maceration and preparation support, and P.M. Lind and three anonymous reviewers for their comments on the manuscript.

Financial support was provided by the Danish Cooperation for Environment in the Arctic and the Commission for Scientific Research in Greenland.

The authors declare they have no competing financial interests.

Received 28 May 2004; accepted 13 September 2004.

up-concentrate in the blubber of ringed seal (*P. hispida*) and bearded seal (*Erignathus barbatus*), which is the primary food of the polar bear (AMAP 2004; de March et al. 1998; Lie et al. 2003; Norstrom et al. 1998). Recent studies of polar bears from Svalbard have indicated that high levels of organochlorines negatively affect levels of retinol (vitamin A) and thyroid hormones (Braathen et al. 2004; Skaare et al. 2001) and possibly also negatively affect cortisols, sex steroids, and reproductive organs (female pseudohermaphrodites), although these latter mechanisms are not clearly understood (Haave et al. 2003; Oskam et al. 2003, 2004; Sonne et al., in press; Wiig et al. 1998). Other studies have associated high levels of organochlorines with low levels of IgG, suggesting possible immunotoxic effects on the IgG levels (Bernhoft et al. 2000; Lie E, Larsen HJS, Larsen S, Johansen GM, Derocher AE, Lunn NJ, et al., unpublished data). Overall, these studies support the notion that organochlorines may cause disruption and thereby potentially affect bone mineral composition.

To determine whether exposure to organochlorines and PBDEs may have adversely affected bone mineral composition in polar bears, we compared BMD in skulls of 41 individual polar bears collected in East Greenland during the supposed pre-polluted period (1892–1932) with 98 polar bear skulls collected during the supposed polluted period (1966–2002). Furthermore, we examined a subset of 58 of the individuals collected during the pollution period to determine if BMD was related to body burden of various organochlorines and PBDEs.

## Materials and Methods

**Sampling and age estimation.** We studied a total of 139 East Greenland polar bear skulls sampled between Skjoldungen at 63°15'N and Danmarks Havn at 76°30'N during 1892–2002. The age determination was carried out by counting the cementum growth layer groups (GLGs) of the lower left incisor ( $I_3$ ) after decalcification, thin sectioning (14  $\mu$ m), and staining (toluidine blue) using the method described by, for example, Hensel and Sorensen (1980) and Dietz et al. (1991). For analyses, the individuals were then categorized into adult males ( $\geq 6$  years of age), adult females ( $\geq 5$  years), and subadults (others) (e.g., Rosing-Asvid et al. 2002). Regarding skull samples from 1892–1987, the sex was available from the expedition files, and in case of absence of this information ( $n = 9$ ), the determination was based on skull morphology.


**Osteodensitometry.** X-Ray osteodensitometry was applied to detect osteopenia (osteoporosis) by use of an X-ray bone densitometer (model XR 26; Norland Corporation, Fort Atkinson, WI, USA), which determined the

BMD (calcium phosphate, hydroxyapatite) using dual X-ray absorptiometry (DXA). The skulls were scanned in “research” mode (speed, 60 mm/sec; resolution,  $3.0 \times 3.0$  mm; width, 24.9 cm) and analyzed using XR software (revision 2.4; Norland Corporation), which generated a picture of the bone segment and calculated the BMD of hydroxyapatite in grams per square centimeter (Figure 1).

To ensure that BMD in the skull represents the mineral status of the skeletal system in general, a study was conducted where the BMD of the skull, one femur, and three lumbar vertebrae were compared in a subset of 13 free-ranging polar bears (3 subadults, 2 adult females, and 8 adult males) from Svalbard and East Greenland. The DXA scanner was calibrated daily using a phantom with known mineral density. In addition, the precision was tested by a  $10\times$  rescanning (mean  $\pm$  SD,  $521.96 \pm 0.60$  g/cm<sup>2</sup>), which from the formula  $[1 - (\text{SD}/\text{mean}) \times 100\%]$  gives a precision of 99.88%. Fragmentation and loss of tooth material caused by handling and lead shot were thought to be a problem. A double determination of the BMD in 2 skulls (numbers 5483 and 2891) with and without incisors, canines, premolars, and molars showed that loss of half or more of the material of the large canines altered the result significantly. Because the canines in the material were not fragmented to such a degree, we did not consider fragmentations a problem.

**Contaminant analyses.** Polar bear subcutaneous adipose tissue samples ( $n = 58$ ) were analyzed for PCBs, DDT and its metabolites, HCHs, CHLs, HCB, dieldrin, and PBDEs as described elsewhere (Dietz et al. 2004; Luross et al. 2002; Sandala et al. 2004). The sum of PCBs ( $\Sigma$ PCB) is the total concentrations of the 51 individual or coeluting congeners (if detected): PCBs 31/28, 52, 49, 44, 42, 64/71, 74, 70, 66/95, 60, 101/84, 99, 97, 87, 110, 151, 149, 118, 146, 153, 105, 141, 179, 138, 158, 129/178, 182/187, 183, 128, 174, 177, 171/202/156, 200, 172, 180, 170/190, 201, 203/196, 195, 194, and 206.  $\Sigma$ DDT is the sum of 4,4'-DDT, 4,4'-DDD (dichlorodiphenyldichloroethane), and 4,4'-DDE (dichlorodiphenyldichloroethylene).  $\Sigma$ HCH is the sum of the  $\alpha$ -,  $\beta$ -, and  $\gamma$ -hexachlorocyclohexane.  $\Sigma$ CHL is the total concentration of oxychlordane, *trans*-chlordane, nonachlor III (MC6), *trans*-nonachlor, *cis*-nonachlor, and heptachlor epoxide.  $\Sigma$ PBDE concentration is the total of 35 individual or coeluting congeners (if detected): PBDE numbers 10, 7, 11, 8, 12/13, 15, 30, 32, 28/33, 35, 37, 75, 71, 66, 47, 49, 77, 100, 119, 99, 116, 85, 155/126, 105, 154, 153, 140, 138, 166, 183, 181, and 190 (Muir DCG, Dietz R, Riget FF, Sonne C, Letcher RJ, Born EWB, unpublished data). All contaminant data are given in nanograms per gram lipid weight (l.w.).

**Statistics.** The BMD showed no deviation from normality (Shapiro-Wilk test), whereas contaminant data were log-transformed (base  $e$ ) before analyses in order to meet the criteria of normality and homogeneity of the variance. The significance level was set to  $p \leq 0.05$ , and a significance level of  $0.05 < p \leq 0.10$  was considered a trend. First, we tested the condylobasal skull length versus age within each group (i.e., subadults of both sexes, adult females, and adult males) in an analysis of covariance (ANCOVA) with skull length as a dependent variable, periods (before and after 1960 respectively) as class variables, age as a covariable, and their first-order interaction links (age  $\times$  period). The result from this analysis showed that the relationship of skull length versus age was the same in the two periods, which justified the use of non-length-corrected skull BMD in the further analyses (all,  $p > 0.26$ ). Second, the relationship of BMD versus age was tested by a linear regression analyses (BMD as a dependent variable and age as an independent variable) for subadults of both sexes, adult females, and adult males. To test for period differences, we used an ANCOVA with BMD as a dependent variable, age/sex (subadult females, subadult males, adult females, and adult males) and period (before and after 1960 respectively) as class variables, age as a covariable, and the first-order interaction links (age  $\times$  period, age  $\times$  age/sex, and age/sex  $\times$  period) between these




**Figure 1.** DXA scanning image of a 12-year-old female East Greenland polar bear skull sampled in 1972. Note the high-density areas of cortical bone tissue and the lower density areas of trabecular bone tissue.

variables. The model was successively reduced for nonsignificant interactions ( $p > 0.05$ ) judged from the type III sum of squares, and the significance of the remaining factors was evaluated from the final model (least square means). A temporal trend over the entire period 1892–2002 was analyzed by a multiple regression analysis with skull BMD as the dependent variable and the individual age and year of kill as explanatory variables for subadults of both sexes, adult females, and adult males, respectively (the relationship was evaluated from the parameter estimate,  $r^2$ , and  $p$ -value). The relationship between age/sex groups and contaminants was analyzed within a one-way analysis of variance on the log-transformed contaminant data, and significant results were tested by Tukey's post hoc test. The skull BMD versus contaminant ( $\Sigma$ PCB,  $\Sigma$ DDT,  $\Sigma$ CHL, HCB,  $\Sigma$ HCH, dieldrin, and  $\Sigma$ PBDE) relationships were explored by multiple regressions with skull BMD as the dependent variable and the age and contaminant concentrations as explanatory variables within age/sex groups (subadults of both sexes, adult females, and adult males). Finally, the relationship between levels of contaminants and BMD was evaluated from the parameter estimate,  $r^2$ , and  $p$ -value.

## Results

We found a highly significant correlation between BMD in skull and femur ( $r = 0.99$ ;  $p < 0.001$ ;  $n = 13$ ), and skull and vertebrae ( $r = 0.97$ ;  $p < 0.001$ ;  $n = 8$ ). These results justified the use of BMD measurements in skull to reflect the status of the skeletal system, although information on body conditions and nutritional stressors, relevant for osteoblastic and osteoclastic activity, was not available.

**Skull BMD and age/sex differences.** BMD was analyzed in 139 skulls representing the period 1892–2002, and consisted of 64 subadults, 40 adult females and 35 adult males. The BMD increased with age in subadults ( $p < 0.001$ ) but not adults (both,  $p > 0.05$ ; Figure 2). BMD differed between males and females ( $p < 0.01$ ) in the order subadult females < subadult males < adult females < adult males. Furthermore, BMD in females 14–23 years of age seemed to decline significantly with age ( $p < 0.04$ ).



**Figure 2.** BMD ( $\text{g}/\text{cm}^2$ ) in skulls from East Greenland polar bears versus individual age.

**Period differences and temporal trends in skull BMD.** Forty-one skulls were available from the supposed pre-pollution period (1892–1932) and 98 from the supposed pollution period (1966–2002) (Table 1). BMD in skulls sampled in the pollution period was significantly lower than BMD sampled in skulls from the pre-pollution period for subadults and adult males ( $p < 0.05$ ) but not for adult females ( $p > 0.9$ ) (Table 1). In addition, the multiple regression analyses of BMD versus individual age and year of kill (1892–2002) showed that BMD decreased over the entire period in adult males ( $p < 0.01$ ), and a similar trend was found for subadults ( $p = 0.07$ ) (Table 2). There was no BMD time trend for adult females ( $p > 0.5$ ).

**Skull BMD and contaminants.** The range and variation of organochlorine and PBDE contaminants (nanograms per gram l.w.) in the 58 polar bears sampled during 1999–2001 are presented in Table 3. Levels of  $\Sigma$ DDT, dieldrin,  $\Sigma$ HCH, and  $\Sigma$ PBDE were not different between subadults, adult females, and adult males (all,  $p > 0.07$ ). However, levels of  $\Sigma$ PCB were higher in adult males when compared with adult females ( $p \leq 0.05$ ). Additional information on the relationship between organochlorines and age, sex, and season in East Greenland polar bears from 1999 through 2002 has been published by Dietz et al. (2004) and Sandala et al. (2004).

BMD was found to be negatively correlated with levels of  $\Sigma$ PCB ( $p < 0.04$ ) and  $\Sigma$ CHL in subadults ( $p < 0.03$ ), whereas BMD was negatively correlated with  $\Sigma$ DDT ( $p < 0.02$ ) and dieldrin ( $p < 0.002$ ) in adult males (Table 4). In addition, a trend of  $\Sigma$ PBDE being negatively correlated to BMD in subadults was found ( $p = 0.06$ ), whereas no significant relations were found for adult females (Table 4).

**Table 1.** Skull BMD [ $\text{g}/\text{cm}^2 \pm \text{SD}$  ( $n$ )] for subadult and adult East Greenland polar bears from 1892 to 2002.

Period	Variable	Subadult females	Subadult males	Adult females	Adult males
1892–1932	BMD	1.67 $\pm$ 0.37 (7)	2.22 $\pm$ 0.19 (5)	1.99 $\pm$ 0.13 (9)	2.73 $\pm$ 0.21 (20)
	Age	2.6 $\pm$ 1.3 (7)	4.4 $\pm$ 1.3 (5)	12.7 $\pm$ 3.7 (9)	11.5 $\pm$ 4.5 (20)
1966–2002	BMD	1.55 $\pm$ 0.3* (17)	1.85 $\pm$ 0.32* (35)	1.98 $\pm$ 0.13 (31)	2.49 $\pm$ 0.24** (15)
	Age	2.8 $\pm$ 1 (17)	3.2 $\pm$ 1.1 (35)	12.1 $\pm$ 6.3 (31)	10.7 $\pm$ 5.5 (15)

Data are divided into two periods: 1892–1932 (supposed organochlorine and PBDE nonpolluted) and 1966–2002 (supposed organochlorine and PBDE polluted). BMD ( $\text{g}/\text{cm}^2$ ) was obtained by DXA scanning of the entire skull, and age (years) was obtained by counting the GLG of the lower  $I_3$  tooth.

\* $p \leq 0.05$  and \*\* $p \leq 0.01$  significantly lower during 1966–2002 compared with the 1892–1932 period for the given age/sex group.

**Table 2.** Significant results from the multiple regression analyses of skull BMD versus age and year of kill in East Greenland polar bears during 1892–2002.

Age/sex group	Equation	$r^2$	$p_{\text{age}}$	$p_{\text{yok}}$	No.
Subadults	$\text{BMD} = 0.193 \times \text{age} - 0.00254 \times \text{yok} + 6.3$	0.64	< 0.001	0.07*	64
Adult males	$\text{BMD} = 0.014 \times \text{age} - 0.00324 \times \text{yok} + 8.8$	0.31	0.2	< 0.01**	35

yok, year of kill. The equation is given as  $[\text{BMD} = A \times \text{age} + B \times \text{yok} + C]$ , with BMD ( $\text{g}/\text{cm}^2$ ) as the dependent variable and age (years) and yok (1892–2002) as the explanatory variables.  $A$ ,  $B$ , and  $C$  are specific parameter estimates;  $r^2$  is the regression coefficient of the model;  $p_{\text{age}}$  is the  $p$ -value for age; and  $p_{\text{yok}}$  is the  $p$ -value for the year of kill.

\*Nonsignificant trend of BMD decline over the entire period 1892–2002 at the  $0.05 < p \leq 0.10$  level. \*\*Significant BMD decline over the entire period 1892–2002 at the  $p \leq 0.01$  level.

## Discussion

**BMD and age/sex differences.** The high correlation in BMD between skull and femur and vertebrae, respectively, is useful because skull samples of polar bears (and other mammals) are present at national zoological museums all over the world, which makes various time-trend bone studies possible. Our results clearly show that skull BMD increased more rapidly in subadults compared with adults, in accordance with previous studies of ringed seals from Northwest Greenland (Sonne-Hansen et al. 2002). Female polar bears usually give birth to two cubs every third year (December) and mobilize and transfer large amounts of calcium and phosphate during gestation and during the postpartum (suckling) period, which lasts up to 2 years (Ramsay and Stirling 1988). In this period, calcium is used for fetal skeletal production and maintenance of the mother's and offspring's calcium phosphate homeostasis (Ramsay and Stirling 1988). Because the female polar bear mobilizes these large amounts of calcium and phosphate, adult females are expected to have a lower BMD compared with adult males. Such a difference was also found in the present study. Similar differences have been found in humans (e.g., Van Langendonck et al. 2002). As suggested for humans, the marked difference in BMD between the sexes could be the result of a higher muscle mass and strength in males, leading to higher biomechanical loading of the bone. This would lead to increased bone formation through the stimulation of the mechanotransduction system in the osteocytes (Van Langendonck et al. 2002).

Earlier studies show that sufficient levels of sex steroids (estrogens and androgens) are important in the development of the human cortical bone structures in boys, girls,

teenagers, adults, and the elderly (Hampson et al. 2002; Juul 2001; Leder et al. 2001; Szulc et al. 2001). On the other hand, high levels of estrogen-active substances (intrinsic, extrinsic) stimulate the expression of secondary sexual characteristics (Hampson et al. 2002; Juul 2001; Leder et al. 2001; Szulc et al. 2001). Therefore, growth delay and osteopenia (osteoporosis) have been associated with hypogonadism and lower estrogen levels in both sexes (Leder et al. 2001; Nelson 2003; Szulc et al. 2001). The age-related decrease in BMD in females in the present study was probably associated with a menopause phase after 15 years of age, but this requires a larger sample size (Figure 2) (Derocher and Stirling 1994).

**Period differences and temporal trends in skull BMD.** In both analyses of subadults of both sexes and adult males, the individuals from the prepollution period had a higher skull BMD compared with those from the polluted period. These results suggest that there is a linkage between decreased BMD for bears from the polluted period and exposure to environmental stressors compared with bears in the prepollution period. Two major environmental stressors could be linked to mineral loss in polar bear skulls: anthropogenic organochlorine compounds and PBDEs and/or climate oscillations (AMAP 2004; de March et al. 1998; Førlund et al. 2002). Concentrations of, for example, ΣPCB in the adipose tissue of East Greenland polar bears have, over the last four decades, reached levels that can elicit adverse biological effects on immunologic parameters and vitamin A levels, which may be linked to the present decrease in skull BMD (stress) (AMAP 2004; de March et al. 1998). However, during the same period global warming has resulted in a reduction in the ice coverage in the East Greenland area (Comiso 2002; Rothcock et al. 1999). Although population ecology has not been studied in East Greenland, the situation is probably similar for polar bears from the Hudson Bay area in Canada (Stirling et al. 1999). A reduction of the sea ice in the Hudson Bay area has reduced the bears' access to ringed seals, resulting in reduced body condition and lowered natality in the polar bears (Stirling et al. 1999).

Temporal differences with respect to potential effects of PCB and DDT exposure on periodontitis and osteoporosis in gray seal and harbor seal was investigated by Bergman et al. (1992), Mortensen et al. (1992), and Schandorff (1997). They found exostosis and periodontitis, often with substantial loss of alveolar bone in mandible and maxilla (osteoporosis). These changes could have been caused by hormonal imbalance potentially induced by PCBs and by DDT and its metabolites, with malformation of the calcium helix structures around the collagen matrix

(DeLillis 1989). These results are further supported by the investigations of Render et al. (2000a, 2000b, 2001). However, it must be noted that the ranges of ΣPCB and ΣDDT levels in the seals were orders of magnitude higher compared with levels in the present polar bears (Blomkvist et al. 1992).

Lind et al. (2003) investigated the BMD in the male gray seals ( $n = 43$ ) reported by Bergman et al. (1992). The method used was peripheral quantitative computed tomography, which made it possible to distinguish between cortical and trabecular bone in os mandibularis and os radius, respectively (DXA scanning used in the present study gives the average of trabecular and cortical bone density). Three sample groups of seals were compared: 1850–1955 (no pollution), 1965–1985 (high pollution), and 1986–1997 (fairly low pollution). They found that radius trabecular BMD was significantly higher in the fairly low pollution period (1986–1997) compared with the high pollution period (1965–1985), whereas mandible cortical BMD was significantly lower in the fairly low pollution period (1986–1997) compared with the no-pollution period (1850–1955). Our study of BMD in East Greenland polar bears supports the findings of Lind et al. (2003).

**BMD levels and contaminants.** Bone density expresses the bone mineral composition determined by the activity of osteoblastic bone formation and osteoclastic bone resorption, which is regulated by androgens and estrogens through cytokines (Manolagas and Jilka 1995; Manolagas et al. 1995). Studies on Svalbard have shown that PCBs

may negatively influence plasma cortisol, estrogen, and testosterone levels (Haave et al. 2003; Oskam et al. 2003, 2004) and plasma retinol and thyroid hormone levels in polar bears (Braathen et al. 2004; Skaare et al. 2001). These studies all indicate that organochlorines in Svalbard polar bears (and likely also East Greenland bears, because the organohalogen compound levels are comparable) potentially affect endocrine homeostasis, which again may lead to bone mineral loss (osteoporosis). Another polar bear study from Svalbard associated high levels of organochlorines with low levels of IgG, suggesting possible immunotoxic effects (Bernhoft et al. 2000; Lie E, Larsen HJS, Larsen S, Johansen GM, Derocher AE, Lunn NJ, et al., unpublished data). This potential effect may lower the immune response and enhance stress with increased cortisol levels, which potentially affects the bone mineral composition (osteoporosis).

The present study indicated that high concentrations of ΣPCB and ΣCHL are associated with reduced skull BMD in subadults and that ΣDDT and dieldrin are associated with reduced skull BMD in adult males. These BMD relationships with ΣPCB, ΣCHL, ΣDDT, and dieldrin concentrations in subadults of both sexes and adult males may suggest endocrine-related effects (e.g., AMAP 2002; Birnbaum 1994; Damstra et al. 2002; de March et al. 1998; Lind et al. 2003, 2004). For example, PCBs and DDT and its metabolites have shown *in vitro* and *in vivo* to be weak agonists/antagonists of estrogen-receptor-mediated activity; organochlorine-

**Table 3.** Concentrations [mean ± SD (median), ng/g l.w.] of various contaminants in subcutaneous adipose tissue of 58 East Greenland polar bears sampled during 1999–2001.

Compound	Subadults (n = 35)	Adult females (n = 14)	Adult males (n = 9)
ΣPCB	6,597 ± 2,726 (6,089)	5,334 ± 2,150* (5,770)	8,637 ± 4,111* (8,280)
ΣCHL	1,598 ± 884 (1,469)	1,379 ± 591 (1,353)	1,055 ± 517 (914)
ΣDDT	392 ± 209 (376)	358 ± 149 (366)	481 ± 331 (496)
ΣHCH	196 ± 68 (172)	195 ± 186 (151)	294 ± 210 (181)
Dieldrin	210 ± 100 (196)	174 ± 70 (154)	177 ± 81 (172)
HCB	99 ± 84 (70)	75 ± 82 (51)	51 ± 28 (48)
ΣPBDE	62 ± 33 (53)	53 ± 17 (53)	52 ± 16 (49)

\*Significant difference between adult females and males at the  $p \leq 0.05$  level.

**Table 4.** Significant results from the multiple regression analyses of skull BMD versus age and contaminant concentrations in East Greenland polar bears sampled during 1999–2001.

Age/sex group	Equation	$r^2$	$p_{age}$	$p_{cont}$	No.
Subadults	$BMD = 0.26 \times age - 0.25 \times [\ln(\Sigma PCB)] + 3.1$	0.59	< 0.001	< 0.04**	35
	$BMD = 0.24 \times age - 0.19 \times [\ln(\Sigma CHL)] + 2.4$	0.6	< 0.001	< 0.03**	35
	$BMD = 0.25 \times age - 0.18 \times [\ln(\Sigma PBDE)] + 1.69$	0.58	< 0.001	0.06*	35
Adult males	$BMD = 0.02 \times age - 0.17 \times [\ln(\Sigma DDT)] + 3.4$	0.69	> 0.08	< 0.02**	9
	$BMD = -0.005 \times age - 0.37 \times [\ln(\text{dieldrin})] + 4.5$	0.85	0.43	< 0.002#	9

The equation is given as  $[BMD = A \times age + B \times \ln(\text{contaminant}) + C]$ , with BMD ( $g/cm^2$ ) as the dependent variable and age (years) and log-transformed contaminant concentration  $[\ln(\text{ng/g l.w.})]$  as the explanatory variables. A, B, and C are specific parameter estimates;  $r^2$  is the regression coefficient of the model;  $p_{age}$  is the  $p$ -value for age; and  $p_{cont}$  is the  $p$ -value for contaminants.

#Nonsignificant trend of a negative correlation between BMD and  $\ln(\Sigma PBDE)$  at the  $0.05 < p \leq 0.10$  level. \*\*Significant negative correlation between BMD and organochlorine contaminant group at the  $p \leq 0.05$  level. \*Significant negative correlation between BMD and organochlorine contaminant group at the  $p \leq 0.01$  level.

mediated induction of cytochrome P450 isozyme activity can affect circulating sex hormone levels (e.g., estrogens) (Navas and Segner 1998), and this is also of relevance in the polar bear (e.g., Letcher et al. 1996). Relationships between 4,4'-DDE concentrations and BMD in humans have been reported (Beard et al. 2000; Glynn et al. 2000). Glynn et al. (2000) found significant negative correlations between 4,4'-DDE and BMD in 68 sedentary women (where concentrations are lower compared with the present polar bears) and concluded that 4,4'-DDE may also have a negative effect on BMD in men (with contaminant levels comparable with those found in the polar bears). Lind et al. (2004) investigated the relationship between DDT and its metabolites and bone composition in juvenile female American alligators (*Alligator mississippiensis*) in Lake Apopka, Florida. Compared with data from a nonpolluted reference alligator subpopulation, the tibial trabecular BMD was increased, and the authors suggested that environmental estrogenic compounds (e.g., DDT and its metabolites) disrupted the normal bone remodeling process (inhibition of osteoclast activity), which had resulted in increased BMD.

Guo et al. (1994) found that children ( $n = 25$ ) of primiparous PCB-contaminated mothers (Yu-Cheng rice oil disease) were significantly smaller and had less total lean mass and less soft tissue mass but not lower BMD compared with a control group. The PCB levels in the children (serum) were 10.3 ng/g l.w., which is lower than the levels in polar bears in the present study. Alveblom et al. (2003) investigated the incidence of osteoporotic fractures in fishermen and their wives from the Baltic Sea (high pollution) and compared these with fishermen from the west coast of Sweden (low pollution) as controls. For vertebral fractures, there was a significantly higher incidence rate ratio for east coast (Baltic) women compared with west coast women, and a similar but nonsignificant tendency was found for men. The PCB concentration (10 congeners) was 2,000 ng/g l.w. (serum), which was significantly higher compared with the west coast population but lower compared with the range in the subcutaneous adipose tissue of East Greenland polar bears. These environmental studies support the findings of negative associations between PCBs/DDT and BMD levels in East Greenland polar bears.

In the present study, we observed a negative correlation between  $\Sigma$ PBDE concentrations in adipose tissue and BMD in subadults. Disturbances in thyroid function and developmental toxicity (histopathology) have been shown to be associated with PBDEs in laboratory rats (e.g., de Wit 2002) as well as in polar bears from Svalbard (Braathen et al. 2004; Skaare et al. 2001).

## Conclusions

Skull BMD increased with age in subadults and was higher in males than in females at all ages. For adult females from 14 years of age, a menopausal BMD decrease was indicated, but further examination requires a larger sample size. BMD in skulls from subadult females, subadult males, and adult males sampled in the supposed pollution period (1966–2002) was significantly lower than BMD in skulls from the period before supposed pollution with organochlorine and PBDE compounds (1892–1932). Furthermore, correlative relationships suggest that  $\Sigma$ PCB,  $\Sigma$ CHL, dieldrin, and  $\Sigma$ DDT exposure negatively influenced BMD in skulls from subadults of both sexes and adult males.

## Correction

In the manuscript originally published online, the years 1892–1960 and 1961–2002 were used to represent the pre- and post-organochlorine/PBDE periods, respectively. These years have been changed throughout to reflect the years in which the skulls were actually collected (1892–1932 and 1966–2002).

## REFERENCES

- Alveblom AK, Rylander L, Johnell O, Hagmar L. 2003. Incidence of hospitalized osteoporotic fractures in cohorts with high dietary intake of persistent organic compounds. *Int Arch Occup Environ Health* 76:246–248.
- AMAP. 2004. Persistent Organic Pollutants in the Arctic. AMAP Assessment 2002. Oslo:Arctic Monitoring and Assessment Programme.
- Beard J, Marshall S, Jong K, Newton R, Triplett-McBride T, Humphries B, et al. 2000. 1,1,1-Trichloro-2,2-bis(*p*-chlorophenyl)-ethane (DDT) and reduced bone mineral density. *Arch Environ Health* 55(3):177–180.
- Bengtsson BE, Bengtsson A, Himberg M. 1985. Fish deformities and pollution in some Swedish waters. *Ambio* 14:32–35.
- Bergman A, Olsson M, Reiland S. 1992. Skull bone lesions in the Baltic grey seal (*Halichoerus grypus*). *Ambio* 21:517–519.
- Bernhoft A, Skaare JU, Wiig Ø, Derocher AE, Larsen HJS. 2000. Possible immunotoxic effects of organochlorines in polar bears (*Ursus maritimus*) at Svalbard. *J Toxicol Environ Health A* 57(7):561–574.
- Bernhoft A, Wiig Ø, Skaare JU. 1997. Organochlorines in polar bears (*Ursus maritimus*) at Svalbard. *Environ Pollut* 96:159–175.
- Birnbaum LS. 1994. Endocrine effects of prenatal exposure to PCBs, dioxins, and other xenobiotics: implications for policy and future research. *Environ Health Perspect* 102:676–679.
- Blomkvist G, Roos A, Jensen S, Bignert A, Olsson M. 1992. Concentrations of sDDT and PCB in seals from Swedish and Scottish waters. *Ambio* 21(8):539–545.
- Braathen M, Derocher AE, Wiig Ø, Sørmo EG, Lie E, Skaare JU, Jenssen BM. 2004. Relationships between PCBs and thyroid hormones and retinol in female and male polar bears. *Environ Health Perspect* 112:826–833.
- Brandt M, Siegel MI. 1978. The effects of stress on cortical bone thickness in rodents. *Am J Phys Anthropol* 49:31–34.
- Colborn T, Vom Saal FA, Soto AM. 1993. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspect* 101:378–384.
- Comiso JC. 2002. A rapidly declining perennial sea ice cover in the Arctic [Letter]. *Geophys Res Lett* 29:1956.
- Damstra T, Barlow S, Bergman A, Kavlock R, Kraak GVD. 2002. Global Assessment of the State-of-the-Science of Endocrine Disruptors. Geneva:World Health Organization.
- de Guise S, Lagace A, Beland P, Girard C, Higgins R. 1995. Nonneoplastic lesions in beluga whales (*Delphinapterus leucas*) and other marine mammals from the St. Lawrence estuary. *J Comp Pathol* 112(3):257–271.
- DeLillis RA. 1989. The endocrine system. In: Robbins Pathological Basis of Disease (Cotran RS, Kumar V, Robbins SL, eds). Philadelphia:W.B. Saunders Co., 1205–1276.
- de March BGE, de Wit C, Muir DCG, Braune B, Gregor DJ, Norstrom RJ. 1998. Persistent organic pollutants. In: AMAP Assessment Report: Arctic Pollution Issues. Oslo:Arctic Monitoring and Assessment Programme, 183–372.
- Derocher AE, Stirling I. 1994. Age-specific reproductive performance of female polar bears (*Ursus maritimus*). *Can J Zool* 72(4):527–536.
- de Wit CA. 2002. An overview of brominated flame retardants in the environment. *Chemosphere* 46(5):583–624.
- Dietz R, Heide-Jørgensen MP, Härkönen T, Teilmann J, Valentin N. 1991. Age determination of European harbour seal (*Phoca vitulina* L.). *Sarsia* 76:17–21.
- Dietz R, Riget FF, Sonne C, Letcher R, Born EW, Muir DCG. 2004. Seasonal and temporal trends in polychlorinated biphenyls and organochlorine pesticides in East Greenland polar bears (*Ursus maritimus*), 1990–2001. *Sci Total Environ* 331:107–124.
- Doyle WJ, Kelley C, Siegel MI. 1977. The effects of audiogenic stress on the growth of long bones in the laboratory rat (*Rattus norvegicus*). *Growth* 41:183–189.
- Feldman EC. 1995. Hyperadrenocorticism. In: Textbook of Veterinary Internal Medicine, Vol. 2 (Ettinger SJ, Feldman EC, eds). Philadelphia:W.B. Saunders Co., 1538–1578.
- Fernie K, Bortolotti G, Drouillard K, Smits J, Marchant T. 2003. Developmental toxicity of in ovo exposure to polychlorinated biphenyls: II. Effects of maternal or paternal exposure on second-generation nestling American kestrels. *Environ Toxicol Chem* 22(11):2688–2694.
- Førland EJ, Hanssen-Bauer I, Jónsson T, Kern-Hansen C, Nordli PØ, Tveito OE, et al. 2002. Twentieth-century variations in temperature and precipitation in the Nordic Arctic. *Polar Rec* 38(206):203–210.
- Ganong WF. 1991. Review of Medical Physiology. 15th ed. Norwalk, CT:Appleton & Lange.
- Glynn AW, Michaësson K, Lind PM, Wolk A, Aune M, Atuma S, et al. 2000. Organochlorines and bone mineral density in Swedish men from the general population. *Osteoporos Int* 11:1036–1042.
- Guo YL, Lin CJ, Yao WJ, Ryan JJ, Hsu CC. 1994. Musculoskeletal changes in children prenatally exposed to polychlorinated-biphenyls and related-compounds (Yu-Cheng children). *J Toxicol Environ Health* 41(1):83–93.
- Haave M, Ropstad E, Derocher AE, Lie E, Dahl E, Wiig Ø, et al. 2003. Polychlorinated biphenyls and reproductive hormones in female polar bears at Svalbard. *Environ Health Perspect* 111:431–436.
- Hampson G, Bhargava N, Cheung J, Vaja S, Seed PT, Fogelman I. 2002. Low circulating estradiol and adrenal androgens concentrations in men on glucocorticoids: a potential contributory factor in steroid-induced osteoporosis. *Metabolism* 51(11):1458–1462.
- Hensel RJ, Sorensen FE. 1980. Age determination of live polar bears. *Int Conf Bear Res Manage* 4:93–100.
- Jamsa T, Viluksela M, Tuomisto JT, Tuomisto J, Tuukkanen J. 2001. Effects of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin on bone in two rats strains with different aryl hydrocarbon receptor structures. *J Bone Miner Res* 16(10):1812–1820.
- Johansson S, Lind PM, Hakansson H, Oxlund H, Örborg J, Melhus H. 2002. Subclinical hypervitaminosis A causes fragile bones in rats. *Bone* 31(6):685–689.
- Johansson S, Melhus H. 2001. Vitamin A antagonizes calcium response to vitamin D in man. *J Bone Miner Res* 16(10):1899–1905.
- Juul A. 2001. The effects of oestrogens on linear bone growth. *Hum Reprod Update* 7(3):303–313.
- Leder BZ, Smith MR, Fallon MA, Lee MLT, Finkelstein JS. 2001. Effects of gonadal steroid suppression on skeletal sensitivity to parathyroid hormone in men. *J Clin Endocr Metab* 86(2):511–516.
- Letcher RJ, Norstrom RJ, Lin S, Ramsay MA, Bandiera SM. 1996. Immunoquantification and microosomal monooxygenase activities of hepatic cytochromes P4501A and P4502B and chlorinated hydrocarbon contaminant levels in polar bear (*Ursus maritimus*). *Toxicol Appl Pharmacol* 137:127–140.
- Lie E, Bernhoft A, Riget FF, Belikov SE, Boltunov AN, Derocher AE, et al. 2003. Geographical distribution of organochlorine pesticides (OCPs) in polar bears (*Ursus*

- maritimus*) in the Norwegian and Russian Arctic. *Sci Total Environ* 306:159–170.
- Lie E, Larsen HJS, Larsen S, Johansen GM, Derocher AE, Lunn NJ, et al. 2004. Does high organochlorine (OC) exposure impair the resistance to infection in polar bears (*Ursus maritimus*)? Part I: Effect of OCs on the humoral immunity? *J Toxicol Environ Health A*(67):555–582.
- Lind PM, Bergman A, Olsson M, Örborg J. 2003. Bone mineral density in male Baltic grey seal. *Ambio* 32(6):385–388.
- Lind PM, Eriksen EF, Sahlin L, Edlund M, Örborg J. 1999. Effects of the antiestrogenic environmental pollutants 3,3',4,4',5-pentachlorobiphenyl (PCB-126) in rat bone and uterus: diverging effects in ovariectomized and intact animals. *Toxicol Appl Pharmacol* 154:236–244.
- Lind PM, Larsson S, Johansson S, Melhus H, Wikström M, Lindhe Ö, et al. 2000b. Bone tissue composition, dimensions and strength in female rats given an increased dietary level of vitamin A or exposed to 3,3',4,4',5-pentachlorobiphenyl (PCB-126) alone or in combination with vitamin C. *Toxicology* 151:11–23.
- Lind PM, Larsson S, Öxlund H, Håkansson H, Nyberg K, Eklund T, et al. 2000a. Change of bone tissue composition and impaired bone strength in rats exposed to 3,3',4,4',5-pentachlorobiphenyl (PCB126). *Toxicology* 150:41–51.
- Lind PM, Milnes MR, Lundberg R, Bermudez D, Örborg J, Guillelte LJ. 2004. Abnormal bone composition in female juvenile American alligators from a pesticide-polluted lake (Lake Apopka, Florida). *Environ Health Perspect* 112:359–362.
- Luroso JM, Alaei M, Sergeant DB, Cannon CM, Whittle DM, Solomon KR, et al. 2002. Spatial distribution of polybrominated diphenyl ethers and polybrominated biphenyls in lake trout from the Laurentian Great Lakes. *Chemosphere* 46:665–672.
- Manolagas SC, Bellido T, Jilka RL. 1995. New insights into the cellular, biochemical and molecular basis of postmenopausal and senile osteoporosis: roles of IL-6 and gp130. *Int J Immunopharmacol* 17(2):109–116.
- Manolagas SC, Jilka RL. 1995. Bone marrow, cytokines, and bone remodeling. Emerging insights into the pathophysiology of osteoporosis. *N Engl J Med* 332(5):305–311.
- Michaelsson K, Lithell H, Vessby B, Melhus H. 2003. Serum retinol levels and the risk of fracture. *New Engl J Med* 348(4):287–294.
- Mooney MP, Siegel MI, Gest TR. 1985. Prenatal stress and increased fluctuating asymmetry in the parietal bones of neonatal rats. *Am J Phys Anthropol* 68:131–134.
- Mortensen PÅ, Bergman A, Bignert A, Hansen HJ, Härkönen T, Olsson M. 1992. Prevalence of skull lesions in harbour seals (*Phoca vitulina*) in Swedish and Danish museum collections: 1835–1988. *Ambio* 21:520–524.
- Navas JM, Segner H. 1998. Antiestrogenic activity of anthropogenic and natural chemicals. *Environ Sci Pollut Res* 5:75–82.
- Nelson HD. 2003. Postmenopausal osteoporosis and estrogen. *Am Fam Physician* 68(4):606–615.
- Nilsson JA. 1994. Energetic stress and the degree of fluctuating asymmetry—implications for a long lasting, honest signal. *Evol Ecol* 8(3):248–255.
- Norstrom RJ, Belikov S, Born EW, Garner GW, Malone B, Olpienski S, et al. 1998. Chlorinated hydrocarbon contaminants in polar bears from eastern Russia, North America, Greenland and Svalbard: biomonitoring of Arctic pollution. *Arch Environ Contam Toxicol* 35(2):354–367.
- Oskam IC, Ropstad E, Dahl E, Lie E, Derocher AE, Wiig Ø, et al. 2003. Organochlorines affect the major androgenic hormone, testosterone, in male polar bears (*Ursus maritimus*) at Svalbard. *J Toxicol Environ Health A* 66(22):2119–2139.
- Oskam IC, Ropstad E, Lie E, Derocher AE, Wiig Ø, Dahl E, et al. 2004. Organochlorines affect the steroid hormone cortisol in free-ranging polar bears (*Ursus maritimus*) at Svalbard, Norway. *J Toxicol Environ Health A* 67:959–977.
- Polischuk S, Ramsay M, Norstrom N. 2002. Body burdens and tissue concentrations of organochlorines in polar bears (*Ursus maritimus*) vary during seasonal fasts. *Environ Pollut* 118:29–39.
- Polischuk SC, Letcher RJ, Norstrom RJ, Ramsay MA. 1995. Preliminary results of fasting on the kinetics of organochlorines in polar bears (*Ursus maritimus*). *Sci Total Environ* 160/161:465–472.
- Promislow JHE, Goodman-Gruen D, Slymen DJ, Barrett-Connor E. 2002. Retinol uptake and bone mineral density in the elderly: the Rancho Bernardo Study. *J Bone Miner Res* 17(8):1349–1358.
- Ramsay MA, Stirling I. 1988. Reproductive biology and ecology of female polar bears (*Ursus maritimus*). *J Zool (Lond)* 214:601–634.
- Render JA, Aulerich RJ, Bursian SJ, Nachreiner RF. 2000a. Proliferation of maxillary and mandibular periodontal squamous cells in mink fed 3,3',4,4',5-pentachlorobiphenyl (PCB 126). *J Vet Diagn Invest* 12(5):477–479.
- Render JA, Bursian SJ, Rosenstein DS, Aulerich RJ. 2001. Squamous epithelial proliferation in the jaws of mink fed diets containing 3,3',4,4',5-pentachlorobiphenyl (PCB 126) or 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD). *Vet Hum Toxicol* 43(1):22–26.
- Render JA, Hochstein JR, Aulerich RJ, Bursian SJ. 2000b. Proliferation of periodontal squamous epithelium in mink fed 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD). *Vet Hum Toxicol* 42(2):85–86.
- Rosing-Asvid A, Born EW, Kingsley MCS. 2002. Age at sexual maturity of males and timing of the mating season of polar bears (*Ursus maritimus*) in Greenland. *Polar Biol* 25:878–883.
- Rothcock DA, Yu Y, Maykut GA. 1999. Thinning of the Arctic sea-ice cover. *Geophys Res Lett* 26(23):3469–3472.
- Sandala GM, Sonne-Hansen C, Dietz R, Muir DCG, Valters K, Bennett ER, et al. 2004. Methyl sulfone and hydroxylated PCB metabolites in adipose and whole blood of polar bear (*Ursus maritimus*) from Scoresby Sound, Greenland. *Sci Total Environ* 2004 331:125–141.
- Sarazin M, Alexandre C, Thomas T. 2000. Influence on bone metabolism of dietary trace elements, protein, fat, carbohydrates and vitamins. *Joint Bone Spine* 67(5):408–418.
- Schandorff S. 1997. Developmental stability and skull lesions in the harbour seal (*Phoca vitulina*) in the 19th and 20th centuries. *Ann Zool Fenn* 34:151–166.
- Selye H. 1973. The evolution of the stress concept. *Am Sci* 61:692–699.
- Siegel MI, Doyle WJ. 1975a. The differential effects of prenatal and postnatal audiogenic stress on fluctuating dental asymmetry. *J Exp Zool* 191:211–214.
- Siegel MI, Doyle WJ. 1975b. Stress and fluctuating limb asymmetry in various species of rodents. *Growth* 39:363–369.
- Siegel MI, Mooney MP. 1987. Perinatal stress and increased fluctuating asymmetry of dental calcium in the laboratory rat. *Am J Phys Anthropol* 73:267–270.
- Siegel MI, Mooney MP, Taylor AB. 1992. Dental and skeletal reduction as a consequence of environmental stress. *Acta Zool Fenn* 191:145–149.
- Siegel P, Siegel MI, Krimmer EC, Doyle WJ, Barry H. 1977. Fluctuating dental asymmetry as an indicator of the stressful prenatal effects of delta<sup>9</sup>-tetrahydrocannabinol in the laboratory rat. *Toxicol Appl Pharmacol* 42:339–344.
- Singh S, Casper RF, Fritz PC, Sukhu B, Ganss B, Girard B, et al. 2000. Inhibition of dioxin effects on bone formation in vitro by a newly described aryl hydrocarbon receptor antagonist, resveratrol. *J Endocrinol* 167(1):183–195.
- Skaare JU, Bernhoft A, Wiig Ø, Norum KR, Haug E, Eide DM, et al. 2001. Relationship between plasma levels of organochlorines, retinol and thyroid hormones from polar bears (*Ursus maritimus*) at Svalbard. *J Toxicol Environ Health A* 62:227–241.
- Sonne C, Leifsson PS, Dietz R, Born EW, Letcher RJ, Kirkegaard M, et al. In press. Enlarged clitoris in wild polar bears (*Ursus maritimus*) can be misdiagnosed as pseudohermaphroditism. *Sci Total Environ*.
- Sonne-Hansen C, Dietz R, Leifsson PS, Hyldstrup L, Riget FF. 2002. Cadmium toxicity to ringed seals (*Phoca hispida*)—an epidemiological study of possible cadmium induced nephropathy and osteodystrophy in ringed seals (*Phoca hispida*) from Qaanaaq in Northwest Greenland. *Sci Total Environ* 295:167–181.
- Stirling I, Lunn NJ, Iacozza J. 1999. Long-term trends in the population ecology of polar bears in western Hudson Bay in relation to climatic change. *Arctic* 52(3):294–306.
- Szulec P, Hofbauer LC, Heufelder AE, Roth S, Delmas PD. 2001. Osteoprotegerin serum levels in men: correlation with age, estrogen and testosterone status. *J Clin Endocr Metab* 86(7):3162–3165.
- Valentine DW, Soulé M. 1973. Effects of *p,p'*-DDT on developmental stability of pectoral fin rays in the grunion (*Leuresthes tenuis*). *Nat Mar Fish Serv Fish Bull* 71:921–925.
- Van Langendonck L, Claessens AL, Lefevre J, Thomis M, Philippaerts R, Delvaux K, et al. 2002. Association between bone mineral density (DXA), body structure, and body composition in middle-aged men. *Am J Hum Biol* 14(6):735–742.
- Wiig Ø, Derocher AE, Cronin MM, Skaare JU. 1998. Female pseudohermaphrodite polar bears at Svalbard. *J Wildlife Dis* 34(4):792–796.
- Zakharov MZ, Yablokov AV. 1990. Skull asymmetry in the Baltic grey seal: effects of environmental pollution. *Ambio* 19(5):266–269.