

**Table 7. Estimated numbers of deaths of persons with AIDS, by year of death and selected characteristics, 2001–2005 and cumulative—United States and dependent areas**

	Year of death					Cumulative <sup>a</sup>
	2001	2002	2003	2004	2005	
<b>Data for 50 states and the District of Columbia</b>						
<b>Age at death (yrs)</b>						
<13	47	25	23	15	7	4,865
13–14	3	9	7	14	14	271
15–19	44	38	38	38	42	1,061
20–24	206	152	163	188	157	8,555
25–29	612	555	531	509	457	43,157
30–34	1,672	1,451	1,328	1,208	1,102	94,260
35–39	3,145	2,922	2,889	2,604	2,129	114,833
40–44	3,714	3,527	3,682	3,655	3,371	101,420
45–49	3,035	3,231	3,355	3,479	3,261	69,832
50–54	2,082	2,279	2,480	2,634	2,635	41,897
55–59	1,098	1,174	1,371	1,508	1,529	23,600
60–64	624	602	732	755	805	13,497
≥65	698	675	805	845	808	13,511
<b>Race/ethnicity</b>						
White, not Hispanic	5,239	5,153	5,263	5,137	5,006	235,879
Black, not Hispanic	9,085	8,927	9,077	9,302	8,562	211,559
Hispanic	2,436	2,306	2,774	2,664	2,444	77,125
Asian/Pacific Islander	99	93	88	113	97	3,383
American Indian/Alaska Native	79	84	75	85	81	1,657
<b>Transmission category</b>						
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	5,995	5,867	6,111	6,078	5,929	260,749
Injection drug use	3,749	3,662	3,759	3,570	3,159	104,450
Male-to-male sexual contact and injection drug use	1,342	1,273	1,354	1,314	1,364	39,920
High-risk heterosexual contact <sup>b</sup>	1,485	1,434	1,554	1,729	1,584	24,655
Other <sup>c</sup>	169	163	156	136	104	9,824
Subtotal	12,740	12,400	12,934	12,826	12,140	439,598
<b>Female adult or adolescent</b>						
Injection drug use	1,829	1,876	1,916	1,959	1,651	41,529
High-risk heterosexual contact <sup>b</sup>	2,258	2,225	2,400	2,531	2,413	40,233
Other <sup>c</sup>	86	84	94	77	64	4,082
Subtotal	4,172	4,185	4,411	4,567	4,128	85,844
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	66	52	53	58	46	4,800
Other <sup>d</sup>	3	4	6	1	1	515
Subtotal	69	56	59	60	48	5,315

**Table 7. Estimated numbers of deaths of persons with AIDS, by year of death and selected characteristics, 2001–2005 and cumulative—United States and dependent areas (cont)**

	Year of death					Cumulative <sup>a</sup>
	2001	2002	2003	2004	2005	
<b>Region of residence</b>						
Northeast	5,091	5,047	5,376	4,904	3,948	174,327
Midwest	1,682	1,675	1,655	1,619	1,541	52,933
South	7,469	7,361	7,776	8,353	8,240	191,845
West	2,738	2,559	2,597	2,577	2,588	111,652
<b>Subtotal for 50 states and the District of Columbia</b>	<b>16,980</b>	<b>16,641</b>	<b>17,404</b>	<b>17,453</b>	<b>16,316</b>	<b>530,756</b>
<b>Data for U.S. dependent areas</b>	<b>724</b>	<b>662</b>	<b>598</b>	<b>628</b>	<b>619</b>	<b>19,355</b>
<b>Total<sup>e</sup></b>	<b>17,726</b>	<b>17,318</b>	<b>18,020</b>	<b>18,099</b>	<b>17,011</b>	<b>550,394<sup>f</sup></b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

<sup>a</sup> From the beginning of the epidemic through 2005.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>d</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>e</sup> Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

<sup>f</sup> Includes 1,162 persons of unknown race or multiple races, 280 persons of unknown state of residence, and 2 persons who were residents of other areas.