

Table 21. Reported AIDS cases for female adults and adolescents, by transmission category and race/ethnicity, 2005 and cumulative—United States and dependent areas

Transmission category	2005		Cumulative ^a	
	No.	%	No.	%
White, not Hispanic				
Injection drug use	506	28	14,519	40
Hemophilia/coagulation disorder	2	0	119	0
High-risk heterosexual contact ^b	749	42	15,094	42
Sex with injection drug user	188	10	5,536	15
Sex with bisexual male	60	3	1,779	5
Sex with person with hemophilia	6	0	325	1
Sex with HIV-infected transfusion recipient	7	0	336	1
Sex with HIV-infected person, risk factor not specified	488	27	7,118	20
Receipt of blood transfusion, blood components, or tissue	20	1	1,871	5
Other/risk factor not reported or identified	517	29	4,653	13
Total	1,794	100	36,256	100
Black, not Hispanic				
Injection drug use	1,130	16	37,977	35
Hemophilia/coagulation disorder	5	0	143	0
High-risk heterosexual contact ^b	3,079	44	46,642	43
Sex with injection drug user	367	5	13,184	12
Sex with bisexual male	115	2	2,190	2
Sex with person with hemophilia	6	0	117	0
Sex with HIV-infected transfusion recipient	12	0	266	0
Sex with HIV-infected person, risk factor not specified	2,579	37	30,885	28
Receipt of blood transfusion, blood components, or tissue	53	1	1,531	1
Other/risk factor not reported or identified	2,711	39	22,158	20
Total	6,978	100	108,451	100
Hispanic				
Injection drug use	371	19	12,407	36
Hemophilia/coagulation disorder	3	0	65	0
High-risk heterosexual contact ^b	990	51	16,964	49
Sex with injection drug user	175	9	6,293	18
Sex with bisexual male	46	2	782	2
Sex with person with hemophilia	1	0	45	0
Sex with HIV-infected transfusion recipient	7	0	130	0
Sex with HIV-infected person, risk factor not specified	761	39	9,714	28
Receipt of blood transfusion, blood components, or tissue	9	0	608	2
Other/risk factor not reported or identified	578	30	4,570	13
Total	1,951	100	34,614	100

Table 21. Reported AIDS cases for female adults and adolescents, by transmission category and race/ethnicity, 2005 and cumulative—United States and dependent areas (cont)

Transmission category	2005		Cumulative ^a	
	No.	%	No.	%
Asian/Pacific Islander				
Injection drug use	9	11	124	12
Hemophilia/coagulation disorder	0	0	7	1
High-risk heterosexual contact ^b	37	44	524	52
Sex with injection drug user	4	5	112	11
Sex with bisexual male	0	0	79	8
Sex with person with hemophilia	0	0	4	0
Sex with HIV-infected transfusion recipient	1	1	23	2
Sex with HIV-infected person, risk factor not specified	32	38	306	30
Receipt of blood transfusion, blood components, or tissue	3	4	93	9
Other/risk factor not reported or identified	36	42	258	26
Total	85	100	1,006	100
American Indian/Alaska Native				
Injection drug use	15	28	267	42
Hemophilia/coagulation disorder	0	0	3	0
High-risk heterosexual contact ^b	24	44	264	41
Sex with injection drug user	8	15	101	16
Sex with bisexual male	0	0	26	4
Sex with person with hemophilia	0	0	2	0
Sex with HIV-infected transfusion recipient	0	0	5	1
Sex with HIV-infected person, risk factor not specified	16	30	130	20
Receipt of blood transfusion, blood components, or tissue	2	4	17	3
Other/risk factor not reported or identified	13	24	88	14
Total	54	100	639	100
Total cases				
Injection drug use	2,047	19	65,534	36
Hemophilia/coagulation disorder	11	0	340	0
High-risk heterosexual contact ^b	4,918	45	79,872	44
Sex with injection drug user	751	7	25,310	14
Sex with bisexual male	223	2	4,876	3
Sex with person with hemophilia	13	0	493	0
Sex with HIV-infected transfusion recipient	27	0	767	0
Sex with HIV-infected person, risk factor not specified	3,904	36	48,426	27
Receipt of blood transfusion, blood components, or tissue	87	1	4,144	2
Other/risk factor not reported or identified	3,881	35	31,879	18
Total	10,944^c	100	181,769^d	100

^a From the beginning of the epidemic through 2005.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes 82 females of unknown race or multiple races.

^d Includes 803 females of unknown race or multiple races.