

Table 11. Estimated numbers of persons living with AIDS at the end of 2005, by race/ethnicity, sex, and transmission category—50 states and the District of Columbia (Revised June 2007)

Transmission category	White, not Hispanic		Black, not Hispanic		Hispanic		Asian/Pacific Islander		American Indian/Alaska Native		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	98,062	75	54,858	44	34,245	56	2,560	72	670	56	191,362	59
Injection drug use	13,194	10	34,680	28	15,125	25	325	9	205	17	63,864	20
Male-to-male sexual contact and injection drug use	11,757	9	9,625	8	4,041	7	174	5	202	17	25,961	8
High-risk heterosexual contact ^b	5,647	4	22,988	19	7,419	12	422	12	99	8	36,784	11
Other ^c	1,805	1	1,651	1	585	1	74	2	17	1	4,154	1
Subtotal	130,464	100	123,803	100	61,416	100	3,556	100	1,193	100	322,125	100
Female adult or adolescent												
Injection drug use	7,313	39	18,714	31	5,059	32	113	16	137	36	31,521	33
High-risk heterosexual contact ^b	10,801	58	39,495	66	10,482	66	539	77	225	60	61,938	65
Other ^c	526	3	1,479	2	412	3	48	7	14	4	2,500	3
Subtotal	18,641	100	59,689	100	15,953	100	700	100	375	100	95,959	100
Child (<13 yrs at diagnosis)												
Perinatal	501	91	2,458	98	650	95	19	94	13	99	3,661	97
Other ^d	50	9	39	2	36	5	1	6	0	1	126	3
Subtotal	551	100	2,497	100	686	100	20	100	13	100	3,787	100
Total^e	149,658		185,988		78,054		4,276		1,581		421,873	

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

^a Includes 2,316 persons of unknown race or multiple races.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^e Includes 3 person of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.