

## Bibliography

1996

---

- F. J. Eppig and J. A. Poisal, "Medicare Beneficiary Information Needs: 1994," Health Care Financing Review, Vol. 18, No. 1 (Fall 1996): 247-252.  
Report presents enrollee information needs and sources of information from the 1994 MCBS.
- M. E. Gornick, P. W. Eggers, T. W. Reilly, R. M. Mentnech, L. K. Fitterman, L. E. Kucken, and B. C. Vladeck, "Effects of Race and Income on Mortality and Use of Services Among Medicare Beneficiaries," New England Journal of Medicine, Vol. 335, No. 11 (September 12, 1996): 791-799.  
1990 census data on median income was combined at the zipcode level with 1993 Medicare administrative data for 26 million beneficiaries 65 years old or older. 1993 MCBS data were used to validate results and determine influenza vaccination rates. Results of calculated age-adjusted mortality rates and age- and sex-adjusted rates for selected diagnoses showed that race and income have substantial effects on mortality and use of services among Medicare beneficiaries.
- James Grana, and Bruce Stuart, "The Impact of Insurance on Access to Physician Services for Elderly People with Arthritis," Inquiry, 33: 326-338 (Winter 1996/97): 326-338.  
Focusing on beneficiaries with arthritis, the study examines the relationship between the degree of health insurance coverage, the likelihood of receiving care, and the amount of care. Finding a positive relationship, the authors discuss the possible effects of increasing insurance coverage.
- L. Gruenberg, E. Kaganova, and M. C. Hornbrook, "Improving the AAPCC With Health-Status Measures From the MCBS," Health Care Financing Review, Vol. 17, No. 3 (Spring 1996): 59-75.  
Data from the 1991 MCBS were used in predictive modeling of 1992 Medicare costs with AAPCC (Adjusted Average Per Capita Cost). The model produces stable unbiased estimates of expenditures on validation samples.
- Mary A. Laschober, and Gary L. Olin, Health & Health Care of the Medicare Population: Data From the 1992 Medicare Current Beneficiary Survey, (Rockville, MD: Westat, Inc., November 1996).  
Presents design of the MCBS and findings for the 1992 Cost and Use file. The first in an annual series of data reports from the MCBS cost and use file. Further reports are available for the 1993 data (Laschober, 1997) and the 1994 data (Olin and Liu, 1998).
- T. L. Mark and L. C. Paramore, "Pneumococcal Pneumonia and Influenza Vaccination: Access to and Use by U. S. Hispanic Medicare Beneficiaries," American Journal of Public Health, Vol. 86, No. 11 (November 1996): 1545-1550.  
1992 MCBS data are used to determine differences in influenza and pneumococcal vaccination rate between Hispanic and non-Hispanic white elderly Medicare

beneficiaries. Speaking Spanish was significantly, negatively associated with influenza vaccination.

Jennifer O'Sullivan, Jason S. Lee, and Benson Yang, Medicare: The Role of Supplemental Health Insurance, (Washington, DC: Congressional Research Service, The Library of Congress, October 10, 1996) Report 96-826 EPW. Medicare covers less than half of beneficiaries' total health care expenses. This Report to the Congress describes supplementary sources of coverage (employer-sponsored, individually purchased, or Medicaid), their role in protecting various segments of the population, and the implications of policy options.

Gerald Riley, Cynthia Tudor, Yen-Pin Chiang, and Melvin Ingber, "Health Status of Medicare Enrollees in HMOs and Fee-for-Service in 1994," Health Care Financing Review, Vol. 17, No. 4 (Summer 1996): 65-76. HMO respondents were less likely to report fair or poor health, functional impairment, or heart disease. The Medicare payment formula for HMOs does not adequately adjust for the better health and consequent lower expected costs of HMO enrollees. The addition of health-status measures would improve payment accuracy.

Linda J. Weiss, and Jan Blustein, "Faithful Patients: The Effect of Long-Term Physician-Patient Relationships on the Costs and Use of Health Care by Older Americans," American Journal of Public Health, Vol. 86, No. 12 (December 1996): 1742-1747. Assesses influence of duration of patient-physician relationship for the amount and type of care received. Most Medicare elderly have long-time ties with a primary physician, and those that do tend to have lower medical care costs.