

ASSESSMENT OF GULF OF ALASKA  
ATKA MACKEREL  
(Partial Chapter)

by  
Sandra A. Lowe and Lowell W. Fritz

EXECUTIVE SUMMARY

Relative to the November 1999 SAFE report, the following substantive changes have been made in the current draft of the Atka mackerel chapter:

Changes in the Input Data

Catch data are updated.

Changes in the Assessment Methodology

There are no changes in the assessment methodology

Changes in Assessment Results

There are no changes in assessment results

Response to SSC comments

There were no SSC comments pertaining to the Atka mackerel assessment from the December 1999 SSC minutes.

In this assessment the following issues are highlighted:

- Rationale for maintaining separate Bering Sea/Aleutian Islands and Gulf of Alaska Atka mackerel assessments is provided.
- There is no reliable estimate of current biomass from the Gulf of Alaska bottom trawl survey. Using Tier 6 criteria, overfishing level is set equal to the average catch from 1978-95, which equals 6,200 mt.
- Given that:
  - a) there is no reliable estimate of current biomass,
  - b) Leslie estimates of local population sizes suggest that abundance has declined significantly in localized areas from 1992-94, and
  - c) the species has exhibited vulnerability to fishing pressure in the past,it is recommended that the ABC for Atka mackerel in the GOA be 600 mt, enough to satisfy only the anticipated bycatch needs of other trawl fisheries, principally those for Pacific cod, rockfish and pollock.

Atka mackerel (*Pleurogrammus monopterygius*) are distributed from the east coast of the Kamchatka peninsula, throughout the Komandorskiye and Aleutian Islands, north to the Pribilof Islands in the eastern Bering Sea, and eastward through the Gulf of Alaska to southeast Alaska. Their center of abundance according to past surveys has been in the Aleutian Islands, particularly from Buldir Island to Seguam Pass.

An Atka mackerel population existed in the Gulf of Alaska primarily in the Kodiak, Chirikof, and Shumagin areas, and supported a large foreign fishery through the early 1980s. By the mid-1980s, this fishery, and presumably the population, had all but disappeared. A further indication that the population had disappeared is the bycatch of Atka mackerel in other fisheries was less than 5 mt prior to their inclusion into the Other Species category in 1988. The dramatic decline of the Atka mackerel fishery in the Gulf of Alaska suggests that the area may be the edge of the species' range and be populated only during periods when recruitment, probably as juveniles, from the Aleutian portion of the range is strong (Ronholt 1989). Recently, Atka mackerel have been detected by the summer trawl surveys only in the Shumagin (Western) area of the Gulf of Alaska.

A morphological and meristic study suggested that there may be separate populations in the Gulf of Alaska and the Aleutian Islands (Levada 1979). This study was based on comparisons of samples collected off Kodiak Island in the central Gulf, and the Rat Islands in the Aleutians. Lee (1985) also conducted a morphological study of Atka mackerel from the Bering Sea, Aleutian Islands and Gulf of Alaska. The data showed some differences (although not consistent by area for each characteristic analyzed), suggesting a certain degree of reproductive isolation. However, results from a genetics study comparing Atka mackerel samples from the western Gulf of Alaska with samples from the Eastern, Central, and Western Aleutians showed no evidence of discrete stocks (Lowe et al. 1998). Between-sample variation was extremely low among the four samples indicating a large amount of gene flow is occurring throughout the range. It is presumed that gene flow is occurring during the larval, pelagic stage, particularly in the Aleutian portion of their range, and that the localized aggregations reflect the distribution of surviving, settled larvae and juveniles. Differences in growth rates consistently observed throughout their Alaskan range are phenotypic characteristics reflecting differences in the local environment.

While genetic information suggests that the Aleutian Island (AI) and Gulf of Alaska (GOA) populations of Atka mackerel could be managed as a unit stock, there are significant differences in population size, distribution, recruitment patterns, and resilience to fishing that suggest otherwise. Bottom trawl surveys and fishery data suggest that the Atka mackerel population in the GOA is smaller and much more patchily distributed than that in the AI, and composed almost entirely of fish > 30 cm in length. There are also more areas of moderate Atka mackerel density in the AI than in the GOA. The lack of small fish in the GOA suggests that Atka mackerel recruit to that region differently than in the AI, perhaps as juveniles moving east from the larger population in the AI rather than from larval settlement in the area. This might also explain the greater sensitivity to fishing depletion in the GOA as shown by both the history of the GOA fishery since the early 1970s and Leslie depletion estimates of catchability (Lowe and Fritz 1996). Catches of Atka mackerel from the GOA peaked in 1975 at about 27,000 mt. Recruitment to the AI population was low from 1980-1985, and catches in the GOA declined to 0 in 1986. Only after a series of large year classes recruited to the AI region in the late 1980s did the population and fishery reestablish in the GOA beginning in the early 1990s. After passage of these year classes through the population, the GOA population, as sampled in the 1999 GOA bottom trawl survey, has declined and is very patchy in its distribution. Leslie depletion analyses using AI and GOA fishery data suggest that catchability increased from one year to the next in the GOA fished areas, but remained the same in the AI areas. These differences in population resilience, size, distribution, and recruitment argue for separate management of GOA and AI stocks despite their genetic similarities.

### 10.2.1 Catch History and Fishery Management

Prior to the mid-1980s, Atka mackerel were fished exclusively by foreign vessels, primarily from the Soviet Union. Landings peaked at 27,777 mt in 1975, then dropped to almost 0 in 1986. Some joint venture operations participated in this fishery from 1983 to 1985. All landings since then have been taken by the domestic fishery.

In 1988, Atka mackerel were combined in the other species category due to low abundance and the absence of a directed fishery for the previous several years. However, beginning in 1990, Atka mackerel were targeted in the western Gulf of Alaska (GOA). From 1990-1993, catches of other species in the GOA were dominated by Atka mackerel, primarily from the Western regulatory area. Atka mackerel were separated from the other species category and became a separate target category in the GOA in 1994 after approval of Amendment 31 to the Fishery Management Plan for the Groundfish Fishery of the Gulf of Alaska. Recent catches of Atka mackerel from the GOA have been:

Year	GOA MANAGEMENT AREA			TOTAL
	Western	Central	Eastern	
1990 <sup>1</sup>	1,416	0	0	1,416
1991 <sup>2</sup>	3,249	9	0	3,258
1992 <sup>2</sup>	13,785	49	0	13,834
1993 <sup>2</sup>	4,867	2,143	0	7,010
1994 <sup>2</sup>	2,661	877	0	3,538
1995 <sup>2</sup>	329	370	2	701
1996 <sup>2</sup>	1,577	9	0	1,586
1997 <sup>2</sup>	321	8	2	331
1998 <sup>2</sup>	279	38	0	317
1999 <sup>3</sup>				262
2000 <sup>3,4</sup>	-	-	-	170

1 Actual observed catch

2 NMFS Blend data

3 Catches not available by regulatory area; TAC is set GOA-wide.

4 NMFS Bulletin Board as of 10/21/00

Total catches of Atka mackerel were small until 1992, when approximately 14,000 mt were taken in the Shumagin area. The 1990 catch of 1,416 mt is a minimum estimate, since this was the tonnage actually observed by domestic observers. The Regional Office's estimate of catch for 1990 is underestimated, as Gulf of Alaska Atka mackerel catches were incorrectly being reported as landed in the Aleutian Islands (pers. comm. Galen Tromble, Regional Office, Juneau, Alaska). For 1995 and 1996, the Council approved an ABC and a TAC of 3,240 mt for Gulf of Alaska Atka mackerel. For purposes of data collection and effort

dispersion, 2,310 mt was allocated to the Western or Shumagin subarea (Area 610) and 925 mt was allocated to the Central, or the combined Chirikof and Kodiak subareas (Areas 620 and 630). The Western subarea (610) was not opened to the directed Atka mackerel fishery in 1995 because the overfishing level for Pacific ocean perch (POP) was nearly reached; Atka mackerel fisheries have had significant bycatch of POP (A. Smoker, NMFS, Juneau, AK, pers. comm.). In 1996, the fishery in the Western subarea was restricted to a 12-h opening on July 1, due again to concerns about the POP catch exceeding TAC and approaching the overfishing level; about 1,300 mt of Atka mackerel was caught. The 1996 POP catch exceeded the Central area POP overfishing level, thus there was no opening for the Atka mackerel fishery in that area. Since 1997 the Atka mackerel fishery has been managed as a bycatch-only fishery with TACs of 1,000 mt in 1997 and 600 mt in 1998, 1999 and 2000.

Scientific research catches are reported in the SAFE reports. Table 10.1 documents annual research catches (1977 - 1998) from NMFS trawl surveys.

### 10.2.2 Description of the Directed Fishery

The patterns of the Atka mackerel fishery generally reflect the behavior of the species: (1) the fishery is highly localized and usually occurs in the same few locations each year; (2) the schooling semi-demersal nature of the species makes it particularly susceptible to trawl gear fished on the bottom; and (3) trawling occurs almost exclusively at depths less than 200 m. Observed Atka mackerel fishery trawl locations in 1990 through 1994 in the Gulf of Alaska are shown in Figure 10.1. In 1990, catches occurred near the edge of the continental shelf on Davidson Bank south of Unimak and Sanak Islands. During 1991-1994, the fishery in the western GOA was concentrated off the southern coast of Umnak Island. However, in 1993 and 1994, areas south and east of the Shumagin Islands were also fished for Atka mackerel, and contributed about 30% of the total GOA catch for the year.

### 10.2.3 Bycatch and Discards

The amounts (mt) of Atka mackerel retained and discarded by target fishery and area in the Gulf of Alaska in 1994 and 1995 are listed below (NMFS/AFSC Blend data for 1994 and 1995):

1994	Area 610			Area 620			Gulf of Alaska		
	Retain	Discard	Total	Retain	Discard	Total	Retain	Discard	Total
Target Fishery									
Atka mackerel	2,232.5	160.1	2,392.6	785.8	85.3	871.0	3,018.3	245.4	3,263.7
Pacific cod, trawl	1.3	0.5	1.8	-	-	-	1.3	0.5	1.8
Rockfish, trawl	230.7	26.1	256.8	0.9	-	0.9	231.6	26.2	257.9
Pelagic pollock	8.2	0.1	8.3	-	0.0	0.0	8.2	0.2	8.3
All others, trawl	-	0.1	0.1	2.4	0.6	3.0	2.4	1.7	4.1
Grand Total	2,472.7	187.0	2,659.7	789.1	85.8	874.9	3,261.8	273.9	3,535.7
1995	Retain	Discard	Total	Retain	Discard	Total	Retain	Discard	Total
Atka mackerel, trawl	0.0	0.0	0.0	161.3	0.0	161.3	161.3	0.0	161.3
Bottom Pollock, trawl	107.0	4.1	111.1	0.0	0.0	0.0	107.0	4.2	111.2
Pacific cod, trawl	3.6	84.0	87.6	2.3	48.7	51.0	5.8	167.5	173.3
Rockfish, trawl	109.1	15.5	124.7	117.1	2.2	119.3	227.2	20.0	247.2
All others	0.2	5.1	5.3	0.0	0.8	0.8	0.2	7.2	7.4
Grand Total	219.9	108.7	328.6	280.7	51.7	332.4	501.5	199.0	700.5

Other than the 1994 Atka mackerel fishery, the only fisheries that caught significant bycatch amounts of Atka mackerel in 1994 and 1995 were the pollock, cod and rockfish trawl fisheries in areas 610 and 620. In 1995, the bottom pollock fishery in area 610 had an Atka mackerel bycatch rate (mt of Atka mackerel per mt of pollock) of less than 1%, as did the cod trawl fisheries in the Western and Central GOA (areas 610, 620 and 630). Calculation of true Atka mackerel bycatch rates for the rockfish fisheries is complicated by certain vessels apparently “topping-off” with Atka mackerel on specific hauls, rather than catching Atka mackerel along with rockfish in the same haul.

The discard rate for the 1994 Gulf Atka mackerel fishery as a whole was about 8%, which was about half that of the Aleutian Islands Atka mackerel fishery in 1994 (over 16%). This most likely reflects the differences in mean sizes of Atka mackerel in the two regions, larger fish being in the GOA. However, direct information about the size or sex composition of the discards is unavailable.

#### 10.2.4 Fishery Length Frequencies

Atka mackerel length distributions from the 1990-1994 fisheries by area and sex are shown in Figures 10.2 and 10.3. Male and female Atka mackerel from the 1990 fishery on Davidson Bank averaged 44.6 cm and 48.7 cm, respectively. These fish were larger than those caught in 1991-94 south of Umnak Island, where males averaged 38.4, 39.9, 41.8, and 44.4 cm in length in the four years, respectively, and females averaged 39.0, 41.6, 42.2, and 44.9 cm. The fishery south of Umnak Island appears to have harvested from the same cohort of fish in each of the last four years, since there have been both slight increases in mean length and little recruitment of younger fish (decreases in numbers between 30 and 35 cm). In 1993 and 1994 in the Shumagin Islands area, Atka mackerel similar in size to those caught on Davidson Bank in 1990 were caught; males averaged 44.4 and 44.9 cm, while females averaged 46.5 and 47.7 cm in the two years, respectively.

#### 10.2.4 Fishery Sex Ratios

In certain areas and months, catches of Atka mackerel were comprised of more females than males (Figures 10.2 and 10.3). The large percentage of female Atka mackerel caught by the fishery on Davidson Bank and near the Shumagin Islands could be a result of segregation of the population by sex during spawning and periods of male nest-guarding in summer and early fall. By contrast, males and females were more evenly represented in the fishery length-frequency samples collected near Umnak Island in 1991-94, regardless of the time of year. This suggests differential habitat utilization by Atka mackerel in areas frequented by the fishery in the Gulf of Alaska, which was also observed between areas in the Aleutian Islands (Fritz and Lowe 1998).

#### 10.2.5 Fishery Age Frequencies

There is only very limited age data available from the 1990 Davidson Bank fishery, the 1992 Umnak Island fishery and the 1994 fishery which operated off Umnak Island, Davidson Bank and Shumagin Bank (Figures 10.4-10.5). In 1990, most Atka mackerel aged were 3-4 year-olds of the 1986 and 1987 year classes, with smaller numbers of 5-7 year-olds from the 1983-1985 year classes. The oldest Atka mackerel from Davidson Bank was 13 years old (1977 year class). In the 1992 Umnak Island fishery, most Atka mackerel aged were 3-4 year-olds from the 1988 and 1989 year classes, and no fish older than 8 years old were aged. The 1988 year class continues to dominate the age distributions in the 1994 fishery (Figure 10.5). The oldest fish aged from the 1994 collection was an 11 year old from Shumagin Bank.

#### 10.2.6 Fishery and Steller Sea Lions

The western stock of Steller sea lions (defined as west of 144°W, or at Cape Suckling east of Prince William Sound) is currently listed as endangered under the Endangered Species Act, and has been listed as threatened since 1990. In 1991-92, 10 nm annual trawl exclusion zones were established around all rookeries west of 150°W; in 1992-93, 20 nm trawl exclusion zones were established around 6 rookeries in the eastern Aleutian Islands that are operational only during the BSAI pollock A-season. In 1993, NMFS designated Steller sea lion critical habitat, which includes a 20 nm aquatic zone around all rookeries and major haulouts west of 144°W, and three foraging areas, one of which contains Shelikof Strait. Sea lion food habits data collected in the Aleutian Islands revealed that Atka mackerel was the most common food item of adults and juveniles in the summer (NMFS 1995).

From 1977 to 1984 and in 1990, 0-11% of the annual Gulf of Alaska Atka mackerel harvest was caught within 20 miles of all Gulf of Alaska sea lion rookeries and major haulouts, reflecting the offshore distribution of the fishery. In 1991-93, however, the fishery moved closer to shore, and this percentage increased to 82-98%, almost all of which was caught between 10-20 nm of Steller sea lion rookeries on Ogchul and Adugak Islands (near Umnak Island), and Atkins and Chernabura Islands in the Shumagin Islands (Figure 10.1).

Leslie depletion estimates of local fishery harvest rates were much greater than estimated Gulf-wide harvest rates (Lowe and Fritz 1996; 1997). This could have adversely affected Steller sea lion foraging success, which raised concerns about how the fishery may have affected food availability and the potential for recovery of the Steller sea lion population. There has not been a directed Gulf of Alaska Atka mackerel fishery since 1996. In June 1998, the Council passed a fishery regulatory amendment which proposed a four-year timetable to temporally and spatially disperse and reduce the level of Atka mackerel fishing within Steller sea lion critical habitat in the Bering Sea/Aleutian Islands. The management of the Bering Sea/Aleutian Islands Atka mackerel fishery is detailed in Lowe and Fritz (1999)

### 10.3.1 Absolute Abundance and Survey Biomass

Bottom trawl surveys of the Gulf of Alaska groundfish community have been conducted every three years since 1984 using an area-depth stratified and area-swept design (Figure 10.6). In 1999, the same GOA survey design was maintained, but effort allocation was shifted to provide more even coverage within depth strata. Atka mackerel are a very difficult species to survey because: (1) they do not have a swim bladder, making them poor targets for hydroacoustic surveys; (2) they prefer hard, rough and rocky bottom which makes sampling with the standard survey bottom trawl gear difficult; and (3) their schooling behavior and patchy distribution makes the species susceptible to large variances in catches which greatly affect area-swept estimates of biomass.

The general groundfish surveys of the Gulf of Alaska are particularly problematic for Atka mackerel given the characteristics described above. In 1996, a meaningful estimate of biomass could not be determined from the data due to extreme variances. Over 98% of the Atka mackerel caught in the 1996 survey were encountered in a single haul within a large stratum (stratum 11, Figure 10.6), which yielded a large stratum biomass with an extremely large confidence interval.

Although estimates of abundance from earlier surveys have been presented in previous assessments, they were also compromised by the problem of large confidence intervals, although not to the same degree as observed in 1996. For instance, as in the 1996 survey, 98% of all the Atka mackerel caught in the 1990 survey were captured in one haul. Atka mackerel have been inconsistently caught in the GOA surveys, appearing in 5%, 28%, 12% and 20% of the hauls in the Shumagin area in the 1990, 1993, 1996, and 1999 GOA surveys, respectively. Examples of the haul-by-haul distribution of Atka mackerel catches in the 1996 and 1999 surveys are shown in Figures 10.7 and 10.8, respectively. What can be concluded from this is that the general groundfish GOA bottom trawl survey, as it has been designed and used since 1984, does not assess Atka mackerel well and the resulting biomass estimates are not considered reliable indicators of absolute abundance or as indices of trend.

Given the problems with assessing GOA Atka mackerel with the bottom trawl survey, there is no reliable estimate of current biomass of Atka mackerel in the GOA. The only indicator of recent trends in abundance comes from analyses of catch-per-unit-effort of the 1992-94 directed Atka mackerel fisheries south of Umnak Island and southeast of the Shumagin Islands (Figure 10.1), which are detailed in the BSAI assessment (Lowe and Fritz 1997) and in the appendix of the 1996 assessment (Lowe and Fritz 1996). These analyses suggest that the Umnak Island Atka mackerel population declined 81% between 1992 and 1994, while the Shumagin Island population declined 58%.

### 10.3.2 Survey Length Frequencies

Length frequency distributions from the 1990, 1993, 1996 and 1999 surveys are shown in Figure 10.9. Mean lengths of males and females, respectively, from each survey were: 37.7 and 36.7 cm in 1990, 42.7 and 44.1 cm in 1993, 45.4 and 47.0 cm in 1996, and 45.4 and 46.8 cm in 1999.

In the 1990 length distribution, Atka mackerel between 20-25 cm were likely 2-year olds from the 1988 year class. The 1991 Aleutian Islands survey also found large numbers of fish from the 1988 year class (Lowe 1992). In the 1993, 1996, and 1999 length distributions, the modes at 41-43 cm, 45-47 cm and 46-51 cm, respectively, are probably composed primarily of 1988 year-class fish. In the 1999 survey there were small numbers of fish less than 37 cm (Figure 10.9)

### 10.3.3 Survey Age Frequencies

Age distributions from the 1993 Gulf of Alaska trawl survey (Figure 10.10) show a mode of 5-year olds from the 1988 year class and a smaller mode of 2-year old fish from the 1990 year class. The age range from survey data is very narrow, basically ages 2-6. Atka mackerel are a summer-fall spawning fish that apparently do not lay down an otolith annulus in the first year (Anderl et. al, 1996.). This is accommodated by adding one year to ages determined by counts of otolith hyaline zones by the Alaska Fisheries Science Center (AFSC) Age and Growth Laboratory. All age data presented in this report have been corrected this way.

## 10.4

## BIOLOGICAL PARAMETERS

### 10.4.1 Natural Mortality, Age of Recruitment, and Maximum Age

A natural mortality rate of 0.3 is assumed for Gulf of Alaska Atka mackerel based on Aleutian Islands Atka mackerel (Lowe 1997).

A qualitative look at the sparse GOA fishery age data shows recruitment patterns similar to the Aleutian Islands fishery. The age of first recruitment appears to be 2 years, and full recruitment at 4 years (Figure 10.4). This pattern becomes somewhat obscured when a strong year class dominates the distributions (Figure 10.5).

The maximum age seen in the Gulf of Alaska fishery is 13 years (1990 fishery). This compares with a maximum age of 15 years for the Aleutian Islands.

### 10.4.2 Length and Weight at Age

Parameters of the von Bertalanffy length-age equation and a weight-length relationship were calculated from the combined 1990, 1992, and 1994 fishery data. Sexes were combined to provide an adequate sample size. The estimated von Bertalanffy growth parameters are:

$$\begin{aligned}L_{\infty} &= 54.56 \text{ cm} \\K &= 0.22 \\t_0 &= -2.78 \text{ yr}\end{aligned}$$

$$\text{Length-age equation: Length (cm)} = L_{\infty}\{1 - \exp[-K(\text{age} - t_0)]\}.$$

The weight-length relationship was determined to be:

$$\text{Weight (kg)} = 4.61\text{E-}05 * \text{Length (cm)}^{2.698}$$

These curves are shown in Figure 10.11 along with the observed mean values. Growth parameters were also estimated from data collected during the 1993 Gulf of Alaska survey. As in the Aleutians, the survey tends to select for smaller fish at age than the fishery. The estimated von Bertalanffy parameters from the 1993 survey are:

$$\begin{aligned}L_{\infty} &= 47.27 \text{ cm} \\K &= 0.610 \\t_0 &= 0.38 \text{ yr}\end{aligned}$$

and the estimated weight-length relationship is:



$$\text{Weight (kg)} = 1.55\text{E-}05 * \text{Length (cm)}^{2.979}$$

These curves are also shown in Figure 10.11 for comparison with the fishery data.

#### 10.4.3 Maturity at Length and Age

Female maturity at length and age were determined for Gulf of Alaska Atka mackerel (McDermott and Lowe 1997). The age at 50% maturity is 3.6 years and length at 50% maturity is 38.3 cm (Figure 10.12). The maturity schedules are given in Table 10.2.

#### 10.4.4 Selectivity at Age

The small amount of age data for Gulf of Alaska Atka mackerel show similar selectivity patterns as seen in the Aleutian survey and fishery data. The fishery data tend to show older fish than the survey samples. The oldest age from the 1993 survey was 9 years old and the age distribution consisted of mostly 2-6 year olds (Figure 10.10). The fishery age data is dominated by fish between 3-8 years of age.

### 10.5 ACCEPTABLE BIOLOGICAL CATCH

There is no reliable estimate of current Atka mackerel biomass in the Gulf of Alaska. In this situation, Tier 6 of Amendment 44 of the BSAI FMP defines the overfishing level (OFL) as the average catch from 1978-95, and that ABC cannot exceed 75% of the OFL. The average annual catch from 1978-95 is 6,200 mt, which is the overfishing level; thus ABC is capped at  $6,200 \times 0.75 = 4,700$  mt. However, we recommend that ABC be set lower than 4,700 mt for the following reasons:

- 1) When ABCs were lower than 4,700 mt, such as in 1994 when the ABC was 3,280 mt, the fishery may have created localized depletions of Atka mackerel in the two primary fished areas, south of Umnak Island and southeast of the Shumagin Islands (see appendix in Lowe and Fritz 1996). The 1994 ABC was set using a 15% harvest rate applied to the 1993 survey biomass estimate of 21,600 mt. The two 1994 fisheries at Umnak and Shumagin combined for over 3,000 mt of the 3,500 mt caught that year, and harvest rates far exceeded the target 15% in each area: at Umnak, the harvest rate was estimated at 85%, and at Shumagin, the harvest rate was estimated at 91%. The 1990 and 1993 surveys also found that Atka mackerel in the GOA were principally congregated in these two areas used by the fishery. These data indicate that the fishery was very efficient in removing fish from these areas and at rates which far surpassed the target Gulf-wide harvest rate.
- 2) Analyses of local fishery CPUEs suggests that the Atka mackerel populations at Umnak and Shumagin Islands declined significantly between 1992 and 1994 (see appendix in Lowe and Fritz 1996). This also reflects the trend of the Aleutian Island Atka mackerel population during that period.
- 3) The GOA Atka mackerel population appears to be particularly vulnerable to fishing pressure because of its very patchy distribution and sporadic recruitment patterns. This is reflected in the Leslie depletion analyses (appendix in Lowe and Fritz 1996) and by the disappearance of the population in the mid-1980s following a period with annual catches as high as 27,000 mt.

For the above reasons, **we recommend a 2001 ABC for GOA Atka mackerel sufficient only to satisfy the bycatch needs of other trawl fisheries**, a recommendation identical to that made since 1997. Catches of Atka mackerel in the GOA in 1997, 1998, 1999, and 2000 were only 331, 291, 316, and 170 mt, respectively, which could represent the natural bycatch of Atka mackerel in other groundfish fisheries. **We recommend a 2001 GOA Atka mackerel ABC of 600 mt**, or approximately double the 1997-1999 catches.

The overfishing level for Gulf Atka mackerel was calculated using Tier 6 criteria, the average catch from 1978-95, since there is no reliable estimate of current biomass. **The average catch from 1978-95 is 6,200 mt, which is the 2001 overfishing level for Atka mackerel in the Gulf of Alaska.**

The Steller sea lion (*Eumetopias jubatus*) was listed as threatened under the Endangered Species Act (ESA) in April 1990. As a result of the listing, trawling was prohibited within 10 nautical miles (nm) of all rookeries in the Central and Western Gulf of Alaska year-round beginning in June 1991; 10 nm no-trawl buffer zones were created around all other Steller sea lion rookeries in the Aleutian Islands and Bering Sea in January 1992. The intent of this action was to exclude trawl fishing activity from areas known to be important for sea lion foraging and reproduction. While there is no proven cause and effect relationship between the decline in Steller sea lion numbers and increases in fishery removals near terrestrial sea lion habitats, NMFS imposed the 10 nm trawl exclusion zones based on general conservation principles in an effort to promote sea lion recovery. Aerial surveys conducted through 1996 revealed that the Steller sea lion population in the western GOA has been relatively stable since 1989, but at about 50% of the size that existed prior to the decline (mid-1970s). This is in contrast to the central GOA, where the sea lion population has declined over 80% in the same period, and continues to decline at about 10% per year (NMFS 1995).

Steller sea lion food habits data (from analysis of scats) from the Aleutian Islands indicates that Atka mackerel is an important part of their diet, at least during summer (NMFS 1995). The prevalence of Atka mackerel and walleye pollock in sea lion scats reflected the distributions of each fish species in the Aleutian Islands region. The percentage occurrence of Atka mackerel was progressively greater in samples taken in the central and western Aleutian Islands, where most of the Atka mackerel biomass in the Aleutian Islands is located. Conversely, the percentage occurrence of pollock was greatest in the eastern Aleutian Islands. Steller sea lion food habits data from the western Gulf of Alaska are relatively sparse, so it is not known how important Atka mackerel is to sea lions in this area. The close proximity of fishery locations to sea lion rookeries in the western Gulf suggests that Atka mackerel could be a prey item at least during the summer. Analyses of fishery CPUE revealed that the fishery may create temporary localized depletions of Atka mackerel, and that these depletions may last for weeks after the vessels have left the area. This supports the argument already made above in the ABC section for a conservative harvest policy for Atka mackerel in the Gulf of Alaska.

$M = 0.30$	$F_{ABC}$	= unknown
Age at full selection = 4 years	$F_{OFL}$	= unknown
2001 exploitable biomass = unknown	2001 Overfishing level	= 6,200 mt
	2001 ABC	= 600 mt

- Anderl, D.M, A. Nishimura and S.A. Lowe. 1996. Is the first annulus on the otolith of the Atka mackerel, *Pleurogrammus monopterygius*, missing? Fish. Bull. 94:163-169
- Fritz, L. W., and S. A. Lowe. 1998. Seasonal distributions of Atka mackerel (*Pleurogrammus monopterygius*) in commercially-fished areas of the Aleutian Islands and the Gulf of Alaska. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-92, 29 p.
- Gorbunova, N.N. 1962. Razmnozhenie i razvite ryb semeistva terpugovykh (Hexagrammidae) (Spawning and development of greenlings (family Hexagrammidae)). Tr. Inst. Okeanol., Akad. Nauk SSSR 59:118-182. In Russian. (Trans. by Isr. Program Sci. Trans., 1970, p. 121-185 in T.S. Rass (editor), Greenlings: taxonomy, biology, interoceanic transplantation; available from the U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA., as TT 69-55097).
- Levada, T.P. 1979. Comparative morphological study of Atka mackerel. Pac. Sci. Res. Inst. Fish. Oceanogr.5(TINRO), Vladivostok, U.S.S.R., Unpublished manuscript.
- Lee, J.U. 1985. Studies on the fishery biology of the Atka mackerel *Pleurogrammus monopterygius* (Pallas) in the north Pacific Ocean. Bull. Fish. Res. Dev. Agency, 34, pp.65-125.
- Lowe, S. A. 1992. Atka Mackerel. In Stock Assessment and Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Regions as Projected for 1993. North Pacific Fisheries Management Council, P.O. Box 103136, Anchorage, Alaska, 99510.
- Lowe, S. A. 1993. Atka Mackerel. In Stock Assessment and Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Regions as Projected for 1994. North Pacific Fisheries Management Council, P.O. Box 103136, Anchorage, Alaska, 99510.
- Lowe, S. A., and L. W. Fritz. 1996. Atka mackerel In Stock Assessment and Evaluation Report for the Groundfish Resources of the Gulf of Alaska as Projected for 1997. North Pacific Fisheries Management Council, P.O. Box 103136, Anchorage, Alaska, 99510.
- Lowe, S. A., and L. W. Fritz. 1997. Atka mackerel. In Stock Assessment and Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Regions as Projected for 1998. North Pacific Fisheries Management Council, P.O. Box 103136, Anchorage, Alaska, 99510.
- Lowe, S. A., and L. W. Fritz. 1999. Atka mackerel. In Stock Assessment and Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Regions as Projected for 2000. North Pacific Fisheries Management Council, P.O. Box 103136, Anchorage, Alaska, 99510.
- Lowe, S. A. and G. G. Thompson. 1993. Exploitation strategies for the Atka mackerel resource of the Aleutian Islands. In Proceedings of the International Symposium on Management for Exploited Fish Populations. G. Kruse, D.M Eggers, R.J. Marasco, C. Pautzke, and T.J. Quinn II (editors). Alaska Sea Grant College Program Report No. 93-02, University of Alaska Fairbanks.
- Lowe, S. A., D. M. Van Doornik, and G. A. Winans. 1998. Geographic variation in genetic and growth patterns of the Atka mackerel (*Pleurogrammus monopterygius*; Hexagrammidae) in the Aleutian Archipelago. Fish. Bull. U. S. 96: 502-515.
- McDermott, S.F. and S.A. Lowe. 1997. The reproductive cycle of Atka mackerel (*Pleurogrammus monopterygius*) in Alaska waters. Fish. Bull. U.S. 95:231-333
- NMFS. 1995. Status review of the United States Steller sea lion (*Eumetopias jubatus*) population. National Marine Mammal Laboratory, Alaska Fishery Science Center, National Marine Fisheries Service, 7600 Sand Point Way, NE, Seattle, WA 98115.
- Ronholt, L.L. 1989. Atka mackerel. In Condition of groundfish resources of the Gulf of Alaska in 1988. T. K. Wilderbuer (editor). NOAA Tech. Memo. NMFS F/NWC-165.

Table 10.1 Research catches of Atka mackerel from NMFS surveys in the Gulf of Alaska.

Year	Research Catches (mt)
1977	0.3
1978	2.8
1979	0.4
1980	3.8
1981	35.3
1982	27.0
1983	0.4
1984	7.5
1985	65.7
1986	
1987	6.2
1988	
1989	
1990	2.7
1991	
1992	
1993	2.4
1994	
1995	
1996	15.1
1997	

Table 10.2. Schedules of age and length specific maturity from McDermott and Lowe (1997).

<u>Length (cm)</u>	<u>Proportion mature</u>	<u>Age</u>	<u>Proportion mature</u>
20	0.00	1	0.00
21	0.00	2	0.04
22	0.00	3	0.22
23	0.00	4	0.69
24	0.00	5	0.94
25	0.00	6	0.99
26	0.00	7	1.00
27	0.00	8	1.00
28	0.00	9	1.00
29	0.00	10	1.00
30	0.00		
31	0.01		
32	0.01		
33	0.02		
34	0.05		
35	0.09		
36	0.17		
37	0.29		
38	0.46		
39	0.63		
40	0.78		
41	0.88		
42	0.93		
43	0.97		
44	0.98		
45	0.99		
46	1.00		
47	1.00		
48	1.00		
49	1.00		
50	1.00		