

SEER Incidence and Delay Adjusted Incidence Rates⁺ All Cancer Sites, By Sex






Figure I-21

⁺ Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and the APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence and Delay Adjusted Incidence Rates+ Both Sexes



+ Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and the APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence and Delay Adjusted Incidence Rates⁺ Males




Figure I-23

+ Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and the APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence and Delay Adjusted Incidence Rates⁺ Females




Figure I-24

+ Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and the APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table II-1
ALL SITES (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINTPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINTPOINTS BY RACE AND SEX

		<u>Jointpoint Segment 1</u>		<u>Jointpoint Segment 2</u>		<u>Jointpoint Segment 3</u>		<u>Jointpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1983	0.9 ^c	1983-1992	1.8 ^c	1992-1995	-1.8	1995-2000	0.1
All Races	Male	1975-1989	1.3 ^c	1989-1992	5.1 ^c	1992-1995	-4.8 ^c	1995-2000	-0.1
All Races	Female	1975-1979	-0.3	1979-1987	1.6 ^c	1987-2000	0.3 ^c		
White	Male and Female	1975-1982	0.8 ^c	1982-1992	1.8 ^c	1992-1995	-1.9	1995-2000	0.2
White	Male	1975-1989	1.4 ^c	1989-1992	4.9 ^c	1992-1995	-5.1 ^c	1995-2000	0.0
White	Female	1975-1979	-0.2	1979-1987	1.7 ^c	1987-2000	0.3 ^c		
Black	Male and Female	1975-1983	1.7 ^c	1983-1989	0.6	1989-1992	4.3 ^c	1992-2000	-1.3 ^c
Black	Male	1975-1981	2.8 ^c	1981-1989	0.6	1989-1992	6.9 ^c	1992-2000	-2.2 ^c
Black	Female	1975-1991	1.1 ^c	1991-2000	-0.2				
<u>SEER Delay-Adjusted Incidence^a</u>									
All Races	Male and Female	1975-1983	0.9 ^c	1983-1992	1.9 ^c	1992-1995	-1.8	1995-2000	0.6
All Races	Male	1975-1989	1.4 ^c	1989-1992	5.2 ^c	1992-1995	-4.8 ^c	1995-2000	0.5
All Races	Female	1975-1979	-0.3	1979-1987	1.5 ^c	1987-2000	0.4 ^c		
White	Male	1975-1989	1.4 ^c	1989-1992	5.1 ^c	1992-1995	-5.1 ^c	1995-2000	0.7 ^c
White	Female	1975-1979	-0.2	1979-1987	1.6 ^c	1987-2000	0.6 ^c		
Black	Male	1975-1981	2.8 ^c	1981-1989	0.7	1989-1992	6.8 ^c	1992-2000	-1.7 ^c
Black	Female	1975-1991	1.2 ^c	1991-2000	0.2				
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1990	0.5 ^c	1990-1994	-0.4	1994-1998	-1.4 ^c	1998-2000	-0.3
All Races	Male	1975-1980	0.9 ^c	1980-1992	0.2 ^c	1992-2000	-1.5 ^c		
All Races	Female	1975-1990	0.6 ^c	1990-1995	-0.2	1995-1998	-1.3 ^c	1998-2000	0.1
White	Male and Female	1975-1990	0.4 ^c	1990-1994	-0.3	1994-1998	-1.3 ^c	1998-2000	-0.1
White	Male	1975-1980	0.8 ^c	1980-1992	0.2 ^c	1992-2000	-1.4 ^c		
White	Female	1975-1990	0.6 ^c	1990-1995	-0.2	1995-1998	-1.3 ^c	1998-2000	0.3
Black	Male and Female	1975-1984	1.3 ^c	1984-1992	0.6 ^c	1992-2000	-1.4 ^c		
Black	Male	1975-1982	1.8 ^c	1982-1990	1.0 ^c	1990-1993	-0.3	1993-2000	-1.9 ^c
Black	Female	1975-1991	1.0 ^c	1991-2000	-0.6 ^c				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ All Cancer Sites, by Race and Sex




Figure II-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table III-1
 BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINTPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINTPOINTS BY RACE AND SEX

		<u>Jointpoint Segment 1</u>		<u>Jointpoint Segment 2</u>		<u>Jointpoint Segment 3</u>		<u>Jointpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1988	1.4 ^c	1988-2000	-0.5 ^c				
All Races	Male	1975-1989	1.2 ^c	1989-2000	-0.5				
All Races	Female	1975-1987	1.7 ^c	1987-2000	-0.6 ^c				
White	Male and Female	1975-1988	1.5 ^c	1988-2000	-0.3				
White	Male	1975-1989	1.3 ^c	1989-2000	-0.3				
White	Female	1975-1987	1.9 ^c	1987-2000	-0.5				
Black	Male and Female	1975-2000	0.1						
Black	Male	1975-2000	0.0						
Black	Female	1975-2000	0.3						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-2000	0.7 ^c						
White	Female	1975-1987	1.8 ^c	1987-2000	-0.2				
Black	Male	1975-2000	0.1						
Black	Female	1975-2000	0.4						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1977	3.9 ^c	1977-1980	-1.2	1980-1991	1.2 ^c	1991-2000	-0.7 ^c
All Races	Male	1975-1977	4.4	1977-1982	-0.4	1982-1990	1.4 ^c	1990-2000	-0.7 ^c
All Races	Female	1975-1992	0.9 ^c	1992-2000	-0.9 ^c				
White	Male and Female	1975-1992	0.9 ^c	1992-2000	-0.7 ^c				
White	Male	1975-1977	5.0	1977-1981	-0.6	1981-1991	1.2 ^c	1991-2000	-0.7 ^c
White	Female	1975-1992	1.0 ^c	1992-2000	-0.8 ^c				
Black	Male and Female	1975-1984	-0.2	1984-1987	4.3	1987-2000	-0.5		
Black	Male	1975-2000	0.1						
Black	Female	1975-1991	1.3 ^c	1991-2000	-0.4				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Jointpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Brain and Other Nervous System Cancer, by Race and Sex




Figure III-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table IV-1
 FEMALE BREAST CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINPOINTS BY RACE AND AGE

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1980	-0.4	1980-1987	3.7 ^c	1987-2000	0.4 ^c		
All Races	Under 50	1975-1980	-1.4	1980-1986	2.9 ^c	1986-2000	-0.3		
All Races	50 and Over	1975-1982	0.8	1982-1986	5.8 ^c	1986-2000	0.7 ^c		
White	All Ages	1975-1980	-0.3	1980-1987	3.8 ^c	1987-2000	0.4 ^c		
White	Under 50	1975-1980	-1.2	1980-1986	2.8 ^c	1986-2000	-0.3 ^c		
White	50 and Over	1975-1980	-0.1	1980-1987	4.3 ^c	1987-2000	0.7 ^c		
Black	All Ages	1975-1992	2.2 ^c	1992-2000	-0.2				
Black	Under 50	1975-1991	1.5 ^c	1991-2000	-0.9				
Black	50 and Over	1975-1995	2.3 ^c	1995-2000	-1.3				
<u>SEER Delay-Adjusted Incidence^a</u>									
All Races	All Ages	1975-1980	-0.6	1980-1986	4.1 ^c	1986-2000	0.6 ^c		
White	All Ages	1975-1980	-0.5	1980-1987	4.1 ^c	1987-1993	-0.2	1993-2000	1.4 ^c
Black	All Ages	1975-1992	2.3 ^c	1992-2000	0.2				
<u>U.S. Cancer Mortality^b</u>									
All Races	All Ages	1975-1990	0.4 ^c	1990-2000	-2.3 ^c				
All Races	Under 50	1975-1991	-0.4 ^c	1991-2000	-3.7 ^c				
All Races	50 and Over	1975-1990	0.6 ^c	1990-2000	-2.0 ^c				
White	All Ages	1975-1990	0.3 ^c	1990-2000	-2.5 ^c				
White	Under 50	1975-1991	-0.7 ^c	1991-2000	-4.2 ^c				
White	50 and Over	1975-1990	0.5 ^c	1990-2000	-2.2 ^c				
Black	All Ages	1975-1991	1.6 ^c	1991-2000	-1.0 ^c				
Black	Under 50	1975-1987	1.7 ^c	1987-1996	-1.1 ^c	1996-2000	-3.6 ^c		
Black	50 and Over	1975-1993	1.6 ^c	1993-2000	-1.2 ^c				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

Table IV-2
 FEMALE BREAST CANCER (*In Situ*)

TRENDS IN SEER INCIDENCE^a USING THE JOINTPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINTPOINTS BY RACE AND AGE

		<u>Jointpoint Segment 1</u>		<u>Jointpoint Segment 2</u>		<u>Jointpoint Segment 3</u>		<u>Jointpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1982	-0.8	1982-1986	32.3 ^b	1986-2000	6.1 ^b		
All Races	Under 50	1975-1982	-2.3	1982-1986	27.9 ^b	1986-2000	3.8 ^b		
All Races	50 and Over	1975-1981	-1.9	1981-1987	26.0 ^b	1987-2000	6.8 ^b		
White	All Ages	1975-1982	-0.7	1982-1986	33.0 ^b	1986-2000	5.9 ^b		
White	Under 50	1975-1982	-3.9	1982-1985	40.5 ^b	1985-2000	4.0 ^b		
White	50 and Over	1975-1981	-1.3	1981-1987	26.1 ^b	1987-2000	6.5 ^b		
Black	All Ages	1975-1981	2.9	1981-1988	18.9 ^b	1988-2000	7.0 ^b		
Black	Under 50	1975-2000	6.9 ^b						
Black	50 and Over	1975-1980	-2.1	1980-1988	20.8 ^b	1988-2000	7.9 ^b		
<u>SEER Delay-Adjusted Incidence^a</u>									
White	All Ages	1975-1982	-0.7	1982-1986	32.8 ^c	1986-2000	6.0 ^c		
Black	All Ages	1975-1981	2.9	1981-1988	18.8 ^c	1988-2000	7.1 ^c		

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Jointpoint Regression Program.

Trends are from the SEER 9 areas.

The APC is significantly different from zero (p<.05).

^a
^b

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Female Breast Cancer, by Race




Figure IV-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence and Delay Adjusted Incidence Rates+ Female Breast Cancer (*In Situ*), by Race




Figure IV-3

+ Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table V-1
CERVIX UTERI CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINPOINTS BY RACE AND AGE

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1981	-4.5 ^c	1981-1998	-1.2 ^c	1998-2000	-9.3		
All Races	Under 50	1975-1981	-3.6 ^c	1981-1998	-0.5	1998-2000	-10.0		
All Races	50 and Over	1975-1983	-4.9 ^c	1983-1991	-0.9	1991-2000	-3.4 ^c		
White	All Ages	1975-1982	-4.8 ^c	1982-1990	0.4	1990-2000	-2.5 ^c		
White	Under 50	1975-1981	-4.7 ^c	1981-1989	1.5	1989-2000	-1.7 ^c		
White	50 and Over	1975-1985	-4.6 ^c	1985-1991	0.3	1991-2000	-3.9 ^c		
Black	All Ages	1975-2000	-3.8 ^c						
Black	Under 50	1975-2000	-3.2 ^c						
Black	50 and Over	1975-2000	-4.2 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	All Ages	1975-1982	-4.8 ^c	1982-1990	0.4	1990-2000	-2.3 ^c		
Black	All Ages	1975-2000	-3.7 ^c						
<u>U.S. Cancer Mortality^b</u>									
All Races	All Ages	1975-1982	-4.4 ^c	1982-1996	-1.6 ^c	1996-2000	-3.8 ^c		
All Races	Under 50	1975-1981	-3.8 ^c	1981-1997	-0.7 ^c	1997-2000	-6.1 ^c		
All Races	50 and Over	1975-1982	-4.6 ^c	1982-2000	-2.3 ^c				
White	All Ages	1975-1983	-4.2 ^c	1983-1996	-1.3 ^c	1996-2000	-3.4 ^c		
White	Under 50	1975-1979	-5.0 ^c	1979-1998	-0.4 ^c	1998-2000	-7.7		
White	50 and Over	1975-1983	-4.8 ^c	1983-2000	-2.1 ^c				
Black	All Ages	1975-1981	-4.8 ^c	1981-1993	-2.4 ^c	1993-2000	-4.7 ^c		
Black	Under 50	1975-2000	-3.2 ^c						
Black	50 and Over	1975-2000	-3.2 ^c						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Cervix Uteri Cancer, by Race




Figure V-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table VI-1
COLON AND RECTUM CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1985	0.8 ^c	1985-1995	-1.8 ^c	1995-1998	1.1	1998-2000	-3.0 ^c
All Races	Male	1975-1986	1.1 ^c	1986-1995	-2.1 ^c	1995-1998	1.0	1998-2000	-3.6
All Races	Female	1975-1985	0.3 ^c	1985-1995	-1.9 ^c	1995-1998	1.7	1998-2000	-3.0
White	Male and Female	1975-1985	0.8 ^c	1985-1995	-2.0 ^c	1995-1998	1.4	1998-2000	-3.6 ^c
White	Male	1975-1986	1.1 ^c	1986-1995	-2.3 ^c	1995-1998	1.2	1998-2000	-4.3 ^c
White	Female	1975-1985	0.3	1985-1993	-2.2 ^c	1993-2000	-0.2		
Black	Male and Female	1975-1980	3.3 ^c	1980-2000	-0.2				
Black	Male	1975-1980	4.6	1980-2000	-0.1				
Black	Female	1975-2000	-0.1						
<u>SEER Delay-Adjusted Incidence^a</u>									
All Races	Male and Female	1975-1985	0.8 ^c	1985-1995	-1.8 ^c	1995-1998	1.2	1998-2000	-2.3
All Races	Male	1975-1986	1.1 ^c	1986-1995	-2.0 ^c	1995-2000	-0.2		
All Races	Female	1975-1985	0.3	1985-1994	-1.8 ^c	1994-2000	0.2		
White	Male	1975-1986	1.1 ^c	1986-1995	-2.3 ^c	1995-1998	1.4	1998-2000	-3.5
White	Female	1975-1985	0.3	1985-1993	-2.2 ^c	1993-2000	0.0		
Black	Male	1975-1980	4.5	1980-2000	-0.0				
Black	Female	1975-2000	-0.0						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1984	-0.5 ^c	1984-2000	-1.7 ^c				
All Races	Male	1975-1979	0.6	1979-1987	-0.6 ^c	1987-2000	-1.9 ^c		
All Races	Female	1975-1984	-1.0 ^c	1984-2000	-1.8 ^c				
White	Male and Female	1975-1984	-0.7 ^c	1984-2000	-1.9 ^c				
White	Male	1975-1985	-0.2	1985-2000	-2.0 ^c				
White	Female	1975-1984	-1.2 ^c	1984-2000	-2.0 ^c				
Black	Male and Female	1975-1990	0.6 ^c	1990-2000	-0.8 ^c				
Black	Male	1975-1990	1.2 ^c	1990-2000	-0.6 ^c				
Black	Female	1975-1985	0.7 ^c	1985-2000	-0.7 ^c				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

Table VI-2
COLON CANCER (Invasive) AND RECTUM CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>
<u>SEER Cancer Incidence for Colon</u>									
All Races	Male and Female	1975-1985	1.1 ^c	1985-1995	-1.6 ^c	1995-2000	-0.5		
All Races	Male	1975-1985	1.7 ^c	1985-2000	-1.5 ^c				
All Races	Female	1975-1983	0.7	1983-2000	-1.1 ^c				
White	Male and Female	1975-1985	1.1 ^c	1985-1995	-1.9 ^c	1995-1998	1.0	1998-2000	-3.6 ^c
White	Male	1975-1985	1.7 ^c	1985-2000	-1.6 ^c				
White	Female	1975-1985	0.6 ^c	1985-1993	-2.0 ^c	1993-2000	-0.3		
Black	Male and Female	1975-1980	3.7 ^c	1980-2000	-0.2				
Black	Male	1975-1989	1.6 ^c	1989-2000	-1.0				
Black	Female	1975-2000	0.0						
<u>SEER Cancer Incidence for Rectum</u>									
All Races	Male and Female	1975-1987	-0.3	1987-1994	-2.4 ^c	1994-2000	0.4		
All Races	Male	1975-1990	-0.3	1990-1993	-4.9	1993-2000	0.1		
All Races	Female	1975-1984	-0.2	1984-1995	-2.2 ^c	1995-1998	3.3	1998-2000	-4.7
White	Male and Female	1975-1987	-0.3	1987-1994	-2.6 ^c	1994-2000	0.4		
White	Male	1975-1990	-0.3	1990-1993	-5.5	1993-2000	0.1		
White	Female	1975-1984	-0.1	1984-1995	-2.4 ^c	1995-1998	3.5	1998-2000	-5.2
Black	Male and Female	1975-2000	-0.1						
Black	Male	1975-2000	0.1						
Black	Female	1975-2000	-0.3						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

*f The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Colon and Rectum Cancer, by Race and Sex




Figure VI-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table VIII-1
ESOPHAGUS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-2000	0.6 ^c						
All Races	Male	1975-2000	0.8 ^c						
All Races	Female	1975-2000	-0.2						
White	Male and Female	1975-1981	-0.6	1981-2000	1.6 ^c				
White	Male	1975-2000	1.7 ^c						
White	Female	1975-2000	0.1						
Black	Male and Female	1975-1986	1.0	1986-2000	-4.3 ^c				
Black	Male	1975-1986	1.2	1986-2000	-4.8 ^c				
Black	Female	1975-1980	9.4	1980-2000	-2.3 ^c				
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-2000	1.7 ^c						
White	Female	1975-2000	0.2						
Black	Male	1975-1986	1.1	1986-2000	-4.8 ^c				
Black	Female	1975-1980	9.3	1980-2000	-2.2 ^c				
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-2000	0.7 ^c						
All Races	Male	1975-1985	0.7 ^c	1985-1994	1.2 ^c	1994-2000	0.5 ^c		
All Races	Female	1975-2000	0.1						
White	Male and Female	1975-1983	0.4 ^c	1983-2000	1.7 ^c				
White	Male	1975-1982	0.4	1982-2000	1.9 ^c				
White	Female	1975-2000	0.4 ^c						
Black	Male and Female	1975-1982	1.1 ^c	1982-1993	-1.6 ^c	1993-2000	-4.7 ^c		
Black	Male	1975-1984	0.8 ^c	1984-1994	-2.0 ^c	1994-2000	-5.5 ^c		
Black	Female	1975-1990	-0.1	1990-2000	-3.1 ^c				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Esophagus Cancer, by Race and Sex




Figure VIII-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XI-1
KIDNEY AND RENAL PELVIS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1990	2.4 ^c	1990-2000	1.2 ^c				
All Races	Male	1975-1989	2.2 ^c	1989-2000	1.0 ^c				
All Races	Female	1975-1992	2.7 ^c	1992-2000	1.0				
White	Male and Female	1975-1990	2.4 ^c	1990-2000	1.1 ^c				
White	Male	1975-2000	1.6 ^c						
White	Female	1975-1992	2.8 ^c	1992-2000	0.7				
Black	Male and Female	1975-2000	2.9 ^c						
Black	Male	1975-2000	2.8 ^c						
Black	Female	1975-2000	3.3 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-2000	1.7 ^c						
White	Female	1975-1991	2.8 ^c	1991-2000	1.3 ^c				
Black	Male	1975-2000	3.0 ^c						
Black	Female	1975-2000	3.5 ^c						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1992	1.1 ^c	1992-2000	-0.3				
All Races	Male	1975-1991	1.1 ^c	1991-2000	-0.2				
All Races	Female	1975-1995	1.1 ^c	1995-2000	-1.6 ^c				
White	Male and Female	1975-1992	1.1 ^c	1992-2000	-0.2				
White	Male	1975-1992	1.0 ^c	1992-2000	-0.3				
White	Female	1975-1995	1.1 ^c	1995-2000	-1.4				
Black	Male and Female	1975-1991	2.7 ^c	1991-2000	-0.1				
Black	Male	1975-1990	3.0 ^c	1990-2000	0.1				
Black	Female	1975-1995	2.4 ^c	1995-2000	-2.3				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Kidney and Renal Pelvis Cancer, by Race and Sex




Figure XI-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XIII-1
LEUKEMIA (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1995	0.0	1995-2000	-1.8 ^c				
All Races	Male	1975-1997	-0.2	1997-2000	-3.4				
All Races	Female	1975-2000	-0.2 ^c						
White	Male and Female	1975-1995	0.1	1995-2000	-1.9 ^c				
White	Male	1975-1997	-0.1	1997-2000	-4.1				
White	Female	1975-2000	-0.1						
Black	Male and Female	1975-2000	-0.6 ^c						
Black	Male	1975-2000	-0.5						
Black	Female	1975-2000	-0.7 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-2000	0.3 ^c						
White	Female	1975-2000	0.5 ^c						
Black	Male	1975-2000	0.0						
Black	Female	1975-2000	-0.1						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1980	0.3	1980-2000	-0.3 ^c				
All Races	Male	1975-2000	-0.3 ^c						
All Races	Female	1975-1980	0.7	1980-2000	-0.4 ^c				
White	Male and Female	1975-2000	-0.3 ^c						
White	Male	1975-2000	-0.3 ^c						
White	Female	1975-2000	-0.3 ^c						
Black	Male and Female	1975-1991	0.6 ^c	1991-2000	-1.0 ^c				
Black	Male	1975-1990	1.1 ^c	1990-2000	-0.9 ^c				
Black	Female	1975-1983	1.3	1983-2000	-0.4 ^c				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Leukemia, by Race and Sex




Figure XIII-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XIV-1
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1984	1.6	1984-2000	4.3 ^c				
All Races	Male	1975-2000	3.7 ^c						
All Races	Female	1975-2000	3.3 ^c						
White	Male and Female	1975-1984	0.9	1984-2000	4.4 ^c				
White	Male	1975-1984	1.3	1984-2000	4.5 ^c				
White	Female	1975-1984	0.7	1984-2000	3.6 ^c				
Black	Male and Female	1975-2000	2.8 ^c						
Black	Male	1975-2000	3.0 ^c						
Black	Female	1975-2000	2.6 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-1987	2.0 ^c	1987-2000	5.2 ^c				
White	Female	1975-1985	0.9	1985-2000	4.0 ^c				
Black	Male	1975-2000	3.2 ^c						
Black	Female	1975-2000	2.8 ^c						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1980	0.2	1980-1987	2.0 ^c	1987-1996	3.8 ^c	1996-2000	0.8
All Races	Male	1975-1985	1.5 ^c	1985-1996	3.7 ^c	1996-2000	1.0		
All Races	Female	1975-1987	0.8 ^c	1987-1995	3.8 ^c	1995-2000	0.3		
White	Male and Female	1975-1978	-1.2	1978-1987	1.7 ^c	1987-1996	3.9 ^c	1996-2000	0.7
White	Male	1975-1984	1.1 ^c	1984-1996	3.7 ^c	1996-2000	1.0		
White	Female	1975-1987	0.6 ^c	1987-1996	3.5 ^c	1996-2000	-0.5		
Black	Male and Female	1975-1988	0.8 ^c	1988-1995	3.2 ^c	1995-2000	0.4		
Black	Male	1975-2000	1.6 ^c						
Black	Female	1975-2000	1.8 ^c						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence and Delay Adjusted Incidence Rates+ Liver and Intrahepatic Bile Duct Cancer, by Race and Sex




Figure XIV-1

+ Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XV-1
LUNG AND BRONCHUS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1982	2.5 ^c	1982-1991	1.0 ^c	1991-2000	-0.9 ^c		
All Races	Male	1975-1982	1.5 ^c	1982-1992	-0.5 ^c	1992-2000	-2.2 ^c		
All Races	Female	1975-1988	4.6 ^c	1988-1998	1.3 ^c	1998-2000	-2.8		
White	Male and Female	1975-1988	1.9 ^c	1988-1998	-0.3	1998-2000	-3.2		
White	Male	1975-1981	1.5 ^c	1981-1991	-0.3	1991-2000	-2.1 ^c		
White	Female	1975-1988	4.8 ^c	1988-1998	1.4 ^c	1998-2000	-2.9		
Black	Male and Female	1975-1984	3.5 ^c	1984-2000	-0.6 ^c				
Black	Male	1975-1984	3.1 ^c	1984-2000	-1.6 ^c				
Black	Female	1975-1990	4.3 ^c	1990-2000	0.4				
<u>SEER Delay-Adjusted Incidence^a</u>									
All Races	Male and Female	1975-1982	2.5 ^c	1982-1991	1.0 ^c	1991-2000	-0.7 ^c		
All Races	Male	1975-1982	1.4 ^c	1982-1991	-0.4	1991-2000	-1.8 ^c		
All Races	Female	1975-1982	5.6 ^c	1982-1991	3.4 ^c	1991-2000	0.7 ^c		
White	Male	1975-1981	1.5 ^c	1981-1990	-0.2	1990-2000	-1.8 ^c		
White	Female	1975-1982	5.6 ^c	1982-1988	4.2 ^c	1988-1998	1.6 ^c	1998-2000	-1.6
Black	Male	1975-1984	3.0 ^c	1984-2000	-1.5 ^c				
Black	Female	1975-1990	4.2 ^c	1990-2000	0.6				
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1982	2.7 ^c	1982-1991	1.6 ^c	1991-2000	-0.7 ^c		
All Races	Male	1975-1982	1.8 ^c	1982-1991	0.4 ^c	1991-2000	-1.8 ^c		
All Races	Female	1975-1983	5.9 ^c	1983-1992	3.8 ^c	1992-2000	0.6 ^c		
White	Male and Female	1975-1982	2.6 ^c	1982-1991	1.7 ^c	1991-2000	-0.6 ^c		
White	Male	1975-1982	1.7 ^c	1982-1991	0.4 ^c	1991-2000	-1.7 ^c		
White	Female	1975-1983	5.9 ^c	1983-1992	3.8 ^c	1992-2000	0.6 ^c		
Black	Male and Female	1975-1982	3.3 ^c	1982-1991	1.7 ^c	1991-2000	-1.3 ^c		
Black	Male	1975-1981	3.1 ^c	1981-1989	1.6 ^c	1989-1993	-0.7	1993-2000	-2.5 ^c
Black	Female	1975-1981	6.7 ^c	1981-1991	4.0 ^c	1991-2000	0.8 ^c		

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

Table XV-2
 SMALL CELL LUNG AND BRONCHUS CANCER (Invasive) AND NON-SMALL CELL LUNG AND BRONCHUS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence for Small Cell Lung and Bronchus^a</u>									
All Races	Male and Female	1975-1982	6.2 ^c	1982-1989	1.2	1989-2000	-2.4 ^c		
All Races	Male	1975-1980	6.4 ^c	1980-1988	0.5	1988-2000	-3.5 ^c		
All Races	Female	1975-1982	9.6 ^c	1982-1991	3.0 ^c	1991-2000	-1.7 ^c		
White	Male and Female	1975-1982	6.3 ^c	1982-1989	1.4 ^c	1989-2000	-2.2 ^c		
White	Male	1975-1980	6.2 ^c	1980-1988	0.6	1988-2000	-3.4 ^c		
White	Female	1975-1982	9.8 ^c	1982-1989	3.9 ^c	1989-2000	-0.9 ^c		
Black	Male and Female	1975-1986	4.6 ^c	1986-2000	-2.1 ^c				
Black	Male	1975-1985	4.8 ^c	1985-2000	-3.2 ^c				
Black	Female	1975-2000	2.1 ^c						
<u>SEER Cancer Incidence for Non-Small Cell Lung and Bronchus^a</u>									
All Races	Male and Female	1975-1988	1.5 ^c	1988-1998	0.0	1998-2000	-3.2		
All Races	Male	1975-1981	1.1 ^c	1981-1992	-0.3	1992-2000	-1.9 ^c		
All Races	Female	1975-1988	4.3 ^c	1988-1998	1.7 ^c	1998-2000	-2.6		
White	Male and Female	1975-1988	1.6 ^c	1988-1998	0.1	1998-2000	-3.3		
White	Male	1975-1990	0.2	1990-2000	-1.8 ^c				
White	Female	1975-1988	4.4 ^c	1988-1998	1.9 ^c	1998-2000	-2.7		
Black	Male and Female	1975-1984	3.2 ^c	1984-2000	-0.5 ^c				
Black	Male	1975-1984	2.8 ^c	1984-2000	-1.4 ^c				
Black	Female	1975-1990	4.2 ^c	1990-2000	0.6				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Lung and Bronchus Cancer, by Race and Sex




Figure XV-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XVI-1
MELANOMA OF THE SKIN (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>	<u>Year</u>	<u>APC</u>
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1981	6.0 ^c	1981-2000	2.7 ^c				
All Races	Male	1975-1985	5.6 ^c	1985-2000	3.1 ^c				
All Races	Female	1975-1981	5.3 ^c	1981-2000	2.1 ^c				
White	Male and Female	1975-1981	6.3 ^c	1981-2000	3.0 ^c				
White	Male	1975-1985	5.8 ^c	1985-2000	3.3 ^c				
White	Female	1975-1981	6.1 ^c	1981-1994	2.0 ^c	1994-1997	5.8	1997-2000	-0.5
Black	Male and Female	1975-2000	0.3						
Black	Male	1975-2000	2.1						
Black	Female	1975-2000	-1.2						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-1980	7.5 ^c	1980-2000	4.3 ^c				
White	Female	1975-1981	5.9 ^c	1981-1993	2.4 ^c	1993-2000	4.6 ^c		
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1989	1.6 ^c	1989-2000	-0.1				
All Races	Male	1975-1987	2.4 ^c	1987-1998	0.8 ^c	1998-2000	-2.9		
All Races	Female	1975-1988	0.8 ^c	1988-2000	-0.6 ^c				
White	Male and Female	1975-1989	1.7 ^c	1989-2000	0.0				
White	Male	1975-1987	2.6 ^c	1987-1998	0.9 ^c	1998-2000	-2.9		
White	Female	1975-1988	1.0 ^c	1988-2000	-0.4				
Black	Male and Female	1975-2000	-0.3						
Black	Male	1975-2000	-1.0 ^c						
Black	Female	1975-2000	0.1						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Melanoma of the Skin, White, by Sex




Figure XVI-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XVIII-1
NON-HODGKIN LYMPHOMA (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1990	3.6 ^c	1990-1995	1.5	1995-2000	-0.7		
All Races	Male	1975-1991	4.3 ^c	1991-2000	-0.1				
All Races	Female	1975-1988	3.0 ^c	1988-1998	1.5 ^c	1998-2000	-2.8		
White	Male and Female	1975-1991	3.7 ^c	1991-2000	0.2				
White	Male	1975-1991	4.3 ^c	1991-2000	-0.2				
White	Female	1975-1990	2.9 ^c	1990-2000	0.9 ^c				
Black	Male and Female	1975-1995	4.6 ^c	1995-2000	-3.8 ^c				
Black	Male	1975-1995	5.6 ^c	1995-2000	-5.3				
Black	Female	1975-2000	2.6 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-1990	4.5 ^c	1990-2000	0.6 ^c				
White	Female	1975-1988	3.1 ^c	1988-2000	1.6 ^c				
Black	Male	1975-1995	5.7 ^c	1995-2000	-4.5				
Black	Female	1975-2000	2.8 ^c						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1991	2.5 ^c	1991-1997	1.6 ^c	1997-2000	-2.9 ^c		
All Races	Male	1975-1996	2.5 ^c	1996-2000	-2.2 ^c				
All Races	Female	1975-1996	2.2 ^c	1996-2000	-2.1 ^c				
White	Male and Female	1975-1996	2.3 ^c	1996-2000	-2.2 ^c				
White	Male	1975-1995	2.5 ^c	1995-2000	-1.2 ^c				
White	Female	1975-1996	2.2 ^c	1996-2000	-2.1 ^c				
Black	Male and Female	1975-1997	2.7 ^c	1997-2000	-6.0 ^c				
Black	Male	1975-1997	2.8 ^c	1997-2000	-6.7				
Black	Female	1975-1997	2.7 ^c	1997-2000	-5.1 ^c				


The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Non-Hodgkin Lymphoma, by Race and Sex



+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XX-1
OVARY CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINTPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINTPOINTS BY RACE AND AGE

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1986	0.0	1986-1989	4.7	1989-2000	-0.7 ^c		
All Races	Under 65	1975-1986	-1.0 ^c	1986-1989	5.3	1989-2000	-0.6 ^c		
All Races	65 and Over	1975-1991	1.7 ^c	1991-2000	-1.1 ^c				
White	All Ages	1975-1986	0.1	1986-1989	4.9	1989-2000	-0.7 ^c		
White	Under 65	1975-1986	-0.8 ^c	1986-1989	5.5	1989-2000	-0.6		
White	65 and Over	1975-1991	1.8 ^c	1991-2000	-0.9				
Black	All Ages	1975-2000	0.4						
Black	Under 65	1975-2000	-0.3						
Black	65 and Over	1975-2000	1.0 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	All Ages	1975-1986	0.1	1986-1989	4.8	1989-2000	-0.4		
Black	All Ages	1975-2000	0.5 ^c						
<u>U.S. Cancer Mortality^b</u>									
All Races	All Ages	1975-1983	-1.1 ^c	1983-1991	0.5	1991-2000	-0.8 ^c		
All Races	Under 65	1975-1980	-3.3 ^c	1980-2000	-1.5 ^c				
All Races	65 and Over	1975-1985	0.7 ^c	1985-1991	1.9 ^c	1991-2000	-0.3		
White	All Ages	1975-1981	-1.5 ^c	1981-1991	0.3	1991-2000	-0.7 ^c		
White	Under 65	1975-1980	-3.3 ^c	1980-2000	-1.4 ^c				
White	65 and Over	1975-1984	0.5 ^c	1984-1991	1.7 ^c	1991-2000	-0.1		
Black	All Ages	1975-1992	0.4 ^c	1992-2000	-1.0				
Black	Under 65	1975-2000	-1.7 ^c						
Black	65 and Over	1975-1992	2.3 ^c	1992-2000	-1.2				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Ovary Cancer, by Race




Figure XX-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XXI-1
PANCREAS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1978	-2.1	1978-1983	1.1	1983-2000	-0.5 ^c		
All Races	Male	1975-2000	-0.7 ^c						
All Races	Female	1975-1984	1.4 ^c	1984-2000	-0.5 ^c				
White	Male and Female	1975-2000	-0.3 ^c						
White	Male	1975-1978	-3.3	1978-2000	-0.6 ^c				
White	Female	1975-1984	1.3 ^c	1984-2000	-0.5 ^c				
Black	Male and Female	1975-2000	-0.2						
Black	Male	1975-2000	-0.5						
Black	Female	1975-2000	0.1						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-1981	-2.3 ^c	1981-1984	1.7	1984-1993	-1.4 ^c	1993-2000	0.7
White	Female	1975-1984	1.2 ^c	1984-2000	-0.4 ^c				
Black	Male	1975-2000	-0.4						
Black	Female	1975-2000	0.2						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-2000	-0.1 ^c						
All Races	Male	1975-1986	-0.8 ^c	1986-2000	-0.3 ^c				
All Races	Female	1975-1984	0.8 ^c	1984-2000	0.1				
White	Male and Female	1975-1997	-0.2 ^c	1997-2000	0.7				
White	Male	1975-1986	-1.0 ^c	1986-2000	-0.2 ^c				
White	Female	1975-1984	0.6 ^c	1984-2000	0.1				
Black	Male and Female	1975-1989	1.2 ^c	1989-2000	-0.6 ^c				
Black	Male	1975-1990	0.6 ^c	1990-2000	-1.0 ^c				
Black	Female	1975-1986	2.3 ^c	1986-2000	-0.1				

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Pancreas Cancer, by Race and Sex




Figure XXI-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XXII-1
PROSTATE CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINTPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINTPOINTS BY RACE AND AGE

		<u>Jointpoint Segment 1</u>		<u>Jointpoint Segment 2</u>		<u>Jointpoint Segment 3</u>		<u>Jointpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1988	2.6 ^c	1988-1992	16.4 ^c	1992-1995	-11.6 ^c	1995-2000	1.6 ^c
All Races	Under 65	1975-1985	2.4 ^c	1985-1989	7.6 ^c	1989-1992	25.8 ^c	1992-2000	3.0 ^c
All Races	65 and Over	1975-1988	2.5 ^c	1988-1992	15.1 ^c	1992-1995	-14.9 ^c	1995-2000	0.6
White	All Ages	1975-1988	2.8 ^c	1988-1992	16.2 ^c	1992-1995	-12.6 ^c	1995-2000	1.7 ^c
White	Under 65	1975-1985	2.5 ^c	1985-1989	8.8 ^c	1989-1992	24.8 ^c	1992-2000	2.6 ^c
White	65 and Over	1975-1988	2.7 ^c	1988-1992	14.8 ^c	1992-1995	-15.8 ^c	1995-2000	0.8
Black	All Ages	1975-1989	2.0 ^c	1989-1992	21.5 ^c	1992-1996	-5.4 ^c	1996-2000	0.5
Black	Under 65	1975-1989	2.0 ^c	1989-1993	25.7 ^c	1993-2000	2.2 ^c		
Black	65 and Over	1975-1989	2.0 ^c	1989-1992	20.2 ^c	1992-1995	-10.3 ^c	1995-2000	-1.7
<u>SEER Delay-Adjusted Incidence^a</u>									
All Races	All Ages	1975-1988	2.6 ^c	1988-1992	16.6 ^c	1992-1995	-11.6 ^c	1995-2000	2.3 ^c
White	All Ages	1975-1988	2.8 ^c	1988-1992	16.5 ^c	1992-1995	-12.4 ^c	1995-2000	3.0 ^c
Black	All Ages	1975-1989	2.0 ^c	1989-1992	22.0 ^c	1992-1996	-5.0 ^c	1996-2000	2.3 ^c
<u>U.S. Cancer Mortality^b</u>									
All Races	All Ages	1975-1987	0.9 ^c	1987-1991	3.1 ^c	1991-1994	-0.6	1994-2000	-4.0 ^c
All Races	Under 65	1975-1987	0.5 ^c	1987-1990	3.1	1990-1996	-3.2 ^c	1996-2000	-5.6 ^c
All Races	65 and Over	1975-1987	0.9 ^c	1987-1991	3.2 ^c	1991-1994	-0.4	1994-2000	-4.0 ^c
White	All Ages	1975-1987	0.8 ^c	1987-1991	3.1 ^c	1991-1994	-0.8	1994-2000	-4.2 ^c
White	Under 65	1975-1987	0.7 ^c	1987-1990	3.7	1990-2000	-4.4 ^c		
White	65 and Over	1975-1987	0.8 ^c	1987-1991	3.2 ^c	1991-1994	-0.6	1994-2000	-4.1 ^c
Black	All Ages	1975-1988	1.9 ^c	1988-1993	3.2 ^c	1993-2000	-2.6 ^c		
Black	Under 65	1975-1992	0.6 ^c	1992-2000	-3.0 ^c				
Black	65 and Over	1975-1988	2.1 ^c	1988-1993	3.4 ^c	1993-2000	-2.5 ^c		

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Jointpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Prostate Cancer, by Race




Figure XXII-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XXIII-1
STOMACH CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-2000	-1.6 ^c						
All Races	Male	1975-1988	-1.1 ^c	1988-2000	-2.1 ^c				
All Races	Female	1975-2000	-1.7 ^c						
White	Male and Female	1975-2000	-1.9 ^c						
White	Male	1975-2000	-1.9 ^c						
White	Female	1975-2000	-2.2 ^c						
Black	Male and Female	1975-2000	-1.1 ^c						
Black	Male	1975-2000	-1.2 ^c						
Black	Female	1975-2000	-0.8 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-2000	-1.8 ^c						
White	Female	1975-2000	-2.1 ^c						
Black	Male	1975-2000	-1.1 ^c						
Black	Female	1975-2000	-0.8 ^c						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1987	-2.5 ^c	1987-1990	-0.4	1990-2000	-2.9 ^c		
All Races	Male	1975-1987	-2.4 ^c	1987-1990	-0.4	1990-2000	-3.3 ^c		
All Races	Female	1975-1987	-2.8 ^c	1987-1990	-0.6	1990-2000	-2.5 ^c		
White	Male and Female	1975-1987	-2.8 ^c	1987-1990	-0.8	1990-2000	-3.2 ^c		
White	Male	1975-1994	-2.3 ^c	1994-2000	-3.9 ^c				
White	Female	1975-1977	-4.7	1977-2000	-2.6 ^c				
Black	Male and Female	1975-1987	-1.6 ^c	1987-1990	0.9	1990-2000	-2.5 ^c		
Black	Male	1975-1991	-1.1 ^c	1991-2000	-2.6 ^c				
Black	Female	1975-2000	-1.2 ^c						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Stomach Cancer, by Race and Sex




Figure XXIII-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table XXVI-1
URINARY BLADDER CANCER (Invasive and In Situ)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-1987	0.7 ^c	1987-2000	0.0				
All Races	Male	1975-1987	0.9 ^c	1987-2000	-0.2				
All Races	Female	1975-2000	0.2 ^c						
White	Male and Female	1975-1987	0.9 ^c	1987-2000	0.1				
White	Male	1975-1987	1.1 ^c	1987-2000	-0.2				
White	Female	1975-2000	0.3 ^c						
Black	Male and Female	1975-2000	0.3						
Black	Male	1975-2000	0.4						
Black	Female	1975-2000	0.5						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-1987	1.2 ^c	1987-1996	-0.5	1996-2000	2.1 ^c		
White	Female	1975-2000	0.4 ^c						
Black	Male	1975-2000	0.5						
Black	Female	1975-2000	0.6						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1977	0.0	1977-1987	-2.1 ^c	1987-2000	-0.2 ^c		
All Races	Male	1975-1982	-1.4 ^c	1982-1987	-2.3 ^c	1987-2000	-0.3 ^c		
All Races	Female	1975-1986	-1.7 ^c	1986-2000	-0.3 ^c				
White	Male and Female	1975-1977	0.1	1977-1987	-2.1 ^c	1987-2000	0.0		
White	Male	1975-1988	-1.7 ^c	1988-2000	-0.2 ^c				
White	Female	1975-1987	-1.7 ^c	1987-2000	-0.1				
Black	Male and Female	1975-2000	-1.1 ^c						
Black	Male	1975-2000	-1.0 ^c						
Black	Female	1975-2000	-0.8 ^c						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Urinary Bladder Cancer, by Race and Sex




Figure XXVI-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).