

Table C1
Adult Self-Reported Current Asthma Prevalence Rate (Percent) and
Prevalence (Number) by State: BRFSS 1999

Note 1: CI designates confidence interval.

State	Sample Size	Prevalence (percent)	Standard Error	Lower 95% CI	Upper 95% CI	Prevalence (number)	Lower 95% CI	Upper 95% CI
AZ	1,738	5.5	0.81	3.9	7.1	189,607	134,159	245,056
AR	2,991	7.2	0.57	6.1	8.4	137,398	115,727	159,068
DC	1,257	6.8	0.77	5.2	8.3	27,452	21,295	33,609
ID	4,937	7.9	0.52	6.9	8.9	68,682	59,617	77,747
IA	3,499	7.7	0.55	6.6	8.8	158,920	136,186	181,653
LA	1,659	4.9	0.64	3.6	6.1	152,370	112,649	192,091
ME	1,662	8.3	0.79	6.7	9.8	76,773	62,099	91,446
MS	2,137	5.0	0.53	3.9	6.0	96,772	76,566	116,978
MO	4,220	7.3	0.55	6.2	8.4	293,634	249,546	337,721
NE	2,709	6.5	0.51	5.5	7.5	75,896	64,117	87,675
NH	1,205	8.8	0.93	7.0	10.6	76,086	60,220	91,953
PA	3,479	6.8	0.51	5.8	7.8	608,516	517,269	699,762
SC	3,395	5.8	0.45	4.9	6.7	161,387	136,626	186,149
UT	3,180	7.5	0.58	6.4	8.7	103,206	87,600	118,813
VT	3,091	8.0	0.58	6.9	9.1	34,752	29,704	39,801
WA	3,525	7.0	0.46	6.1	7.9	292,596	255,856	329,336
WI	2,166	7.3	0.67	6.0	8.6	281,522	230,367	332,677
WY	2,395	7.1	0.56	6.0	8.2	24,221	20,480	27,961

Footnote 1: If values for the standard error and the 95% confidence intervals are not provided, the normal approximation to the binomial distribution does not apply because of the small sample size.

Footnote 2: When the sample size is less than 50, estimates are not precise and should be interpreted with caution.