

Table C2.1
Adult Self-Reported Current Asthma Prevalence Rate (Percent)
by Sex and State: BRFSS 1999

Note 1: CI designates confidence interval.

State	Male					Female				
	Sample Size	Prevalence (percent)	Standard Error	Lower 95% CI	Upper 95% CI	Sample Size	Prevalence (percent)	Standard Error	Lower 95% CI	Upper 95% CI
AZ	710	4.5	1.17	2.2	6.8	1,028	6.4	1.11	4.2	8.6
AR	1,165	5.3	0.82	3.7	6.9	1,826	9.0	0.79	7.5	10.6
DC	456	5.0	1.12	2.8	7.2	801	8.3	1.06	6.2	10.4
ID	2,084	6.8	0.79	5.3	8.4	2,853	8.9	0.67	7.6	10.2
IA	1,375	6.2	0.75	4.7	7.6	2,124	9.1	0.79	7.5	10.6
LA	659	3.9	1.00	1.9	5.8	1,000	5.8	0.82	4.2	7.4
ME	732	7.2	1.13	5.0	9.5	930	9.2	1.11	7.0	11.4
MS	835	4.3	0.84	2.6	5.9	1,302	5.6	0.66	4.3	6.9
MO	1,696	6.4	0.88	4.7	8.1	2,524	8.1	0.69	6.7	9.5
NE	1,074	5.2	0.72	3.8	6.6	1,635	7.6	0.72	6.2	9.1
NH	485	5.2	1.15	2.9	7.5	720	12.1	1.42	9.4	14.9
PA	1,388	5.3	0.80	3.8	6.9	2,091	8.1	0.65	6.8	9.3
SC	1,365	4.4	0.67	3.1	5.7	2,030	7.0	0.61	5.8	8.2
UT	1,368	5.9	0.80	4.3	7.5	1,812	9.1	0.85	7.4	10.7
VT	1,245	6.2	0.76	4.7	7.7	1,846	9.7	0.87	8.0	11.4
WA	1,570	4.8	0.56	3.7	5.9	1,955	9.2	0.71	7.9	10.6
WI	912	5.8	0.87	4.1	7.5	1,254	8.7	1.00	6.7	10.6
WY	1,061	5.2	0.71	3.8	6.6	1,334	8.9	0.86	7.2	10.6

Footnote 1: If values for the standard error and the 95% confidence intervals are not provided, the normal approximation to the binomial distribution does not apply because of the small sample size.

Footnote 2: When the sample size is less than 50, estimates are not precise and should be interpreted with caution.