

Table L1
Adult Self-Reported Lifetime Asthma Prevalence Rate (Percent) and
Prevalence (Number) by State: BRFSS 1999

Note 1: CI designates confidence interval.

State	Sample Size	Prevalence (percent)	Standard Error	Lower 95% CI	Upper 95% CI	Prevalence (number)	Lower 95% CI	Upper 95% CI
AZ	1,739	7.7	0.94	5.9	9.6	267,148	202,499	331,798
AR	2,996	11.3	0.67	10.0	12.6	214,197	188,561	239,832
DC	1,258	11.7	0.99	9.7	13.6	47,400	39,528	55,273
ID	4,951	11.3	0.58	10.2	12.5	99,047	88,698	109,397
IA	3,517	11.6	0.66	10.3	12.9	241,051	213,363	268,739
LA	1,660	9.1	0.85	7.5	10.8	284,982	231,926	338,038
ME	1,664	12.7	0.97	10.8	14.6	118,730	100,592	136,868
MS	2,142	9.4	0.76	7.9	10.9	183,093	153,483	212,702
MO	4,228	11.1	0.69	9.7	12.4	445,840	390,179	501,501
NE	2,712	9.6	0.62	8.4	10.9	113,530	99,091	127,968
NH	1,213	14.3	1.23	11.9	16.7	124,914	103,063	146,765
PA	3,489	10.7	0.62	9.5	11.9	965,473	852,342	1,078,604
SC	3,405	9.7	0.60	8.5	10.9	272,048	238,455	305,641
UT	3,185	12.4	0.76	11.0	13.9	171,211	150,865	191,557
VT	3,094	11.7	0.71	10.4	13.2	51,170	44,847	57,494
WA	3,539	11.9	0.62	10.7	13.1	497,677	446,562	548,793
WI	2,171	10.2	0.76	8.7	11.7	393,247	334,849	451,644
WY	2,399	10.5	0.67	9.2	11.8	36,015	31,487	40,542

Footnote 1: If values for the standard error and the 95% confidence intervals are not provided, the normal approximation to the binomial distribution does not apply because of the small sample size.

Footnote 2: When the sample size is less than 50, estimates are not precise and should be interpreted with caution.