

Table 1. Estimated numbers of cases of HIV/AIDS, by year of diagnosis and selected characteristics, 2001–2005—33 states and U.S. dependent areas with confidential name-based HIV infection reporting (Revised June 2007)

	Year of diagnosis				
	2001	2002	2003	2004	2005
Data for 33 states					
Age at diagnosis (yrs)					
<13	368	294	212	179	168
13–14	53	46	51	35	43
15–19	1,010	1,007	1,007	1,012	1,213
20–24	3,184	3,076	3,223	3,486	3,876
25–29	4,482	4,325	4,064	4,207	4,581
30–34	6,557	6,007	5,274	5,031	5,123
35–39	7,956	7,305	6,584	6,025	6,123
40–44	6,441	6,074	5,956	5,662	6,054
45–49	4,564	4,234	4,170	4,063	4,396
50–54	2,604	2,506	2,524	2,499	2,790
55–59	1,356	1,267	1,310	1,416	1,535
60–64	711	724	665	735	768
≥65	659	608	592	664	660
Race/ethnicity					
White, not Hispanic	10,925	10,723	10,329	10,639	11,559
Black, not Hispanic	20,868	19,372	18,087	17,389	18,121
Hispanic	7,400	6,654	6,534	6,251	6,782
Asian/Pacific Islander	266	305	344	351	417
American Indian/Alaska Native	162	192	182	188	195
Transmission category					
Male adult or adolescent					
Male-to-male sexual contact	16,167	16,037	15,693	16,469	18,296
Injection drug use	4,837	4,048	3,726	3,340	3,441
Male-to-male sexual contact and injection drug use	1,532	1,432	1,330	1,311	1,324
High-risk heterosexual contact ^a	4,938	4,657	4,407	4,107	4,255
Other ^b	161	155	139	125	139
Subtotal	27,635	26,329	25,295	25,352	27,455
Female adult or adolescent					
Injection drug use	2,769	2,269	2,104	1,942	1,851
High-risk heterosexual contact ^a	9,014	8,433	7,871	7,413	7,734
Other ^b	158	147	147	125	124
Subtotal	11,941	10,849	10,122	9,481	9,708
Child (<13 yrs at diagnosis)					
Perinatal	317	249	188	155	142
Other ^c	51	45	26	25	26
Subtotal	368	294	214	180	168
Subtotal for 33 states	39,944	37,472	35,631	35,012	37,331
Data for U.S. dependent areas	62	57	46	41	36
Total^d	40,007	37,529	35,678	35,053	37,367

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

Data include persons with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later diagnosis of AIDS, or concurrent diagnoses of HIV infection and AIDS.

See Technical Notes for the list of areas that have had laws or regulations requiring confidential name-based HIV infection reporting since at least 2001.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.