

Table 4. Estimated numbers of AIDS cases in children <13 years of age, by year of diagnosis and selected characteristics, 2001–2005 and cumulative—50 states and the District of Columbia (Revised June 2007)


	Year of diagnosis					Cumulative ^a
	2001	2002	2003	2004	2005	
Race/ethnicity						
White, not Hispanic	13	14	12	6	6	1,613
Black, not Hispanic	85	72	46	31	46	5,631
Hispanic	23	16	11	9	13	1,738
Asian/Pacific Islander	1	1	0	1	1	54
American Indian/Alaska Native	0	1	0	1	0	32
Transmission category						
Hemophilia/coagulation disorder	0	0	0	0	0	226
Mother with documented HIV infection or 1 of the following risk factors	118	103	71	50	67	8,460
Injection drug use	14	11	8	6	4	3,198
Sex with injection drug user	10	4	6	2	1	1,388
Sex with bisexual male	4	2	0	2	1	203
Sex with person with hemophilia	1	0	0	0	0	36
Sex with HIV-infected transfusion recipient	0	0	0	0	0	22
Sex with HIV-infected person, risk factor not specified	40	36	20	20	29	1,508
Receipt of blood transfusion, blood components, or tissue	1	2	1	0	0	143
Has HIV infection, risk factor not specified	49	47	35	20	31	1,964
Receipt of blood transfusion, blood components, or tissue	0	2	0	0	0	372
Other/risk factor not reported or identified	3	0	0	0	1	42
Total^b	121	105	71	50	68	9,101

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

^a From the beginning of the epidemic through 2005.

^b Includes children of unknown race or multiple races. Cumulative total includes 33 children of unknown race or multiple races. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Figure 1. Estimated numbers of AIDS cases in children <13 years of age, by year of diagnosis, 1992–2005—50 states and the District of Columbia (Revised June 2007)



Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays, but not for incomplete reporting.