

**Table 2. Time to an AIDS diagnosis after a diagnosis of HIV infection, by selected characteristics, 2004—33 states and U.S. dependent areas with confidential name-based HIV infection reporting (Revised June 2007)**

	≥12 Months after diagnosis of HIV infection <sup>a</sup>		<12 Months after diagnosis of HIV infection <sup>b</sup>		Total
	No.	(%) <sup>c</sup>	No.	(%) <sup>c</sup>	No.
<b>Data for 33 states</b>					
<b>Age at diagnosis (yrs)</b>					
<13	149	84	29	16	179
13–14	28	79	7	21	35
15–19	838	83	172	17	1,010
20–24	2,707	78	777	22	3,484
25–29	2,914	69	1,291	31	4,204
30–34	3,232	64	1,787	36	5,019
35–39	3,561	59	2,463	41	6,024
40–44	3,173	56	2,482	44	5,655
45–49	2,062	51	1,994	49	4,056
50–54	1,233	49	1,262	51	2,495
55–59	654	46	760	54	1,414
60–64	330	45	404	55	735
≥65	298	45	366	55	664
<b>Race/ethnicity</b>					
White, not Hispanic	6,675	63	3,953	37	10,627
Black, not Hispanic	10,485	60	6,890	40	17,375
Hispanic	3,577	57	2,661	43	6,238
Asian/Pacific Islander	194	56	155	44	350
American Indian/Alaska Native	110	59	78	41	188
<b>Transmission category</b>					
<b>Male adult or adolescent</b>					
Male-to-male sexual contact	10,305	63	6,147	37	16,452
Injection drug use	1,663	50	1,671	50	3,334
Male-to-male sexual contact and injection drug use	779	59	531	41	1,310
High-risk heterosexual contact <sup>d</sup>	2,180	53	1,922	47	4,102
Other <sup>e</sup>	59	47	66	53	125
Subtotal	14,986	59	10,337	41	25,323
<b>Female adult or adolescent</b>					
Injection drug use	1,157	60	782	40	1,939
High-risk heterosexual contact <sup>d</sup>	4,829	65	2,577	35	7,405
Other <sup>e</sup>	57	46	68	54	125
Subtotal	6,043	64	3,427	36	9,470
<b>Child (&lt;13 yrs at diagnosis)</b>					
Perinatal	126	81	29	19	155
Other <sup>f</sup>	24	95	1	5	25
Subtotal	149	83	30	17	180
<b>Subtotal for 33 states</b>	<b>21,178</b>	<b>61</b>	<b>13,795</b>	<b>39</b>	<b>34,973</b>
<b>Data for U.S. dependent areas</b>	<b>21</b>	<b>52</b>	<b>19</b>	<b>48</b>	<b>41</b>
<b>Total<sup>g</sup></b>	<b>21,200</b>	<b>61</b>	<b>13,814</b>	<b>39</b>	<b>35,014<sup>h</sup></b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

See Technical Notes for the list of areas that have had laws or regulations requiring confidential name-based HIV infection reporting since at least 2001.

Data exclude 39 persons whose month of diagnosis of HIV infection is unknown.

<sup>a</sup> Includes persons in whom AIDS has not developed.

<sup>b</sup> Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

<sup>c</sup> Percentages represent proportions of the total number of diagnoses of HIV/AIDS made during 2004 for the corresponding group (see row entries).

<sup>d</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>e</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>f</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>g</sup> Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

<sup>h</sup> Includes 194 persons of unknown race or multiple races.