

Active Bacterial Core Surveillance (ABCs) Report Emerging Infections Program Network



Streptococcus pneumoniae, 2006

ABCs Areas

California (San Francisco County and children < 5 years in Alameda and Contra Costa counties); Colorado (5 county Denver area); Connecticut; Georgia (20 county Atlanta area); Maryland (6 county Baltimore area); Minnesota; New Mexico; New York (15 county Rochester and Albany areas and children <5 years in Erie county); Oregon (3 county Portland area); Tennessee (11 urban counties)

ABCs Population

The surveillance areas represent 28,204,455 persons. Source: National Center for Health Statistics bridged-race vintage 2006 postcensal file

ABCs Case Definition

Invasive pneumococcal disease: isolation of *Streptococcus pneumoniae* from normally sterile site in resident of a surveillance area in 2006.

ABCs Methodology

ABCs personnel routinely contacted all microbiology laboratories serving acute care hospitals in their area to identify cases. Standardized case report forms that include information on demographic characteristics, clinical syndrome, and outcome of illness were completed for each identified case. Pneumococcal isolates were collected and sent to reference laboratories for susceptibility testing using CLSI methods and serotyping. Regular laboratory audits assessed completeness of active surveillance and detected additional cases.

Rates of invasive pneumococcal disease were calculated using population estimates for 2006. For national estimates, race- and age-specific rates of disease were applied from the aggregate surveillance area to the age and racial distribution of the 2006 U.S. population. Cases with unknown race were distributed by area based on reported race distribution for known cases within the eight age categories.

Reported ABCs Profiles

Race	No.	(Rate*)
White	2,614	(11.8)
Black	1,095	(25.1)
Other	213	(12.8)

Unknown race (n=684) distributed amongst known

Citation

Centers for Disease Control and Prevention. 2007. Active Bacterial Core Surveillance Report, Emerging Infections Program Network, *Streptococcus pneumoniae*, 2006. Available via the Internet:

http://www.cdc.gov/ncidod/dbmd/abcs/survreports/spneu06.pdf

		ases	D	eaths	
Age (years)	No.	(Rate*)	No.	(Rate*)	
< 1		(33.8)	3	(0.71)	
1	156	(36.7)	1	(0.24)	
2-4	149	(11.7)	5	(0.39)	
5-17	162	(3.3)	4	(0.08)	
18-34	265	(4.1)	15	(0.23)	
35-49	787	(12.0)	74	(1.13)	
50-64	993	(20.0)	118	(2.38)	
≥ 65	1,268	(40.2)	238	(7.55)	
Total	3,922	(13.9)	458	(1.62)	

^{*} Cases or deaths per 100,000 population for ABCs areas

Syndrome	No.	(% *)
Meningitis	241	(6.1)
Bacteremia without focus	904	(23.0)
Pneumonia with bacteremia	2,607	(66.5)

^{*} Percent of cases

Antibiotic	\mathbf{s}^*	$\mathbf{I}^{\boldsymbol{\dagger}}$	R‡
Susceptibility	%	%	%
Penicillin	74.4	15.2	10.4
Cefotaxime	93.8	5.0	1.2
Erythromycin	78.4	0.12	21.5
TMP/Sulfa	77.3	6.8	15.9
Tetracycline	88.7	0.21	11.1
Levofloxacin	99.5	0.06	0.44
Vancomycin	100.0	0.0	0.0

Based on reference lab testing of 3,402 isolates

National Estimates of Invasive Disease

Cases: 41,400 (13.8/100,000) Deaths: 5,000 (1.7/100,000)

Healthy People 2010 Update

Objective: Decrease the incidence of invasive pneumococcal infections to 46 per 100,000 persons less than 5 years of age and to 42 per 100,000 persons aged 65 and older.

Age (year)	2010 Objective	2006 Rate **
< 5	46/100,000	20.4/100,000
≥ 65	42/100,000	40.1/100,000

^{*} Cases per 100,000 U.S. population < 5 years or ≥ 65 years

For more information, visit our web site:

http://www.cdc.gov/abcs

^{*} Cases per 100,000 population for ABCs areas

^{*} Susceptible; † Intermediate; ‡ Resistant based on year 2006 CLSI definitions