

Table 36. Culture-Positive Tuberculosis Cases with Drug Susceptibility Results, by Monodrug (INH) or Multidrug Resistance: 59 Reporting Areas, 2003

Reporting Area	Total Culture Positive Cases	Cases with Initial Drug Susceptibility Testing Performed		Resistance ¹			
				Isoniazid		Isoniazid and Rifampin	
		No.	(%)	No.	(%)	No.	(%)
United States	11,704	10,549	(90.1)	869	(8.2)	114	(1.1)
Alabama	205	191	(93.2)	7	(3.7)	0	(0.0)
Alaska	48	45	(93.8)	4	(8.9)	2	(4.4)
Arizona	228	222	(97.4)	14	(6.3)	2	(0.9)
Arkansas	97	81	(83.5)	1	(1.2)	0	(0.0)
California	2,498	2,325	(93.1)	238	(10.2)	33	(1.4)
Colorado	72	71	(98.6)	7	(9.9)	1	(1.4)
Connecticut	105	94	(89.5)	8	(8.5)	1	(1.1)
Delaware	28	26	(92.9)	2	(7.7)	0	(0.0)
District of Columbia	71	67	(94.4)	3	(4.5)	0	(0.0)
Florida	852	785	(92.1)	79	(10.1)	7	(0.9)
Georgia	396	374	(94.4)	20	(5.3)	2	(0.5)
Hawaii	80	67	(83.8)	11	(16.4)	4	(6.1)
Idaho	10	8	(80.0)	4	(50.0)	0	(0.0)
Illinois	480	410	(85.4)	21	(5.1)	3	(0.7)
Indiana	119	114	(95.8)	6	(5.3)	1	(0.9)
Iowa	35	20	(57.1)	—	—	—	—
Kansas	54	54	(100.0)	8	(14.8)	1	(1.9)
Kentucky	119	110	(92.4)	5	(4.5)	1	(0.9)
Louisiana	206	188	(91.3)	10	(5.3)	1	(0.5)
Maine	20	16	(80.0)	0	(0.0)	0	(0.0)
Maryland	208	170	(81.7)	18	(10.6)	2	(1.2)
Massachusetts	197	185	(93.9)	11	(5.9)	0	(0.0)
Michigan	204	188	(92.2)	14	(7.4)	3	(1.6)
Minnesota	172	157	(91.3)	27	(17.2)	5	(3.2)
Mississippi	96	79	(82.3)	6	(7.6)	0	(0.0)
Missouri	108	105	(97.2)	6	(5.7)	0	(0.0)
Montana	7	7	(100.0)	1	(14.3)	1	(14.3)
Nebraska	22	18	(81.8)	3	(16.7)	0	(0.0)
Nevada	89	88	(98.9)	7	(8.0)	0	(0.0)
New Hampshire	12	12	(100.0)	1	(8.3)	0	(0.0)
New Jersey	400	396	(99.0)	40	(10.1)	6	(1.5)
New Mexico	44	44	(100.0)	4	(9.1)	0	(0.0)
New York State ²	254	251	(98.8)	22	(8.8)	6	(2.4)
New York City	873	864	(99.0)	81	(9.4)	19	(2.2)
North Carolina	294	275	(93.5)	17	(6.2)	0	(0.0)
North Dakota	4	2	(50.0)	—	—	—	—
Ohio	172	130	(75.6)	5	(3.8)	0	(0.0)
Oklahoma	126	96	(76.2)	7	(7.3)	0	(0.0)
Oregon	87	75	(86.2)	6	(8.0)	0	(0.0)
Pennsylvania	273	225	(82.4)	25	(11.1)	4	(1.8)
Rhode Island	30	21	(70.0)	—	—	—	—
South Carolina	183	171	(93.4)	13	(7.6)	0	(0.0)
South Dakota	17	17	(100.0)	0	(0.0)	0	(0.0)
Tennessee	225	163	(72.4)	—	—	—	—
Texas	1,296	1,011	(78.0)	53	(5.2)	5	(0.5)
Utah	29	24	(82.8)	2	(8.3)	1	(4.2)
Vermont	7	5	(71.4)	—	—	—	—
Virginia	262	242	(92.4)	23	(9.5)	2	(0.8)
Washington	215	189	(87.9)	12	(6.3)	0	(0.0)
West Virginia	19	16	(84.2)	1	(6.3)	0	(0.0)
Wisconsin	54	54	(100.0)	6	(11.1)	0	(0.0)
Wyoming	2	1	(50.0)	—	—	—	—
American Samoa ³
Fed. States of Micronesia ³
Guam ³	40	29	(72.5)	—	—	—	—
N. Mariana Islands ³	27	26	(96.3)	3	(11.5)	1	(3.8)
Puerto Rico ³	108	92	(85.2)	4	(4.3)	1	(1.1)
Republic of Palau ³
U.S. Virgin Islands ³

¹Isolates may be resistant to other drugs. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for $\geq 75\%$ of cases.

²Excludes New York City.

³Not included in U.S. totals.

Note: Ellipses indicate data not available.