


Figure 8.1. Prevalence of current smoking among adults aged 18 years and over: United States, 1997-2004




NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded persons with unknown smoking status (about 1% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000-02 were recalculated using weights derived from the 2000 census. See "About This Release" and table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997-2004 National Health Interview Surveys.

- In 2004, 20.9% (95% confidence interval = 20.3%-21.5%) of adults aged 18 years and over were current smokers, continuing a decline in the prevalence of current smoking among adults in the United States.
- The prevalence of current smoking among U.S. adults has declined over time from 24.7% in 1997 to 20.9% in 2004.

Figure 8.2. Percent distribution of smoking status among adults aged 18 years and over, by sex: United States, 2004




NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded 331 persons (1.1%) with unknown smoking status.

DATA SOURCE: Sample Adult Core component of the 2004 National Health Interview Survey.

■ The percentage of current smokers was higher for men (23.4%) than for women (18.5%).

■ The percentage of former smokers was higher for men than for women, and the percentage of those who had never smoked was higher for women than for men.

Figure 8.3. Prevalence of current smoking among adults aged 18 years and over, by age group and sex: United States, 2004




NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded 331 persons (1.1%) with unknown smoking status.

DATA SOURCE: Sample Adult Core component of the 2004 National Health Interview Survey.

- For both sexes combined, the percentage of adults who were current smokers was lower among adults aged 65 years and over (8.8%) than among adults aged 18-44 years (23.7%) and 45-64 years (22.4%). This pattern in current smoking by age group was seen in both men and women.
- For all three age groups, men were more likely than women to be current smokers.

Figure 8.4. Age- sex-adjusted prevalence of current smoking among adults aged 18 years and over, by race/ethnicity: United States, 2004



NOTES: Current smokers were defined as those who smoked more than 100 cigarettes in their lifetime and now smoke every day or some days. The analyses excluded 331 persons (1.1%) with unknown smoking status. Estimates are age- sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

DATA SOURCE: Sample Adult Core component of the 2004 National Health Interview Survey.

- The age- sex-adjusted prevalence of current smoking was 14.2% for Hispanic persons, 22.7% for non-Hispanic white persons, and 20.0% for non-Hispanic black persons.
- Non-Hispanic white adults and non-Hispanic black adults were more likely than Hispanic adults to be current smokers.

Data tables for figures 8.1-8.4:

Data table for figure 8.1. Prevalence of current smoking among adults aged 18 years and over: United States, 1997-2004

Year	Percent (95% confidence interval)	
	Crude ¹	Age-adjusted ²
1997	24.7 (24.1-25.3)	24.6 (24.0-25.1)
1998	24.1 (23.5-24.7)	24.0 (23.1-24.8)
1999	23.5 (22.9-24.1)	23.3 (22.7-24.0)
2000	23.2 (22.5-23.8)	23.1 (22.5-23.7)
2001	22.7 (22.1-23.3)	22.6 (22.0-23.2)
2002	22.4 (21.7-23.0)	22.3 (21.7-22.9)
2003	21.6 (21.0-22.2)	21.5 (20.9-22.1)
2004	20.9 (20.3-21.5)	20.8 (20.2-21.4)

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

Data table for figure 8.2. Percent distribution of smoking status among adults aged 18 years and over, by sex: United States, 2004

Smoking status and sex	Percent	95% confidence interval
Never		
Total	57.7	56.9-58.4
Male	51.7	50.7-52.8
Female	63.2	62.2-64.1
Former		
Total	21.4	20.9-22.0
Male	24.8	24.0-25.7
Female	18.3	17.6-19.0
Current		
Total	20.9	20.3-21.5
Male	23.4	22.6-24.3
Female	18.5	17.8-19.3

Data table for figure 8.3. Prevalence of current smoking among adults aged 18 years and over, by age group and sex: United States, 2004

Age and sex	Percent	95% confidence interval
18-44 years		
Total	23.7	22.8-24.7
Male	26.1	24.8-27.4
Female	21.4	20.3-22.5
45-64 years		
Total	22.4	21.4-23.4
Male	25.0	23.5-26.6
Female	19.8	18.7-21.1
65 years and over		
Total	8.8	8.0-9.7
Male	9.8	8.5-11.3
Female	8.1	7.2-9.1
18 years and over: crude¹		
Total	20.9	20.3-21.5
Male	23.4	22.6-24.3
Female	18.5	17.8-19.3
18 years and over: age-adjusted²		
Total	20.8	20.2-21.4
Male	23.0	22.2-23.9
Female	18.7	17.9-19.5

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

Data table for figure 8.4. Adjusted prevalence of current smoking among adults aged 18 years and over, by race/ethnicity: United States, 2004

Race/ethnicity	Percent (95% confidence interval)	
	Age- sex-adjusted ¹	Age-adjusted ²
Hispanic or Latino	14.2 (13.1-15.4)	14.3 (13.2-15.5)
Not Hispanic or Latino:		
White, single race	22.7 (21.9-23.5)	22.7 (21.9-23.4)
Black, single race	20.0 (18.3-21.8)	19.7 (18.1-21.5)

Age- sex-adjusted estimates are presented in the figure. Estimates are age- sex-adjusted to the 2000 projected U.S. standard population using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted to the 2000 projected U.S. standard population using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.