

Table 33. Tuberculosis Cases and Percentages by Noninjecting Drug Use,¹ Age ≥15: Reporting Areas, 2007

Reporting Area	Total Cases	Cases with Information on Noninjecting Drug Use		Cases Reporting Noninjecting Drug Use	
		No.	(%)	No.	(%)
United States	12,518	12,103	(96.7)	965	(8.0)
Alabama	168	167	(99.4)	12	(7.2)
Alaska	47	44	(93.6)	6	(13.6)
Arizona	278	242	(87.1)	21	(8.7)
Arkansas	95	95	(100.0)	4	(4.2)
California	2,592	2,516	(97.1)	159	(6.3)
Colorado	104	102	(98.1)	5	(4.9)
Connecticut	105	102	(97.1)	5	(4.9)
Delaware	18	18	(100.0)	0	(0.0)
District of Columbia	58	58	(100.0)	7	(12.1)
Florida	928	925	(99.7)	113	(12.2)
Georgia	437	430	(98.4)	49	(11.4)
Hawaii	119	39	(32.8)	--	--
Idaho	9	9	(100.0)	1	(11.1)
Illinois	504	449	(89.1)	44	(9.8)
Indiana	123	120	(97.6)	12	(10.0)
Iowa	42	42	(100.0)	3	(7.1)
Kansas	55	55	(100.0)	6	(10.9)
Kentucky	116	115	(99.1)	6	(5.2)
Louisiana	207	179	(86.5)	44	(24.6)
Maine	17	17	(100.0)	0	(0.0)
Maryland	257	255	(99.2)	11	(4.3)
Massachusetts	209	207	(99.0)	6	(2.9)
Michigan	205	194	(94.6)	13	(6.7)
Minnesota	216	214	(99.1)	8	(3.7)
Mississippi	128	128	(100.0)	8	(6.3)
Missouri	115	114	(99.1)	15	(13.2)
Montana	11	11	(100.0)	0	(0.0)
Nebraska	23	23	(100.0)	0	(0.0)
Nevada	93	89	(95.7)	3	(3.4)
New Hampshire	11	10	(90.9)	0	(0.0)
New Jersey	437	437	(100.0)	14	(3.2)
New Mexico	49	48	(98.0)	3	(6.3)
New York State ²	245	215	(87.8)	10	(4.7)
New York City	888	860	(96.8)	82	(9.5)
North Carolina	320	318	(99.4)	44	(13.8)
North Dakota	7	6	(85.7)	0	(0.0)
Ohio	239	236	(98.7)	23	(9.7)
Oklahoma	128	120	(93.8)	13	(10.8)
Oregon	87	82	(94.3)	10	(12.2)
Pennsylvania	249	239	(96.0)	14	(5.9)
Rhode Island	37	36	(97.3)	0	(0.0)
South Carolina	205	203	(99.0)	26	(12.8)
South Dakota	13	13	(100.0)	0	(0.0)
Tennessee	224	224	(100.0)	45	(20.1)
Texas	1,406	1,406	(100.0)	93	(6.6)
Utah	37	37	(100.0)	1	(2.7)
Vermont	3	3	(100.0)	0	(0.0)
Virginia	291	291	(100.0)	10	(3.4)
Washington	269	266	(98.9)	11	(4.1)
West Virginia	24	24	(100.0)	1	(4.2)
Wisconsin	68	68	(100.0)	4	(5.9)
Wyoming	2	2	(100.0)	0	(0.0)
American Samoa ³	3	3	(100.0)	0	(0.0)
Fed. States of Micronesia ³	49	35	(71.4)	--	--
Guam ³	61	56	(91.8)	0	(0.0)
Marshall Islands ³	93	90	(96.8)	0	(0.0)
N. Mariana Islands ³	41	37	(90.2)	0	(0.0)
Puerto Rico ³	94	94	(100.0)	15	(16.0)
Republic of Palau ³	11	9	(81.8)	0	(0.0)
U.S. Virgin Islands ³

¹Noninjecting drug use within past 12 months of TB diagnosis. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

²Excludes New York City.

³Not included in U.S. totals.

Note: Ellipses indicate data not available.