

References

- Andersen KE, Boman A, Hamann K, Wahlberg JE. 1984. Guinea pig maximization tests with formaldehyde releasers. Results from two laboratories. *Contact Dermatitis* 10(5):257-266.
- Basketter DA, Scholes EW. 1992. Comparison of the local lymph node assay with the guinea-pig maximization test for the detection of a range of contact allergens. *Food and Chemical Toxicology* 30(1): 65-69.
- Basketter DA, Scholes EW, Kimber I. 1994. The performance of the local lymph node assay with chemicals identified as contact allergens in the human maximization test. *Food and Chemical Toxicology* 32(6): 543-547.
- Basketter DA, Gerberick GF, Kimber I. 1998. Strategies for identifying false positive responses in predictive skin sensitization tests. *Food and Chemical Toxicology* 36(4): 327-333.
- Basketter DA, Lea LJ, Cooper K, Stocks J, Dickens A, Pate I, et al. 1999a. Threshold for classification as a skin sensitizer in the local lymph node assay: A statistical evaluation. *Food and Chemical Toxicology* 37(12): 1167-1174.
- Basketter DA, Lea LJ, Cooper KJ, Ryan CA, Gerberick GF, Dearman RJ, et al. 1999b. Identification of metal allergens in the local lymph node assay. *American Journal of Contact Dermatitis* 10(4): 207-212.
- Basketter DA, Clapp C, Jefferies D, Safford B, Ryan CA, Gerberick F, et al. 2005. Predictive identification of human skin sensitization thresholds. *Contact Dermatitis* 53(5): 260-267.
- Gad SC, Dunn BJ, Dobbs DW. 1986. Development and validation of an alternative dermal sensitization test: The mouse ear swelling test (MEST). *Toxicology and Applied Pharmacology* 84(1): 93-114.
- Gerberick GF, House RV, Fletcher ER, Ryan CA. 1992. Examination of the local lymph node assay for use in contact sensitization risk assessment. *Fundamental and Applied Toxicology* 19(3): 438-445.
- Gerberick GF, Ryan CA, Kern PS, Schlatter H, Dearman RJ, Kimber I, et al. 2005. Compilation of historical local lymph node data for evaluation of skin sensitization alternative methods. *Dermatitis* 16(4): 157-202.
- Griem P, Goebel C, Scheffler H. 2003. Proposal for a risk assessment methodology for skin sensitization based on sensitization potency data. *Regulatory Toxicology and Pharmacology* 38(3): 269-290.
- Guillot JP, Gonnet JF, Clement C, Faccini JM. 1983. Comparative study of methods chosen by the Association Francaise de Normalisation (AFNOR) for evaluating sensitizing potential in the albino guinea-pig. *Food and Chemical Toxicology* 21(6): 795-805.
- Hicks R, Hewitt PJ, Lam HF. 1979. An investigation of the experimental induction of hypersensitivity in the guinea pig by material containing chromium, nickel and cobalt

from arc welding fumes. *International Archives of Allergy and Applied Immunology* 59(3): 265-272.

Hindsen M, Bruze M, Christensen OB. 1999. Individual variation in nickel patch test reactivity. *American Journal of Contact Dermatitis* 10(2): 62-67.

Kligman AM. 1966. The identification of contact allergens by human assay. 3. The maximization test: a procedure for screening and rating contact sensitizers. *Journal of Investigative Dermatology* 47(5): 393-409.

Lea LJ, Warbrick EV, Dearman RJ, Kimber I, Basketter DA. 1999. The impact of vehicle on assessment of relative skin sensitization potency of 1,4-dihydroquinone in the local lymph node assay. *American Journal of Contact Dermatitis* 10(4): 213-218.

Magnusson B, Kligman AM. 1969. The identification of contact allergens by animal assay. The guinea pig maximization test. *Journal of Investigative Dermatology* 52(3): 268-276.

Marzulli FN, Maibach HI. 1974. The use of graded concentrations in studying skin sensitizers: experimental contact sensitization in man. *Food and Cosmetics Toxicology* 12(2): 219-227.

Ryan CA, Gerberick GF, Cruse LW, Basketter DA, Lea L, Blaikie L, et al. 2000. Activity of human contact allergens in the murine local lymph node assay. *Contact Dermatitis* 43(2): 95-102.

Ryan CA, Cruse LW, Skinner RA, Dearman RJ, Kimber I, Gerberick GF. 2002. Examination of a vehicle for use with water soluble materials in the murine local lymph node assay. *Food and Chemical Toxicology* 40(11): 1719-1725.

Schneider K, Akkan Z. 2004. Quantitative relationship between the local lymph node assay and human skin sensitization assays. *Regulatory Toxicology and Pharmacology* 39(3): 245-255.

Uter W, Fuchs T, Hausser M, Ippen H. 1995. Patch test results with serial dilutions of nickel sulfate (with and without detergent), palladium chloride, and nickel and palladium metal plates. *Contact Dermatitis* 32(3): 135-142.

Wahlberg JE, Boman A. 1985. Guinea pig maximization test. *Current problems in dermatology* 14: 59-106.