

# **Post-Chernobyl Thyroid Cancer in Exposed Children**


**Maureen Hatch, Ph.D.  
Radiation Epidemiology Branch**

**Epidemiology Course  
May 16, 2007**

# The Chernobyl Accident – 26 April 1986

- 10 days of releases into the atmosphere
- Widespread and spotty fallout due to rain and changing wind directions
- Iodine 131 the principal contaminant
- Over 200,000 evacuated, 5 million living in contaminated regions

# Chernobyl Fallout



## Contaminated\* Areas in European Countries Following the Chernobyl Accident (Izrael et al. 1996)

Country	Area in deposition-density ranges (km <sup>2</sup> )			
	37-185 kBqm <sup>-2</sup>	185-555 kBqm <sup>-2</sup>	555-1480 kBqm <sup>-2</sup>	>1480 kBqm <sup>-2</sup>
Russia	49,800	5,700	2,100	300
Belarus	29,900	10,200	4,200	2,200
Ukraine	37,200	4,200	900	600
Sweden	12,000	-	-	-
Finland	11,500	-	-	-
Austria	8,600	-	-	-
Norway	5,200	-	-	-
Bulgaria	4,800	-	-	-
Switzerland	1,300	-	-	-
Greece	1,200	-	-	-
Slovenia	300	-	-	-
Italy	300	-	-	-
Moldova	80	-	-	-

\* The contaminated areas are defined as those where the <sup>137</sup>Cs deposition density resulting from the Chernobyl accident was greater than 37 kBqm<sup>-2</sup>



# Exposure to Radioactive Iodine from Chernobyl

- $^{131}\text{I}$  concentrates in the thyroid (thyroid dose much greater than average body dose)
- Can be inhaled and ingested (mainly in milk)
- Children received the highest doses (small thyroid mass, high milk consumption)

# **Iodine Deficiency in Contaminated Areas**

- **Possible risk factor for thyroid cancer**
- **Increases uptake of radioiodines**
- **May stimulate thyroid cell proliferation**
- **May increase effect of radioiodines**

# **Radiation and Thyroid Cancer: What was Known before Chernobyl**

- **Atomic bomb**
  - **Biggest increase in children**
- **X-ray exposures: medical uses**
  - **Increase following exposure in childhood**
- **$^{131}\text{I}$ : dx and tx**
  - **No obvious increase in adults but data sparse in children**

# Data from Chernobyl will contribute to:

- Understanding of  $^{131}\text{I}$  carcinogenesis
- Effective handling of future nuclear events
- Safe use of radioiodines in clinical practice



# **Exposure in Childhood to Fallout: Radioiodines from the Nevada Test Site ,1985-1986**

- **3545 schoolchildren screened and interviewed, 2473 (2496) analyzed**
- **Doses based on diet and deposition**
- **Mean dose=170 (120) mGy**
- **Significant excess of thyroid neoplasms (n=19, 23)**

**Kerber R, et al., JAMA 1993; Lyon et al., Epidemiol 2006**

# Thyroid Disease in Those Exposed as Children to Iodine 131 from the Hanford Nuclear Plant


- 5199 children from contaminated areas (1944-1957), 3440 screened and analyzed
- Thyroid radiation dose estimated through specially designed computer program
- Mean dose 174 mGy
- No dose-response relationship with benign or malignant thyroid disease

# Thyroid Cancer in Contaminated Areas of Ukraine, 1981-1990

Year	Thyroid Cancer (No.)
1981	0
1982	0
1983	0
1984	0
1985	0
1986	0
1987	0
1988	0
1989	0
1990	3


Prisyazhiuk A, et al., The Lancet 1991

# Cases of Childhood (■) and adolescent (□) thyroid carcinoma in Belarus, registered yearly from 1986- 1995







# Age Distribution at the time of diagnosis of thyroid cancer patients from Belarus and from Italy and France



(Pacini et al. 1997)

# Papillary cancer, solid subtype



**Real effect  
or  
Screening effect?**

# Case-Control Study in Belarus

- Belarus, ages 0-16, dx in 1987-1992
- 107 cases, 214 controls (**same opportunity for diagnosis**)
- exposure assessment ecological



# Case-Control Study in Belarus

Dose (Gy)	Cases	Controls	OR(95% CI)
< 0.3	64	88	1.00
0.3-0.9	26	15	2.38 (1.2, 4.9)
1 +	17	4	5.84 (2.0, 17.3)

Astakhova L, et al., Radiat Res 1998

# Thyroid cancer in Bryansk region of the Russian Federation

- Population-based case-control study (0-19)
- 26 cases, diagnosed before Oct 1, 1997
- 52 matched controls from Russian national registry
- Doses estimated from semi-empirical model

# Odds Ratios and 95% Confidence Intervals for Thyroid Cancer by Median Radiation Dose, Russian Federation

Median Dose (mGy)	No. of Cases	No. of Controls	OR (95% CI)
23	4	16	1.00 (ref.)
139	5	14	1.65 (0.32-8.50)
427	4	16	3.05 (0.42-22.1)
1049	13	6	44.7 (3.30-604)

Davis S, et al., Rad Res 2004

# **Radiation Dose and Iodine Status: Belarus and Russian Federation, 1992- 1998**

- **Population-based case-control study (<15)**
- **276 cases, 1300 matched controls**
- **Stable iodine status based on settlement soil levels**
- **Consumption of potassium iodide from interview**



# Risk of Thyroid Cancer at 1 Gy, Belarus and Russian Federation

- Median dose in controls = 245 mGy

OR at 1 Gy (95% CI),	5.5 (3.1, 9.5) -
from different models	8.4 (4.1, 17.3)

---

# Radiation Dose and Iodine Status: Belarus and Russian Federation, 1992- 1998

OR at 1 Gy (95% CI)

---

---

Potassium  
iodide

Highest two tertiles  
of soil iodine

Lowest tertiles  
of soil iodine

---

---

No

3.5 (1.8, 7.0)

10.8 (5.6, 20.8)

Yes

1.1 (0.3, 3.6)

3.3 (1.9, 10.6)

---

---

Cardis E, et al., JNCI 2005

# **Joint Effect of Iodine Deficiency and Radiation Dose: Bryansk region of the Russian Federation, 1996**

- **3070 individuals in 78 settlements**
- **2590 ages 6-18**
- **Urinary iodine measurements**
- **34 histologically confirmed cancers**
- **Dose estimated from semi-empirical models**

**Shakhtarin V, et al., IJE 2003**

# Iodine Levels and Radiation Dose: Bryansk region, Russian Federation, 1996

Urinary Iodine Excretion ( $\mu\text{g}/\text{dl}$ )	ERR per Gy Estimate	95% CI
< 5.0	24.1	(1.7, 78.31)
5.0 – 7.49	18.3	(10.7, 28.6)
7.5 – 9.99	16.2	(0.8, 49.3)
$\geq 10$	13.0	(-11.0, 71.2)

Shakhtarin V, et al., IJE 2003



# Thyroid Cancer Risk in Areas of Ukraine

- Ecological study of 301,907 (1-18) in 1,293 rural settlements
- 24% with individual dose estimates; 76% with 'individualized' estimates
- 232 histologically confirmed thyroid cancers through 12/01
- ERR/Gy=8.0 (95% CI 4.6-15)

# Ukrainian-American Thyroid Study Belarusian-American Thyroid Study



A collaboration between scientists from  
Ukraine, Belarus, NCI and Columbia  
University



# Approach

- **Cohort study of 25,000 exposed children**
- **Biennial screening examinations of the thyroid gland, 1998 - present**
  - **Palpation**
  - **Ultrasound**
  - **Fine Needle Aspiration as indicated**
  - **Thyroid hormone, thyroid antibody and iodine excretion measurements**

# Study Endpoints

- **Thyroid cancer**
- Benign neoplasm
- Diffuse goiter
- Nodular goiter
- **Autoimmune thyroiditis (AIT)**
- Thyrotoxicosis (hyperthyroidism)
- Hypothyroidism
- Hyperparathyroidism
- Hypoparathyroidism
- **Iodine deficiency**

# Dosimetry



- Direct measurements
- Questionnaire data
- Radioecologic modelling

**Mean in Ukraine**      **0.78 Gy**

**Median**                      **0.30 Gy**



# Approximate Mean Doses From Selected Low Dose Radiation Exposures<sup>†</sup>

Source	Approximate mean individual dose, mGy
Thyroid dose from Chernobyl (Ukraine, <18 y at exposure)	780
Breast dose to scoliosis patients	100
Pediatric CT scan (stomach dose from abdominal scan)	25
Single screening mammogram	3

<sup>†</sup> Adapted from DJ Brenner et al. 2003


## Odds Ratios <sup>a</sup> and 95% Confidence Intervals (95% CI) by Thyroid Dose

Dose Categories (Gy)	Cases (n)	Odds Ratio (95% CI)
0 – 0.24	9	Ref.
0.25 – 0.74	9	2.31 (0.91 – 5.88)
0.75 – 1.49	10	6.25 (2.50 – 15.6)
1.50 – 2.99	8	8.97 (3.39 – 23.7)
3.00+	9	15.30 (5.88 – 40.0)

<sup>a</sup> all odds ratios adjusted for gender, age at screening p (trend) = <0.0001

Tronko M, Howe G, et al., JNCI, 2006

# Plot of the Odds Ratio Estimates and the Corresponding 95% Confidence Intervals from the Categorical Analysis and a Fitted Dose-Response Line



# Models of Excess Relative Risk per Gray (ERR) and Interactions of Dose, Gender and Age at Exposure

Variable	Parameter and Estimates		
<b>Dose</b>	<b>ERR = 5.25*</b>		
<b>Gender: Male</b>	<b>RR = 2.21</b>	<b>p = 0.14</b>	
<b>Female</b>	<b>RR = 16.57</b>		
<b>Dose</b>			
<b>Age at exposure: 0 – 4</b>	<b>RR = 9.08</b>	<b>p = 0.58</b>	
<b>5 – 9</b>	<b>RR = 7.00</b>		
<b>10+</b>	<b>RR = 3.39</b>		

\* Statistically significant

# Conclusions from Cohort Study

- **Strong, linear relationship between dose and response**
- **Not confounded by screening**
- **Suggestive modification by gender and age at exposure, but not by iodine**
- **ERR = 5.25 v. ERR = 7.7 (pooled analysis of external radiation)**

# General Conclusions

- Consistent results from analytic studies (**5-6 fold excess overall**)
- Strong dose-response
- Magnitude of risk similar to external radiation



# Questions Still Remain About....

- **Age and gender as modifiers of thyroid cancer risk in children**
- **Role of iodine deficiency**
- **Risk of thyroid cancer in exposed adults**
- **Risk in those exposed in utero**
- **Specific molecular features**
- **Changes in tumor characteristics**

# Thyroid Cancer Morbidity and Mortality Due to Chernobyl

- ~ 5,000 cases of thyroid cancer through 2002
- 15 thyroid cancer deaths

# Thyroid Cancer Morbidity and Mortality Due to Chernobyl

- **Variable estimates of lifetime excess**
  - 4,000 – 9,000 deaths (WHO, 2005)
  - 30,000-60,000 cancer deaths (Greens/EFA Party, 2006)
  - 93,000 cancer deaths (Greenpeace, 2006)

# Postscript

- Cohort of 110,645 Ukrainian Male Cleanup Workers
- Nested case-control study of leukemia and related disorders
- 70 confirmed, analyzable leukemias, five matched controls (age, residence)
- Individual RADRUE (time and motion-based) dose estimate