Supplemental Materials #2: Rates and Trends by Site and Sex for 1992-1999

Incidence SEER 12 Areas	Both Sexes	Male	Female	Both Sexes	Male	Female
All Races	Rate	Rate	Rate	APC	APC	APC
All Sites	475.8	570.5	413.6	-0.9 *	-2.3 *	
Brain and Other Nervous System	6.5	7.7	5.4	-0.4	-0.7	-0.1
Breast	0.0		132.1	0.1	0.,	1.1 *
Cervix Uteri			10.2			-2.3 *
Colon and Rectum	54.3	64.6	46.7	-0.6	-1.0 *	
Corpus and Uterus, NOS			24.5			0.3
Esophagus	4.5	7.5	2.1	0.5	0.6	-0.4
Hodgkin Lymphoma	2.7	3.0	2.4	-0.1	-0.4	0.1
Kidney and Renal Pelvis	10.7	14.8	7.4	0.9 *	0.5	1.1 *
Larynx	4.2	7.5	1.6	-2.6 *	-3.5 *	
Leukemias	12.2	16.0	9.4	-1.6 *	-1.7 *	-1.5
Liver and Intrahepatic Bile Duct	5.3	8.1	3.1	4.3 *	3.8 *	4.5 *
Lung and Bronchus	63.9	84.5	49.2	-1.3 *	-2.5 *	0.1
Melanoma of the Skin	15.5	19.5	12.7	2.8 *	2.9 *	2.7 *
Myeloma	5.5	6.9	4.6	-1.1	-1.3	-1.3
Non-Hodgkin's Lymphomas	19.0	23.7	15.2	0.1	-0.6	1.1 *
Oral Cavity and Pharynx	11.4	17.0	6.7	-2.1 *	-2.6 *	-1.3
Ovary			17.1			-0.9 *
Pancreas	11.0	12.5	9.8	-0.6	-0.7	-0.7
Prostate		178.9			-4.2 *	
Stomach	9.3	13.5	6.3	-1.3 *	-2.1 *	-0.4
Testis		5.0			1.1	
Thyroid	6.3	3.5	9.0	3.2 *	1.2	3.9 *
Urinary Bladder	20.2	35.5	9.2	-0.4	-0.6 *	-0.6

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard population by 5-year age groups.

[~] Statistic could not be calculated.

[^] Statistic not displayed due to less than 25 cases.

^{*} The APC is significantly different from zero (p<0.05).