

Assays Using Rat Uterine Cytosol

Reference	Acton et al. (1983)	Allen et al. (1980)	Anstead et al. (1989)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	Rats (otherwise unspecified)	Rats (otherwise unspecified)	Rats (otherwise unspecified)
<i>Age of animals</i>	Mature	n.p.	n.p.
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	n.p.	estradiol benzoate treated 3 x 0.16 µg	n.p.
<i>Buffer for preparation of cytosol</i>	n.p.	TED (10 mM Tris, 1.5 mM EDTA, 0.5 mM dithiothreitol)	n.p.
<i>Dilution of tissue with buffer</i>	n.p.	8 uteri in 4 mL TED buffer	n.p.
<i>Protein concentration of cytosol</i>	n.p.	n.p.	n.p.
Competitive binding assay			
<i>Volume and concentration of radiolabelled 17β-estradiol</i>	volume n.p.; 1 nM	50 µL; 7x10 ⁻⁹ mol/L	n.p.
<i>Specific activity of radioligand</i>	n.p.	58 Ci/mmol	n.p.
<i>Solvent used to dissolve competing ligand</i>	ethanol	TED buffer	n.p.
<i>Concentration range of competing ligand</i>	10 ⁻³ to 10 µg/mL	10 ⁻⁹ to 10 ^{-5.5} mol/L	10 ⁻³ to 10 µg/mL
<i>Volume of ER prep used</i>	n.p.	150 µL	n.p.
<i>No. of replicates</i>	n.p.	n.p.	n.p.
<i>No. of times assay repeated</i>	n.p.	n.p.	n.p.
<i>Time of incubation</i>	16 hours	30 mins	n.p.
<i>Temperature of incubation</i>	3°C	30°C	n.p.
<i>Measure of nonspecific binding(y/n) and concentration</i>	y, 10 ⁻³ to 10 µg/mL	n.p.	n.p.
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	Dextran-coated charcoal	200 µL dextran-coated charcoal (250mg% Norit A, 25mg% dextran in TED buffer)	Dextran-coated charcoal
<i>Incubation time and temperature</i>	n.p.	20 min. in ice cold water	n.p.
<i>Centrifugation speed</i>	n.p.	2000xg	n.p.
<i>Centrifugation time and temperature</i>	n.p.	4°C for 5 min.	n.p.
Data calculations			
<i>Program or method used for calculating data</i>	n.p.	n.p.	n.p.
<i>Data plotted as</i>	n.p.	% specific ct./min. to controls vs. Conc. ligand in incubate (mol/L)	n.p.
<i>Data format in paper (e.g., IC₅₀, K_i)</i>		graphical	n.p.
<i>Calculation of RBA</i>	IC ₅₀ E ₂ /IC ₅₀ ligand (estimated)	IC ₅₀ E ₂ /IC ₅₀ competitor x100	n.p.

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity

Assays Using Rat Uterine Cytosol

Reference	Ashby et al. (1999)	Blair et al. (2000)	Connor et al. (1997)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	AP Rats	Sprague Dawley rats	Sprague Dawley rats
<i>Age of animals</i>	21-25 days old	245±18 days old	24 days
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	n.p.	10 days prior to sacrifice	n.p.
<i>Buffer for preparation of cytosol</i>	TEGM (10 mM Tris, 1.5 mM EDTA, MgCl ₂ 3 mM, 10% glycerol, pH 7.6)	TEDG (10 mM Tris, 1.5 mM EDTA, 10 mM D, 10% glycerol, pH 7.4)	TESHMo (10 mM Tris-HCl, pH 7.4; 1.5 mM EDTA, 15 mM thioglycerol; 10 mM sodium molybdate)
<i>Dilution of tissue with buffer</i>	50 mg/mL buffer	100 mg/mL buffer	50 mg/mL buffer
<i>Protein concentration of cytosol</i>	n.p.	n.p.	n.p.
Competitive binding assay			
<i>Volume and concentration of radiolabelled 17β -estradiol</i>	volume n.p.; 5 nM - 500 μM	10 μL; 1 nM	volume n.p.; 10 nM
<i>Specific activity of radioligand</i>	n.p.	141 Ci/mmol	130 Ci/mmol
<i>Solvent used to dissolve competing ligand</i>	n.p.	100% ethanol	n.p.
<i>Concentration range of competing ligand</i>	5 nM - 500 μM	n.p.	1 mM - 0.1 μM
<i>Volume of ER prep used</i>	100 μL	50 μL	n.p.
<i>No. of replicates</i>	n.p.	2	3
<i>No. of times assay repeated</i>	n.p.	2	n.p.
<i>Time of incubation</i>	18 hours	20 hours	8 hours
<i>Temperature of incubation</i>	4°C	4°C	4°C
<i>Measure of nonspecific binding(y/n) and concentration</i>	n.p.	y	n.p.
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	Hydroxyapatite in TEGM buffer	750 μl cold hydroxyapatite in 50mM Tris pH 7.4	0.1 volume Dextran-coated charcoal
<i>Incubation time and temperature</i>	n.p.	20 min. at 4°C	n.p.
<i>Centrifugation speed</i>	n.p.	600xg	8000xg
<i>Centrifugation time and temperature</i>	n.p.	4°C for 5 min.	10 min.; temp. n.p.
Data calculations			
<i>Program or method used for calculating data</i>	n.p.	n.p.	n.p.
<i>Data plotted as</i>	% Control vs. Concentration (M)	% [³ H]-E ₂ bound vs. Competitor concentration (M)	n.p.
<i>Data format in paper (e.g., IC₅₀, K_i)</i>	IC ₅₀	IC ₅₀	IC ₅₀
<i>Calculation of RBA</i>	n.p.	IC ₅₀ E ₂ /IC ₅₀ competitor x 100	IC ₅₀ E ₂ /IC ₅₀ competitor

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity

Assays Using Rat Uterine Cytosol

Reference	Elsby et al. (2000)	Fang et al. (2001)	Gabbard and Segaloff (1983)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	AP rats	Sprague Dawley rats	AXC rats
<i>Age of animals</i>	21-25 days old	245±18 days old	Mature
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	n.p.	10 days prior to sacrifice	5 days prior to sacrifice
<i>Buffer for preparation of cytosol</i>	TEGM (10 mM Tris, 1.5 mM EDTA, MgCl ₂ 3 mM, 10% glycerol, pH 7.6)	TEDG (10 mM Tris, 1.5 mM EDTA, 10 mM D, 10% glycerol, pH 7.4)	1.5 mM Tris, 1.0 mM EDTA, 20 mM sodium molybdate, pH 7.4
<i>Dilution of tissue with buffer</i>	50 mg/mL buffer	17 mg/mL	10 mg/mL buffer
<i>Protein concentration of cytosol</i>	n.p.	n.p.	6 mg/mL
Competitive binding assay			
<i>Volume and concentration of radiolabelled 17β -estradiol</i>	volume n.p.; 5 nM - 500 μM	10 μL; 1 nM	n.p.
<i>Specific activity of radioligand</i>	111 Ci/mmol	141 Ci/mmol	53 Ci/mmol
<i>Solvent used to dissolve competing ligand</i>	n.p.	100% ethanol	ethanol
<i>Concentration range of competing ligand</i>	5 nM - 500 μM	n.p.	1 nM to 1 μM
<i>Volume of ER prep used</i>	100 μL	50 μL	100 μL
<i>No. of replicates</i>	n.p.	2	3
<i>No. of times assay repeated</i>	n.p.	2	n.p.
<i>Time of incubation</i>	18 hours	20 hours	2 hours
<i>Temperature of incubation</i>	4°C	4°C	4°C
<i>Measure of nonspecific binding(y/n) and concentration</i>	n.p.	n.p.	n.p.
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	250 μL 60% Hydroxyapatite in TEGM buffer	750 μl cold hydroxyapatite in 50mM Tris pH 7.4	Dextran-coated charcoal (Norite-A + dextran + human gamma globulin)
<i>Incubation time and temperature</i>	n.p.	20 min. at 4°C	15 min. at 4°C
<i>Centrifugation speed</i>	1000xg	600xg	3500xg
<i>Centrifugation time and temperature</i>	10 min. at room temp.	4°C for 5 min.	10 min. at 0°C
Data calculations			
<i>Program or method used for calculating data</i>	n.p.	n.p.	n.p.
<i>Data plotted as</i>	% Control vs. Concentration (M)	n.p.	% Bound radioactivity vs. Log concentration
<i>Data format in paper (e.g., IC₅₀, K_i)</i>	IC ₅₀	RBA	relative displacing activity (RDA)
<i>Calculation of RBA</i>	IC ₅₀ E ₂ /IC ₅₀ competitor x 100	IC ₅₀ E ₂ /IC ₅₀ competitor x 100	IC ₅₀ E ₂ /IC ₅₀ competitor x 100

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity

Assays Using Rat Uterine Cytosol

Reference	Jaimez et al. (2000)	Jordan et al. (1986)	Laws et al. (1996)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	Rats (otherwise unspecified)	Sprague Dawley rats	Long Evans rats
<i>Age of animals</i>	Immature	18-21 days	Adult (60 days)
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	n.p.	n.p.	11 days prior to sacrifice
<i>Buffer for preparation of cytosol</i>	TEDM (20mM Tris-HCl, 1.5 mM EDTA, 0.25 mM dithiothreitol, 10 µg/mL leupeptine, 10% glycerol, pH 7.4)	TED (10 mM Tris, 1.5 mM EDTA, 0.5 mM dithiothreitol, pH 7.4)	TE-G30%-MTG (50 mM Tris, 0.9 mM EDTA, 30% (v/v) glycerol, 0.15% (v/v) monothio glycerol, pH 7.4)
<i>Dilution of tissue with buffer</i>	1:6 (w/v) tissue to buffer ratio	2 uteri/mL buffer	50 mg/mL buffer
<i>Protein concentration of cytosol</i>	n.p.	n.p.	n.p.
Competitive binding assay			
<i>Volume and concentration of radiolabelled 17β -estradiol</i>	volume n.p.; 1 nM	100 µL; 5x10 ⁻⁹ mol/L	volume n.p.; 1 nM
<i>Specific activity of radioligand</i>	n.p.	51 Ci/mmol	111Ci/nmol
<i>Solvent used to dissolve competing ligand</i>	n.p.	ethanol	20% glycerol/ethanol; TE-G30%-MTG buffer
<i>Concentration range of competing ligand</i>	n.p.	n.p.	0.0001-1000 µM
<i>Volume of ER prep used</i>	n.p.	200 µL	200 µL
<i>No. of replicates</i>	n.p.	n.p.	n.p.
<i>No. of times assay repeated</i>	n.p.	n.p.	n.p.
<i>Time of incubation</i>	18 hours	18 hours or 30 min.	30 min.
<i>Temperature of incubation</i>	4°C	30°C or 4°C	30°C
<i>Measure of nonspecific binding(y/n) and concentration</i>	n.p.	y	y
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	Dextran-coated charcoal (250 mg Norite-A + 25 mg Dextran T-70 in 100 mL TEDM buffer)	Dextran-coated charcoal (0.25% Norit A and 0.025% dextran in TED buffer)	250 µL 60% hydroxyapatite in TEG-MTG buffer
<i>Incubation time and temperature</i>	n.p.	20 min. at 4°C	15 min. at room temp.
<i>Centrifugation speed</i>	800xg	2000xg	1000xg
<i>Centrifugation time and temperature</i>	15 min. at 4°C	10 min. at 4°C	10 min. at 4°C
Data calculations			
<i>Program or method used for calculating data</i>	n.p.	n.p.	Graph Pad Prism
<i>Data plotted as</i>	n.p.	% specific [³ H]-E ₂ binding in controls vs. log concentration (mol/L)	% Specific binding vs. Competitor (M)
<i>Data format in paper (e.g., IC₅₀, K_i)</i>	RBA	IC ₅₀	K _i calculated from EC ₅₀
<i>Calculation of RBA</i>	n.p.	IC ₅₀ E ₂ /IC ₅₀ competitor x 100	n.p.

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity

Assays Using Rat Uterine Cytosol

Reference	Laws et al. (2000)	Leibl and Spona (1982)	Liu et al. (1994)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	Long Evans rats	Sprague Dawley rats	Sprague Dawley rats
<i>Age of animals</i>	Adult (60 days)	Adult (60 - 80 days)	30 days
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	11 days prior to sacrifice	10 days prior to sacrifice	n.p.
<i>Buffer for preparation of cytosol</i>	TE-G30%-MTG (50 mM Tris, 0.9 mM EDTA, 30% (v/v) glycerol, 0.15% (v/v) monothioglycerol, pH 7.4)	TMK buffer (10 mM Tris, 1.5 mM MgCl ₂ , 10 mM KCl, pH 7.2)	TEGD (10 mM Tris, 1.5 mM EDTA, 1 mM dithiothreitol, 10% (v/v) glycerol)
<i>Dilution of tissue with buffer</i>	50 mg/mL buffer	5 uteri/4 mL buffer	n.p.
<i>Protein concentration of cytosol</i>	n.p.	n.p.	n.p.
Competitive binding assay			
<i>Volume and concentration of radiolabelled 17β-estradiol</i>	volume n.p.; 1 nM	volume n.p.; 1 nM	volume n.p.; 1 nM
<i>Specific activity of radioligand</i>	111Ci/nmol	58 Ci/mmol	147 Ci/mmol
<i>Solvent used to dissolve competing ligand</i>	20% glycerol/ethanol; TE-G30%-MTG buffer	ethanol	n.p.
<i>Concentration range of competing ligand</i>	0.0001-1000 μ M	1 pM -1 μ M	1 nM - 10 μ M
<i>Volume of ER prep used</i>	200 μ L	n.p.	200 μ g protein
<i>No. of replicates</i>	n.p.	2	n.p.
<i>No. of times assay repeated</i>	n.p.	n.p.	3
<i>Time of incubation</i>	30 min.	18 hours	2 hours
<i>Temperature of incubation</i>	30°C	4°C	22°C
<i>Measure of nonspecific binding(y/n) and concentration</i>	y	n.p.	y
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	250 μ L 60% hydroxyapatite in TEG-MTG buffer	0.25 mL dextran-coated charcoal (0.6% charcoal Norit A, 0.06% dextran T-60)	100 μ L dextran-coated charcoal (5% charcoal and 0.5% dextran in TEGD buffer)
<i>Incubation time and temperature</i>	15 min. at room temp.	20 min., 4°C	20 min., 4°C
<i>Centrifugation speed</i>	1000xg	3000xg	1500xg
<i>Centrifugation time and temperature</i>	10 min. at 4°C	10 min.; temp. n.p.	10 min. at 4°C
Data calculations			
<i>Program or method used for calculating data</i>	Graph Pad Prism	n.p.	n.p.
<i>Data plotted as</i>	% Specific binding vs. Competitor (M)	% Bound vs. Concentration competitor	[³ H]-E ₂ complex (%) vs. ligand (nM)
<i>Data format in paper (e.g., IC₅₀, K_i)</i>	K _i calculated from EC ₅₀	graphical	graphical (EC ₅₀ estimated)
<i>Calculation of RBA</i>	n.p.	n.p.	n.p.

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity

Assays Using Rat Uterine Cytosol

Reference	McBlain (1987)	Nelson et al. (1973)	Olea et al. (1996)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	Sprague Dawley rats	Sprague Dawley rats	Rats (otherwise unspecified)
<i>Age of animals</i>	4-5 weeks	2-5 months	Immature
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	n.p.	n.p.	n.p.
<i>Buffer for preparation of cytosol</i>	(10 mM Tris, 1.5 mM EDTA, 12 mM monothioglycerol, 10 mM sodium molybdate, 10% (v/v) glycerol, pH 7.4)	10 mM Tris-HCL+1.5 mM EDTA, pH 7.4	Phosphate buffer
<i>Dilution of tissue with buffer</i>	Uteri powdered under liquid N ₂	Uteri from 3-4 mice in 10 mL	n.p.
<i>Protein concentration of cytosol</i>	n.p.	n.p.	2 mg/mL
Competitive binding assay			
<i>Volume and concentration of radiolabelled 17β-estradiol</i>	volume n.p.; 2 nM	volume n.p.; 2 nM	volume n.p.; 3 nM
<i>Specific activity of radioligand</i>	n.p.	48 Ci/mmol	103 Bq/mmol
<i>Solvent used to dissolve competing ligand</i>	ethanol (final conc. 1.5%)	absolute ethanol	ethanol
<i>Concentration range of competing ligand</i>	0.2 nM-20 μ M	0.1 to 500 μ M	0.1 nM to 100 μ M
<i>Volume of ER prep used</i>	n.p.	400 μ g protein	n.p.
<i>No. of replicates</i>	n.p.	n.p.	n.p.
<i>No. of times assay repeated</i>	n.p.	3	n.p.
<i>Time of incubation</i>	18 hours	1 hour	16 hours
<i>Temperature of incubation</i>	4°C	4°C	0-4°C
<i>Measure of nonspecific binding(y/n) and concentration</i>	y	n.p.	y
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	0.5 mL dextran-coated charcoal + 1 mg/mL BSA	0.5 mL activated charcoal and 0.5m% Dextran T40 in Tris/HCL	Dextran + charcoal
<i>Incubation time and temperature</i>	10 min.	15 min. at 4°C	n.p.
<i>Centrifugation speed</i>	12,800xg	2000xg for 5 min.	n.p.
<i>Centrifugation time and temperature</i>	5 min.; temp. n.p.	5 min.; temp. n.p.	n.p.
Data calculations			
<i>Program or method used for calculating data</i>	n.p.	n.p.	n.p.
<i>Data plotted as</i>	[³ H]-E ₂ Bound (% of control) vs. Molar excess of competitor	% Inhibition of [³ H]-E ₂ binding vs. Concentration (μ M)	% Specific binding vs. Concentration (M)
<i>Data format in paper (e.g., IC₅₀, K_i)</i>	IC ₂₅ , IC ₅₀ , K _i	graphical	graphical, RBA
<i>Calculation of RBA</i>	n.p.	n.p.	IC ₅₀ E ₂ /IC ₅₀ competitor x 100

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity

Assays Using Rat Uterine Cytosol

Reference	Perez et al. (1998)	Qian and Abul-Hajj (1990)	Rijks et al. (1996)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	Rats (otherwise unspecified)	Sprague Dawley rats	Sprague Dawley rats
<i>Age of animals</i>	Immature	Immature	Mature
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	n.p.	n.p.	n.p.
<i>Buffer for preparation of cytosol</i>	Phosphate buffer	n.p.	ER buffer (10 mM Tris.HCl, 1.0 mM EDTA, 1 mM dithiothreitol, 10 mM sodium molybdate, 0.25M sucrose, pH 7.4)
<i>Dilution of tissue with buffer</i>	n.p.	n.p.	200 mg/mL buffer
<i>Protein concentration of cytosol</i>	2 mg/mL	3.5 mg/mL	3-4 mg/mL
Competitive binding assay			
<i>Volume and concentration of radiolabelled 125I-estradiol</i>	volume n.p.; 3 nM	50 μ l; concentration n.p.	50 μ l; concentration n.p.
<i>Specific activity of radioligand</i>	103 Bq/mmol	91 Ci/mmol	4.26 TBq/mmol
<i>Solvent used to dissolve competing ligand</i>	ethanol	n.p.	n.p.
<i>Concentration range of competing ligand</i>	10 pM to 100 μ M	1 nM to 3 μ M	10 pM to 2 μ M
<i>Volume of ER prep used</i>	n.p.	150 μ l	50 μ L
<i>No. of replicates</i>	n.p.	2	n.p.
<i>No. of times assay repeated</i>	n.p.	n.p.	2
<i>Time of incubation</i>	16 hours	3 hours	18 hours
<i>Temperature of incubation</i>	4°C	4°C	4°C
<i>Measure of nonspecific binding(y/n) and concentration</i>	y	y	n.p.
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	Dextran + charcoal	Dextran-coated charcoal (10 mM Trizma base, 1.0 mM EDTA, 250 mM sucrose, 0.05% dextran, 0.5% charcoal)	n.p.
<i>Incubation time and temperature</i>	n.p.	15 min. 4°C	n.p.
<i>Centrifugation speed</i>	n.p.	2000xg	n.p.
<i>Centrifugation time and temperature</i>	n.p.	0°C for 5 min.	n.p.
Data calculations			
<i>Program or method used for calculating data</i>	n.p.	n.p.	LIGAND computer program
<i>Data plotted as</i>	% Specific binding vs. Concentration (M)	n.p.	% Bound vs. Concentration competitor (nM)
<i>Data format in paper (e.g., IC₅₀, K_i)</i>	graphical, RBA	RBA	K _i
<i>Calculation of RBA</i>	IC ₅₀ E ₂ /IC ₅₀ competitor x 100	IC ₅₀ E ₂ /IC ₅₀ competitor x 100	K _i reference steroid/ K _i competitor x 100

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity

Assays Using Rat Uterine Cytosol

Reference	Routledge et al. (1998)	Waller et al. (1996)	Zacharewski et al. (1998)
Preparation of receptor			
<i>Species/strain from which receptor obtained</i>	Rats (otherwise unspecified)	Long Evans rats	Sprague Dawley rats
<i>Age of animals</i>	8-10 weeks	Adult (60 days)	22 day old
<i>Source of receptor</i>	Uterus	Uterus	Uterus
<i>Isolated preparation</i>	Cytosol	Cytosol	Cytosol
<i>When ovariectomized</i>	n.p.	11 days prior to sacrifice	n.p.
<i>Buffer for preparation of cytosol</i>	TEGM (10 mM Tris, 1.5 mM EDTA, 3 mM MgCl ₂ , 10% glycerol, pH 7.4)	TE-G30%-MTG (50 mM Tris, 0.9 mM EDTA, 30% (v/v) glycerol, 0.15% (v/v) monothioglycerol, pH 7.4)	TEGD buffer (10 mM Tris base 1.5 mM EDTA, 1 mM dithiothreitol, 10% glycerol, pH 7.6)
<i>Dilution of tissue with buffer</i>	50 mg/mL	50 mg/mL buffer	200 mg/mL buffer
<i>Protein concentration of cytosol</i>	n.p.	n.p.	2 mg/mL
Competitive binding assay			
<i>Volume and concentration of radiolabelled 17β-estradiol</i>	volume n.p.; 5 nM	volume n.p.; 1 nM	30 μ L; 1 nM
<i>Specific activity of radioligand</i>	110 Ci/mmol	111 Ci/nmol	84 Ci/mmol
<i>Solvent used to dissolve competing ligand</i>	n.p.	20% glycerol/ethanol; TE-G30%-MTG buffer	dimethyl sulfoxide
<i>Concentration range of competing ligand</i>	0.5 nM -500 μ M	0.0001-1000 μ M	1-1000 μ M
<i>Volume of ER prep used</i>	100 μ L	200 μ L	240 μ L
<i>No. of replicates</i>	2	n.p.	2
<i>No. of times assay repeated</i>	n.p.	n.p.	3
<i>Time of incubation</i>	18 hours	30 min.	30 min.
<i>Temperature of incubation</i>	4°C	30°C	30°C
<i>Measure of nonspecific binding(y/n) and concentration</i>	n.p.	y	y, 30 μ L
Separation of ligand			
<i>Type of slurry (hydroxyapatite, charcoal, protamine sulfate)</i>	250 μ L 60% hydroxyapatite	250 μ L 60% hydroxyapatite in TEG-MTG buffer	125 μ L 60% hydroxyapatite in TEGD buffer
<i>Incubation time and temperature</i>	n.p.	15 min. at room temp.	n.p.
<i>Centrifugation speed</i>	1,000xg	1000xg	n.p.
<i>Centrifugation time and temperature</i>	10 min.	10 min. at room temp.	n.p.
Data calculations			
<i>Program or method used for calculating data</i>	Ligand Competition Analysis Software (Lundon Software, Chagrin Falls, OH)	one side competitive binding curves (Graph Pad Prism)	n.p.
<i>Data plotted as</i>	% Control vs. Molarity	% Specific binding vs. Competitor (M)	³ H-E ₂ Bound vs. Log concentration of unlabeled competitor (M)
<i>Data format in paper (e.g., IC₅₀, K_i)</i>	graphical	K _i calculated from EC ₅₀	IC ₅₀
<i>Calculation of RBA</i>	n.p.	n.p.	n.p.

Abbreviations: n.p. = not provided; n.a. = not applicable; RBA = relative binding affinity