


Partition of Trend in Incidence Rates for the Time Period 1992-2002 All Races, Both Sexes


Overall Decreasing Regression Coefficient : -2.75



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Percents may not add to 100 due to rounding.

Partition of Trend in Incidence Rates for the Time Period 1992-2002 All Races, Males


Overall Decreasing Regression Coefficient : -7.90



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Percents may not add to 100 due to rounding.

Partition of Trend in Incidence Rates for the Time Period 1992-2002 All Races, Females


Overall Increasing Regression Coefficient : 0.28



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Percents may not add to 100 due to rounding.

Partition of Trend in Death Rates for the Time Period 1992-2002 All Races, Both Sexes


Overall Decreasing Regression Coefficient : -2.13



Source: NCHS public-use file for the total US.
Percents may not add to 100 due to rounding.

Partition of Trend in Death Rates for the Time Period 1992-2002 All Races, Males


Overall Decreasing Regression Coefficient : -3.87



Source: NCHS public-use file for the total US.
Percents may not add to 100 due to rounding.

Partition of Trend in Death Rates for the Time Period 1992-2002 All Races, Females

Overall Decreasing Regression Coefficient : -1.26



Source: NCHS public-use file for the total US.
Percents may not add to 100 due to rounding.