


## 2,4-Diaminotoluene

### CAS No. 95-80-7

Reasonably anticipated to be a human carcinogen  
First Listed in the *Second Annual Report on Carcinogens* (1981)



### Carcinogenicity

2,4-Diaminotoluene is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of carcinogenicity in experimental animals (NCI 1979, IARC 1978, 1986). When administered in the diet, 2,4-diaminotoluene induced hepatocellular carcinomas or neoplastic nodules in rats of both sexes and fibromas of the subcutaneous tissue in male rats; it also increased the incidence of carcinomas or adenomas of the mammary gland in female rats. When administered in the diet, 2,4-diaminotoluene induced hepatocellular carcinomas and may have increased the incidence of lymphomas in female mice. No tumors occurred at significantly increased incidences in male mice. When injected subcutaneously, the compound induced local sarcomas in rats of both sexes (IARC 1978, 1986).

No adequate data were available to evaluate the carcinogenicity of 2,4-diaminotoluene in humans (IARC 1978).

### Properties

2,4-Diaminotoluene is a colorless to brown, needle-shaped crystalline solid or powder that is soluble in water, ethanol, ether, and benzene. When heated to decomposition, it emits toxic fumes of nitrogen oxides. 2,4-Diaminotoluene is available in the United States as a commercial grade; however, the primary use of this chemical is as an intermediate without isolation, and it is typically produced as part of a mixture of approximately 80% 2,4- and 20% 2,6-diaminotoluene. It is also produced in smaller amounts as part of a mixture containing approximately 65% 2,4- and 35% 2,6-diaminotoluene (HSDB 2000).

### Use

The primary use of 2,4-diaminotoluene is as an intermediate in the production of toluene diisocyanate, which is used to produce polyurethane. Small amounts of 2,4-diaminotoluene are used to produce dyes for textiles, leathers, furs, and wood and biological stains. 2,4-Diaminotoluene can be used in the production of approximately 60 dyes, 28 of which are believed to be produced in significant amounts worldwide. The following eight dyes, believed to be produced commercially from 2,4-diaminotoluene, were produced in the United States in 1975: C.I. Basic Brown 4, Basic Orange 1, Direct Brown 154, Direct Black 4, Direct Black 9, Leuco Sulfur Brown 10, Leuco Sulfur Brown 26, and Sulfur Black 2. These dyes are generally used to color silk, wool, paper, and leather. Some are also used to dye cotton fibers and cellulosic fibers, in spirit varnishes and wood stains, as indicators in the manufacture of pigments, and as biological stains (IARC 1978).

2,4-Diaminotoluene is used as a developer for direct dyes, particularly to obtain black, dark-blue, and brown shades, and to obtain navy-blue and black colors on leather. It is also used in dyeing furs. In the U.S., 2,4-diaminotoluene was used in hair dye formulations until this use was banned in 1971 (IARC 1978). Other applications include its use for the preparation of impact resins, polyamides with superior wire coating properties, antioxidants, hydraulic fluids, polyurethane foams, fungicide stabilizers, and as a photographic developer (HSDB 2000).

### Production

In 1978, the IARC reported that 2,4-diaminotoluene had been produced commercially in the United States for over 50 years (IARC 1978); recent U.S. production data for only the 2,4-isomer could not be found in the published literature. Chem Sources (2001) identified seven U.S. suppliers of 2,4-diaminotoluene. In 1977, eight companies were reported to have produced more than 240 million lb and three companies imported 5,500 lb (TSCA 1979). No current data on imports or exports were available.

### Exposure

The primary routes of potential human exposure to 2,4-diaminotoluene are dermal contact and inhalation. The National Occupational Exposure Survey (1981-1983) indicated that 8,513 workers, including 395 women, were potentially exposed to 2,4-diaminotoluene (NIOSH 1990). EPA's Toxic Chemical Release Inventory (TRI) reported that a total of 664 lb of 2,4-diaminotoluene was released to the environment, only to the atmosphere, from five facilities (TRI99 2001).

Potential consumer exposure may occur as a result of the presence of trace contaminants in products that contain 2,4-diaminotoluene-based dyes (e.g., furs, leather, silk, textiles, and wool) (IARC 1978). Furthermore, the compound is a degradation product of the polyester urethane foam used in Meme silicone breast implants (Luu *et al.* 1998). Elevated levels were detected in the urine and plasma of patients up to two years post-operation; the levels ranged from 0.4 to 6 ng/mL (Sepai *et al.* 1995, Luu *et al.* 1998).

### Regulations

#### EPA

##### Clean Air Act

NESHAP: Listed as a Hazardous Air Pollutant (HAP)

NSPS: Manufacture of substance is subject to certain provisions for the control of Volatile Organic Compound (VOC) emissions

##### Comprehensive Environmental Response, Compensation, and Liability Act

Reportable Quantity (RQ) = 10 lb

##### Emergency Planning and Community Right-to-Know Act

Toxics Release Inventory: Listed substance subject to reporting requirements

##### Resource Conservation and Recovery Act

Listed as a Hazardous Constituent of Waste

### REFERENCES

- ChemSources. 2001. Chemical Sources International, Inc. <http://www.chemsources.com>.
- HSDB. 2000. Hazardous Substances Data Base. National Library of Medicine. <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>.
- IARC. 1979. Some Monomers, Plastics, and Synthetic Elastomers, and Acrolein. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 19. Lyon, France: International Agency for Research on Cancer. 513 pp.
- IARC. 1986. Some Chemicals Used in Plastics and Elastomers. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 39. Lyon, France: International Agency for Research on Cancer. 403 pp.
- Luu, H. M., J. C. Hutter and H. F. Bushar. 1998. A physiologically based pharmacokinetic model for 2,4-toluenediamine leached from polyurethane foam-covered breast implants. *Environ Health Perspect* 106(7): 393-400.
- NCI. 1979. Bioassay of 2,4-Diaminotoluene for Possible Carcinogenicity (CAS no. 95-80-7). Technical Report Series No 162. DHEW (NIH) Publication No. 79-1718. Bethesda, MD: National Institute of Health. 122 pp.
- NIOSH. 1990. National Occupational Exposure Survey (1981-83). Unpublished provisional data as of 7/1/90. Cincinnati, OH: U. S. Department of Health and Human Services.
- Sepai, O., D. Henschler, S. Czech, P. Eckert and G. Sabbioni. 1995. Exposure to toluenediamines from polyurethane-covered breast implants. *Toxicol Lett* 77(1-3): 371-8.
- TRI99. 2001. Toxic Chemical Release Inventory 1999. Data contained in the Toxic Chemical Release Inventory (TRI). National Library of Medicine. <http://www.epa.gov/triexplorer/>.
- TSCA. 1979. Toxic Substances Control Act, Chemical Substances Inventory.