

**Table 1 (continued). Cancer incidence trends by joinpoint analyses\* for 1973 through 1999 for the most common cancers by race, sex, and age, SEER†, 1973-1999.**

Site/Race/Sex		Age		Joinpoint Analyses* (1973-1999)									
				Trend 1		Trend 2		Trend 3		Trend 4			
				Range of Years	APC‡	Range of Years	APC‡	Range of Years	APC‡	Range of Years	APC‡		
<b>Female Breast</b>													
<b>All Races</b>		<b>All ages</b>		1973-80	-0.6	1980-87	3.7 §	1987-99	0.5 §				
		<b>&lt;50</b>		1973-80	-1.5	1980-86	2.9 §	1986-99	-0.3				
		<b>50-64</b>		1973-80	-1.2	1980-85	4.1 §	1985-99	1.6 §				
		<b>65-74</b>		1973-78	-1.2	1978-87	4.6 §	1987-99	0.4				
		<b>75+</b>		1973-82	0.8	1982-87	5.2 §	1987-99	0.0				
<b>White</b>		<b>All ages</b>		1973-80	-0.5	1980-87	3.8 §	1987-99	0.4				
		<b>&lt;50</b>		1973-80	-1.4 §	1980-86	2.9 §	1986-99	-0.4				
		<b>50-64</b>		1973-80	-1.1	1980-85	4.4 §	1985-99	1.5 §				
		<b>65-74</b>		1973-78	-1.0	1978-87	4.7 §	1987-99	0.3				
		<b>75+</b>		1973-82	0.8	1982-87	5.4 §	1987-99	0.0				
<b>Black</b>		<b>All ages</b>		1973-79	-0.7	1979-86	3.9 §	1986-99	0.9 §				
		<b>&lt;50</b>		1973-91	1.4 §	1991-99	-1.1						
		<b>50-64</b>		1973-99	2.1 §								
		<b>65-74</b>		1973-99	2.2 §								
		<b>75+</b>		1973-99	1.7 §								

\* Joinpoint analysis of trends allowed for up to three joinpoints based on rates per 100,000 that were age-adjusted to the 2000 U.S. standard population.

† SEER (Surveillance, Epidemiology, and End Results) registries covering 10% of the U.S. population: metropolitan areas of San Francisco, Detroit, Atlanta, and Seattle-Puget Sound; and States of Connecticut, Hawaii, Iowa, Utah, and New Mexico.

‡ APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population using the joinpoint regression program.

§ APC is statistically significantly different from zero (two-sided p<.05).

‡ Joinpoint results are not shown if less than 10 cases in any year.