

Some Environmental Contaminants Influence Motor and Feeding Behaviors in the Ornate Wrasse (*Thalassoma pavo*) via Distinct Cerebral Histamine Receptor Subtypes

Giuseppina Giusi,¹ Rosa Maria Facciolo,¹ Raffaella Alò,¹ Antonio Carelli,¹ Maria Madeo,¹ Pietro Brandmayr,² and Marcello Canonaco¹

¹Comparative Neuroanatomy Laboratory, and ²Zoocenoses Laboratory, Ecology Department, University of Calabria, Cosenza, Italy

Common environmental contaminants such as heavy metals and pesticides pose serious risks to behavioral and neuroendocrine functions of many aquatic organisms. In the present study, we show that the heavy metal cadmium and the pesticide endosulfan produce such effects through an interaction of specific cerebral histamine receptor subtypes in the teleost ornate wrasse (*Thalassoma pavo*). Treatment of this teleost with toxic cadmium levels for 1 week was sufficient to induce abnormal swimming movements, whereas reduced feeding behaviors were provoked predominantly by elevated endosulfan concentrations. In the brain, these environmental contaminants caused neuronal degeneration in cerebral targets such as the mesencephalon and hypothalamus, damage that appeared to correlate with altered binding levels of the three major histamine receptors (subtypes 1, 2, and 3). Although cadmium accounted for reduced binding activity of all three subtypes in most brain regions, it was subtype 2 that seemed to be its main target, as shown by a very great ($p < 0.001$) down-regulation in mesencephalic areas such as the stratum griseum central layer. Conversely, endosulfan provided very great and great ($p < 0.01$) up-regulating effects of subtype 3 and 1 levels, respectively, in preoptic-hypothalamic areas such as the medial part of the lateral tuberal nucleus, and in the suprachiasmatic nucleus. These results suggest that the neurotoxicant-dependent abnormal motor and feeding behaviors may well be tightly linked to binding activities of distinct histamine subtypes in localized brain regions of the *Thalassoma pavo*. **Key words:** cadmium, diencephalon, endosulfan, histamine receptor subtype, mesencephalon, teleost fish. *Environ Health Perspect* 113:1522–1529 (2005). doi:10.1289/ehp.7983 available via <http://dx.doi.org/> [Online 14 July 2005]

A number of neurotoxic environmental contaminants, recognized as endocrine disruptors, have aroused much interest in the field of neuroendocrinology (Pillai et al. 2003). In particular, polycyclic aromatic hydrocarbons, polychlorinated biphenyls, and many heavy metals such as arsenic, cobalt, and mercury in the organic form appear to impair growth, development, and sociosexual behavior in vertebrates (Beauvais et al. 2001; Bisson and Hontela 2002). Cadmium is one of the heavy metals that pose an increasing health threat to ecologic communities and humans (Harvey et al. 1999). Because of widespread industrial applications such as the use of alloys for metal coatings and nickel-cadmium batteries as well as the burning of fossil fuels, urban traffic, and waste incineration, this pollutant is emitted into the atmosphere (Liao and Freedman 1998). It is taken up readily by humans and other mammals not only via inhalation but also via the food web (Waalkes et al. 1992). By binding to cysteine residues or generating reactive oxygen species (Risso-de Faverney et al. 2001), Cd has been shown to influence genomic and postgenomic processes in liver, kidney, lung, and brain (Minami et al. 2001). Similar alterations appear to be linked with neuronal dysfunctions in the hypothalamic–pituitary–testicular axis and inadequate neurosecretory activities of pituitary cells (Lafuente et al. 2001). Interestingly, these same degenerative processes in the preferential

brain region, that is, the olfactory bulb, appear to corroborate evidence that foraging and aggression controlled by this region constitute vital behaviors in fish (Griffiths and Armstrong 2000; Scherer et al. 1997).

The persistence and accumulation of the insecticide endosulfan (6,7,8,9,10,10-hexachloro-1,5,5a,6,9,9a-hexahydro-6,9-methano-2,4,3-benzodioxathiepin-3-oxide) in human tissues are of concern (Martinez et al. 1997). Its resistance to biologic degradation and its low water solubility not only favor binding to soil particles and persistence within surface waters but also prompt bioconcentrations resulting in up to 600 times ambient water concentration in some species (Miller and Llados 1999). The purpose of this insecticide was to protect economically important crops such as tobacco and cotton; little attention was paid to neurologic risks to humans or other animals in terrestrial and aquatic ecosystems. Recent studies show that similar pesticides alter aggressive and reproductive behaviors in teleost fish (Carlson et al. 1998; Roex et al. 2001) through the interference of cerebral neuromediating systems (Clark 1997) such as the histaminergic system. This system is actively involved in blocking chemical-dependent stressful conditions such as writhing and consequently modifies responses to cold and immobilization states (Ferretti et al. 1998; Ito 2000).

The histamine (HA) receptor complex is one of the main neurosignaling systems that, in addition to “allergic” and anti-inflammatory processes, has been recognized for its role in many neurologic functions such as attention, arousal, cognition, movement, and feeding in mammalian species (Lin et al. 1996). Structurally, the histaminergic neuronal fibers, originating from the hypothalamus, are projected extensively throughout the central nervous system and promote their actions via three distinct receptor subtypes denoted H_nR (H₁R, H₂R, H₃R) (Hill et al. 1997). All but H₃R subtypes are postsynaptically located and are coupled positively to adenylyl cyclase and phospholipase C, whereas H₃R has been isolated at both the presynaptic and postsynaptic levels (Brown et al. 2001). Recently, Spieler et al. (1999) observed that inhibition of the H₁R site is linked to the improvement of appetitive reversal learning and memory tasks in the goldfish *Carassius auratus*—a relationship that was further supported by the detection, via the application of their specific and selective antagonists, of H₃R (Peitsaro et al. 2000) and H₁R (Choich et al. 2004) in brain areas of the zebrafish *Danio rerio* and the tilapia *Oreochromis niloticus*, respectively. Moreover, identification of these subtypes in other vertebrates such as amphibians and reptiles (Brodin et al. 1990; Inagaki et al. 1991) is consistent with their highly conserved profile throughout vertebrate phylogeny (Kaslin and Panula 2001). On the basis of the above data, the intention of our study was to establish whether Cd and endosulfan neurotoxicologic effects on behavioral activities could be controlled through cerebral histaminergic neuronal mechanisms in the ornate wrasse. In fish, adaptations to environmental variables, including chemical–physical water properties such as temperature, photoperiod, ionic balance, and

Address correspondence to M. Canonaco, Comparative Neuroanatomy Laboratory, Ecology Department, University of Calabria, 87030 Rende (CS), Italy. Telephone: 0039-984-492974. Fax: 0039-984-492986. E-mail: canonaco@unical.it

This study was supported partially by the contract sponsor MEMO-BIOMAR research program and COFIN (Cofinancial Projects) of MIUR (Italian University Research Ministry).

The authors declare they have no competing financial interests.

Received 1 February 2005; accepted 14 July 2005.

pollutants (Huang et al. 2004), are involved in the acceleration of the diandric-protogynic physiologic state, which makes fish valuable models for the investigation of neurologic adaptation mechanisms (Bass and Grober 2001; Larson et al. 2003). Furthermore, the host-cleaning symbiosis of the ornate wrasse *Thalassoma pavo* not only is critical for the stability of ecosystems (Zander and Sörje 2002) but also improves environmental conditions for commercially valuable fish and thus makes the ornate wrasse an important ecologic biomarker species.

Materials and Methods

Animals. We collected 32 young female ornate wrasses (body weight, 20–25 g; length, ~16–18 cm) from the Tyrrhenian Sea; they were acclimated to laboratory conditions for 1 week. During this period, fish were maintained in flow-through tanks containing 150 L of seawater (19–21°C, pH 7.8) under a 12/12-hr light/dark photoperiod and fed daily (10 g/kg body weight) with commercial food (Morubel, Milan, Italy) corresponding to 2% of biomass in the tank. Animal maintenance and experimental procedures were in accordance with the *Guide for Use and Care of Laboratory Animals* (European Communities Council Directive 1986), and efforts were made to minimize animal suffering and reduce number of specimens used.

Experimental treatments and behavioral observations. In the first part of the study, fish ($n = 12$) were exposed for 1 week to one of the two concentrations of Cd ($\text{CdCl}_2 \cdot 2\text{H}_2\text{O}$; Sigma, Milan, Italy): a sublethal concentration (2.26 mg/L; $n = 6$) or a calculated maximum

acceptable toxicant (MAT) concentration (11.32 mg/L; $n = 6$), which were both less than the 96-hr LC_{50} (lethal median concentration) value of 28.68 mg/L (Giusi et al. 2004) and the 96-hr LC_{50} value of 20.12 mg/L obtained in the white sea bass, *Lates calcarifer* (Thophon et al. 2003). This contaminant was prepared by dissolving $\text{CdCl}_2 \cdot 2\text{H}_2\text{O}$ in appropriate volumes of seawater. Other fish ($n = 12$) were exposed for the same length of time to two concentrations of endosulfan (Sigma): a sublethal concentration (0.2 $\mu\text{g/L}$; $n = 6$) and a calculated MAT concentration (1.3 $\mu\text{g/L}$; $n = 6$) that were both less than the 96-hr LC_{50} value of 3.30 $\mu\text{g/L}$ (Giusi et al. 2004), and the 96-hr LC_{50} value varying between 1.4 and 1.5 $\mu\text{g/L}$ for freshwater fish (Johnson and Finley 1980), sublethal and MAT concentrations that fall within the ranges reported in surface waters (0.039–0.205 $\mu\text{g/L}$) and after runoff water events (0.01–1.3 $\mu\text{g/L}$) from agricultural areas, respectively (Naqvi and Vaishnavi 1993). The pesticide was dissolved in seawater. We compared both treatment groups with controls ($n = 8$) consisting of fish maintained under identical conditions except that only vehicle was added to the tanks. During the entire experiment, accumulation of wastes and pathogens was avoided by replacing the tanks with fresh seawater every day. To reduce to a minimum the stressful conditions of this operation, we rapidly transferred the fish with a small fishing net to the tanks containing fresh seawater and either one of the two contaminants in an effort to achieve the intended nominal concentrations. Throughout the behavioral sessions, we checked the feeding habits of the ornate wrasse to ensure that the fish ingested water containing

the neurotoxicant, although the uptake of these compounds primarily depends on their passage through the gill system (Thophon et al. 2003).

We assessed the behavior and mortality for all fish that received either Cd or endosulfan in four 1-hr sessions each day for 1 week. The motor and feeding behaviors that we analyzed included hyperactive movements consisting in either swimming toward the surface, swimming in the same direction, or “bumping” into each other or against the tank; assuming a “relaxed” position or simply being inactive; and hyperventilation, which is defined as the number of times that the operculum opens and closes in a 1-hr observation session. We also recorded feeding frequency and quantity (milligrams) of food ingested during each observation session. The quantity of food ingested was determined after the residual food recovered at the bottom of the tank was dried and weighed. The above motor activities and feeding behaviors of both treatment and control groups, estimated as mean activity per 24 hr \pm SE, were recorded with a digital video camera (model TR 7000 E; Sony, Tokyo, Japan) and elaborated at a personal computer (Microsoft Windows XP; Microsoft Corp., Redmond, WA) using EthoLog software (version 2.2.5; Visual Basic, São Paulo, Brazil) for behavioral analyses.

Amino cupric silver staining. To establish whether abnormal behavioral activities were related to neuronal damage, fish treated in the same manner as in the behavioral study with sublethal ($n = 3$) and MAT ($n = 3$) concentrations of Cd and endosulfan were decapitated and their brains quickly removed (within 30 sec) and stored at -40°C according to common cryostat procedures for unfixed




Figure 1. Assessment of effects of Cd and endosulfan on some motor activities of *Thalassoma pavo*: (A) hyperactive movements, (B) movements toward surface, (C) feeding frequency, (D) quantity (mg) of food ingested, and (E) hyperventilation activity. For these behaviors a sublethal concentration of endosulfan and Cd as well as a MAT concentration of the two contaminants, respectively, were compared with controls. Values (means of activities/24 hr \pm SE) of motor activities and feeding behaviors were estimated daily during four 1-hr observations for 1 week, as described in “Materials and Methods.” The behavioral data were analyzed by one-way ANOVA followed where necessary by post hoc Neuman-Keuls multiple-range test.

* $p < 0.05$; *** $p < 0.001$.

brains (Canonaco et al. 1997). Brains were mounted on a freezing stage of a sliding cryostat (Microm-HM505E; Zeiss, Wallford, Germany), and a serial set of representative coronal sections (30 μm) was selected at an interval of 240 μm for amino cupric silver

staining procedures according to previously published methods (de Olmos et al. 1994); we adapted the exposition time (25 min) to neutral red for the different brain sections of our fish species. This method, which has been used mainly for mammalian brain studies, proved to

be appropriate for cerebral neuronal fields that have undergone degeneration processes. The brain sections were rinsed with distilled H_2O , placed into dishes containing the preimpregnating solution (silver nitrate [AgNO_3], distilled H_2O , D,L-alanine, copper nitrate [$\text{Cu}(\text{NO}_3)_2$], cadmium nitrate [$\text{Cd}(\text{NO}_3)_2$], lanthanum nitrate [$\text{La}(\text{NO}_3)_2$], neutral red, pyridine triethanolamine, isopropanol), heated in a microwave oven (45–50°C) for 50 min, and cooled at room temperature for 3 hr. The sections were then rinsed in distilled H_2O , and after a quick rinse in acetone they were placed in an impregnating solution AgNO_3 , distilled H_2O , ethanol, acetone, lithium hydroxide (LiOH), ammonium hydroxide (NH_4OH) for 50 min, followed by a 25-min fixation in a reducer solution (formalin, citric acid monohydrate, ethanol, distilled H_2O) at a temperature range of 32–35°C. These sections were left overnight in distilled H_2O , and the next day they were placed in a first bleaching solution [potassium ferricyanide in potassium chlorate solution, lactic acid] for 60 sec at room temperature. Afterward, they were bleached in a second bleaching solution (potassium permanganate, sulfuric acid) for 60 sec and rinsed in distilled H_2O . For the stabilization phase, sections were transferred in sodium thiosulfate solution and rinsed again in distilled H_2O . Finally, they were immersed in a rapid fixer solution for 5 min and counterstained with 0.5% neutral red solution (Carlo Erba, Milan, Italy) for 25 min, dehydrated in ethanol (50–100%) and xylene, and mounted with DPX (*p*-xylene-bis[*N*-pyridinium bromide]; Sigma) for observations with a bright-field Dialux EB 20 microscope (Leitz, Stuttgart, Germany). The effects of both neurotoxicants on the argyrophilic reaction at the different brain levels were compared with controls that consisted of fish maintained under conditions identical to those of the two treatment groups except that only vehicle was added to the tanks. Because the same negative results were obtained at all brain levels, only two representative posterior areas were illustrated and compared with the different brain areas of the two treatment groups.

Effects of Cd and endosulfan on H_1R – H_3R

The neurotoxic actions of Cd and endosulfan were also correlated with the type of distribution pattern of the H_1R neuronal system. Fish treated with the sublethal ($n = 4$) and MAT ($n = 4$) concentrations of Cd and endosulfan along with their control ($n = 6$) were used. The brains were removed and quickly frozen for storage at -40°C , and brain sections (14 μm thick) were thawed, dried at room temperature, and then handled according to *in vitro* binding studies for mammals (Ryu et al. 1996) that were adapted for fish brain sections (Peitsaro et al. 2000). Briefly, we incubated sections in 150 mM sodium potassium phosphate buffer




Figure 2. Photomicrographs showing the amino cupric silver staining pattern in rostral (i), middle (j), and posterior (k) brain areas of the *Thalassoma pavo*, treated with a MAT concentration of (a–c) Cd or (e–g) endosulfan. The effects of Cd ($n = 6$; arrows) were mostly observed in telencephalic and in mesencephalic areas such as Dm2 (a) and in the piriform SGC neurons of the optic tectum (b), respectively, and in the pretectal NGa (c), compared with control ($n = 8$); controls gave comparable results at all brain levels for both neurotoxicants as described in “Materials and Methods,” and so these same controls (d, h) were also used for the effects of endosulfan. In the case of endosulfan ($n = 6$), the major effects (arrows) were detected in the interneurons of the entopeduncular nucleus (e) and in the preoptic NSC area (f) and NLTm (g) of the hypothalamic lobe.

Abbreviations: CP, central posterior thalamic nucleus; Dc2, central part of dorsal telencephalon, subdivision 2; DI, lateral part of the dorsal telencephalon; Dm2–Dm4, medial part of the dorsal telencephalon, subdivisions 2–4; DP, dorsal posterior thalamic nucleus; E, entopeduncular nucleus; NAP, anterior periventricular nucleus; NAT, anterior tubular nucleus; NGa, anterior part of the nucleus glomerulosus; NH, habenular nucleus; NLTm, medial part of lateral tubular nucleus; NLTv, ventral part of lateral tubular nucleus; NPGm, medial preglomerular nucleus; NPO, preoptic nucleus; NPP, posterior periventricular nucleus; NSC, supra-chiasmatic nucleus; NT, nucleus taenia; OT, optic tectum; POA, preoptic area; PSp, parvocellular superficial pretectal nucleus; SCO, subcommissural organ; TLo, torus longitudinalis; VCe, cerebellum valvula; VM, ventromedial thalamic nucleus; Vot, ventral optic tract; Vp, postcommissural nucleus of the ventral telencephalon.

(Sigma) 2 mM MgCl₂ and 100 μM dithiothreitol pH 7.4 (Roche Diagnostic, Milan, Italy) containing different concentrations (0.5–20 nM) of [³H]-*N*-α-methyl-HA (NAMH; PerkinElmer Life Sciences, Boston, MA, USA). Some sections were incubated with 10 nM [³H]-NAMH using a wipe assay procedure. This concentration displayed the greatest affinity for H₁R in the presence of the different values (1 μM–1 nM) of the following specific HA antagonists (Sigma): H₁R antagonist pyrilamine, H₂R antagonist cimetidine, and H₃R antagonist thioperamide. Other sections were incubated with 10 nM [³H]-NAMH plus 500 μM of their corresponding antagonist for nonspecific binding values that proved to be similar to that of the background; subsequently an autoradiographic film (Hyperfilm; Amersham, Piscataway, NJ, USA) was apposed to dried sections and to slides containing plastic standards.

After an exposure period of 6 weeks (25°C), we developed the autoradiographic films according to previous methods (Canonaco et al. 1997) and we evaluated the different H₁R–H₃R binding densities, expressed in femtomole per milligram wet tissue weight, with a Panasonic Telecamera (objective lens FD; 50 mm, 1:3.5; Canon, Milan, Italy) attached to a Macintosh computer-assisted image analyzer system running Scion-Image 2.0 (National Institutes of Health Image, Bethesda, MD, USA). We stained labeled sections with cresyl violet acetate to identify the diencephalic, mesencephalic, and telencephalic brain regions, using the perch fish atlas (Cerdá-Reverter et al. 2001a, 2001b).

Statistical analysis. For the receptor binding study, Scatchard analyses of saturation binding data, which were fitted by a one-site and/or two-site model [based on the significance of extrasum squares using a LIGAND program (Munson and Rodbard 1980)] supplied relative affinity states and maximal binding receptor binding densities. To compare behavioral observations and histaminergic receptor

binding data, we compared the treatment groups using a one-way analysis of variance (ANOVA) when there was a significant *p*-value ≤ 0.05, according to the Neuman-Keuls multiple-range post hoc test.

Results

Behavioral analysis. Treatment of the ornate wrasse with Cd and endosulfan accounted for a net differentiation in the type of behavior responses. The MAT concentration of both stressors—11.32 mg/L and 1.3 μg/L, respectively—induced stereotype motor behaviors during the entire experimental session. Fish treated with Cd at 11.32 mg/L exhibited greater (*p* < 0.001; Figure 1A) hyperactive swimming activities such as moving in only a vertical direction and/or “bumping” against each other or against the glass tanks, in contrast to controls, which were often inactive and spent most of their time along the bottom of the tank. Fish treated with a concentration of 2.26 mg/L Cd displayed only moderate stereotype behaviors (Figure 1B), including hyperactive movements that consisted of swimming mainly in a vertical direction toward the surface of the water, whereas controls exhibited more random movements. Conversely, endosulfan caused a significant increase of some hyperactive movements (*p* < 0.05; Figure 1A,B) such as swimming in a vertical direction, whereas “bumping” type of swimming behaviors occurred in a less significant manner. This pesticide markedly reduced feeding, even at the lower concentration (0.2 μg/L). With both concentrations of endosulfan tested, feeding behavior was irregular, and overall, treated animals ate less food than did controls (Figure 1C,D). The MAT concentrations of both contaminants caused an excessive production of mucus on the operculum surface and, after 24 hr, hyperventilation became increasingly more severe up to the end of the study (Figure 1E).

Analysis of amino cupric silver-stained tissue. From the amino cupric silver staining

analysis, it was possible to correlate these abnormal behaviors with evident neurodegeneration processes in telencephalic and mesencephalic regions. In particular, a MAT concentration of Cd supplied damaged external pyramidal neuron, as exhibited by a typically argyrophilic dark neuronal perikarya and often by a shrunken and folded appearance compared with little or no damage in controls (Figure 2d,h). This feature was limited mainly to the medial dorsal part of the telencephalon, subdivision 2 (Dm2; Figure 2a), and the pyramidal layer of the mesencephalic stratum griseum central (SGC; Figure 2b), which showed consistent dark axonal processes. The effects of Cd seemed to extend to other areas of the brain, namely, the anterior part of the nucleus glomerulosus (NGa; Figure 2c) of the diencephalic pretectal region that is involved, via mesencephalic circuits, with the regulation of visual motor functions in teleosts (Kaslin and Panula 2001). With endosulfan, substantial neurodegeneration was present in ventral telencephalic regions such as the entopeduncular nucleus (e; Figure 2e) plus the diencephalic suprachiasmatic nucleus (NSC; Figure 2f) and the medial part of lateral tuberal nucleus (NLTm; Figure 2g). In these brain regions endosulfan produced an altered pattern of neurons defined as an “interrupted string of pearls” as noted with degeneration of interneurons of mammals (Siegel et al. 1999).

Effects of Cd and endosulfan on H₁R–H₃R

When the regional distribution of HA receptors was determined in the presence of Cd and endosulfan, we observed a peculiar pattern of histaminergic expressing neurons in the same above brain regions of *Thalassoma pavo*. Such a relationship was based on a similar optimal [³H]-NAMH binding constant (Figure 3) in both treated and control fish with respect to that of rodents (unpublished data). Overall, the




Figure 3. (A) A saturation curve of [³H]-NAMH binding (fmol/mg wet tissue weight ± SE), using wipe assays, was determined for the preoptic area of the *Thalassoma pavo* treated with MAT concentrations of Cd and endosulfan and compared with controls as described in “Materials and Methods.” (B) From the linear Scatchard plot, the negative slope was calculated to provide the mean dissociation constant (nM), whereas the intercept of the curve at the abscissa provided the maximal number of binding sites. Evaluation of saturation-binding study supplied similar results in three separate trials.




Figure 4. Displacement curves of [³H]-NAMH (% of total binding) in preoptic area of the *Thalassoma pavo* (*n* = 6) were generated in the presence of different concentrations (1 μM to 1 nM) of cold NAMH and of selective HA antagonists thioperamide, pyrilamine, and cimetidine as described in “Materials and Methods.” Each point represents mean ± SE of three separate tests.

highest ($> 140 < 200$ fmol/mg wet tissue weight) HA binding densities were shown to be typical of rostral areas such as the preoptic nucleus (NPO) as well as the torus longitudinalis (TL_o) and SGC of midbrain regions, whereas lower ($> 70 < 110$ fmol/mg wet tissue weight) binding densities were reported for the central nucleus of the ventral telencephalon and molecular stratum of the cerebellum. Application of the selective HA receptor antagonists enabled us to demonstrate that it was the diencephalic region that proved to be a preferential target of the major distribution differences of all subtypes (H_1R-H_3R), as displayed by notable displacement capacities of these subtypes in the preoptic area (Figure 4), as well as high H_1R and H_2R levels in areas such as NPO (45%) and in the nucleus of the saccus vasculosus (NSV; 43%), respectively (Figure 5). The subtype H_3R was predominantly higher in some regions and especially in Dm2 (45%) of the telencephalon and in TL_o (44%) of the mesencephalon.

It is noteworthy that fish treated with a MAT Cd dose showed a down-regulating effect of H_2R -expressing neurons, as displayed by the low binding densities in some midbrain regions of the representative autoradiograms (Figure 6). Of all the regions examined, SGC (−115%; $p < 0.001$) and NGa (−90%) of the mesencephalon (Figure 7A) seemed to contain the greatest down-regulating activities of H_2R -producing neurons, whereas a moderate ($p < 0.05$) reduction was evident in the habenular nucleus (NH; −45%). A similar reduction that appeared to be also maintained for H_1R -producing neurons and precisely a very great and great ($p < 0.01$) reduction of H_1R levels in TL_o (−105%) and the central posterior thalamic nucleus (CP; −65%), respectively, whereas a moderate up-regulating activity was instead detected for NLTm (+38%; Figure 7B). The subtype H_3R did not appear to be a major target of Cd (Figure 7C) aside from the moderately higher levels (+40%) obtained in the external cellular layer (ECL) of the olfactory bulb.

The effects of endosulfan appeared instead to be preferentially directed toward H_3R -producing neurons, as shown by the greater binding densities in the representative autoradiograms of midbrain regions (Figure 6b). The diencephalic region (Figure 8A) provided very great up-regulating effects, especially in the NLTm (+110%) and the nucleus of the posterior hypothalamic recess (NRP; +78%). By contrast, greatly decreased levels were detected in another hypothalamic area, that is, the ventromedial thalamic nucleus (VM; −70%). Moreover, the H_1R -producing neurons of this brain region were a preferred target for endosulfan effects (Figure 8B), as indicated by the greatly increased levels in the NSC (+68%) plus moderately higher levels in the NSV (+40%). Conversely, the other subtype (H_2R) did not prove to be a preferred target of this pesticide (Figure 8C) despite the moderately higher H_2R levels in NPO (+60%).

Discussion

We describe here for the first time neurotoxicologic effects of the heavy metal Cd and the insecticide endosulfan that are responsible for abnormal motor and feeding behaviors through histaminergic mechanisms in the ornate wrasse. A first abnormal behavior consisted of stereotype motor activities such as swimming in a constant direction or “bumping” against each other and/or against the glass tank, especially when the fish were treated with a MAT Cd concentration. These abnormal behaviors, as reported previously for *Thalassoma pavo* observed under field conditions (Giusi et al. 2004) and in a wide variety of fauna ranging from terrestrial vertebrates such as rodents (Lafuente et al. 2001) to aquatic species such as amphibians (James et al. 2004) and Chordata Ascidaacea (Bellas et al. 2001), should not be surprising because of wide distribution of this heavy metal in the different ecosystems. This condition appears mainly in aquatic communities because Cd readily accumulates in the different tissues after

uptake via the calcium transport pathway of gill's chloride cells (Wood 2001), above all in the olfactory structures that are considered to be its preferential target (Tallkvist et al. 2002). In this context the interference of such sensory communicating structures in the ornate wrasse may offset normal responses to olfaction-mediated stimuli such as migration and physical contact with other fish, which is in accordance with the irregular responses to alarm signals as well as modification of aggressive social relationships that have been reported in rainbow trout treated with toxic Cd doses (Scott et al. 2003).

In pesticides both sublethal and MAT concentrations of endosulfan caused abnormal feeding behaviors, whereas altered swimming movements were less evident than in Cd-treated animals. As a consequence, the consumption of food at an asynchronous rhythm and at different time intervals is in good agreement with other pesticides, accounting for feeding difficulties via neuronal functional hindrances in the goldfish (Bretaud et al. 2000). Similar difficulties obtained even under sublethal concentrations tends to suggest that sensorimotor threshold activities are susceptible to this contaminant, as shown by “startled” motor behaviors being tightly associated with the olfactory-dependent neuromodiation of




Figure 5. Percentage binding levels (of total) ± SE of H_1R , H_2R , and H_3R sites in diencephalic (A) and extradiencephalic (B) regions of the *Thalassoma pavo* ($n = 6$) were determined in the presence of their respective selective antagonists as described in “Materials and Methods.”

Abbreviations: Dm2, medial part of the dorsal telencephalon, subdivision 2; NPO, preoptic nucleus; NRP, nucleus of the posterior hypothalamic recess; NSC, supra-chiasmatic nucleus; NSV, nucleus of the saccus vasculosus; SGC, stratum griseum centrale; TL_o, torus longitudinalis; VM, ventromedial thalamic nucleus.




Figure 6. Representative binding autoradiograms displaying distinct receptor densities (black line) of H_2R in the posterior regions of the *Thalassoma pavo* treated with a MAT concentration of Cd (a; $n = 4$) and of H_3R in the same brain regions of animals that, instead, received a MAT concentration of endosulfan (b; $n = 4$), with respect to their corresponding (e, f) controls ($n = 6$). Binding pattern of these two subtypes appeared to be highly specific as shown by very similar background levels reported for [3H]-NAMH in presence of a 500× concentration of the selective antagonists cimetidine (c) and thioperamide (d), respectively, as described in “Materials and Methods.”

Abbreviations: NAT, anterior tubular nucleus; NGa, anterior part of the nucleus glomerulosus; NLTm, medial part of lateral tubular nucleus; OT, optic tectum; TL_o, torus longitudinalis; VM, ventromedial thalamic nucleus.

optomotor responses such as predation, foraging, and orientation toward food odor (Pan and Dutta 1998). These olfactory-dependent responses appear to be determining elements for feeding behaviors throughout the various biologic phases of the fish, as demonstrated by both young and adult Japanese killifish being neither attracted to nor able to consume food after receiving similar endosulfan doses (Gormley and Teather 2003).

When the toxicologic actions of both environmental contaminants were assessed at the structural level of the brain, notable neurodegenerative events were observed, as shown by the diffused amino cupric silver staining of neurons in the different brain regions. With this method it was possible not only to immediately and clearly detect the precise location of neuronal trauma (Siegel et al. 1999) but also to distinguish between somata and axonal damage in some diencephalic, mesencephalic, and telencephalic sites of *Thalassoma pavo*. Of the brain regions exposed to MAT Cd concentrations, the telencephalic Dm2 displayed the greatest axonal fiber damage and interstitial edema. This condition fits nicely with the infiltration properties of the heavy metal in mammalian telencephalic regions such as the hippocampus (Mendez-Armenta et al. 2001), which is involved in analogous functions such

as learning, spatial memory, and motor behaviors that are controlled by Dm2 in fish (Portavella et al. 2004). Even SGC and NGa of mesencephalic and pretectal areas, respectively, which are related to the modulation of multisensorial inputs (visual, acoustic, and electroreceptive signals), supplied perturbed dendritic spine formation and deformed soma in a fashion similar to that of mesencephalic trigeminal neurons of rodents exposed to Cd (Yoshida 2001). The effects of endosulfan were instead involved mainly with axonal deformations of interneurons in diencephalic areas such as NSC and preoptic area of the hypothalamus, an effect that tends to overlap cellular alterations and interstitial infiltration events induced by the pesticide carbofuran in teleosts (Ram et al. 2001). Ram et al. (2001) also showed that such a contaminant was responsible for a reduction in number and size of neurons and consequently altered neurotransmission functions in this same brain region.

Interestingly, the neuronal alterations provoked by both environmental contaminants in the present study seemed to coincide with changes of the histaminergic transcriptional activities in some telencephalic and mesencephalic regions and in the anterior and posterior areas of the hypothalamus. The hypothalamic area is considered to be a key

production site of HA not only in mammals (Pillot et al. 2002) but also in amphibians (Airaksinen and Panula 1990) and in some fish species such as the zebrafish (Kaslin and Panula 2001). Because toxicologic effects of Cd and endosulfan occur in distinct and localized brain regions seems to support strongly a behavior-linked relationship of these neurotoxins, as demonstrated by Cd being preferentially directed toward the motor-controlling cerebral regions and endosulfan being involved predominantly on endocrine-dependent activities of hypothalamic areas. The effects of Cd exposure on pretectal and tegmental areas are characterized primarily by a down-regulatory activity of H₂R-expressing neurons, whereas a similar activity of H₁R-expressing neurons was detected for TLo and CP. Moreover, on the basis of the low levels of H₂R occurring not only in key motor telencephalic areas but also in mesencephalic and cerebellar regions of the mormyrid electric fish (Han et al. 2000) and of other vertebrates (Minami et al. 2001), it appears that a down-regulation of this subtype might represent an important condition of the histaminergic inhibitory effects on locomotor behaviors (Santos et al. 2003). The inhibitory effects may be accomplished by the regulation of parameters such as swimming velocity, location of objects, and overall vestibular activities




Figure 7. The effects of both sublethal and MAT concentrations of Cd on H₂R (A), H₁R (B), and H₃R (C) with respect to their controls ($n = 6$) were expressed as a percentage binding level \pm SE in the different brain regions of the *Thalassoma pavo*, as described in "Materials and Methods." The levels were compared using one-way ANOVA followed where necessary by a post hoc Neuman-Keuls multiple-range test.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Abbreviations: CP, central posterior thalamic nucleus; E, entopeduncular nucleus; ECL, external cellular layer of olfactory bulb; NGa, anterior part of the nucleus glomerulosus; NH, habenular nucleus; NLTm, medial part of lateral tubular nucleus; NPO, preoptic nucleus; NSC, suprachiasmatic nucleus; NSV, nucleus of the saccus vasculosus; SGC, stratum griseum central; TLo, torus longitudinalis; VM, ventromedial thalamic nucleus.




Figure 8. The effects of both sublethal and MAT concentrations of endosulfan on H₃R (A), H₁R (B), and H₂R (C) with respect to their controls ($n = 6$) were expressed as a percentage binding level \pm SE in the different brain regions of the *Thalassoma pavo*, as described in "Materials and Methods." The levels were compared using one-way ANOVA followed where necessary by a post hoc Neuman-Keuls multiple-range test.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. Abbreviations: CP, central posterior thalamic nucleus; Dm2, medial part of the dorsal telencephalon, subdivision 2; NH, habenular nucleus; NLTm, medial part of lateral tubular nucleus; NPO, preoptic nucleus; NRP, nucleus of the posterior hypothalamic recess; NSC, suprachiasmatic nucleus; NSV, nucleus of the saccus vasculosus; SGC, stratum griseum central; TLo, torus longitudinalis; VM, ventromedial thalamic nucleus.

that are controlled by these same brain regions (Meek 1990; Xue et al. 2003).

Conversely, endosulfan appeared predominantly to promote enhanced levels of H₃R-expressing neurons mainly in hypothalamic areas such as NLTm and NRP as well as of H₁R-expressing neurons in NSC. This relationship appears to be strengthened by decreased swimming and feeding behaviors obtained immediately (after 2 hr) in the Chinook salmon when treated with the organophosphate pesticide diazinon (Scholz et al. 2000). The finding that the diencephalic region is a major target of pesticide toxic effects should not be surprising because polychlorinated biphenyls interfere with other hypothalamic activities, including the regulation of body temperature and the activities of the hypothalamic–pituitary–gonadal circuits, with severe consequences on reproductive and hormone-releasing activities (Bloomquist 2003; Cooper et al. 2000). It is noteworthy that high levels of H₃R-expressing neurons have been correlated with a reduction of food intake through the suppression of appetite and energy expenditure in the same hypothalamic areas (Takahashi et al. 2002). In addition, the high levels of H₁R-expressing neurons in other hypothalamic sites of the ornate wrasse plus the inhibition of these subtypes accounting for improved feeding habits in the goldfish (Spieler et al. 1999) appear to be consistent with an important inhibitory role of H₁R and H₃R, at least in hypothalamic nuclei of this teleost.

In conclusion, these results provide direct evidence that the toxicologic risks of endosulfan and Cd on the motor and feeding behavior of *Thalassoma pavo*, as shown by evident morphologic neuronal damages and distinct H_nR-expressing patterns, appear to be very strongly correlated with histaminergic neurosignaling mechanisms. Although most research to date has mostly considered the physiologic risks of the environmental toxicants, here we show that the abnormal behaviors could be linked to specific HA subtype interactions operating in some cerebral regions, at least in the ornate wrasse. Consequently, the motor activities appear to be tightly linked to Cd via variations of mainly H₂R-expressing neurons in the mesencephalic and telencephalic regions, whereas modified feeding behaviors induced by endosulfan seem to be related to the differences of H₁R- and H₃R-expressing neurons mainly in hypothalamic areas. We are still at the beginning of this research, but molecular neuronal interests directed toward the role of environmental disruptors on aquatic organisms could provide further insights regarding not only the behavioral hazards of these contaminants but also neurotoxic mechanisms operating during the entire development cycle of fish, with the intent of minimizing ecologic and commercial risks of this very important class of vertebrates.

REFERENCES

- Airaksinen MS, Panula P. 1990. Comparative neuroanatomy of the histaminergic system in the brain of the frog *Xenopus laevis*. *J Comp Neurol* 292:412–423.
- Bass AH, Grober MS. 2001. Social and neural modulation of sexual plasticity in teleost fish. *Brain Behav Evol* 57:293–300.
- Beauvais SL, Jones SB, Parris JT, Brewer SK, Little EE. 2001. Cholinergic and behavioral neurotoxicity of carbaryl and cadmium to larval rainbow trout (*Oncorhynchus mykiss*). *Ecotoxicol Environ Saf* 49:84–90.
- Bellas J, Vasquez E, Beiras R. 2001. Toxicity of Hg, Cu and Cr on early developmental stages of *Ciona intestinalis* (Chordata, Ascidiacea) with potential application marine water quality assessment. *Water Res* 35(12):2905–12.
- Bisson M, Hontela A. 2002. Cytotoxic and endocrine-disrupting potential of atrazine, diazinon, endosulfan, and mancozeb in adrenocortical steroidogenic cells of rainbow trout exposed *in vitro*. *Toxicol Appl Pharmacol* 180:110–117.
- Bloomquist JR. 2003. Chloride channels as tools for developing selective insecticides. *Arch Insect Biochem Physiol* 54:145–156.
- Bretaud S, Toutant JP, Saglio P. 2000. Effects of carbofuran, diuron and nicosulfuron on acetylcholinesterase activity in goldfish (*Carassius auratus*). *Ecotoxicol Environ Saf* 47:117–124.
- Brodin L, Hökfelt T, Grillner S, Panula P. 1990. Distribution of histaminergic neurons in the brain of the lamprey *Lampetra fluviatilis* as revealed by histamine-immunoreactivity. *J Comp Neurol* 292:435–442.
- Brown RE, Stevens RS, Haas HL. 2001. The physiology of brain histamine. *Prog Neurobiol* 63:637–642.
- Canonaco M, Tavolara R, Facciolo RM. 1997. Dimorphic distribution of the two main GABA_A binding sites in cortical and limbic areas of a rodent living in natural environmental conditions. *J Comp Neurol* 380:423–434.
- Carlson RW, Bradbury SP, Drummond RA, Hammermeister DE. 1998. Neurological effects on startle responses and escape from predation by medaka exposed to organic chemicals. *Aquat Toxicol* 43:51–68.
- Cerdá-Reverter JM, Zanuy S, Muñoz-Cueto JA. 2001a. Cytoarchitectonic study of the brain of a perciform species, the sea bass *Dicentrarchus labrax*. I. The telencephalon. *J Morphol* 247:217–228.
- Cerdá-Reverter JM, Zanuy S, Muñoz-Cueto JA. 2001b. Cytoarchitectonic study of the brain of a perciform species, the sea bass *Dicentrarchus labrax*. II: The diencephalon. *J Morphol* 247:229–251.
- Choich JA, El-Nabawi A, Silbergeld EK. 2004. Evidence of histamine receptors in fish brain using an *in vivo* [¹⁴C]-2-deoxyglucose autoradiographic method and an *in vitro* receptor-binding autoradiographic method. *Environ Res* 94:86–93.
- Clark JM. 1997. Insecticides as a tool in probing vital receptors and enzymes in excitable membranes. *Pest Biochem Physiol* 57:235–254.
- Cooper RL, Stoker TE, Tyrey L, Goldman JM, McElroy WK. 2000. Atrazine disrupts the hypothalamic control of pituitary-ovarian function. *Toxicol Sci* 53:297–307.
- de Olmos JS, Beltramino CA, De Lorenzo SDO. 1994. Use of amino-cupric silver technique for the detection of early and semiacute neuronal degeneration caused by neurotoxins, hypoxia, and physical trauma. *Neurotoxicol Teratol* 16(6):545–561.
- European Communities Council Directive. 1986. Guide for use and care of laboratory animals. *Off J Eur Commun* L358:1–29.
- Ferretti C, Blengio M, Ghi P, Adage T, Portaleone P, Ricci Gamalero S. 1998. Hypothalamic histamine release in normal and stressed rats is affected by sex and aging. *Pharmacol Biochem Behav* 59:225–260.
- Giusi G, Madeo M, Alò R, Carelli A, Facciolo RM, Canonaco M, et al. 2004. Effects of environmental stressors on neuronal histaminergic and behavioral activities of the teleost *Thalassoma pavo*. *Horm Behav* 46:106.
- Gormley KL, Teather KL. 2003. Developmental, behavioural, and reproductive effects experience by Japanese medaka (*Oryzias latipes*) in response to short-term exposure to endosulfan. *Ecotoxicol Environ Saf* 54:330–338.
- Griffiths SW, Armstrong JD. 2000. Differential responses of kin and nonkin salmon to patterns of water flow: does recirculation influence aggression? *Anim Behav* 59:1019–1023.
- Han VZ, Grant K, Bell CC. 2000. Rapid activation of GABAergic interneurons and possible calcium independent release in the mormyrid electrosensory lobe. *J Neurophysiol* 83(3):1592–1604.
- Harvey PW, Rush KC, Cockburn A. 1999. Endocrine and hormonal toxicology: an integrated mechanistic and target systems approach. In: *Endocrine and Hormonal Toxicology* (Harvey PW, Rush KC, Cockburn A, eds). Chichester, UK:Wiley, 1–11.
- Hill SJ, Ganellin CR, Timmermann H, Schwartz JC, Shankley NP, Young JM, et al. 1997. International union of pharmacology. XIII: Classification of histamine receptors. *Pharmacol Rev* 49:253–278.
- Huang YS, Wang LH, Tsai CL. 2004. Photoperiod affects the development of central neurotransmitter systems of tilapia, *Oreochromis mossambicus*. *Neurosci Lett* 355:201–204.
- Inagaki N, Panula P, Yamatodani A, Wada H. 1991. Organization of the histaminergic system in the brain of the teleost *Trachurus trachurus*. *J Comp Neurol* 310:94–102.
- Ito C. 2000. The role of brain histamine in acute and chronic stresses. *Biomol Pharmacol* 54:263–267.
- James SM, Little EE, Semlitsch RD. 2004. The effect of soil composition and hydration on the bioavailability and toxicity of cadmium to hibernating juvenile American toads. *Environ Pollut* 132:523–532.
- Johnson WW, Finley MT. 1980. *Handbook of Acute Toxicity of Chemicals to Fish and Aquatic Invertebrates*. Resource 137. Washington, DC:U.S. Department of Interior, Fish and Wildlife Service.
- Kaslin J, Panula P. 2001. Comparative anatomy of the histaminergic and other aminergic system in zebrafish (*Danio rerio*). *J Comp Neurol* 440:342–377.
- Lafuente A, Fenández-Rey E, Seara R, Pérez-Lorenzo M, Esquifino AI. 2001. Alternate cadmium exposure differentially affects amino acid metabolism within the hypothalamus, median eminence, striatum and prefrontal cortex of male rats. *Neurochem Internat* 39:187–192.
- Larson ET, Norris DO, Summers CH. 2003. Monoaminergic changes associated with socially induced sex reversal in the saddleback wrasse. *Neuroscience* 119:251–263.
- Liao VH, Freedman JH. 1998. Cadmium-regulated genes from the nematode *Caenorhabditis elegans*. Identification and cloning of new cadmium-responsive genes by differential display. *J Biol Chem* 273:31962–31970.
- Lin JS, Hou Y, Sakai K, Jouvet M. 1996. Histaminergic descending inputs to the mesopontine tegmentum and their role in the control of cortical activation and wakefulness in the cat. *J Neurosci* 16:1523–1537.
- Martinez MP, Angulo R, Pozo R, Jodral M. 1997. Organochlorine pesticides in pasteurized milk and associated health risks. *Food Chem Toxicol* 35:621–624.
- Meek J. 1990. Tectal morphology: connections, neurones and synapses. In: *The Visual System of Fish* (Douglas R, Djamgoz M, eds). London:Chapman and Hall, 239–277.
- Mendez-Armenta M, Barroso-Moguel R, Villeda-Hernandez J, Nava-Ruiz C, Rios C. 2001. Histopathological alterations in the brain regions of rats after prenatal combined treatment with cadmium and dexamethasone. *Toxicology* 161(3):189–199.
- Miller LL, Lladós R. 1999. *Toxicological Profile for Endosulfan*. Draft Report 4.3. Atlanta, GA:U.S. Agency for Toxic Substance and Disease Registry, 162–163, 169.
- Minami A, Takeda A, Nishibaba D, Takefuta S, Oku N. 2001. Cadmium toxicity in synaptic neurotransmission in the brain. *Brain Res* 894:336–339.
- Munson PJ, Rodbard D. 1980. LIGAND: a versatile computerize approach for characterization of ligand-binding systems. *Anal Biochem* 107:220–239.
- Naqvi SM, Vaishnavi C. 1993. Bioaccumulation potential and toxicity of endosulfan insecticide to non-target animals. *Comp Biochem Physiol C* 105:347–361.
- Pan G, Dutta HM. 1998. The inhibition of brain acetylcholinesterase activity of juvenile largemouth bass *Micropterus salmoides* by sublethal concentrations of diazinon. *Environ Res A* 79:133–137.
- Peitsaro N, Anichtchik OV, Panula P. 2000. Identification of a histamine H₂-like receptor in the zebrafish (*Danio rerio*) brain. *J Neurochem* 75:718–724.
- Pillai A, Priya L, Gupta S. 2003. Effects of combined exposure to lead and cadmium on the hypothalamic-pituitary axis function in proestrous rats. *Food Chem Toxicol* 41:379–384.
- Pillot C, Heron A, Cochois V, Tardivel-Lacombe J, Ligneau X, Schwartz JC, et al. 2002. A detailed mapping of the histamine H₃ receptor and its gene transcript in rat brain. *Neuroscience* 114(1): 173–193.
- Portavella M, Torres B, Salas C. 2004. Avoidance response in goldfish: emotional and temporal involvement of medial and lateral telencephalic pallium. *J Neurosci* 24(9):2335–2342.

- Ram RN, Singh IJ, Singh DV. 2001. Carbofuran induced impairment in the hypothalamo-neurohypophyseal-gonadal complex in the teleost, *Channa punctatus* (Bloch). *J Environ Biol* 22(3):193–200.
- Risso-de Faverney C, Devaux A, Lafaurie M, Girard JP, Bailly B, Rahmani R. 2001. Cadmium induced apoptosis and genotoxicity in rainbow trout hepatocytes through generation of reactive oxygen species. *Aquat Toxicol* 53:65–76.
- Roex EWM, Giovannangelo M, van Gestel CAM. 2001. Reproductive impairment in the zebrafish *Danio rerio*, upon chronic exposure to 1,2,3-trichlorobenzene. *Ecotoxicol Environ Saf* 48:196–201.
- Ryu JH, Yanai K, Zhao XL, Watanabe T. 1996. The effect of dopamine D₁ receptor stimulation on the up-regulation of histamine H₂-receptors following destruction of the ascending dopaminergic neurones. *Br J Pharmacol* 118:585–592.
- Santos NR, Huston JP, Brandão ML. 2003. Blockade of histamine H₂ receptors of the periaqueductal gray and inferior colliculus induces fear-like behaviors. *Pharmacol Biochem Behav* 75:25–33.
- Scherer E, McNicol RE, Evans RE. 1997. Impairment of lake trout foraging by chronic exposure to cadmium: a black-box experiment. *Aquat Toxicol* 37:1–7.
- Scholz NL, Truelove NK, French BL, Berejikian BA, Quinn TP, Casillas E, et al. 2000. Diazinon disrupt antipredator and homing behaviours in Chinook salmon (*Oncorhynchus tshawytscha*). *Can J Fish Aquat Sci* 57:1911–1918.
- Scott GR, Sloman KA, Rouleau C, Wood CM. 2003. Cadmium disrupts behavioural and physiological responses to alarm substance in juvenile rainbow trout (*Oncorhynchus mykiss*). *J Exp Biol* 206:1779–1790.
- Siegel JM, Nienhuis R, Gulyani S, Ouyang S, Wu MF, Mignot E, et al. 1999. Neuronal degeneration in canine narcolepsy. *J Neurosci* 19(1):248–257.
- Spieler RE, Nelson CA, Huston JP, Mattioli R. 1999. Post-trial administration of H₁ histamine receptor blocker improves appetitive reversal learning and memory in goldfish, *Carassius auratus*. *Neurosci Lett* 277:5–8.
- Takahashi K, Suwa H, Ishikawa T, Kotani H. 2002. Targeted disruption of H₃ receptors results in changes in brain histamine tone leading to an obese phenotype. *J Clin Invest* 110(12):1791–1800.
- Talkvist J, Persson E, Henriksson J, Tiälve H. 2002. Cadmium-metallothionein interactions in the olfactory pathways of rats and pikes. *Toxicol Sci* 67:108–113.
- Thophon S, Kruatrachue M, Upatham ES, Pokethitiyook S, Jaritkhuan S. 2003. Histopathological alterations of white seabass *Lates calcarifer*, in acute and subchronic cadmium exposure. *Environ Pollut* 121:307–320.
- Waalkes MP, Coogan TP, Barter RA. 1992. Toxicological principles of metal carcinogenesis with special emphasis on cadmium. *Crit Rev Toxicol* 22:175–201.
- Wood CM. 2001. Toxic responses of the gill. In: *Target Organ Toxicity in Marine and Freshwater Teleosts* (Schlenk D, Benson WH, eds). London: Taylor & Francis, 1–87.
- Xue HG, Yamamoto N, Yang CY, Kerem G, Yoshimoto M, Imura K, et al. 2003. Fiber connections of the torus longitudinalis and optic tectum in holocentrid teleosts. *J Comp Neurol* 462: 194–212.
- Yoshida S. 2001. Re-evaluation of acute neurotoxic effects of Cd²⁺ on mesencephalic trigeminal neurons of the adult rat. *Brain Res* 892(1):102–110.
- Zander CD, Sötje I. 2002. Seasonal and geographical differences in cleaner fish activity in the Mediterranean Sea. *Helgol Mar Res* 55:232–241.

Southeast Asia that will allow natural solar ultraviolet-B radiation (UV-B) to suppress the virus before infection occurs.

Influenza viruses and various non-pigmented bacteria are killed by UV-B wavelengths in sunlight (Hollaender and Oliphant 1944). Biomass smoke significantly suppresses natural levels of UV-B, and severe smoke pollution reduced UV-B by up to 95% during the burning seasons in Brazil in 1995 (Mims 1996) and 1997 (Mims FM III, White B, unpublished data). Reduced UV-B on 6 days in August 1997 was well correlated ($r^2 = 0.83$) with an increase in the ratio of nonpigmented bacteria vulnerable to UV-B to pigmented bacteria that are protected from UV-B (Mims and White 1998). Although airborne influenza viruses were not measured, 1997 hospital admission records at Alta Floresta, Brazil, showed that influenza incidence was highest during the burning season (de Castro GC, personal communication).

Human cases of avian influenza in Thailand and Vietnam peaked during the winter burning seasons of 2003 and 2004 (Thailand Ministry of Public Health 2005). Assuming similar optical properties of biomass smoke in Southeast Asia and Brazil, where UV-B and optical depth are highly correlated, optical depth measurements over Thailand and Vietnam by NASA's Terra and Aqua satellites suggest highly suppressed UV-B during these avian influenza outbreaks (Mims FM III, unpublished data).

Human cases of avian influenza in Thailand and Vietnam since December 2003

have peaked during both the rainy season and the burning season. Thus, periods of prolonged cloudiness and severe smoke pollution could play a role in initiating avian and other influenza outbreaks by attenuating the solar UV-B that might otherwise suppress influenza viruses in outdoor air exposed to sunlight. The transmission of avian influenza to people during these periods is enhanced by the fact that poultry raised for human consumption are often kept within several meters of where people live (World Health Organization 2004).

The author declares he has no competing financial interests.

Forrest M. Mims III
Geronimo Creek Observatory
Seguin, Texas
E-mail: forrest.mims@ieec.org

REFERENCES

- Hollaender A, Oliphant J. 1944. The inactivating effect of monochromatic ultraviolet radiation on influenza virus. *J Bacteriol* 48:447–454.
- Longini IM Jr, Nizam A, Xu S, Ungchusak K, Hanshaoworakul W, Cummings DAT, et al. 2005. Containing pandemic influenza at the source. *Science* 309:1083–1087.
- Mims FM III. 1996. Significant reduction in UV-B caused by smoke from biomass burning in Brazil. *Photochem Photobiol* 64:123–125.
- Thailand Ministry of Public Health. 2005. Avian Influenza Surveillance in Human as of July 1, 2005. Available: http://thaigcd.ddc.moph.go.th/AI_case_report_050701.html [accessed 25 August 2005].
- Washam C. 2005. On hens and needles. *Environ Health Perspect* 113:A370.
- World Health Organization. 2004. Avian Influenza A(H5) in Rural Areas in Asia: Food Safety Considerations. Available: <http://www.who.int/foodsafety/micro/avian2/en/> [accessed 25 August 2005].

eHP Published by the National Institute of Environmental Health Sciences

There's Another Option Before Your Very Eyes


Post your career opportunity ad where the most qualified applicants will see it—right here!

Call: 1-919-337-2028
E-mail: ehpads@niehs.nih.gov


ehponline.org Environmental Health PERSPECTIVES

ERRATA

There was an error in Figure 2 of Zeman et al. [*Environ Health Perspect* 110:817–822 (2002)]: the y -axis should have been labeled “Nitrite” instead of “Nitrate.” The corrected figure appears below.



In Giusi et al. [*Environ Health Perspect* 113:1522–1529 (2005)], the colors were incorrect in the key to Figure 1. The corrected figure appears below.



EHP regrets the errors.