

NTP REPORT
ON THE
TOXICITY STUDIES OF ACESULFAME POTASSIUM
(CAS NO. 55589-62-3)

IN FVB/N-TgN(v-*Ha-ras*)Led (Tg.AC)
HEMIZYGOUS MICE

AND CARCINOGENICITY STUDIES OF
ACESULFAME POTASSIUM

IN B6.129-*Trp53*^{tm1Brd} (N5)
HAPLOINSUFFICIENT MICE

(FEED STUDIES)

NATIONAL TOXICOLOGY PROGRAM
P.O. Box 12233
Research Triangle Park, NC 27709

October 2005

NTP GMM 2

NIH Publication No. 06-4460

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
National Institutes of Health

FOREWORD

The National Toxicology Program (NTP) is made up of four charter agencies of the U.S. Department of Health and Human Services (DHHS): the National Cancer Institute (NCI), National Institutes of Health; the National Institute of Environmental Health Sciences (NIEHS), National Institutes of Health; the National Center for Toxicological Research (NCTR), Food and Drug Administration; and the National Institute for Occupational Safety and Health (NIOSH), Centers for Disease Control and Prevention. In July 1981, the Carcinogenesis Bioassay Testing Program, NCI, was transferred to the NIEHS. The NTP coordinates the relevant programs, staff, and resources from these Public Health Service agencies relating to basic and applied research and to biological assay development and validation.

The NTP develops, evaluates, and disseminates scientific information about potentially toxic and hazardous chemicals. This knowledge is used for protecting the health of the American people and for the primary prevention of disease.

The studies described in this Report were performed under the direction of the NIEHS and were conducted in compliance with NTP laboratory health and safety requirements and must meet or exceed all applicable federal, state, and local health and safety regulations. Animal care and use were in accordance with the Public Health Service Policy on Humane Care and Use of Animals. The prechronic and chronic studies were conducted in compliance with Food and Drug Administration (FDA) Good Laboratory Practice Regulations, and all aspects of the chronic studies were subjected to retrospective quality assurance audits before being presented for public review.

The studies described in this Report series were designed and conducted to characterize the toxicologic potential, including carcinogenic activity, of selected agents in laboratory animals that have been genetically modified. These genetic modifications may involve inactivation of selected tumor suppressor functions or activation of oncogenes that are commonly observed in human cancers. This may result in a rapid onset of cancer in the genetically modified animal when exposure is to agents that act directly or indirectly on the affected pathway. An absence of a carcinogenic response may reflect either an absence of carcinogenic potential of the agent or that the selected model does not harbor the appropriate genetic modification to reduce tumor latency and allow detection of carcinogenic activity under the conditions of these subchronic studies. Chemicals selected for NTP toxicology and carcinogenesis studies are chosen primarily on the bases of human exposure, level of production, and chemical structure. The interpretive conclusions presented in this Report are based only on the results of these NTP studies. Extrapolation of these results to other species and quantitative risk analyses for humans require wider analyses beyond the purview of these studies. Selection per se is not an indicator of a chemical's carcinogenic potential.

Details about ongoing and completed NTP studies, abstracts of all NTP Reports, and full versions of the completed reports are available at the NTP's World Wide Web site: <http://ntp.niehs.nih.gov>. In addition, printed copies of these reports are available from NTP as supplies last by contacting (919) 541-3419.

NTP REPORT
ON THE
TOXICITY STUDIES OF ACESULFAME POTASSIUM
(CAS NO. 55589-62-3)

IN FVB/N-TgN(v-*Ha-ras*)Led (Tg.AC)
HEMIZYGOUS MICE

AND CARCINOGENICITY STUDIES OF
ACESULFAME POTASSIUM

IN B6.129-*Trp53*^{tm1Brd} (N5)
HAPLOINSUFFICIENT MICE

(FEED STUDIES)



NATIONAL TOXICOLOGY PROGRAM
P.O. Box 12233
Research Triangle Park, NC 27709

October 2005

NTP GMM 2

NIH Publication No. 06-4460

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
National Institutes of Health

CONTRIBUTORS

National Toxicology Program

Evaluated and interpreted results and reported findings

R.D. Irwin, Ph.D., Study Scientist
 D.E. Malarkey, D.V.M., Ph.D., Study Pathologist
 D.W. Bristol, Ph.D.
 J.R. Bucher, Ph.D.
 J.E. French, Ph.D.
 J.R. Hailey, D.V.M.
 J.K. Haseman, Ph.D.
 R.A. Herbert, D.V.M., Ph.D.
 R.R. Maronpot, D.V.M.
 J.C. Peckham, D.V.M., M.S., Ph.D.
 J.H. Roycroft, Ph.D.
 C.S. Smith, Ph.D.
 G.S. Travlos, D.V.M.
 M.K. Vallant, B.S., M.T.
 K.L. Witt, M.S.

BioReliance Corporation

Conducted studies and evaluated pathology findings

M.L. Wenk, Ph.D., Principal Investigator
 L.L. Lanning, D.V.M.

Experimental Pathology Laboratories, Inc.

Provided pathology review

J.F. Hardisty, D.V.M., Principal Investigator
 G. Willson, B.V.M. & S.

Dynamac Corporation

Prepared quality assurance audits

S. Brecher, Ph.D., Principal Investigator

NTP Pathology Working Group

*Evaluated slides and prepared pathology report
 (December 18, 2001)*

C.A. Cummings, D.V.M., Ph.D., Chairperson
 Pathology Associates, A Charles River Company
 Laboratories, Inc.
 J. Mahler, D.V.M.
 National Toxicology Program

Constella Group, Inc.

Provided statistical analyses

P.W. Crockett, Ph.D., Principal Investigator
 L.J. Betz, M.S.
 K.P. McGowan, M.B.A.
 J.T. Scott, M.S.

Biotechnical Services, Inc.

Prepared Report

S.R. Gunnels, M.A., Principal Investigator
 B.F. Hall, M.S.
 L.M. Harper, B.S.
 D.C. Serbus, Ph.D.
 R.A. Willis, B.A., B.S.

CONTENTS

| | |
|--|------------|
| ABSTRACT | 5 |
| EXPLANATION OF LEVELS OF EVIDENCE OF CARCINOGENIC ACTIVITY | 8 |
| TECHNICAL REPORTS REVIEW SUBCOMMITTEE | 9 |
| SUMMARY OF TECHNICAL REPORTS REVIEW SUBCOMMITTEE COMMENTS | 10 |
| INTRODUCTION | 11 |
| MATERIALS AND METHODS | 17 |
| RESULTS | 23 |
| DISCUSSION AND CONCLUSIONS | 37 |
| REFERENCES | 41 |
| APPENDIX A Summary of Lesions in Male Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 45 |
| APPENDIX B Summary of Lesions in Female Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 59 |
| APPENDIX C Summary of Lesions in Male p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 73 |
| APPENDIX D Summary of Lesions in Female p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 85 |
| APPENDIX E Genetic Toxicology | 97 |
| APPENDIX F Chemical Characterization and Dose Formulation Studies | 101 |
| APPENDIX G Feed and Compound Consumption in the 9-Month Feed Studies of Acesulfame Potassium | 109 |

SUMMARY

Background

Acesulfame potassium is an artificial sweetener widely used in beverages and foods. We tested if acesulfame potassium could cause cancer in two different strains of genetically modified mice.

Methods

We fed groups of male and female Tg.AC mice and male and female p53 mice diets containing up to 3% acesulfame potassium for 9 months. Animals given feed with no sweetener added served as the control groups. Tissues from 15 sites were examined for every animal.


Results

Exposure to acesulfame potassium had no effect on the survival of any of the animal groups. No increases in tumors were seen in males or females from either strain of mice.

Conclusions

We conclude that acesulfame potassium did not cause cancer in the genetically modified mice used in these studies.

ABSTRACT



ACESULFAME POTASSIUM

CAS No. 55589-62-3

Chemical Formula: $C_4H_4KNO_4S$ Molecular Weight: 201.25

Synonyms: ASK; HOE-095K; 6-methyl-1,2,3-oxathiazine-4(3-H)-one-2,2-dioxide potassium salt

Trade names: Sunette, Sweet One

Acesulfame potassium is an artificial sweetener used throughout the world in food and beverages. Acesulfame potassium was nominated by The Center for Science in the Public Interest because of its widespread use. Male and female Tg.AC hemizygous and p53 haploinsufficient mice were exposed to acesulfame potassium (at least 99% pure) in feed for 9 months. Genetic toxicology studies were conducted in mouse peripheral blood erythrocytes.

9-MONTH STUDY IN Tg.AC HEMIZYGOUS MICE

Groups of 15 male and 15 female Tg.AC hemizygous mice were fed diets containing 0%, 0.3%, 1%, or 3% acesulfame potassium (equivalent to average daily doses of approximately 420, 1,400, or 4,500 mg acesulfame potassium/kg body weight to males and 520, 1,700, or 5,400 mg/kg to females) for 40 weeks. Exposure to acesulfame potassium had no effect on survival or mean

body weights. Feed consumption by the exposed groups was similar to that by the control groups throughout the study. There were no neoplasms or nonneoplastic lesions that were attributed to exposure to acesulfame potassium.

9-MONTH STUDY IN p53 HAPLOINSUFFICIENT MICE

Groups of 15 male and 15 female p53 haploinsufficient mice were fed diets containing 0%, 0.3%, 1%, or 3% acesulfame potassium (equivalent to average daily doses of approximately 475, 1,500, or 4,700 mg/kg to males and 570, 1,800, or 5,700 mg/kg to females) for 40 weeks. Exposure to acesulfame potassium had no effect on survival or mean body weights. Feed consumption by the exposed groups was similar to that by the control groups throughout the study. There were no neoplasms or nonneoplastic lesions that were attributed to exposure to acesulfame potassium.

GENETIC TOXICOLOGY

Acesulfame potassium did not increase the frequency of micronucleated erythrocytes in peripheral blood of male or female Tg.AC hemizygous mice administered 0.3% to 3% in dosed feed. A similar study was conducted in p53 haploinsufficient mice, and a significant exposure concentration-related increase in the frequency of micronucleated erythrocytes was noted in males but not females.

CONCLUSIONS

Under the conditions of this 9-month feed study, there was *no evidence of carcinogenic activity** of acesulfame potassium in male or female p53 haploinsufficient mice exposed to 0.3%, 1%, or 3%.

* Explanation of Levels of Evidence of Carcinogenic Activity is on page 8. A summary of the Technical Report Review Subcommittee comments and public discussion on this technical report appears on page 9.

Summary of the 9-Month Carcinogenesis and Genetic Toxicology Studies of Acesulfame Potassium

| | Male p53 Haploinsufficient Mice | Female p53 Haploinsufficient Mice |
|---|---|--|
| Concentrations in feed | 0%, 0.3%, 1%, or 3% (0, 3,000, 10,000, or 30,000 ppm) | 0%, 0.3%, 1%, or 3% |
| Body weights | Exposed groups similar to the control group | Exposed groups similar to the control group |
| Survival rates | 14/15, 15/15, 15/15, 14/15 | 14/15, 14/15, 14/15, 14/15 |
| Nonneoplastic effects | None | None |
| Neoplastic effects | None | None |
| Level of evidence of carcinogenic activity | No evidence | No evidence |
| Genetic toxicology | | |
| Micronucleated erythrocytes | | |
| Mouse peripheral blood <i>in vivo</i> : | Positive in p53 haploinsufficient males; negative in Tg.AC hemizygous males and females and p53 haploinsufficient females | |

EXPLANATION OF LEVELS OF EVIDENCE OF CARCINOGENIC ACTIVITY

The National Toxicology Program describes the results of individual experiments on a chemical agent and notes the strength of the evidence for conclusions regarding each study. Negative results, in which the study animals do not have a greater incidence of neoplasia than control animals, do not necessarily mean that a chemical is not a carcinogen, inasmuch as the experiments are conducted under a limited set of conditions. Positive results demonstrate that a chemical is carcinogenic for laboratory animals under the conditions of the study and indicate that exposure to the chemical has the potential for hazard to humans. Other organizations, such as the International Agency for Research on Cancer, assign a strength of evidence for conclusions based on an examination of all available evidence, including animal studies such as those conducted by the NTP, epidemiologic studies, and estimates of exposure. Thus, the actual determination of risk to humans from chemicals found to be carcinogenic in laboratory animals requires a wider analysis that extends beyond the purview of these studies.

Five categories of evidence of carcinogenic activity are used in the Technical Report series to summarize the strength of the evidence observed in each experiment: two categories for positive results (**clear evidence and some evidence**); one category for uncertain findings (**equivocal evidence**); one category for no observable effects (**no evidence**); and one category for experiments that cannot be evaluated because of major flaws (**inadequate study**). These categories of interpretative conclusions were first adopted in June 1983 and then revised in March 1986 for use in the Technical Report series to incorporate more specifically the concept of actual weight of evidence of carcinogenic activity. For each separate experiment (male rats, female rats, male mice, female mice), one of the following five categories is selected to describe the findings. These categories refer to the strength of the experimental evidence and not to potency or mechanism.

- **Clear evidence** of carcinogenic activity is demonstrated by studies that are interpreted as showing a dose-related (i) increase of malignant neoplasms, (ii) increase of a combination of malignant and benign neoplasms, or (iii) marked increase of benign neoplasms if there is an indication from this or other studies of the ability of such tumors to progress to malignancy.
- **Some evidence** of carcinogenic activity is demonstrated by studies that are interpreted as showing a chemical-related increased incidence of neoplasms (malignant, benign, or combined) in which the strength of the response is less than that required for clear evidence.
- **Equivocal evidence** of carcinogenic activity is demonstrated by studies that are interpreted as showing a marginal increase of neoplasms that may be chemical related.
- **No evidence** of carcinogenic activity is demonstrated by studies that are interpreted as showing no chemical-related increases in malignant or benign neoplasms.
- **Inadequate study** of carcinogenic activity is demonstrated by studies that, because of major qualitative or quantitative limitations, cannot be interpreted as valid for showing either the presence or absence of carcinogenic activity.

For studies showing multiple chemical-related neoplastic effects that if considered individually would be assigned to different levels of evidence categories, the following convention has been adopted to convey completely the study results. In a study with clear evidence of carcinogenic activity at some tissue sites, other responses that alone might be deemed some evidence are indicated as “were also related” to chemical exposure. In studies with clear or some evidence of carcinogenic activity, other responses that alone might be termed equivocal evidence are indicated as “may have been” related to chemical exposure.

When a conclusion statement for a particular experiment is selected, consideration must be given to key factors that would extend the actual boundary of an individual category of evidence. Such consideration should allow for incorporation of scientific experience and current understanding of long-term carcinogenesis studies in laboratory animals, especially for those evaluations that may be on the borderline between two adjacent levels. These considerations should include:

- adequacy of the experimental design and conduct;
- occurrence of common versus uncommon neoplasia;
- progression (or lack thereof) from benign to malignant neoplasia as well as from preneoplastic to neoplastic lesions;
- some benign neoplasms have the capacity to regress but others (of the same morphologic type) progress. At present, it is impossible to identify the difference. Therefore, where progression is known to be a possibility, the most prudent course is to assume that benign neoplasms of those types have the potential to become malignant;
- combining benign and malignant tumor incidence known or thought to represent stages of progression in the same organ or tissue;
- latency in tumor induction;
- multiplicity in site-specific neoplasia;
- metastases;
- supporting information from proliferative lesions (hyperplasia) in the same site of neoplasia or in other experiments (same lesion in another sex or species);
- presence or absence of dose relationships;
- statistical significance of the observed tumor increase;
- concurrent control tumor incidence as well as the historical control rate and variability for a specific neoplasm;
- survival-adjusted analyses and false positive or false negative concerns;
- structure-activity correlations; and
- in some cases, genetic toxicology.

**NATIONAL TOXICOLOGY PROGRAM BOARD OF SCIENTIFIC COUNSELORS
TECHNICAL REPORTS REVIEW SUBCOMMITTEE**

The members of the Technical Reports Review Subcommittee who evaluated the draft NTP Technical Report on acesulfame potassium on May 22, 2003, are listed below. Subcommittee members serve as independent scientists, not as representatives of any institution, company, or governmental agency. In this capacity, subcommittee members have five major responsibilities in reviewing the NTP studies:

- to ascertain that all relevant literature data have been adequately cited and interpreted,
- to determine if the design and conditions of the NTP studies were appropriate,
- to ensure that the Technical Report presents the experimental results and conclusions fully and clearly,
- to judge the significance of the experimental results by scientific criteria, and
- to assess the evaluation of the evidence of carcinogenic activity and other observed toxic responses.

Mary Anna Thrall, D.V.M., Chairperson
Department of Pathology
College of Veterinary Medicine and Biomedical Sciences
Colorado State University
Fort Collins, CO

Kim Boekelheide, M.D., Ph.D.
Division of Biology and Medicine
Department of Pathology and Laboratory Medicine
Brown University
Providence, RI

Hillary M. Carpenter, III, Ph.D.
Minnesota Department of Health
St. Paul, MN

Samuel M. Cohen, M.D., Ph.D.*
Department of Pathology and Microbiology
University of Nebraska Medical Center
Omaha, NE

Michael R. Elwell, D.V.M., Ph.D., Principal Reviewer
Pfizer Global Research and Development
Groton, CT

Walter W. Piegorsch, Ph.D., Principal Reviewer
Department of Statistics
University of South Carolina
Columbia, SC

Stephen M. Roberts, Ph.D.
Department of Physiological Sciences
College of Veterinary Medicine
University of Florida
Gainesville, FL

Richard D. Storer, M.P.H., Ph.D., Principal Reviewer
Department of Genetic and Cellular Toxicology
Merck Research Laboratories
West Point, PA

Mary Vore, Ph.D.
Graduate Center for Toxicology
University of Kentucky
Lexington, KY

Cheryl Lyn Walker, Ph.D.
M.D. Anderson Cancer Center
The University of Texas
Smithville, TX

* Did not attend

SUMMARY OF TECHNICAL REPORTS REVIEW SUBCOMMITTEE COMMENTS

On May 22, 2003, the draft Report on the toxicology and carcinogenesis studies of acesulfame potassium received public review by the National Toxicology Program's Board of Scientific Counselor's Technical Reports Review Subcommittee. The review meeting was held at the National Institute of Environmental Health Sciences, Research Triangle Park, NC.

Dr. R. Irwin, NIEHS, introduced the studies of acesulfame potassium in genetically modified mice by describing the uses of the sweetener, its nomination as a presumptive negative control for evaluation of the transgenic mouse models, the study design, and the survival, body weight, and histopathology observations. The proposed conclusion was:

Under the conditions of this 9-month feed study, there was *no evidence of carcinogenic activity* of acesulfame potassium in male or female p53 haploinsufficient mice exposed to 0.3%, 1%, or 3%. Because this is a new model, there is uncertainty whether the study possessed sufficient sensitivity to detect a carcinogenic effect.


Dr. Storer, the first principal reviewer, agreed with the conclusions. He felt it more appropriate to characterize the Tg.AC mouse model as a zeta-globin promoter activation model with an oncogenic H-*ras* reporter phenotype than as a gain of oncogenic function H-*ras* model.

Dr. Elwell, the second principal reviewer, agreed with the conclusions and inquired if the mice might have been able to tolerate a higher dose. Dr. Irwin replied that in another study with CD-1 mice a mild weight gain depression was seen at this top dose.

Dr. Piegorsch, the third principal reviewer, also agreed with the conclusions.

Dr. Storer moved, and Dr. Elwell seconded, that the conclusions be accepted as written. The motion was then approved unanimously with eight votes.

INTRODUCTION



ACESULFAME POTASSIUM

CAS No. 55589-62-3

Chemical Formula: $C_4H_4KNO_4S$ Molecular Weight: 201.25

Synonyms: ASK; HOE-095K; 6-methyl-1,2,3-oxathiazine-4(3-H)-one-2,2-dioxide potassium salt

Trade names: Sunette, Sweet One

CHEMICAL AND PHYSICAL PROPERTIES

Acesulfame potassium is an odorless, colorless, crystalline powder approximately 200 times sweeter than sugar. Acesulfame potassium has a decomposition point of 225° C, is stable over a broad range of pH, and has a density of 1.81 g/cm³ (*Merck Index*, 1996; HSDB, 2003; Parchem, 2003). Acesulfame potassium is also highly soluble in water, has a long shelf life, and is synergistic when used in combination with other sweeteners (Parchem, 2003).

PRODUCTION, USE, AND HUMAN EXPOSURE

Acesulfame potassium is an artificial sweetener used throughout the world. It is currently available as a dry powder for use in food and beverages and is present as a flavoring agent in numerous food items and diet drinks; therefore, there is extensive human exposure.

ABSORPTION, DISTRIBUTION, AND EXCRETION

After oral administration, acesulfame potassium is rapidly and completely absorbed then rapidly eliminated in the urine of rats, dogs, and humans. After oral administration to rats, blood concentrations peaked within 30 minutes then declined with a $t_{1/2}$ of 4.8 hours. Following oral administration to humans (30 mg), blood levels peaked after 1 to 1.5 hours then declined with a $t_{1/2}$ of 2.5 hours.

Acesulfame potassium is rapidly distributed after oral administration and rapidly eliminated with no evidence of bioaccumulation in any tissue. At the time of maximum blood concentration, the highest levels were present in the gastrointestinal tract, bile, kidneys, and urinary bladder. Following oral administration of [¹⁴C]-labeled acesulfame potassium to rats, 97.5% of the administered dose was excreted in the urine as parent compound within 24 hours; in humans, 98.4% was excreted in the

urine. Similar results were obtained for groups of rats dosed for 8 or 60 days prior to a histopathologic examination (Volz *et al.*, 1991).

TOXICITY

Experimental Animals

One published prechronic study of acesulfame potassium was found in the literature (Sinkeldam *et al.*, 1991a). Groups of 10 male and 10 female Wistar rats received acesulfame potassium in the diet at concentrations of 0%, 1%, 3%, or 10% for 13 weeks. Body weights and food and water consumption were measured. Urine samples were collected, and blood was drawn from the tip of the tail for hematology during week 13; blood was also collected at necropsy for determination of serum enzyme activities. At the end of the study, rats were necropsied, and adrenal gland, brain, cecal weights, heart, kidney, liver, ovary, spleen, testis, thymus, and thyroid gland weights were recorded. Tissues were collected, processed, and a histopathologic examination was conducted. There were no biologically significant differences in hematology or clinical chemistry parameters among the groups. There were slight decreases in mean relative spleen (males) and thymus (females) weights in rats that received 1% acesulfame potassium and in relative spleen weights of 3% females, which were small in magnitude and did not appear to be biologically significant. The only change that was biologically significant was the increase in relative cecal weights in males and females in the 10% groups. No exposure-related histologic lesions were observed.

Humans

No information on toxicity studies of acesulfame potassium in humans was found in the literature.

CARCINOGENICITY

Experimental Animals

Groups of 60 male and 60 female Wistar rats received diets that contained 0, 3,000, 10,000, or 30,000 ppm acesulfame potassium for 120 (males) or 123 (females) weeks (Sinkeldam *et al.*, 1991b). Body weights were recorded weeks 1 to 5, 8, 10, and 12, and at 4-week intervals thereafter. Feed consumption was measured at regular intervals throughout the study, and water

consumption was measured during week 62 and 63. At weeks 33, 53, 78, 102, and 119, blood was collected from 10 males and 10 females in each group for hematology. Blood was drawn from 10 rats per group at weeks 34, 52, and 104 for blood urea nitrogen and glucose and from 10 rats per group for serum chemistry at weeks 34, 52, and 109. At weeks 35, 52, 78, and 104, urine was collected from 10 rats per group for urinalysis. Survival of all exposed groups was similar to the control groups. Mean body weights of 3,000 and 10,000 ppm males and females were similar to the controls throughout the study. The mean body weight of 30,000 ppm males was slightly lower than the controls up to week 80, after which there appeared to be no differences between any of the male groups. The mean body weight of 30,000 ppm females was lower than that of the controls, but the difference was no more than 5%. The only potential exposure-related response was the increased incidences of mammary gland tumors in females. The incidences of fibroadenoma were greater in exposed groups of females (43%, 48%, 48%) than that in control group (30%), and the incidences of adenocarcinoma were greater in exposed females (3%, 5%, 5%) than that in controls (2%). The authors concluded that the increased incidences of adenocarcinoma were not exposure-related because mammary gland adenocarcinomas are common in Wistar rats (historical control rate is 5%), and the incidence in the concurrent control group was relatively low compared to historical control data.

Groups of 100 male and 100 female Swiss mice received diets that contained 0, 3,000, 10,000, or 30,000 ppm acesulfame potassium for 80 weeks (Beems *et al.*, 1991). Body weights were recorded at weeks 1, 2, 4, and at 4-week intervals thereafter. Survival of all exposed groups was similar to that of the controls. Mean body weights of 30,000 ppm males and females were slightly lower than those of the other groups, but the differences were marginal. Mean liver weights of all exposed groups of males were less than that of the control group; kidney weights of exposed males and liver and kidney weights of exposed females were similar to the controls. No exposure-related lesions were observed.

Humans

No epidemiology studies of acesulfame potassium in humans were found in the literature.

GENETIC TOXICITY

Two studies measured cytogenetic effects in bone marrow cells of male Swiss albino mice administered acesulfame potassium by gavage. Mukherjee and Chakrabarti (1997) measured induction of chromosomal aberrations in mice treated with 15 to 2,250 mg/kg; significant increases in chromatid breaks were observed, and the increases in chromosomally abnormal cells were related to dose. A followup study by Mukhopadhyay *et al.* (2000) investigated induction of chromosomal aberrations in mice administered a mixture of acesulfame potassium (1.5, 15, and 150 mg/kg) and aspartame (3.5, 35, and 350 mg/kg); doses were chosen based on proposed limits of the two sweeteners together in soft drinks (150 ppm acesulfame potassium and 350 ppm aspartame); no induction of chromosomal aberrations was observed.

Jung *et al.* (1991) conducted a battery of genetic toxicity tests on acesulfame potassium. At doses from 4 to 10,000 µg/plate, acesulfame potassium failed to increase the frequency of revertants in *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, and TA1538 with or without S9. Over the same dose range, no increase in the frequency point mutations leading to tryptophane independence was observed in *Escherichia coli* WP2uvrA. At concentrations of 10 to 1,000 µg/mL culture media, acesulfame potassium did not increase the frequency of azaguanine-resistant colonies of Chinese hamster V79 cells.

Incubation of freshly isolated hepatocytes from F344 rats for 20 hours in culture medium containing 25 to 5,000 µg/mL, acesulfame potassium did not increase incorporation of H3-thymidine into nuclear DNA, a measure of unscheduled DNA synthesis (Jung *et al.*, 1991).

No increase in the frequency of micronuclei was observed in the bone marrow of male or female NMRI mice after an acute gavage exposure to 0, 450, 1,500, or 4,500 mg acesulfame potassium/kg body weight (Jung *et al.*, 1991). There was no increase in the frequency of chromosomal aberrations in the bone marrow of male or female Chinese hamsters that received the same dose concentrations for 5 days.

BACKGROUND

ON GENETICALLY ALTERED MICE

Mutation and/or deletions of tumor suppressor genes or activation of protooncogenes can disrupt cell function and predispose an animal to cancer. In the current studies, two genetically altered mouse models with either a loss of heterozygosity in a critical cancer gene (Trp53) or a gain of oncogene function (Ha-ras) were used to determine how these animals would respond to acesulfame potassium exposure. These mouse models are susceptible to the rapid development of cancer. The Tg.AC hemizygous and p53 haploinsufficient mice are being evaluated by the National Institute of Environmental Health Sciences (NIEHS) as models for identifying chemical toxicity and/or chemical carcinogenic processes (Tennant *et al.*, 1996).

FVB/N-TgN(v-Ha-ras)Led (Tg.AC) Hemizygous Mouse Model

The Tg.AC mouse (on an FVB/N background) was developed by Leder *et al.* (1990) by introduction via pronuclear injection of a tripartite transgene composed of the promoter of the mouse embryonic zeta-globin gene, through the v-Ha-ras coding sequence, with point mutations in codons 12 and 59, and an SV40 polyadenylation sequence.

The Tg.AC transgenic mouse model has been evaluated as a reporter phenotype (skin papillomas) in response to either genotoxic or nongenotoxic carcinogens, including tumor promoters (Spalding *et al.*, 1993, 1999; Tennant *et al.*, 1999). Tg.AC mice are hemizygous for a mutant v-Ha-ras transgene. The model was developed by Leder *et al.* (1990) with an inducible zeta-globin promoter driving the expression of a mutated v-Ha-ras oncogene and is regarded as a genetically initiated model. With the exception of bone marrow, constitutive expression of the transgene cannot be detected in adult tissues. The transgene is transcriptionally silent until activated by certain treatments including full-thickness wounding, ultraviolet irradiation, or exposure to some chemicals (Cannon *et al.*, 1997; Trempus *et al.*, 1998). The Tg.AC hemizygous mouse develops a high incidence of skin papillomas in response to topical application of 12-*O*-tetradecanoyl-phorbol-13-acetate (TPA), and TPA has been used as a positive control in NIEHS Tg.AC mouse studies (Spalding *et al.*, 1993). Point mutations

in the *Ha-ras* gene are believed to be early events in the induction of skin papillomas and malignancies. Topical application of carcinogens to the shaved dorsal surface of Tg.AC mice induces epidermal squamous cell papillomas or carcinomas, a reporter phenotype that defines the activity of the chemical. The oral route of administration can also generate tumor responses in the skin of Tg.AC mice and lead to squamous cell papillomas and/or carcinomas of the forestomach. To date, the appearance of either spontaneous or induced tumors has been shown to require activation of transgene expression. However, the mechanism of response by the Tg.AC model to chemical carcinogens is not yet understood.

In NIEHS studies, mice are exposed beginning at 2 months of age for a total of 6 to 9 months. Cutaneous papillomas at various sites have been reported at 10% and 7% incidence in 26-week-old control male and female Tg.AC mice, respectively (Mahler *et al.*, 1998). Cutaneous papillomas occurring at sites such as the lip, pinnae, prepuce, and vulva suggest a possible relationship to grooming and chronic irritation. Up to 32% of Tg.AC homozygous and heterozygous male or female mice can develop odontogenic tumors as early as 26 weeks (Wright *et al.*, 1995; Mahler *et al.*, 1998). A number of different tumor types occur in untreated Tg.AC hemizygous mice at an incidence of greater than 3% including odontogenic tumors, forestomach papillomas, cutaneous papillomas, alveolar/bronchiolar adenomas, salivary gland duct carcinomas, and erythro-leukemia (Mahler *et al.*, 1998). In the FVB mouse (the background strain for the Tg.AC hemizygous mouse), alveolar/bronchiolar neoplasms occur at 14 months of age (Mahler *et al.*, 1996).

The Tg.AC hemizygous mouse model was used in the current studies of acesulfame potassium and in the companion studies of aspartame (NTP, 2005) because this model has been reported to detect both nongenotoxic and genotoxic carcinogens (Spalding *et al.*, 1993; Tennant *et al.*, 1995, 1996; Pritchard *et al.*, 2003).

B6.129-Trp53^{tm1Brd} (N5) Haploinsufficient Mouse Model

The heterozygous B6.129-Trp53 (N12)^{tm1Brd(+/-)} mouse (on a B6.129S7 background) was developed by Donehower *et al.* (1992). A null mutation was introduced into one p53 allele by homologous recombination in murine embryonic stem cells. Insertion of a neo cassette resulted in deletion of a 450-base pair gene fragment containing 106 nucleotides of exon 5 and approximately 350 nucleotides of intron 4.

Trp53, a nuclear protein, plays an essential role in the regulation of the cell cycle, specifically in the transition from G₀ to G₁, as well as G₂ to M, and the spindle apparatus. The p53 protein has a short half-life and exists at a very low concentration under normal cell physiological conditions. However, in DNA damaged cells that are able to replicate, p53 is expressed in high amounts with a significant increase in half-life due to post-translational modification (phosphorylation or acetylation). Mutations in p53 may also increase the protein half-life and alter functions that may contribute to transformation and development of the malignant phenotype. p53 is a DNA-binding protein containing DNA-binding, oligomerization, and transcription activation domains. Many amino acid residues in different p53 domains may be phosphorylated or acetylated, which may determine specific p53 functions. It is postulated to bind as a tetramer to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion or promote apoptosis, functioning as a tumor suppressor. This protein is critical to tumor suppression in humans and rodents. Mutants of p53 that fail to bind the consensus DNA binding site, and hence are unable to function as tumor suppressors, frequently occur in human cancers. Alterations of the *Trp53* gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome.

The mouse heterozygous for a p53 null allele (+/-) has only a single functional wild-type p53 allele which provides a target for mutagens. The p53 tumor suppressor gene is one of the most common sites for mutations and gene alterations in human cancer (Harris, 1996a,b,c).

Heterozygous p53^(+/-) transgenic mice develop normally, and like humans and other mammals, develop cancer (primarily lymphomas or sarcomas) with age, but often with decreased latency and increased susceptibility.

STUDY RATIONALE

Acesulfame potassium was nominated by The Center for Science in the Public Interest (CSPI) in 1996. At that time, the FDA was considering a food additive petition for the use of acesulfame potassium as an artificial sweetener in nonalcoholic beverages which would have substantially increased human exposure to acesulfame potassium. CSPI asked that the NTP review the data being submitted to support the petition, and if necessary, conduct additional testing or recommend to the FDA that additional testing be conducted by the sponsor. A review of the carcinogenicity data indicated that an

apparently adequate conventional 80-week mouse carcinogenicity study had been conducted with acesulfame potassium (Beems *et al.*, 1991). Because acesulfame potassium is not biotransformed after absorption and was negative in the 80-week mouse carcinogenicity study, it was considered a candidate for a negative control in the evaluation of Tg.AC and p53 mouse models.

The conventional rodent bioassay has been used for over three decades and is credible in identifying carcinogens thought to pose risks to humans (Tomatis *et al.*, 1997). An ongoing goal of the NIEHS and the NTP is to seek other model systems for toxicology and carcinogenesis studies, especially those that can provide mechanistic information that will assist in understanding an agent's mode of action. The use of genetically altered models holds promise for improving the accuracy and efficacy of experimental assessment of the carcinogenic potential of chemicals. Genetically altered mouse models carry activated oncogenes or inactivated tumor suppressor genes known to be involved in neoplastic processes in humans and rodents. This trait may allow them to respond to carcinogens more quickly than conventional

rodent strains. In addition, the neoplastic effects of agents can be observed in genetically altered models within a time frame in which few, if any, spontaneous tumors would arise. The high incidences of spontaneous or background tumors, which occur most often late in the 2-year rodent studies, can hinder interpretation of the findings and their implications for human health. The use of target or reporter genes allows for direct molecular and cellular analysis of a chemical's effects in genetically modified mouse models and can provide additional mechanistic information about the mode of action.

For the past few years, the NIEHS and the NTP have been actively evaluating genetically altered strains in toxicologic testing strategies. Based on completed evaluations, two models, the Tg.AC hemizygous (*v-Ha-ras*) and p53-deficient (p53 haploinsufficient) mice, have shown potential usefulness in identifying carcinogens (Pritchard *et al.*, 2003). Acesulfame potassium was one of the test agents selected for the continued evaluation of the genetically modified Tg.AC hemizygous and p53 haploinsufficient strains.

MATERIALS AND METHODS

PROCUREMENT AND CHARACTERIZATION OF ACESULFAME POTASSIUM

Acesulfame potassium was obtained from Riedel-deHaen (St. Louis, MO) via Research Triangle Institute (RTI) (Research Triangle Park, NC) in one lot (8415-76-02 RTI), which was used in the 9-month studies. Identity and purity analyses were conducted by the analytical chemistry laboratory, RTI, and the study laboratory; the study laboratory (BioReliance Corporation, Rockville, MD) also conducted stability analyses (Appendix F). Reports on analyses performed in support of the acesulfame potassium studies are on file at the National Institute of Environmental Health Sciences.

Lot 8415-76-02 RTI, a white, odorless, crystalline powder, was identified as acesulfame potassium by infrared and proton nuclear magnetic resonance spectroscopy. All spectra were consistent with the structure of acesulfame potassium. The purity of lot 8415-76-02 RTI was determined using high-performance liquid chromatography (HPLC) and confirmed by the study laboratory using major peak comparisons versus a reference sample from the same lot (stored at -20°C); HPLC indicated one major peak and no impurities. The purity was found to be greater than 99%.

To ensure stability, the bulk chemical was stored at room temperature under a headspace of inert gas, protected from light, in polyethylene bags inside plastic pails. Stability of the bulk chemical was monitored during the study by the study laboratory using HPLC. No degradation of the bulk chemical was observed.

PREPARATION AND ANALYSIS OF DOSE FORMULATIONS

The dose formulations were prepared once every two to four weeks by mixing acesulfame potassium with feed (Table F1). Homogeneity studies of the 0.1% and 5% acesulfame potassium formulations and stability studies of the 0.1% formulations were performed by the

analytical chemistry laboratory with HPLC. Homogeneity studies of the 0.3% and 3% formulations were performed by the study laboratory using HPLC. Homogeneity was confirmed, and stability was confirmed for at least 35 days for dose formulations stored in amber glass bottles at -20°C , -5°C , and 25°C , and when exposed to light and air for 7 days.

Periodic analyses of the dose formulations of acesulfame potassium were conducted by the study laboratory using HPLC. During the 9-month studies, the dose formulations were analyzed five times; all 15 of the dose formulations analyzed were within 10% of the target concentrations (Table F2). Animal room samples of these dose formulations were also analyzed, and 12 of the 15 samples were within 10% of the target concentrations (Table F2).

9-MONTH STUDIES

Study Design

Groups of 15 male and 15 female mice were fed diets containing 0%, 0.3%, 1%, or 3% (0, 3,000, 10,000, or 30,000 ppm) acesulfame potassium for 40 weeks.

Source and Specification of Animals

Male and female FVB/N-TgN(v-Ha-ras)Led (Tg.AC) hemizygous and B6.129-*Trp53*^{tm1Brd} (N5) haploinsufficient mice were obtained from Taconic Farms, Inc. (Germantown, NY), for use in the 9-month studies. Tg.AC hemizygous mice were quarantined for 11 days and p53 haploinsufficient mice were quarantined for 13 days before the beginning of the studies. Five male and five female mice in the control and 3% groups were randomly selected for parasite evaluation and gross observation of disease. Tg.AC hemizygous mice were 5 to 6 weeks old and p53 haploinsufficient mice were 6 to 7 weeks old at the beginning of the studies. The health of the mice was monitored during the studies according to the protocols of the NTP Sentinel Animal Program; all results were negative.

Animal Maintenance

Mice were housed individually. Feed and water were available *ad libitum*, and feed consumption was measured weekly. The feed was irradiated to reduce potential microbial contamination. Cages were changed weekly, and racks were changed and rotated every 2 weeks. Further details of animal maintenance are given in Table 1.

Clinical Examinations and Pathology

All animals were observed twice daily. Clinical findings and body weights were recorded initially, weekly, and at the end of the studies.

Complete necropsies were performed on all mice, except the control positive mice treated with TPA. At necropsy, all organs and tissues were examined for grossly visible lesions, and all major tissues were fixed and preserved in 10% neutral buffered formalin, processed and trimmed, embedded in paraffin, sectioned to a thickness of 4 to 6 μm , and stained with hematoxylin and eosin for microscopic examination. For all paired organs (e.g., adrenal gland, kidney, ovary), samples from each organ were examined. Complete microscopic examinations were performed on the control and 3% groups; the lung was

examined in all groups of male Tg.AC hemizygous mice. Tissues examined microscopically are listed in Table 1.

Microscopic evaluations were completed by the study laboratory pathologist, and the pathology data were entered into the Toxicology Data Management System. Upon completion of the laboratory pathologist's histologic evaluation, the slides, paraffin blocks, and residual wet tissues were sent to the NTP Archives for inventory, slide/block match, and wet tissue audit. The slides, individual animal data records, and pathology tables were evaluated by an independent pathology laboratory where quality assessment (QA) was performed. The QA pathologist reviewed all neoplasms from all groups and all slides from all animals in the control and high dose groups in Tg.AC hemizygous and p53 haploinsufficient mice. Results were reviewed by the NTP Pathology Working Group (PWG) Chairperson and two NTP pathologists. Minimal discrepancies were identified and the final diagnoses represent a consensus of the three contractors and two NTP pathologists. Details of these review procedures have been described, in part, by Maronpot and Boorman (1982) and Boorman *et al.* (1985).

TABLE 1
Experimental Design and Materials and Methods in the 9-Month Feed Studies of Acesulfame Potassium

| Tg.AC Hemizygous Mice | p53 Haploinsufficient Mice |
|--|---|
| Study Laboratory BioReliance Corporation (Rockville, MD) | BioReliance Corporation (Rockville, MD) |
| Strain FVB/N-TgN(v-Ha-ras)Led (Tg.AC) hemizygous | B6.129-Trp53 ^{tm1Brd} (N5) haploinsufficient |
| Animal Source Taconic Farms, Inc. (Germantown, NY) | Taconic Farms, Inc. (Germantown, NY) |
| Time Held Before Studies 11 days | 13 days |
| Average Age When Studies Began 5 to 6 weeks | 6 to 7 weeks |
| Date of First Exposure January 31, 2000 (males) February 1, 2000 (females) | February 2, 2000 (males) February 3, 2000 (females) |
| Duration of Exposure 40 weeks | 40 weeks |
| Date of Last Exposure October 30-31, 2000 | November 1-2, 2000 |
| Necropsy Dates October 30-31, 2000 | November 1-2, 2000 |
| Average Age at Necropsy 44 to 45 weeks | 45 to 46 weeks |
| Size of Study Groups 15 males and 15 females | 15 males and 15 females |
| Method of Distribution Animals were distributed randomly into groups of approximately equal initial mean body weights. | Animals were distributed randomly into groups of approximately equal initial mean body weights. |
| Animals per Cage 1 | 1 |
| Method of Animal Identification Tail tattoo | Tail tattoo |
| Diet Irradiated NTP-2000 open formula meal (Zeigler Brothers, Inc., Gardners, PA), available <i>ad libitum</i> | Same as Tg.AC hemizygous mouse study |
| Water Tap water (Washington Suburban Sanitary Commission Potomac Plant) via automatic watering system (Edstrom Industries, Inc., Waterford, WI), available <i>ad libitum</i> | Same as Tg.AC hemizygous mouse study |

TABLE 1
Experimental Design and Materials and Methods in the 9-Month Feed Studies of Acesulfame Potassium

| Tg.AC Hemizygous Mice | p53 Haploinsufficient Mice |
|--|---|
| Cages | |
| Polycarbonate cages (Lab Products, Inc., Seaford, DE), changed weekly | Same as Tg.AC hemizygous mouse study |
| Bedding | |
| Irradiated Sani-Chips® (P.J. Murphy Forest Produces, Montville, NJ), changed weekly | Same as Tg.AC hemizygous mouse study |
| Cage Filters | |
| Remay 2016 (Snow Filtration, West Chester, OH), changed every 2 weeks | Same as Tg.AC hemizygous mouse study |
| Racks | |
| Stainless steel (Lab Products, Inc., Seaford, DE), changed and rotated every 2 weeks | Same as Tg.AC hemizygous mouse study |
| Animal Room Environment | |
| Temperature: 72° ± 3° F | Temperature: 72° ± 3° F |
| Relative humidity: 50% ± 15% | Relative humidity: 50% ± 15% |
| Room fluorescent light: 12 hours/day | Room fluorescent light: 12 hours/day |
| Room air changes: 10/hour | Room air changes: 10/hour |
| Exposure Concentrations | |
| 0%, 0.3%, 1%, or 3% acesulfame potassium in feed, available <i>ad libitum</i> | 0%, 0.3%, 1%, or 3% acesulfame potassium in feed, available <i>ad libitum</i> |
| Type and Frequency of Observation | |
| Observed twice daily; animals were weighed and clinical observations were recorded initially, weekly, and at the end of the studies. Feed consumption was recorded weekly. | Observed twice daily; animals were weighed and clinical observations were recorded initially, weekly, and at the end of the studies. Feed consumption was recorded weekly. |
| Method of Sacrifice | |
| Carbon dioxide asphyxiation | Carbon dioxide asphyxiation |
| Necropsy | |
| Necropsy was performed on all mice. | Necropsy was performed on all mice. |
| Histopathology | |
| Complete histopathology was performed on all mice that died early and 0% and 3% mice. In addition to gross lesions and tissue masses, the following tissues were examined: adrenal gland, bone with marrow, brain, clitoral gland, esophagus, eye, gallbladder, harderian gland, heart and aorta, large intestine (cecum, colon, rectum), small intestine (duodenum, jejunum, ileum), kidney, liver, lung and mainstem bronchi, lymph nodes (mandibular and mesenteric), mammary gland, nose, ovary, pancreas, parathyroid gland, pituitary gland, preputial gland, prostate gland, salivary gland, skin, spleen, stomach (forestomach and glandular), testis (with epididymis and seminal vesicle), thymus, thyroid gland, tongue, trachea, urinary bladder, uterus, vagina, and Zymbal's gland. In addition, the lung was examined in the remaining groups of males. | Complete histopathology was performed on all mice that died early and 0% and 3% mice. In addition to gross lesions and tissue masses, the following tissues were examined: adrenal gland, bone with marrow, brain, clitoral gland, esophagus, eye, gallbladder, harderian gland, heart and aorta, large intestine (cecum, colon, rectum), small intestine (duodenum, jejunum, ileum), kidney, larynx, liver, lung and mainstem bronchi, lymph nodes (mandibular and mesenteric), mammary gland, nose, ovary, pancreas, parathyroid gland, pituitary gland, preputial gland, prostate gland, salivary gland, skin, spleen, stomach (forestomach and glandular), testis (with epididymis and seminal vesicle), thymus, thyroid gland, tongue, trachea, urinary bladder, uterus, vagina, and Zymbal's gland. |

STATISTICAL METHODS

Survival Analyses

The probability of survival was estimated by the product-limit procedure of Kaplan and Meier (1958) and is presented in the form of graphs. Animals found dead of other than natural causes were censored from the survival analyses; animals dying from natural causes were not censored. Statistical analyses for possible dose-related effects on survival used Cox's (1972) method for testing two groups for equality and Tarone's (1975) life table test to identify dose-related trends. All reported P values for the survival analyses are two sided.

Calculation of Incidence

The incidences of neoplasms or nonneoplastic lesions are presented in Tables A1, A4, B1, B4, C1, C4, D1, and D4 as the numbers of animals bearing such lesions at a specific anatomic site and the numbers of animals with that site examined microscopically. For calculation of statistical significance, the incidences of most neoplasms (Tables A3, B3, C3, and D3) and all nonneoplastic lesions are given as the numbers of animals affected at each site examined microscopically. However, when macroscopic examination was required to detect neoplasms in certain tissues (e.g., hardierian gland, intestine, mammary gland, and skin) before microscopic evaluation, or when neoplasms had multiple potential sites of occurrence (e.g., leukemia or lymphoma), the denominators consist of the number of animals on which a necropsy was performed. Tables A3, B3, C3, and D3 also give the survival-adjusted neoplasm rate for each group and each site-specific neoplasm. This survival-adjusted rate (based on the Poly-3 method described below) accounts for differential mortality by assigning a reduced risk of neoplasm, proportional to the third power of the fraction of time on study, only to site-specific lesion free animals that do not reach terminal sacrifice.

Analysis of Neoplasm and Nonneoplastic Lesion Incidences

The Poly-k test (Bailer and Portier, 1988; Portier and Bailer, 1989; Piegorsch and Bailer, 1997) was used to assess neoplasm and nonneoplastic lesion prevalence. This test is a survival-adjusted quantal-response procedure that modifies the Cochran-Armitage linear trend test to take survival differences into account. More specifically, this method modifies the denominator in the

quantal estimate of lesion incidence to approximate more closely the total number of animal years at risk. For analysis of a given site, each animal is assigned a risk weight. This value is one if the animal had a lesion at that site or if it survived until terminal sacrifice; if the animal died prior to terminal sacrifice and did not have a lesion at that site, its risk weight is the fraction of the entire study time that it survived, raised to the kth power.

This method yields a lesion prevalence rate that depends only upon the choice of a shape parameter for a Weibull hazard function describing cumulative lesion incidence over time (Bailer and Portier, 1988). Unless otherwise specified, a value of $k=3$ was used in the analysis of site-specific lesions. This value was recommended by Bailer and Portier (1988) following an evaluation of neoplasm onset time distributions for a variety of site-specific neoplasms in control F344 rats and B6C3F₁ mice (Portier *et al.*, 1986). Bailer and Portier (1988) showed that the Poly-3 test gave valid results if the true value of k was anywhere in the range from 1 to 5. A further advantage of the Poly-3 method is that it does not require lesion lethality assumptions. Variation introduced by the use of risk weights, which reflect differential mortality, was accommodated by adjusting the variance of the Poly-3 statistic as recommended by Bieler and Williams (1993).

Tests of significance included pairwise comparisons of each exposed group with controls and a test for an overall exposure-related trend. Continuity-corrected Poly-3 tests were used in the analysis of lesion incidence, and reported P values are one sided. The significance of lower incidences or decreasing trends in lesions is represented as 1-P with the letter N added (e.g., $P=0.99$ is presented as $P=0.01N$).

Analysis of Continuous Variables

Body weight data, which historically have approximately normal distributions, were analyzed with the parametric multiple comparison procedures of Dunnett (1955) and Williams (1971, 1972). Average severity values were analyzed for significance with the Mann-Whitney U test (Hollander and Wolfe, 1973).

QUALITY ASSURANCE METHODS

The 9-month studies were conducted in compliance with Food and Drug Administration Good Laboratory Practice Regulations (21 CFR, Part 58). In addition, as records from the 9-month studies were submitted to the NTP Archives, these studies were audited

retrospectively by an independent quality assurance contractor. Separate audits covered completeness and accuracy of the pathology data, pathology specimens, final pathology tables, and a draft of this NTP Technical Report. Audit procedures and findings are presented in the reports and are on file at NIEHS. The audit findings were reviewed and assessed by NTP staff, and all comments were resolved or otherwise addressed during the preparation of this Technical Report.

GENETIC TOXICOLOGY

The genetic toxicity of acesulfame potassium was assessed by testing the ability of the chemical to induce increases in the frequency of micronucleated erythrocytes in mouse peripheral blood. The protocol for this study and the results are given in Appendix E.

The genetic toxicity studies have evolved from an earlier effort by the NTP to develop a comprehensive database permitting a critical anticipation of a chemical's carcinogenicity in experimental animals based on numerous considerations, including the molecular structure of the chemical and its observed effects in short-term *in vitro* and *in vivo* genetic toxicity tests (structure-activity relationships). The short-term tests were originally developed to clarify proposed mechanisms of chemical-induced DNA damage based on the relationship between electrophilicity and mutagenicity (Miller and Miller, 1977) and the somatic mutation theory of cancer (Straus, 1981; Crawford, 1985). However, it should be noted that not all cancers arise through genotoxic mechanisms.

DNA reactivity combined with *Salmonella* mutagenicity is highly correlated with induction of carcinogenicity in multiple species/sexes of rodents and at multiple tissue sites (Ashby and Tennant, 1991). A positive response in the *Salmonella* test was shown to be the most predictive *in vitro* indicator for rodent carcinogenicity (89% of the *Salmonella* mutagens are rodent carcinogens) (Tennant *et al.*, 1987; Zeiger *et al.*, 1990). Other tests, although less predictive of rodent carcinogenicity, can provide useful information on the types of DNA and chromosomal damage induced by the chemical under investigation.

The predictivity for carcinogenicity of a positive response in acute *in vivo* bone marrow chromosome aberration or micronucleus tests appears to be less than that in the *Salmonella* test (Shelby *et al.*, 1993; Shelby and Witt, 1995). However, clearly positive results in long-term peripheral blood micronucleus tests have high predictivity for rodent carcinogenicity (Witt *et al.*, 2000); negative results in this assay do not correlate well with either negative or positive results in rodent carcinogenicity studies. Because of the theoretical and observed associations between induced genetic damage and adverse effects in somatic and germ cells, the determination of *in vivo* genetic effects is important to the overall understanding of the risks associated with exposure to a particular chemical. Most organic chemicals that are identified by the International Agency for Research on Cancer as human carcinogens, other than hormones, are genotoxic. The vast majority of these are detected by both the *Salmonella* assay and rodent bone marrow cytogenetics tests (Shelby, 1988; Shelby and Zeiger, 1990).

RESULTS

9-MONTH STUDY IN Tg.AC HEMIZYGOUS MICE

Positive Control Tg.AC Hemizygous Mice

12-*O*-Tetradecanoylphorbol-13-acetate (TPA) (1.25 µg) was dermally administered to groups of 15 males and 15 females three times weekly for up to 16 weeks. All males and 87% of the females developed more than 20 papillomas each by week 16 (data not shown). This is consistent with historical rates found in other studies (Tennant *et al.*, 2001).

Survival

Estimates of 9-month survival probabilities for male and female mice are shown in Table 2 and in the Kaplan-Meier survival curves (Figure 1). Survival of all exposed groups was similar to that of the control groups.

TABLE 2
Survival of Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|----------|---------|----------|----------|
| Male | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Moribund | 2 | 4 | 2 | 1 |
| Animals surviving to study termination | 13 | 11 | 13 | 14 |
| Percent probability of survival at end of study ^a | 87 | 73 | 87 | 93 |
| Mean survival (days) ^b | 258 | 249 | 268 | 267 |
| Survival analysis ^c | P=0.430N | P=0.638 | P=1.000N | P=0.951N |
| Female | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Moribund | 1 | 0 | 2 | 2 |
| Natural deaths | 2 | 3 | 1 | 1 |
| Animals surviving to study termination | 12 | 12 | 12 | 12 |
| Percent probability of survival at end of study | 80 | 80 | 80 | 80 |
| Mean survival (days) | 266 | 255 | 267 | 262 |
| Survival analysis | P=1.000 | P=1.000 | P=1.000 | P=1.000 |

^a Kaplan-Meier determinations

^b Mean of all deaths (uncensored, censored, and terminal sacrifice).

^c The result of the life table trend test (Tarone, 1975) is in the control column, and the results of the life table pairwise comparisons (Cox, 1972) with the controls are in the exposed group columns. A negative trend or lower mortality in an exposed group is indicated by N.




FIGURE 1
Kaplan-Meier Survival Curves for Male and Female Tg.AC Hemizygous Mice
Exposed to Acesulfame Potassium in Feed for 9 Months

Body Weights, Feed and Compound Consumption, and Clinical Findings

Mean body weights of all exposed groups of males were similar to those of the control group throughout the study; mean body weights of all exposed groups of females were less than the control group after week 35 (Tables 3 and 4; Figure 2). Because the body weights of females did not decrease with increasing exposure concentration, the body weight effect was not considered to be treatment related. Feed consumption by the exposed

groups was similar to that by the control groups (Tables G1 and G2). Dietary concentrations of 0.3%, 1%, and 3% acesulfame potassium delivered average daily doses 420, 1,400, or 4,500 mg acesulfame potassium/kg body weight to males and 520, 1,700, or 5,400 mg/kg to females. Clinical findings included jaw masses, malocclusions, and cutaneous papillomas, but these findings did not occur in a pattern that would suggest a relationship to acesulfame potassium exposure.

TABLE 3
Mean Body Weights and Survival of Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Weeks on Study | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|----------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|
| | Av. Wt. (g) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors |
| 1 | 21.2 | 15 | 21.2 | 100 | 15 | 21.3 | 101 | 15 | 21.3 | 101 | 15 |
| 2 | 22.6 | 15 | 22.0 | 97 | 15 | 22.7 | 100 | 15 | 22.5 | 100 | 15 |
| 3 | 23.3 | 15 | 23.6 | 101 | 15 | 23.7 | 102 | 15 | 23.7 | 102 | 15 |
| 4 | 24.2 | 15 | 24.9 | 103 | 15 | 24.8 | 103 | 15 | 24.7 | 102 | 15 |
| 5 | 24.9 | 15 | 25.7 | 103 | 15 | 25.3 | 102 | 15 | 25.1 | 101 | 15 |
| 6 | 25.5 | 15 | 26.3 | 103 | 15 | 26.4 | 104 | 15 | 25.6 | 100 | 15 |
| 7 | 26.3 | 15 | 26.5 | 101 | 15 | 26.8 | 102 | 15 | 26.9 | 102 | 15 |
| 8 | 25.9 | 15 | 27.2 | 105 | 15 | 27.5 | 106 | 15 | 27.4 | 106 | 15 |
| 9 | 27.4 | 15 | 27.6 | 101 | 15 | 27.7 | 101 | 15 | 27.7 | 101 | 15 |
| 10 | 27.5 | 15 | 27.6 | 100 | 15 | 28.0 | 102 | 15 | 27.7 | 101 | 15 |
| 11 | 27.0 | 15 | 28.2 | 104 | 14 | 28.4 | 105 | 15 | 27.6 | 102 | 15 |
| 12 | 28.0 | 15 | 29.0 | 104 | 14 | 29.2 | 104 | 15 | 28.8 | 103 | 15 |
| 13 | 28.2 | 15 | 28.9 | 103 | 14 | 29.4 | 104 | 15 | 29.0 | 103 | 15 |
| 14 | 28.8 | 15 | 29.8 | 104 | 14 | 29.7 | 103 | 15 | 29.3 | 102 | 15 |
| 15 | 29.3 | 15 | 29.9 | 102 | 14 | 30.0 | 102 | 15 | 29.9 | 102 | 15 |
| 16 | 29.5 | 15 | 29.8 | 101 | 14 | 30.3 | 103 | 15 | 30.1 | 102 | 15 |
| 17 | 29.6 | 15 | 29.8 | 101 | 14 | 30.5 | 103 | 15 | 30.4 | 103 | 15 |
| 18 | 29.2 | 15 | 30.5 | 105 | 14 | 30.7 | 105 | 15 | 30.6 | 105 | 15 |
| 19 | 29.4 | 15 | 30.4 | 103 | 14 | 30.5 | 104 | 15 | 30.7 | 104 | 15 |
| 20 | 29.5 | 15 | 29.7 | 101 | 14 | 30.5 | 103 | 15 | 30.6 | 104 | 15 |
| 21 | 29.5 | 15 | 30.1 | 102 | 14 | 31.2 | 106 | 15 | 31.1 | 105 | 15 |
| 22 | 29.5 | 15 | 30.9 | 105 | 14 | 31.6 | 107 | 15 | 31.5 | 107 | 15 |
| 23 | 30.4 | 14 | 32.0 | 105 | 13 | 31.4 | 103 | 15 | 31.8 | 105 | 15 |
| 24 | 31.5 | 13 | 31.1 | 99 | 13 | 31.8 | 101 | 15 | 31.9 | 101 | 15 |
| 25 | 31.6 | 13 | 32.2 | 102 | 13 | 32.2 | 102 | 15 | 32.1 | 102 | 15 |
| 26 | 31.8 | 13 | 32.2 | 101 | 13 | 32.2 | 101 | 15 | 32.8 | 103 | 14 |
| 27 | 32.0 | 13 | 32.2 | 101 | 13 | 32.2 | 101 | 15 | 32.3 | 101 | 14 |
| 28 | 32.0 | 13 | 32.6 | 102 | 13 | 32.9 | 103 | 15 | 32.8 | 103 | 14 |
| 29 | 32.4 | 13 | 32.8 | 101 | 13 | 33.1 | 102 | 14 | 32.5 | 100 | 14 |
| 30 | 32.5 | 13 | 33.3 | 103 | 13 | 33.4 | 103 | 14 | 33.0 | 102 | 14 |
| 31 | 32.5 | 13 | 33.3 | 103 | 13 | 33.1 | 102 | 14 | 33.0 | 102 | 14 |
| 32 | 32.5 | 13 | 33.2 | 102 | 13 | 33.6 | 103 | 14 | 33.7 | 104 | 14 |
| 33 | 32.3 | 13 | 32.5 | 101 | 13 | 33.2 | 103 | 14 | 33.0 | 102 | 14 |
| 34 | 32.2 | 13 | 32.7 | 102 | 13 | 33.4 | 104 | 14 | 33.4 | 104 | 14 |
| 35 | 32.6 | 13 | 32.8 | 101 | 12 | 33.4 | 103 | 14 | 33.4 | 103 | 14 |
| 36 | 33.0 | 13 | 32.7 | 99 | 12 | 33.7 | 102 | 14 | 33.4 | 101 | 14 |
| 37 | 32.7 | 13 | 33.5 | 102 | 12 | 34.1 | 104 | 14 | 33.3 | 102 | 14 |
| 38 | 32.9 | 13 | 33.5 | 102 | 12 | 33.4 | 102 | 14 | 33.2 | 101 | 14 |
| 39 | 33.2 | 13 | 32.9 | 99 | 12 | 34.4 | 104 | 13 | 33.9 | 102 | 14 |
| 40 | 33.2 | 13 | 33.6 | 101 | 11 | 34.1 | 103 | 13 | 33.5 | 101 | 14 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 25.5 | | 26.1 | 102 | | 26.2 | 103 | | 26.0 | 102 | |
| 14-40 | 31.3 | | 31.9 | 102 | | 32.2 | 103 | | 32.1 | 103 | |

TABLE 4
Mean Body Weights and Survival of Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Weeks on Study | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|----------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|
| | Av. Wt. (g) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors |
| 1 | 17.9 | 15 | 17.8 | 99 | 15 | 18.0 | 101 | 15 | 17.8 | 99 | 15 |
| 2 | 17.5 | 15 | 18.2 | 104 | 15 | 18.2 | 104 | 15 | 18.1 | 103 | 15 |
| 3 | 17.8 | 15 | 18.5 | 104 | 15 | 18.3 | 103 | 15 | 18.3 | 103 | 15 |
| 4 | 19.6 | 15 | 19.5 | 100 | 15 | 19.3 | 99 | 15 | 19.7 | 101 | 15 |
| 5 | 19.7 | 15 | 19.6 | 100 | 15 | 19.8 | 101 | 15 | 19.7 | 100 | 15 |
| 6 | 19.7 | 15 | 19.5 | 99 | 15 | 19.6 | 100 | 15 | 19.4 | 99 | 15 |
| 7 | 20.3 | 15 | 20.1 | 99 | 15 | 20.5 | 101 | 15 | 20.6 | 102 | 15 |
| 8 | 20.7 | 15 | 20.8 | 101 | 15 | 21.2 | 102 | 15 | 20.8 | 101 | 15 |
| 9 | 21.3 | 15 | 21.6 | 101 | 15 | 21.5 | 101 | 15 | 21.2 | 100 | 15 |
| 10 | 21.7 | 15 | 21.6 | 100 | 15 | 21.9 | 101 | 15 | 21.4 | 99 | 15 |
| 11 | 22.1 | 15 | 21.9 | 99 | 15 | 22.2 | 101 | 15 | 22.3 | 101 | 15 |
| 12 | 22.5 | 15 | 22.1 | 98 | 14 | 22.6 | 100 | 15 | 21.9 | 97 | 15 |
| 13 | 22.2 | 15 | 22.4 | 101 | 14 | 22.7 | 102 | 15 | 21.9 | 99 | 15 |
| 14 | 22.3 | 15 | 22.7 | 102 | 14 | 22.7 | 102 | 15 | 22.1 | 99 | 15 |
| 15 | 22.5 | 15 | 23.1 | 103 | 14 | 22.9 | 102 | 15 | 22.6 | 100 | 15 |
| 16 | 23.5 | 15 | 23.3 | 99 | 14 | 23.2 | 99 | 15 | 23.4 | 100 | 15 |
| 17 | 23.7 | 15 | 23.6 | 100 | 14 | 23.2 | 98 | 15 | 23.0 | 97 | 15 |
| 18 | 24.0 | 15 | 23.4 | 98 | 14 | 23.7 | 99 | 15 | 23.3 | 97 | 15 |
| 19 | 23.8 | 15 | 23.3 | 98 | 14 | 23.9 | 100 | 15 | 23.3 | 98 | 15 |
| 20 | 24.4 | 15 | 23.5 | 96 | 14 | 23.9 | 98 | 15 | 23.7 | 97 | 15 |
| 21 | 24.6 | 15 | 23.8 | 97 | 14 | 24.4 | 99 | 15 | 23.5 | 96 | 15 |
| 22 | 24.9 | 15 | 24.0 | 96 | 14 | 24.6 | 99 | 15 | 23.3 | 94 | 15 |
| 23 | 25.0 | 15 | 24.1 | 96 | 14 | 24.2 | 97 | 15 | 24.0 | 96 | 15 |
| 24 | 25.4 | 15 | 24.5 | 97 | 14 | 24.8 | 98 | 15 | 24.7 | 97 | 14 |
| 25 | 25.3 | 15 | 24.5 | 97 | 14 | 24.7 | 98 | 15 | 25.1 | 99 | 14 |
| 26 | 25.4 | 15 | 24.8 | 98 | 14 | 25.0 | 98 | 15 | 25.2 | 99 | 14 |
| 27 | 26.7 | 15 | 25.3 | 95 | 14 | 25.2 | 94 | 15 | 25.1 | 94 | 14 |
| 28 | 26.5 | 15 | 25.5 | 96 | 14 | 25.1 | 95 | 15 | 25.1 | 95 | 14 |
| 29 | 26.8 | 15 | 25.2 | 94 | 13 | 25.1 | 94 | 15 | 24.8 | 93 | 14 |
| 30 | 27.1 | 15 | 25.6 | 95 | 13 | 25.4 | 94 | 15 | 25.7 | 95 | 14 |
| 31 | 26.7 | 15 | 25.5 | 96 | 13 | 25.5 | 96 | 14 | 26.1 | 98 | 14 |
| 32 | 26.4 | 15 | 25.7 | 97 | 13 | 26.0 | 99 | 14 | 26.1 | 99 | 14 |
| 33 | 25.9 | 14 | 25.8 | 100 | 13 | 25.5 | 99 | 14 | 25.8 | 100 | 14 |
| 34 | 26.0 | 14 | 26.1 | 100 | 13 | 25.4 | 98 | 14 | 25.6 | 99 | 14 |
| 35 | 26.0 | 13 | 24.9 | 96 | 13 | 25.9 | 100 | 13 | 26.4 | 102 | 13 |
| 36 | 28.3 | 12 | 25.9 | 92 | 13 | 26.3 | 93 | 13 | 26.2 | 93 | 13 |
| 37 | 28.4 | 12 | 26.4 | 93 | 13 | 26.6 | 94 | 13 | 26.3 | 93 | 13 |
| 38 | 27.8 | 12 | 26.7 | 96 | 12 | 26.6 | 96 | 13 | 26.6 | 96 | 12 |
| 39 | 28.5 | 12 | 26.6 | 93 | 12 | 26.8 | 94 | 13 | 26.7 | 94 | 12 |
| 40 | 28.5 | 12 | 26.3 | 92 | 12 | 27.0 | 95 | 12 | 26.6 | 93 | 12 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 20.2 | | 20.3 | 100 | | 20.4 | 101 | | 20.2 | 100 | |
| 14-40 | 25.7 | | 24.8 | 96 | | 24.9 | 97 | | 24.8 | 96 | |




FIGURE 2
Growth Curves for Male and Female Tg.AC Hemizygous Mice
Exposed to Acesulfame Potassium in Feed for 9 Months

Pathology and Statistical Analyses

This section describes the statistically significant or biologically noteworthy changes in the incidences of neoplasms of the jaw and skin. Summaries of the incidences of neoplasms and nonneoplastic lesions, individual animal tumor diagnoses, and statistical analyses of primary neoplasms that occurred with an incidence of at least 5% in at least one animal group are presented in Appendix A for males and Appendix B for females.

Jaw: Gross observations at necropsy included jaw masses in females. The incidence of odontogenic tumor in 1% females was slightly increased (0%, 2/15; 0.3%, 0/15; 1%, 6/15; 3%, 3/15; Table B3); however, the increased incidence was considered due to biologic variability and not to acesulfame potassium exposure.

Skin: Gross observations at necropsy included skin masses (papillomas) at several locations, particularly the face, head, and genital region; most masses occurred on the lip. Based on the microscopic evaluation, there were no significant increases in the incidences of squamous cell papilloma at any specific site in males or females (Tables A1 and B1). However, when papillomas at all sites were combined, there was an increase in the incidence of squamous cell papilloma in 1% males (3/15, 4/15, 8/15, 5/15; Table A3). With the exception of one male in the 3% group and one female in the 1% group, each of which had two papillomas at different locations, all other animals had a single papilloma (Tables A2 and B2).

9-MONTH STUDY IN p53 HAPLOINSUFFICIENT MICE

Survival

Estimates of 9-month survival probabilities for male and female mice are shown in Table 5 and in the Kaplan-Meier survival curves (Figure 3). Survival of all exposed groups was similar to that of the control groups.

Body Weights, Feed and Compound Consumption, and Clinical Findings

Mean body weights of 1% males were greater than those of the controls after week 26; however, this trend likely reflected biologic variability and sample size rather than

a treatment effect (Table 6 and Figure 4). Mean body weights of 0.3% and 3% males and of all exposed groups of females were similar to those of the control groups (Table 7 and Figure 4). Feed consumption by the exposed groups was similar to that by the control groups (Tables G3 and G4). Dietary concentrations of 0.3%, 1%, and 3% acesulfame potassium delivered average daily doses of approximately 475, 1,500, or 4,700 mg/kg to males and 570, 1,800, or 5,700 mg/kg to females. There were no clinical findings related to acesulfame potassium exposure.

TABLE 5
Survival of p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|----------|----------|----------|----------|
| Male | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Moribund | 1 | 0 | 0 | 1 |
| Animals surviving to study termination | 14 | 15 | 15 | 14 |
| Percent probability of survival at end of study ^a | 93 | 100 | 100 | 93 |
| Mean survival (days) ^b | 268 | 274 | 274 | 271 |
| Survival analysis ^c | P=1.000 | P=1.000N | P=1.000N | P=1.000N |
| Female | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Moribund | 1 | 0 | 1 | 0 |
| Natural deaths | 0 | 1 | 0 | 1 |
| Animals surviving to study termination | 14 | 14 | 14 | 14 |
| Percent probability of survival at end of study | 93 | 93 | 93 | 93 |
| Mean survival (days) | 267 | 268 | 264 | 272 |
| Survival analysis | P=1.000N | P=1.000N | P=1.000 | P=1.000N |

^a Kaplan-Meier determinations

^b Mean of all deaths (uncensored, censored, and terminal sacrifice).

^c The result of the life table trend test (Tarone, 1975) is in the control column, and the results of the life table pairwise comparisons (Cox, 1972) with the controls are in the exposed group columns. A negative trend or lower mortality in an exposed group is indicated by N.





FIGURE 3
Kaplan-Meier Survival Curves for Male and Female p53 Haploinsufficient Mice Exposed to Acesulfame Potassium in Feed for 9 Months

TABLE 6
Mean Body Weights and Survival of Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Weeks on Study | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|----------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|
| | Av. Wt. (g) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors |
| 1 | 20.9 | 15 | 20.5 | 98 | 15 | 20.5 | 98 | 15 | 21.1 | 101 | 15 |
| 2 | 21.9 | 15 | 21.7 | 99 | 15 | 22.0 | 101 | 15 | 22.0 | 101 | 15 |
| 3 | 22.1 | 15 | 21.9 | 99 | 15 | 22.3 | 101 | 15 | 22.1 | 100 | 15 |
| 4 | 22.9 | 15 | 22.5 | 98 | 15 | 22.8 | 100 | 15 | 22.7 | 99 | 15 |
| 5 | 23.6 | 15 | 23.6 | 100 | 15 | 23.9 | 101 | 15 | 24.1 | 102 | 15 |
| 6 | 24.1 | 15 | 23.9 | 99 | 15 | 24.4 | 101 | 15 | 24.4 | 101 | 15 |
| 7 | 24.5 | 15 | 24.2 | 99 | 15 | 25.0 | 102 | 15 | 24.7 | 101 | 15 |
| 8 | 25.3 | 15 | 24.8 | 98 | 15 | 25.5 | 101 | 15 | 25.2 | 100 | 15 |
| 9 | 25.4 | 15 | 24.9 | 98 | 15 | 25.6 | 101 | 15 | 25.4 | 100 | 15 |
| 10 | 25.6 | 15 | 25.1 | 98 | 15 | 26.1 | 102 | 15 | 25.6 | 100 | 15 |
| 11 | 25.9 | 15 | 25.4 | 98 | 15 | 26.8 | 104 | 15 | 25.7 | 99 | 15 |
| 12 | 26.5 | 15 | 25.6 | 97 | 15 | 27.1 | 102 | 15 | 26.3 | 99 | 15 |
| 13 | 26.6 | 15 | 25.9 | 97 | 15 | 26.9 | 101 | 15 | 26.5 | 100 | 15 |
| 14 | 27.0 | 15 | 26.7 | 99 | 15 | 27.2 | 101 | 15 | 27.0 | 100 | 15 |
| 15 | 27.5 | 15 | 27.2 | 99 | 15 | 28.3 | 103 | 15 | 27.9 | 102 | 15 |
| 16 | 28.0 | 15 | 27.9 | 100 | 15 | 28.4 | 101 | 15 | 28.1 | 100 | 15 |
| 17 | 28.5 | 15 | 27.7 | 97 | 15 | 28.9 | 101 | 15 | 28.2 | 99 | 15 |
| 18 | 29.4 | 15 | 28.8 | 98 | 15 | 30.1 | 102 | 15 | 29.1 | 99 | 15 |
| 19 | 29.2 | 15 | 28.3 | 97 | 15 | 29.8 | 102 | 15 | 28.9 | 99 | 15 |
| 20 | 29.4 | 15 | 28.6 | 97 | 15 | 29.2 | 99 | 15 | 29.1 | 99 | 15 |
| 21 | 29.7 | 15 | 28.8 | 97 | 15 | 30.0 | 101 | 15 | 29.4 | 99 | 15 |
| 22 | 29.5 | 15 | 29.0 | 98 | 15 | 29.9 | 101 | 15 | 29.5 | 100 | 15 |
| 23 | 29.2 | 15 | 28.5 | 98 | 15 | 29.6 | 101 | 15 | 29.5 | 101 | 15 |
| 24 | 29.6 | 15 | 29.2 | 99 | 15 | 30.3 | 102 | 15 | 30.3 | 102 | 15 |
| 25 | 29.6 | 15 | 28.9 | 98 | 15 | 30.5 | 103 | 15 | 29.9 | 101 | 15 |
| 26 | 29.6 | 15 | 29.4 | 99 | 15 | 30.5 | 103 | 15 | 30.3 | 102 | 15 |
| 27 | 29.0 | 15 | 29.0 | 100 | 15 | 30.9 | 107 | 15 | 30.4 | 105 | 15 |
| 28 | 29.3 | 15 | 29.8 | 102 | 15 | 31.9 | 109 | 15 | 31.0 | 106 | 15 |
| 29 | 30.3 | 14 | 30.2 | 100 | 15 | 32.1 | 106 | 15 | 30.5 | 101 | 15 |
| 30 | 30.4 | 14 | 30.5 | 100 | 15 | 32.4 | 107 | 15 | 31.1 | 102 | 15 |
| 31 | 30.1 | 14 | 30.0 | 100 | 15 | 32.2 | 107 | 15 | 31.1 | 103 | 15 |
| 32 | 29.9 | 14 | 29.3 | 98 | 15 | 32.0 | 107 | 15 | 30.9 | 103 | 15 |
| 33 | 29.6 | 14 | 29.5 | 100 | 15 | 31.1 | 105 | 15 | 30.4 | 103 | 14 |
| 34 | 29.2 | 14 | 29.2 | 100 | 15 | 31.3 | 107 | 15 | 30.2 | 103 | 14 |
| 35 | 29.2 | 14 | 29.3 | 100 | 15 | 30.8 | 106 | 15 | 29.6 | 101 | 14 |
| 36 | 29.1 | 14 | 29.3 | 101 | 15 | 31.0 | 107 | 15 | 29.4 | 101 | 14 |
| 37 | 29.1 | 14 | 29.7 | 102 | 15 | 31.1 | 107 | 15 | 29.7 | 102 | 14 |
| 38 | 29.1 | 14 | 29.5 | 101 | 15 | 31.3 | 108 | 15 | 29.9 | 103 | 14 |
| 39 | 29.2 | 14 | 29.9 | 102 | 15 | 31.7 | 109 | 15 | 30.1 | 103 | 14 |
| 40 | 29.5 | 14 | 30.1 | 102 | 15 | 31.6 | 107 | 15 | 29.9 | 101 | 14 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 24.3 | | 23.8 | 98 | | 24.5 | 101 | | 24.3 | 100 | |
| 14-40 | 29.2 | | 29.0 | 99 | | 30.5 | 104 | | 29.7 | 102 | |

TABLE 7
Mean Body Weights and Survival of Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Weeks on Study | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|----------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|----------------|------------------------|---------------------|
| | Av. Wt. (g) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors | Av. Wt. (g) | Wt. (% of controls) | No. of Survivors |
| 1 | 16.4 | 15 | 17.0 | 104 | 15 | 17.0 | 104 | 15 | 16.4 | 100 | 15 |
| 2 | 18.5 | 15 | 17.9 | 97 | 15 | 18.4 | 100 | 15 | 17.8 | 96 | 15 |
| 3 | 18.4 | 15 | 18.7 | 102 | 15 | 19.2 | 104 | 15 | 18.5 | 101 | 15 |
| 4 | 19.6 | 15 | 19.5 | 100 | 15 | 19.9 | 102 | 15 | 19.1 | 97 | 15 |
| 5 | 20.2 | 15 | 20.7 | 103 | 15 | 20.6 | 102 | 15 | 20.1 | 100 | 15 |
| 6 | 20.9 | 15 | 21.1 | 101 | 15 | 21.2 | 101 | 15 | 21.0 | 101 | 15 |
| 7 | 21.0 | 15 | 21.2 | 101 | 15 | 21.5 | 102 | 15 | 21.1 | 101 | 15 |
| 8 | 21.4 | 15 | 21.5 | 101 | 15 | 21.5 | 101 | 15 | 20.8 | 97 | 15 |
| 9 | 21.4 | 15 | 21.5 | 101 | 15 | 21.7 | 101 | 15 | 21.3 | 100 | 15 |
| 10 | 21.8 | 15 | 21.6 | 99 | 15 | 21.9 | 101 | 15 | 21.6 | 99 | 15 |
| 11 | 22.4 | 15 | 22.4 | 100 | 15 | 22.4 | 100 | 15 | 21.8 | 97 | 15 |
| 12 | 22.6 | 15 | 22.3 | 99 | 15 | 22.6 | 100 | 15 | 22.3 | 99 | 15 |
| 13 | 23.0 | 15 | 23.1 | 100 | 15 | 23.2 | 101 | 15 | 22.8 | 99 | 15 |
| 14 | 23.2 | 15 | 23.6 | 102 | 15 | 23.5 | 101 | 15 | 22.9 | 99 | 15 |
| 15 | 23.6 | 15 | 24.2 | 103 | 15 | 23.7 | 100 | 15 | 22.9 | 97 | 15 |
| 16 | 24.1 | 15 | 25.2 | 105 | 15 | 24.2 | 100 | 15 | 23.6 | 98 | 15 |
| 17 | 24.3 | 15 | 25.1 | 103 | 15 | 24.2 | 100 | 15 | 23.7 | 98 | 15 |
| 18 | 24.7 | 15 | 25.6 | 104 | 15 | 25.0 | 101 | 15 | 24.1 | 98 | 15 |
| 19 | 24.8 | 15 | 25.8 | 104 | 15 | 24.9 | 100 | 14 | 24.6 | 99 | 15 |
| 20 | 25.1 | 15 | 26.1 | 104 | 15 | 25.5 | 102 | 14 | 24.9 | 99 | 15 |
| 21 | 25.4 | 15 | 26.4 | 104 | 15 | 25.7 | 101 | 14 | 25.3 | 100 | 15 |
| 22 | 25.6 | 15 | 26.8 | 105 | 15 | 26.1 | 102 | 14 | 25.5 | 100 | 15 |
| 23 | 26.1 | 15 | 27.0 | 103 | 15 | 26.3 | 101 | 14 | 25.6 | 98 | 15 |
| 24 | 26.1 | 15 | 26.9 | 103 | 15 | 26.6 | 102 | 14 | 26.0 | 100 | 15 |
| 25 | 26.4 | 14 | 26.9 | 102 | 15 | 26.7 | 101 | 14 | 26.3 | 100 | 15 |
| 26 | 26.5 | 14 | 27.1 | 102 | 14 | 26.5 | 100 | 14 | 26.3 | 99 | 15 |
| 27 | 26.7 | 14 | 27.3 | 102 | 14 | 27.2 | 102 | 14 | 26.9 | 101 | 15 |
| 28 | 26.9 | 14 | 27.8 | 103 | 14 | 27.4 | 102 | 14 | 27.4 | 102 | 15 |
| 29 | 27.0 | 14 | 28.0 | 104 | 14 | 28.0 | 104 | 14 | 27.1 | 100 | 15 |
| 30 | 28.2 | 14 | 28.6 | 101 | 14 | 28.5 | 101 | 14 | 28.0 | 99 | 15 |
| 31 | 27.8 | 14 | 28.6 | 103 | 14 | 28.5 | 103 | 14 | 27.9 | 100 | 15 |
| 32 | 28.1 | 14 | 27.9 | 99 | 14 | 28.5 | 101 | 14 | 27.5 | 98 | 15 |
| 33 | 27.5 | 14 | 28.0 | 102 | 14 | 28.4 | 103 | 14 | 27.9 | 102 | 15 |
| 34 | 27.7 | 14 | 27.8 | 100 | 14 | 28.3 | 102 | 14 | 27.5 | 99 | 15 |
| 35 | 27.6 | 14 | 27.3 | 99 | 14 | 27.7 | 100 | 14 | 26.8 | 97 | 15 |
| 36 | 27.3 | 14 | 28.1 | 103 | 14 | 28.2 | 103 | 14 | 27.1 | 99 | 14 |
| 37 | 27.5 | 14 | 28.0 | 102 | 14 | 28.0 | 102 | 14 | 27.6 | 100 | 14 |
| 38 | 27.7 | 14 | 28.1 | 101 | 14 | 27.9 | 101 | 14 | 27.6 | 100 | 14 |
| 39 | 27.5 | 14 | 28.3 | 103 | 14 | 28.2 | 103 | 14 | 28.0 | 102 | 14 |
| 40 | 28.3 | 14 | 28.8 | 102 | 14 | 28.8 | 102 | 14 | 28.3 | 100 | 14 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 20.6 | | 20.7 | 100 | | 20.9 | 101 | | 20.4 | 99 | |
| 14-40 | 26.4 | | 27.0 | 102 | | 26.8 | 102 | | 26.2 | 99 | |




FIGURE 4
Growth Curves for Male and Female p53 Haploinsufficient Mice
Exposed to Acesulfame Potassium in Feed for 9 Months

Pathology and Statistical Analyses

There were no neoplasms or nonneoplastic lesions in p53 haploinsufficient mice that were attributed to exposure to acesulfame potassium. Summaries of the incidences of neoplasms and nonneoplastic lesions, individual animal tumor diagnoses, and statistical analyses of primary neoplasms that occurred with an incidence of at least 5% in at least one animal group are presented in Appendix C for males and Appendix D for females.

GENETIC TOXICOLOGY

Acesulfame potassium did not increase the frequency of micronucleated normochromatic erythrocytes in peripheral blood of male or female Tg.AC hemizygous mice administered 0.3% to 3% of the chemical in feed for 9 months (Table E1). A similar study was conducted in p53 haploinsufficient mice, and in this study, a significant exposure-related increase in the frequency of micronucleated normochromatic erythrocytes was noted in males but not females; micronucleus frequencies in the 1% and 3% male groups were significantly elevated over the control group (Table E2). There was no significant alteration in the percentage of polychromatic erythrocytes among total erythrocytes in either Tg.AC hemizygous or p53 haploinsufficient mice.

DISCUSSION AND CONCLUSIONS

Acesulfame potassium is an oxathiazinone dioxide intense sweetener developed by Hoechst and currently marketed under the commercial names Sweet One and Sunett. Acesulfame is rapidly absorbed after oral administration but undergoes no biotransformation and is rapidly eliminated in the urine as parent compound within 24 hours of ingestion (Volz *et al.*, 1991). The carcinogenicity of acesulfame potassium has been evaluated in Wistar rats (Sinkeldam *et al.*, 1991a) and Swiss mice (Beems *et al.*, 1991). Both studies yielded negative results.

In the current studies, acesulfame potassium was evaluated in Tg.AC hemizygous and p53 haploinsufficient mice as a presumptive negative control to provide additional information regarding the use of these mice as alternative models for evaluation of carcinogenic activity. Specifically, these studies were designed to examine whether extending the study duration to 39 weeks to increase the opportunity for a positive response would lead to a significant increase in background tumors in control mice and to evaluate the forestomach in Tg.AC hemizygous mice as a possible target organ for papilloma induction for studies that use the oral route of administration.

No prechronic studies of acesulfame potassium were conducted by the NTP. Prechronic studies in rats conducted by Hoechst and the 80-week study in Swiss mice were used as a basis for dose selection in the current studies. During the 80-week mouse study, mean body weights of males and females that received 30,000 ppm acesulfame potassium in the diet were reduced, and mean liver weights of all exposed groups were significantly lower than those of the controls at the end of the study (Beems *et al.*, 1991). Based on these results, doses of 0.3%, 1%, and 3% (3,000, 10,000, or 30,000 ppm) administered in feed were selected for the current 9-month studies.

In the current studies, these exposure concentrations appeared to be well tolerated by both strains of mice. Survival and mean body weights of males and females

were unaffected by exposure, and there were no exposure concentration-related clinical findings or other indications of toxicity. Therefore, it is possible that higher dietary concentrations may have been tolerated.

Cutaneous squamous cell papillomas occurred on the lip, prepuce, and pinna of exposed male Tg.AC hemizygous mice. Small epidermal squamous cell papillomas begin to occur at sites of chronic grooming (e.g., ears, nose, lips, paws, and the ano-urogenital areas) when the mice reach 20 to 26 weeks of age (Tenant *et al.*, 2001); however, the incidence is low (0% to 2%) in Tg.AC hemizygous mice and higher (up to 17%) in Tg.AC homozygous mice. The incidences in 34-week old control Tg.AC hemizygous mice was 3.7% in males and 3.8% in female (Mahler *et al.*, 1998). While historical control rates for 45-week old Tg.AC mice used in 39-week studies were not identified, they are expected to be higher; squamous cell papillomas occurred in 20% of the control males and females in the current study. Because spontaneous skin papillomas are relatively common in aging Tg.AC mice, especially in areas of grooming and/or licking, and the increase in the incidence of squamous cell papilloma occurred only in the 1% group of males with no increase in multiplicity, the slight increase was not considered related to acesulfame potassium.

The incidence of odontogenic tumors was slightly increased in Tg.AC hemizygous females that received 1% acesulfame potassium. The incidences of odontogenic tumors were not increased in any group of males or in females that received 3%. Therefore, the increase was not considered exposure related. Odontogenic tumors have occurred at incidences of 18% to 27% in control Tg.AC mice in 6-month studies (Tennant *et al.*, 2001).

Pritchard *et al.* (2003) reported as part of a global review of transgenic cancer model findings that the Tg.AC model had an overall accuracy of 74% in correctly predicting chemicals that are either listed by the International Agency for Research on Cancer, and/or the

NTP in their respective listings of chemicals as known or suspected human carcinogens. Chemicals studied in the Tg.AC line and found negative also contributed to this accuracy score if the chemicals were considered to be noncarcinogens by virtue of not appearing in these listings. The vast majority of the Tg.AC studies examined by Pritchard *et al.* (2003) used the dermal route of exposure and skin papillomas as the reporter phenotype. The pattern of incorrect assignments included both “false positives” and “false negatives.” This level of accuracy is not significantly different from that achieved by the p53 haploinsufficient model or the *Hras2* model in the Pritchard analysis. However, based on discussions at a recent NTP workshop (NTP, 2003) there is a lack of acceptance within the scientific/regulatory community that a positive result in the Tg.AC model represents a true cancer response. Therefore, the confidence in this model to identify carcinogens appears to be less than that in either the p53 haploinsufficient or *Hras2* model.

There were no increases in the incidences of neoplasms in p53 haploinsufficient mice exposed to acesulfame potassium. There was a significant increase in the frequency of micronucleated normochromatic erythrocytes at the end of the exposure period in males that received 1% or 3% acesulfame potassium. Although the increase was statistically significant, the actual magnitude of the increase was only 1 or 1.2 micronuclei/1,000 cells. Moreover, no increase was observed in females. In a review of the results of NTP prechronic and chronic studies in which peripheral blood micronuclei were evaluated in B6C3F₁ mice, Witt *et al.* (2000) considered a small increase in magnitude of response and/or a response in only one sex as a weak response and concluded that weak responses did not correlate with rodent carcinogenicity and were of uncertain biological significance. By contrast, a strong response (determined by magnitude of increase in frequency and response in males and females) strongly correlated with rodent carcinogenicity. The small increase in magnitude of response observed in male p53 haploinsufficient mice in

the current study and the lack of response in females would classify this as a weak response and, therefore, is of uncertain biological significance.

In a study that examined micronuclei formation after acute exposure, Jung *et al.* (1991) administered 0, 450, 1,500, or 4,500 mg acesulfame potassium/kg body weight in distilled water by gavage to groups of 10 NMRI mice. Mice were dosed twice during a 24-hour period and then sacrificed 6 hours after the second dose; bone marrow was collected, and smears were prepared. Examination of 2,000 polychromatic erythrocytes per mouse revealed no increase in the frequency of micronuclei. The positive control, cyclophosphamide, produced a strong response.

The review and analysis of the performance of the p53 haploinsufficient model by Pritchard *et al.* (2003) included data and conclusions as reported by the authors of studies published up to mid-2002. They compared the responses in the p53 haploinsufficient mouse model to a group of 12 known human carcinogens, 19 suspected human carcinogens, and 28 probable human noncarcinogens. In this analysis, the p53 haploinsufficient model had an overall accuracy of 81%. The high accuracy was in large part due to the lack of “false positives” (positive outcomes for probable human noncarcinogens). They concluded that the p53 haploinsufficient model correctly identified 10 of 12 (83%) known human carcinogens, 11 of 19 (58%) suspected carcinogens, and was correctly negative for 27 of 28 (96%) putative noncarcinogens under the conditions of these short-term cancer bioassays. Thus, two known and eight suspected human carcinogens were not detected by the p53 haploinsufficient model. These authors also acknowledged that a number of procedural decisions may have increased the apparent accuracy scores for the mouse models studied. These included such things as accepting a single positive study in a given model as representing a positive response even if other studies of the same chemical were negative or gave equivocal findings in that model. Clearly, the strength of the p53 haploinsufficient model is in its specificity for correctly identifying known or suspected

carcinogens, but its sensitivity to detect carcinogens is a weakness. Because this model either did not identify or only equivocally identified a significant number of known or suspected human carcinogens, negative findings with this model as shown in the current study are of uncertain value.

In summary, there was no evidence of a positive response for papilloma formation in the forestomach or

for tumors at other sites in male or female Tg.AC hemizygous mice administered acesulfame potassium in the diet at concentrations up to 3% for 9 months.

CONCLUSIONS

Under the conditions of this 9-month feed study, there was *no evidence of carcinogenic activity** of acesulfame potassium in male or female p53 haploinsufficient mice exposed to 0.3%, 1%, or 3%.

* Explanation of Levels of Evidence of Carcinogenic Activity is on page 8. A summary of the Technical Report Review Subcommittee comments and public discussion on this technical report appears on page 9.

REFERENCES

- Ashby, J., and Tennant, R.W. (1991). Definitive relationships among chemical structure, carcinogenicity and mutagenicity for 301 chemicals tested by the U.S. NTP. *Mutat. Res.* **257**, 229-306.
- Bailer, A.J., and Portier, C.J. (1988). Effects of treatment-induced mortality and tumor-induced mortality on tests for carcinogenicity in small samples. *Biometrics* **44**, 417-431.
- Beems, H.P., Til, H.P., Newman, A.J., and Mayer, D.G. (1991). Carcinogenicity Study of Acesulfame-K in Mice. In *Acesulfame-K*, pp. 59-70. Marcel Dekker, New York.
- Bieler, G.S., and Williams, R.L. (1993). Ratio estimates, the delta method, and quantal response tests for increased carcinogenicity. *Biometrics* **49**, 793-801.
- Boorman, G.A., Montgomery, C.A., Jr., Eustis, S.L., Wolfe, M.J., McConnell, E.E., and Hardisty, J.F. (1985). Quality assurance in pathology for rodent carcinogenicity studies. In *Handbook of Carcinogen Testing* (H.A. Milman and E.K. Weisburger, Eds.), pp. 345-357. Noyes Publications, Park Ridge, NJ.
- Cannon, R.E., Spalding, J.W., Trempus, C.S., Szczesniak, C.J., Virgil, K.M., Humble, M.C., and Tennant, R.W. (1997). Kinetics of wound-induced v-Ha-ras transgene expression and papilloma development in transgenic Tg.AC mice. *Mol. Carcinog.* **20**, 108-114.
- Code of Federal Regulations (CFR) **21**, Part 58.
- Cox, D.R. (1972). Regression models and life-tables. *J.R. Stat. Soc.* **B34**, 187-220.
- Crawford, B.D. (1985). Perspectives on the somatic mutation model of carcinogenesis. In *Advances in Modern Environmental Toxicology. Mechanisms and Toxicity of Chemical Carcinogens and Mutagens* (M.A. Mehlman, W.G. Flamm, and R.J. Lorentzen, Eds.), pp. 13-59. Princeton Scientific Publishing Co., Inc., Princeton, NJ.
- Donehower, L.A., Harvey, M., Slagle, B.L., McArthur, M.J., Montgomery, C.A., Jr., Butel, J.S., and Bradley, A. (1992). Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* **356**, 215-221.
- Dunnnett, C.W. (1955). A multiple comparison procedure for comparing several treatments with a control. *J. Am. Stat. Assoc.* **50**, 1096-1121.
- Harris, C.C. (1996a). Structure and function of the p53 tumor suppressor gene: Clues for rational cancer therapeutic strategies. *J. Natl. Cancer Inst.* **88**, 1442-1455.
- Harris, C.C. (1996b). p53 Tumor suppressor gene: From the basic research laboratory to the clinic—an abridged historical perspective. *Carcinogenesis* **17**, 1187-1198.
- Harris, C.C. (1996c). The 1995 Walter Hubert Lecture – molecular epidemiology of human cancer: Insights from the mutational analysis of the p53 tumour-suppressor gene. *Brit. J. Cancer* **73**, 261-269.
- Hazardous Substances Data Bank (HSDB) (2003). National Institute for Occupational Safety and Health, HSDB database available through the National Library of Medicine TOXNET System.

- Hollander, M., and Wolfe, D.A. (1973). *Nonparametric Statistical Methods*, pp. 120-123. John Wiley and Sons, New York.
- Integrated Laboratory Systems (ILS) (1990). Micronucleus Data Management and Statistical Analysis Software, Version 1.4. ILS, P.O. Box 13501, Research Triangle Park, NC 27707.
- Jung, R., Kreiling, R., and Mayer, D.G. (1991). Acesulfame-K: Studies for Genotoxic Effects. In *Acesulfame-K*, pp. 87-104. Marcel Dekker, New York.
- Kaplan, E.L., and Meier, P. (1958). Nonparametric estimation from incomplete observations. *J. Am. Stat. Assoc.* **53**, 457-481.
- Kaplan, E.L., and Meier, P. (1958). Nonparametric estimation from incomplete observations. *J. Am. Stat. Assoc.* **53**, 457-481.
- Leder, A., Kuo, A., Cardiff, R.D., Sinn, E., and Leder, P. (1990). v-Ha-ras transgene abrogates the initiation step in mouse skin tumorigenesis: Effects of phorbol esters and retinoic acid. *Proc. Natl. Acad. Sci.* **87**, 9178-9182.
- MacGregor, J.T., Wehr, C.M., Henika, P.R., and Shelby, M.D. (1990). The *in vivo* erythrocyte micronucleus test: Measurement at steady state increases assay efficiency and permits integration with toxicity studies. *Fundam. Appl. Toxicol.* **14**, 513-522.
- Mahler, J.F., Stokes, W., Mann, P.C., Takaoka, M., and Maronpot, R.R. (1996). Spontaneous lesions in aging FVB/N mice. *Toxicol. Pathol.* **24**, 710-716.
- Mahler, J.F., Flagler, N.D., Malarkey, D.E., Mann, P.C., Haseman, J.K., and Eastin, W. (1998). Spontaneous and chemically induced proliferative lesions in Tg.AC transgenic and p53-heterozygous mice. *Toxicol. Pathol.* **26**, 501-511.
- Maronpot, R.R., and Boorman, G.A. (1982). Interpretation of rodent hepatocellular proliferative alterations and hepatocellular tumors in chemical safety assessment. *Toxicol. Pathol.* **10**, 71-80.
- The Merck Index* (1996). 12th ed. (S. Budavari, Ed.), p. 8. Merck and Company Inc., Whitehouse Station, NJ.
- Miller, J.A., and Miller, E.C. (1977). Ultimate chemical carcinogens as reactive mutagenic electrophiles. In *Origins of Human Cancer* (H.H. Hiatt, J.D. Watson, and J.A. Winsten, Eds.), pp. 605-627. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Mukherjee, A., and Chakrabarti, J. (1997). *In vivo* cytogenetic studies on mice exposed to acesulfame-K — a non-nutritive sweetener. *Food Chem. Toxicol.* **35**, 1177-1179.
- Mukhopadhyay, M., Mukherjee, A., and Chakrabarti, J. (2000). *In vivo* cytogenetic studies on blends of aspartame and acesulfame-K. *Food Chem. Toxicol.* **38**, 75-77.
- National Toxicology Program (NTP) (2003). NTP Workshop. Genetically Modified Rodent Models for Cancer Hazard Identification: Selecting Substances for Study and Interpreting and Communicating Results (<http://ntp-server.niehs.nih.gov/Meetings/2003/2003FebTgWkshop.html>).
- National Toxicology Program (NTP) (2005). Toxicology Studies of Aspartame (CAS No. 22839-47-0) in FVB Tg.AC Hemizygous Mice and Carcinogenicity Studies of Aspartame in B6/129 (N5) p53 Haploinsufficient Mice (Feed Studies). Technical Report Series No. 523. NIH Publication No. 06-4459. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, Research Triangle Park, NC (in press).
- Parchem (2003). Acesulfame Potassium. Parchem database available through Parchem Trading Ltd. (http://www.parchem.com/mainf_asp?product=9&print=1)
- Piegorsch, W.W., and Bailer, A.J. (1997). *Statistics for Environmental Biology and Toxicology*, Section 6.3.2. Chapman and Hall, London.
- Portier, C.J., and Bailer, A.J. (1989). Testing for increased carcinogenicity using a survival-adjusted quantal response test. *Fundam. Appl. Toxicol.* **12**, 731-737.

- Portier, C.J., Hedges, J.C., and Hoel, D.G. (1986). Age-specific models of mortality and tumor onset for historical control animals in the National Toxicology Program's carcinogenicity experiments. *Cancer Res.* **46**, 4372-4378.
- Pritchard, J.B., French, J.E., Davis, B.J., and Haseman, J.K. (2003). The role of transgenic mouse models in carcinogen identification. *Environ. Health Perspect.* **111**, 444-454.
- Shelby, M.D. (1988). The genetic toxicity of human carcinogens and its implications. *Mutat. Res.* **204**, 3-15.
- Shelby, M.D., and Witt, K.L. (1995). Comparison of results from mouse bone marrow chromosome aberration and micronucleus tests. *Environ. Mol. Mutagen.* **25**, 302-313.
- Shelby, M.D., and Zeiger, E. (1990). Activity of human carcinogens in the Salmonella and rodent bone-marrow cytogenetics tests. *Mutat. Res.* **234**, 257-261.
- Shelby, M.D., Erexson, G.L., Hook, G.J., and Tice, R.R. (1993). Evaluation of a three-exposure mouse bone marrow micronucleus protocol: Results with 49 chemicals. *Environ. Mol. Mutagen.* **21**, 160-179.
- Sinkeldam, E.J., Kuper, C.F., Beems, R.B., Newman, A.J., and Feron, V.J. (1991a). Combined Chronic Toxicity and Carcinogenicity Study of Acesulfame-K in Rats. In *Acesulfame-K*, pp. 43-58. Marcel Dekker, New York.
- Sinkeldam, E.J., Til, H.P., Groot, A.P., Willems, M.I., Kreiling, R., and Mayer, D.G. (1991b). Toxicity Studies of Acesulfame-K: A New High Intensity Non-Caloric Sweetener. In *Acesulfame-K*, pp. 27-42. Marcel Dekker, New York.
- Spalding, J.W., Momma, J., Elwell, M.R., and Tennant, R.W. (1993). Chemically induced skin carcinogenesis in a transgenic mouse line (TG.AC) carrying a v-HA-ras gene. *Carcinogenesis* **14**, 1335-1341.
- Spalding, J.W., French, J.E., Tice, R.R., Furedi-Machacek, M., Haseman, J.K., and Tennant, R.W. (1999). Development of a transgenic mouse model for carcinogenesis bioassays: Evaluation of chemically induced skin tumors in Tg.AC mice. *Toxicol. Sci.* **49**, 241-254.
- Straus, D.S. (1981). Somatic mutation, cellular differentiation, and cancer causation. *JNCI* **67**, 233-241.
- Tarone, R.E. (1975). Tests for trend in life table analysis. *Biometrika* **62**, 679-682.
- Tennant, R.W., Margolin, B.H., Shelby, M.D., Zeiger, E., Haseman, J.K., Spalding, J., Caspary, W., Resnick, M., Stasiewicz, S., Anderson, B., and Minor, R. (1987). Prediction of chemical carcinogenicity in rodents from in vitro genetic toxicity assays. *Science* **236**, 933-941.
- Tennant, R.W., French, J.E., and Spalding, J.W. (1995). Identifying chemical carcinogens and assessing potential risk in short-term bioassays using transgenic mouse models. *Environ. Health Perspect.* **103**, 942-950.
- Tennant, R.W., Spalding, J., and French, J.E. (1996). Evaluation of transgenic mouse bioassays for identifying carcinogens and noncarcinogens. *Mutat. Res.* **365**, 119-127.
- Tennant, R.W., Stasiewicz, S., Mennear, J., French, J.E., and Spalding, J.W. (1999). Genetically altered mouse models for identifying carcinogens. *IARC Sci. Publ.* **146**, 123-150.
- Tennant, R.W., Stasiewicz, S., Eastin, W.C., Mennear, J.H., and Spalding, J.W. (2001). The Tg.AC (v-Ha-ras) transgenic mouse: Nature of the model. *Toxicol. Pathol.* **29**, 51-59.
- Tomatis, L., Huff, J., Hertz-Picciotto, I., Sandler, D.P., Bucher, J., Boffetta, P., Axelson, O., Blair, A., Taylor, J., Stayner, L., and Barrett, J.C. (1997). Avoided and avoidable risks of cancer. *Carcinogenesis* **18**, 97-105.

- Trempus, C.S., Mahler, J.F., Ananthaswamy, H.N., Loughlin, S.M., French, J.E., and Tennant, R.W. (1998). Photocarcinogenesis and susceptibility to UV radiation in the v-Ha-ras transgenic Tg.AC mouse. *J. Invest. Dermatol.* **111**, 445-451.
- Volz, M., Christ, O., Eckert, H.G., Herok, J., Kellner, M., and Rupp, W. (1991). Kinetics and Biotransformation of Acesulfame-K. In *Acesulfame-K*, pp. 7-26. Marcel Dekker, New York.
- Williams, D.A. (1971). A test for differences between treatment means when several dose levels are compared with a zero dose control. *Biometrics* **27**, 103-117.
- Williams, D.A. (1972). The comparison of several dose levels with a zero dose control. *Biometrics* **28**, 519-531.
- Witt, K.L., Knapton, A., Wehr, C.M., Hook, G.J., Mirsalis, J., Shelby, M.D., and MacGregor, J.T. (2000). Micronucleated erythrocyte frequency in peripheral blood of B6C3F₁ mice from short-term, prechronic, and chronic studies of the NTP Carcinogenesis Bioassay Program. *Environ. Mol. Mutagen.* **36**, 163-194.
- Wright, J.T., Hansen, L., Mahler, J., Szczesniak, C., and Spalding, J.W. (1995). Odontogenic tumours in the v-HA-ras (TG•AC) transgenic mouse. *Arch. Oral Biol.* **40**, 631-638.
- Zeiger, E., Haseman, J.K., Shelby, M.D., Margolin, B.H., and Tennant, R.W. (1990). Evaluation of four in vitro genetic toxicity tests for predicting rodent carcinogenicity: Confirmation of earlier results with 41 additional chemicals. *Environ. Mol. Mutagen.* **16** (Suppl. 18), 1-14.

APPENDIX A
SUMMARY OF LESIONS
IN MALE Tg.AC HEMIZYGOUS MICE
IN THE 9-MONTH FEED STUDY
OF ACESULFAME POTASSIUM

| | | |
|-----------------|--|-----------|
| TABLE A1 | Summary of the Incidence of Neoplasms in Male Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 46 |
| TABLE A2 | Individual Animal Tumor Pathology of Male Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 48 |
| TABLE A3 | Statistical Analysis of Primary Neoplasms in Male Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 54 |
| TABLE A4 | Summary of the Incidence of Nonneoplastic Lesions in Male Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 56 |

TABLE A1
Summary of the Incidence of Neoplasms in Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|----------------------------------|----------|----------|---------|---------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 2 | 4 | 2 | 1 |
| Survivors | | | | |
| Terminal sacrifice | 13 | 11 | 13 | 14 |
| Animals examined microscopically | 15 | 15 | 15 | 15 |
| Alimentary System | | | | |
| Liver | (15) | | (1) | (15) |
| Stomach, forestomach | (15) | (6) | (3) | (15) |
| Squamous cell papilloma | 5 (33%) | 4 (67%) | 2 (67%) | 6 (40%) |
| Tooth | (4) | (3) | | (2) |
| Odontogenic tumor | 4 (100%) | 3 (100%) | | 1 (50%) |
| Cardiovascular System | | | | |
| None | | | | |
| Endocrine System | | | | |
| Thyroid gland | (15) | | | (15) |
| Adenoma | 1 (7%) | | | |
| General Body System | | | | |
| None | | | | |
| Genital System | | | | |
| None | | | | |
| Hematopoietic System | | | | |
| Bone marrow | (15) | | | (15) |
| Spleen | (15) | (1) | (1) | (15) |
| Integumentary System | | | | |
| Skin | (15) | (7) | (8) | (15) |
| Squamous cell papilloma | 1 (7%) | 1 (14%) | 2 (25%) | 4 (27%) |
| Dermis, mast cell tumor benign | 1 (7%) | | | |
| Lip, squamous cell papilloma | 2 (13%) | 3 (43%) | 5 (63%) | 2 (13%) |
| Pinna, squamous cell papilloma | | | 1 (13%) | |
| Prepuce, squamous cell papilloma | 1 (7%) | | | |
| Musculoskeletal System | | | | |
| None | | | | |

TABLE A1
Summary of the Incidence of Neoplasms in Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|---|--------|--------|--------|---------|
| Nervous System | | | | |
| None | | | | |
| Respiratory System | | | | |
| Lung | (15) | (15) | (15) | (15) |
| Alveolar/bronchiolar adenoma | 1 (7%) | 1 (7%) | | 2 (13%) |
| Special Senses System | | | | |
| Harderian gland | (15) | | | (15) |
| Adenoma | | | | 1 (7%) |
| Urinary System | | | | |
| None | | | | |
| Systemic Lesions | | | | |
| Multiple organs ^b | (15) | (15) | (15) | (15) |
| Leukemia erythrocytic | | | 1 (7%) | 1 (7%) |
| Neoplasm Summary | | | | |
| Total animals with primary neoplasms ^c | 12 | 10 | 10 | 10 |
| Total primary neoplasms | 16 | 12 | 11 | 17 |
| Total animals with benign neoplasms | 9 | 8 | 9 | 9 |
| Total benign neoplasms | 12 | 9 | 10 | 15 |
| Total animals with malignant neoplasms | | | 1 | 1 |
| Total malignant neoplasms | | | 1 | 1 |
| Total animals with uncertain neoplasms- | | | | |
| benign or malignant | 4 | 3 | | 1 |
| Total uncertain neoplasms | 4 | 3 | | 1 |

^a Number of animals examined microscopically at the site and the number of animals with neoplasm

^b Number of animals with any tissue examined microscopically

^c Primary neoplasms: all neoplasms except metastatic neoplasms

TABLE A2
Individual Animal Tumor Pathology of Male Tg.AC Hemizygous Mice in the 9-Month Feed Study
of Acesulfame Potassium: 0%

| Number of Days on Study | 1 | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
|----------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 4 | 5 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| | 8 | 7 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| Carcass ID Number | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 2 | 6 | 1 | 2 | 3 | 4 | 5 | 7 | 8 | 9 | 0 | 1 | 3 | 4 | 5 | | | | |
| Hematopoietic System | | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Lymph node, mandibular | + | + | + | + | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + |
| Lymph node, mesenteric | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | M | + | + | + |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Thymus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Integumentary System | | | | | | | | | | | | | | | | | | | |
| Mammary gland | M | M | M | M | M | M | M | M | M | M | M | M | M | M | M | M | M | M | M |
| Skin | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Squamous cell papilloma | | | | X | | | | | | | | | | | | | | | |
| Dermis, mast cell tumor benign | | | | | | | | | | X | | | | | | | | | |
| Lip, squamous cell papilloma | | | | X | | | | X | | | | | | | | | | | |
| Prepuce, squamous cell papilloma | | | | | | | | | | | | | | | | X | | | |
| Musculoskeletal System | | | | | | | | | | | | | | | | | | | |
| Bone | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Skeletal muscle | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Nervous System | | | | | | | | | | | | | | | | | | | |
| Brain | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Respiratory System | | | | | | | | | | | | | | | | | | | |
| Larynx | I | + | + | + | + | + | + | + | + | + | + | + | + | + | + | M | + | + | + |
| Lung | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Alveolar/bronchiolar adenoma | | | | | | | | | | | | | | | X | | | | |
| Nose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Trachea | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Special Senses System | | | | | | | | | | | | | | | | | | | |
| Eye | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Harderian gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Zymbal's gland | M | + | + | + | + | + | + | M | M | M | M | M | M | I | + | M | | | |
| Urinary System | | | | | | | | | | | | | | | | | | | |
| Kidney | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Urinary bladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Systemic Lesions | | | | | | | | | | | | | | | | | | | |
| Multiple organs | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |

Total
Tissues/
Tumors

15
14
14
15
15

15
1
1
2
1

15
15

15

13
15
1
15
15

15
15
7

15
15

15

TABLE A3
Statistical Analysis of Primary Neoplasms in Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|---|------------|----------------|----------------|------------|
| Lung: Alveolar/bronchiolar Adenoma | | | | |
| Overall rate ^a | 1/15 (7%) | 1/15 (7%) | 0/15 (0%) | 2/15 (13%) |
| Adjusted rate ^b | 7.5% | 7.8% | 0.0% | 14.0% |
| Terminal rate ^c | 1/13 (8%) | 1/11 (9%) | 0/13 (0%) | 2/14 (14%) |
| First incidence (days) ^d | 274 (T) | 274 (T) | — ^e | 274 (T) |
| Poly-3 test | P=0.351 | P=0.752 | P=0.487N | P=0.523 |
| Skin: Squamous Cell Papilloma | | | | |
| Overall rate | 3/15 (20%) | 4/15 (27%) | 8/15 (53%) | 5/15 (33%) |
| Adjusted rate | 22.5% | 31.4% | 55.8% | 35.1% |
| Terminal rate | 3/13 (23%) | 4/11 (36%) | 7/13 (54%) | 5/14 (36%) |
| First incidence (days) | 274 (T) | 274 (T) | 264 | 274 (T) |
| Poly-3 test | P=0.429 | P=0.473 | P=0.075 | P=0.381 |
| Stomach (Forestomach): Squamous Cell Papilloma | | | | |
| Overall rate | 5/15 (33%) | 4/15 (27%) | 2/15 (13%) | 6/15 (40%) |
| Adjusted rate | 37.5% | 30.4% | 14.1% | 42.1% |
| Terminal rate | 5/13 (39%) | 3/11 (27%) | 2/13 (15%) | 6/14 (43%) |
| First incidence (days) | 274 (T) | 233 | 274 (T) | 274 (T) |
| Poly-3 test | P=0.359 | P=0.511N | P=0.164N | P=0.554 |
| Tooth: Odontogenic Tumor | | | | |
| Overall rate | 4/15 (27%) | 3/15 (20%) | 0/15 (0%) | 1/15 (7%) |
| Adjusted rate | 26.7% | 20.0% | 0.0% | 7.0% |
| Terminal rate | 2/13 (15%) | 0/11 (0%) | 0/13 (0%) | 1/14 (7%) |
| First incidence (days) | 148 | 71 | — | 274 (T) |
| Poly-3 test | P=0.118N | P=0.501N | P=0.051N | P=0.179N |
| All Organs: Erythrocytic Leukemia | | | | |
| Overall rate | 0/15 (0%) | 0/15 (0%) | 1/15 (7%) | 1/15 (7%) |
| Adjusted rate | 0.0% | 0.0% | 6.7% | 6.7% |
| Terminal rate | 0/13 (0%) | 0/11 (0%) | 0/13 (0%) | 0/14 (0%) |
| First incidence (days) | — | — ^f | 192 | 172 |
| Poly-3 test | P=0.352 | — | P=0.522 | P=0.523 |
| All Organs: Benign Neoplasms | | | | |
| Overall rate | 9/15 (60%) | 8/15 (53%) | 9/15 (60%) | 9/15 (60%) |
| Adjusted rate | 67.4% | 60.9% | 62.7% | 63.2% |
| Terminal rate | 9/13 (69%) | 7/11 (64%) | 8/13 (62%) | 9/14 (64%) |
| First incidence (days) | 274 (T) | 233 | 264 | 274 (T) |
| Poly-3 test | P=0.561N | P=0.521N | P=0.555N | P=0.564N |
| All Organs: Malignant Neoplasms | | | | |
| Overall rate | 0/15 (0%) | 0/15 (0%) | 1/15 (7%) | 1/15 (7%) |
| Adjusted rate | 0.0% | 0.0% | 6.7% | 6.7% |
| Terminal rate | 0/13 (0%) | 0/11 (0%) | 0/13 (0%) | 0/14 (0%) |
| First incidence (days) | — | — | 192 | 172 |
| Poly-3 test | P=0.352 | — | P=0.522 | P=0.523 |

TABLE A3
Statistical Analysis of Primary Neoplasms in Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|-------------|-------------|-------------|-------------|
| All Organs: Benign or Malignant Neoplasms | | | | |
| Overall rate | 12/15 (80%) | 10/15 (67%) | 10/15 (67%) | 10/15 (67%) |
| Adjusted rate | 80.0% | 66.8% | 66.7% | 66.7% |
| Terminal rate | 10/13 (77%) | 7/11 (64%) | 8/13 (62%) | 9/14 (64%) |
| First incidence (days) | 148 | 71 | 192 | 172 |
| Poly-3 test | P=0.392N | P=0.346N | P=0.343N | P=0.343N |

(T) Terminal sacrifice

^a Number of neoplasm-bearing animals/number of animals examined. Denominator is number of animals examined microscopically for lung; for other tissues, denominator is number of animals necropsied.

^b Poly-3 estimated neoplasm incidence after adjustment for intercurrent mortality

^c Observed incidence at terminal kill

^d Beneath the control incidence is the P value associated with the trend test. Beneath the exposed group incidence are the P values corresponding to pairwise comparisons between the controls and that exposed group. The Poly-3 test accounts for the differential mortality in animals that do not reach terminal sacrifice. A negative trend or a lower incidence in an exposed group is indicated by N.

^e Not applicable; no neoplasms in animal group

^f Value of statistic cannot be computed

TABLE A4
Summary of the Incidence of Nonneoplastic Lesions in Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|--|----------|------|----------|----------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 2 | 4 | 2 | 1 |
| Survivors | | | | |
| Terminal sacrifice | 13 | 11 | 13 | 14 |
| Animals examined microscopically | 15 | 15 | 15 | 15 |
| Alimentary System | | | | |
| Liver | (15) | | (1) | (15) |
| Hematopoietic cell proliferation | 1 (7%) | | | |
| Inflammation, chronic | 1 (7%) | | | |
| Hepatocyte, necrosis, focal | 1 (7%) | | | |
| Mesentery | (2) | | | (2) |
| Fat, necrosis, focal | 2 (100%) | | | 2 (100%) |
| Pancreas | (15) | | | (15) |
| Inflammation, chronic active | 1 (7%) | | | |
| Cardiovascular System | | | | |
| None | | | | |
| Endocrine System | | | | |
| Adrenal cortex | (14) | | | (15) |
| Atrophy | 10 (71%) | | | 11 (73%) |
| Hypertrophy, focal | 9 (64%) | | | 5 (33%) |
| General Body System | | | | |
| None | | | | |
| Genital System | | | | |
| Preputial gland | (15) | | (1) | (15) |
| Cyst | 1 (7%) | | 1 (100%) | |
| Testes | (15) | | | (15) |
| Germinal epithelium, degeneration, focal | 1 (7%) | | | 1 (7%) |

^a Number of animals examined microscopically at the site and the number of animals with lesion

TABLE A4
Summary of the Incidence of Nonneoplastic Lesions in Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|-----------|----------|------|----------|
| Hematopoietic System | | | | |
| Bone marrow | (15) | | | (15) |
| Hyperplasia | | | | 1 (7%) |
| Lymph node, mandibular | (14) | | | (15) |
| Hyperplasia | 2 (14%) | | | 2 (13%) |
| Spleen | (15) | (1) | (1) | (15) |
| Atrophy | 1 (7%) | | | |
| Hematopoietic cell proliferation | 10 (67%) | 1 (100%) | | 7 (47%) |
| Thymus | (15) | | | (14) |
| Atrophy, diffuse | 3 (20%) | | | 2 (14%) |
| Atrophy, focal | 2 (13%) | | | |
| Integumentary System | | | | |
| Skin | (15) | (7) | (8) | (15) |
| Sebaceous gland, hyperplasia, focal | 1 (7%) | | | |
| Subcutaneous tissue, pinna, inflammation, acute, focal | | 1 (14%) | | |
| Musculoskeletal System | | | | |
| None | | | | |
| Nervous System | | | | |
| None | | | | |
| Respiratory System | | | | |
| Lung | (15) | (15) | (15) | (15) |
| Alveolar epithelium, alveolus, inflammation, chronic active, focal | | | | 1 (7%) |
| Nose | (15) | | | (15) |
| Glands, hyperplasia | 1 (7%) | | | |
| Special Senses System | | | | |
| Eye | (15) | | | (15) |
| Retina, atrophy | 15 (100%) | | | 14 (93%) |
| Urinary System | | | | |
| Kidney | (15) | (1) | | (15) |
| Infiltration cellular, plasma cell | | 1 (100%) | | |
| Renal tubule, dilatation, diffuse | | | | 1 (7%) |
| Renal tubule, dilatation, focal | | | | 2 (13%) |
| Urinary bladder | (15) | | | (15) |
| Edema | 1 (7%) | | | |

APPENDIX B
SUMMARY OF LESIONS
IN FEMALE Tg.AC HEMIZYGOUS MICE
IN THE 9-MONTH FEED STUDY
OF ACESULFAME POTASSIUM

| | | |
|-----------------|--|-----------|
| TABLE B1 | Summary of the Incidence of Neoplasms in Female Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 60 |
| TABLE B2 | Individual Animal Tumor Pathology of Female Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 62 |
| TABLE B3 | Statistical Analysis of Primary Neoplasms in Female Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 69 |
| TABLE B4 | Summary of the Incidence of Nonneoplastic Lesions in Female Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 71 |

TABLE B1
Summary of the Incidence of Neoplasms in Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|--|----------|---------|----------|---------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 1 | | 2 | 2 |
| Natural deaths | 2 | 3 | 1 | 1 |
| Survivors | | | | |
| Terminal sacrifice | 12 | 12 | 12 | 12 |
| Animals examined microscopically | 15 | 14 | 14 | 15 |
| Alimentary System | | | | |
| Liver | (15) | (2) | | (15) |
| Stomach, forestomach | (14) | (9) | (10) | (14) |
| Squamous cell papilloma | 7 (50%) | 6 (67%) | 9 (90%) | 8 (57%) |
| Squamous cell papilloma, multiple | 2 (14%) | 1 (11%) | | |
| Tongue | (15) | (1) | | (15) |
| Squamous cell carcinoma, metastatic, uncertain primary site | 1 (7%) | | | |
| Tooth | (2) | (1) | (6) | (4) |
| Odontogenic tumor | 2 (100%) | | 6 (100%) | 3 (75%) |
| Cardiovascular System | | | | |
| None | | | | |
| Endocrine System | | | | |
| Pituitary gland | (12) | | | (14) |
| General Body System | | | | |
| None | | | | |
| Genital System | | | | |
| Uterus | (13) | (2) | (1) | (14) |
| Sarcoma stromal | | | | 1 (7%) |
| Hematopoietic System | | | | |
| Spleen | (15) | | (3) | (15) |
| Integumentary System | | | | |
| Skin | (15) | (5) | (6) | (15) |
| Squamous cell papilloma | 1 (7%) | 2 (40%) | 3 (50%) | 1 (7%) |
| Lip, squamous cell papilloma | | | 2 (33%) | |
| Vulva, squamous cell papilloma | 2 (13%) | 2 (40%) | 2 (33%) | 3 (20%) |

TABLE B1
Summary of the Incidence of Neoplasms in Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|---------|--------|------|--------|
| Musculoskeletal System | | | | |
| None | | | | |
| Nervous System | | | | |
| None | | | | |
| Respiratory System | | | | |
| Lung | (15) | | | (15) |
| Alveolar/bronchiolar adenoma | 1 (7%) | | | 1 (7%) |
| Special Senses System | | | | |
| Harderian gland | (14) | (1) | | (15) |
| Urinary System | | | | |
| Kidney | (15) | (1) | | (15) |
| Urinary bladder | (13) | | | (15) |
| Transitional epithelium, papilloma | | | | 1 (7%) |
| Systemic Lesions | | | | |
| Multiple organs | (15) | (14) | (14) | (15) |
| Leukemia erythrocytic | 2 (13%) | 1 (7%) | | 1 (7%) |
| Neoplasm Summary | | | | |
| Total animals with primary neoplasms ^c | 12 | 10 | 13 | 13 |
| Total primary neoplasms | 17 | 12 | 22 | 19 |
| Total animals with benign neoplasms | 11 | 9 | 10 | 10 |
| Total benign neoplasms | 13 | 11 | 16 | 14 |
| Total animals with malignant neoplasms | 2 | 1 | | 2 |
| Total malignant neoplasms | 2 | 1 | | 2 |
| Total animals with metastatic neoplasms | 1 | | | |
| Total metastatic neoplasms | 1 | | | |
| Total animals with malignant neoplasms of uncertain primary site | 1 | | | |
| Total animals with uncertain neoplasms-benign or malignant | 2 | | 6 | 3 |
| Total uncertain neoplasms | 2 | | 6 | 3 |

^a Number of animals examined microscopically at the site and the number of animals with neoplasm

^b Number of animals with any tissue examined microscopically

^c Primary neoplasms: all neoplasms except metastatic neoplasms

TABLE B2
Individual Animal Tumor Pathology of Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium: 0%

| Number of Days on Study | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|----------------------|
| | 2 | 3 | 4 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| | 4 | 6 | 0 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| Carcass ID Number | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 7 | 7 | 7 | 7 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 9 |
| | 9 | 7 | 6 | 8 | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 0 | | | | |
| | | | | | | | | | | | | | | | | | | | Total Tissues/Tumors |
| Alimentary System | | | | | | | | | | | | | | | | | | | |
| Esophagus | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| Gallbladder | A | A | + | + | + | + | + | + | + | + | + | + | + | + | M | + | + | + | 12 |
| Intestine large, colon | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Intestine large, rectum | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Intestine large, cecum | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Intestine small, duodenum | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Intestine small, jejunum | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Intestine small, ileum | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Mesentery | | | | | | + | | | | + | | | | | | | | | 2 |
| Pancreas | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| Salivary glands | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| Stomach, forestomach | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| Squamous cell papilloma | | | | X | | | X | | X | X | X | | X | X | | | | | 7 |
| Squamous cell papilloma, multiple | X | | | | | | | | | | | | | | | | | X | 2 |
| Stomach, glandular | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Tongue | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Squamous cell carcinoma, metastatic, uncertain primary site | | | | X | | | | | | | | | | | | | | | 1 |
| Tooth | | + | | | | | | | | + | | | | | | | | | 2 |
| Odontogenic tumor | | X | | | | | | | | X | | | | | | | | | 2 |
| Cardiovascular System | | | | | | | | | | | | | | | | | | | |
| Blood vessel | + | + | + | + | + | M | + | + | + | + | + | + | + | + | M | M | + | + | 12 |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Endocrine System | | | | | | | | | | | | | | | | | | | |
| Adrenal cortex | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| Adrenal medulla | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| Islets, pancreatic | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| Parathyroid gland | M | A | + | M | M | + | + | M | + | + | M | M | M | M | M | | | | 5 |
| Pituitary gland | + | I | + | + | + | + | + | I | I | + | + | + | + | + | + | + | + | + | 12 |
| Thyroid gland | + | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 14 |
| General Body System | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | |
| Genital System | | | | | | | | | | | | | | | | | | | |
| Clitoral gland | M | A | + | + | + | + | + | M | I | + | + | M | + | M | + | | | | 9 |
| Ovary | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Uterus | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |
| Vagina | A | A | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 13 |

+: Tissue examined microscopically
A: Autolysis precludes examination

M: Missing tissue
I: Insufficient tissue

X: Lesion present
Blank: Not examined

TABLE B2
Individual Animal Tumor Pathology of Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium: 0.3%

| Number of Days on Study | 0 | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
|-----------------------------------|-----|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|-----------------------------|
| Carcass ID Number | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 0 | 9 | 0 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 9 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 3 | 9 | 1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 0 | 2 | 4 | 5 | | | |
| | | | | | | | | | | | | | | | | | | Total Tissues/ Tumors |
| Alimentary System | | | | | | | | | | | | | | | | | | |
| Esophagus | + | | | | | | | | | | | | | | | | | 1 |
| Gallbladder | A | | | | | | | | | | | | | | | | | |
| Intestine large, colon | A | | | | | | | | | | | | | | | | | |
| Intestine large, rectum | A | | | | | | | | | | | | | | | | | |
| Intestine large, cecum | A | | | | | | | | | | | | | | | | | |
| Intestine small, duodenum | A | | | | | | | | | | | | | | | | | |
| Intestine small, jejunum | A | | | | | | | | | | | | | | | | | |
| Intestine small, ileum | A | | | | | | | | | | | | | | | | | |
| Liver | + + | | | | | | | | | | | | | | | | | 2 |
| Pancreas | + | | | | | | | | | | | | | | | | | 1 |
| Salivary glands | + | | | | | | | | | | | | | | | | | 1 |
| Stomach, forestomach | A | A | | + | + | | | + | + | + | + | + | + | + | + | | | + |
| Squamous cell papilloma | | | | X | X | | | X | X | | | | | X | | | | X |
| Squamous cell papilloma, multiple | | | | | | | | | | | | | | X | | | | |
| Stomach, glandular | A | | | | | | | | | | | | | | | | | |
| Tongue | + | | | | | | | | | | | | | | | | | 1 |
| Tooth | | | | + | | | | | | | | | | | | | | |
| Cardiovascular System | | | | | | | | | | | | | | | | | | |
| Blood vessel | + | | | | | | | | | | | | | | | | | 1 |
| Heart | A | | | | | | | | | | | | | | | | | |
| Endocrine System | | | | | | | | | | | | | | | | | | |
| Adrenal cortex | + | | | | | | | | | | | | | | | | | 1 |
| Adrenal medulla | + | | | | | | | | | | | | | | | | | 1 |
| Islets, pancreatic | + | | | | | | | | | | | | | | | | | 1 |
| Parathyroid gland | M | | | | | | | | | | | | | | | | | |
| Pituitary gland | M | | | | | | | | | | | | | | | | | |
| Thyroid gland | A | | | | | | | | | | | | | | | | | |
| General Body System | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | |
| Genital System | | | | | | | | | | | | | | | | | | |
| Clitoral gland | + | | | | | | | | | | | | | | | | | 1 |
| Ovary | A | | A | | | | | | | | | | | | | | | + |
| Uterus | A | | | | + | | + | | | | | | | | | | | |
| Vagina | A | | | | | | | | | | | | | | | | | 2 |
| Hematopoietic System | | | | | | | | | | | | | | | | | | |
| Bone marrow | + | | | | | | | | | | | | | | | | | 1 |
| Lymph node, mandibular | + | | | | | | | | | | | | | | | | | 1 |
| Lymph node, mesenteric | A | | | | | | | | | | | | | | | | | |
| Spleen | A | | | | | | | | | | | | | | | | | |
| Thymus | + | | | | | | | | | | | | | | | | | 1 |

TABLE B2
Individual Animal Tumor Pathology of Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium: 1%

| Number of Days on Study | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | | |
|--------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|-----------------------------|----|
| Carcass ID Number | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | Total Tissues/ Tumors | |
| Alimentary System | | | | | | | | | | | | | | | | | | | | |
| Mesentery | | | | | | | | | | | | | | | | | | | | 1 |
| Stomach, forestomach | | | | | | | | | | | | | | | | | | | | 10 |
| Squamous cell papilloma | | | | | | | | | | | | | | | | | | | | 9 |
| Tooth | | | | | | | | | | | | | | | | | | | | 6 |
| Odontogenic tumor | | | | | | | | | | | | | | | | | | | | 6 |
| Cardiovascular System | | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | | |
| Endocrine System | | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | | |
| General Body System | | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | | |
| Genital System | | | | | | | | | | | | | | | | | | | | |
| Uterus | | | | | | | | | | | | | | | | | | | | 1 |
| Hematopoietic System | | | | | | | | | | | | | | | | | | | | |
| Spleen | | | | | | | | | | | | | | | | | | | | 3 |
| Integumentary System | | | | | | | | | | | | | | | | | | | | |
| Skin | | | | | | | | | | | | | | | | | | | | 6 |
| Squamous cell papilloma | | | | | | | | | | | | | | | | | | | | 3 |
| Lip, squamous cell papilloma | | | | | | | | | | | | | | | | | | | | 2 |
| Vulva, squamous cell papilloma | | | | | | | | | | | | | | | | | | | | 2 |
| Musculoskeletal System | | | | | | | | | | | | | | | | | | | | |
| Bone | | | | | | | | | | | | | | | | | | | | 1 |
| Nervous System | | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | | |
| Respiratory System | | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | | |
| Special Senses System | | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | | |
| Urinary System | | | | | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | | | | | |
| Systemic Lesions | | | | | | | | | | | | | | | | | | | | |
| Multiple organs | | | | | | | | | | | | | | | | | | | | 14 |

TABLE B2
Individual Animal Tumor Pathology of Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium: 3%

| | | |
|------------------------------------|---------------------------------|--------------------------|
| Number of Days on Study | 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | |
| | 5 3 5 7 7 7 7 7 7 7 7 7 7 7 7 | |
| | 6 2 5 4 4 4 4 4 4 4 4 4 4 4 4 | |
| Carcass ID Number | 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | Total Tissues/ Tumors |
| | 2 2 2 2 2 2 2 2 2 3 3 3 3 3 3 | |
| | 4 1 9 2 3 5 6 7 8 0 1 2 3 4 5 | |
| Integumentary System | | |
| Mammary gland | + + + + + + + + + + + + + + + | 15 |
| Skin | + + + + + + + + + + + + + + + | 15 |
| Squamous cell papilloma | | 1 |
| Vulva, squamous cell papilloma | | 3 |
| Musculoskeletal System | | |
| Bone | + + + + + + + + + + + + + + + | 15 |
| Skeletal muscle | + + + + + + + + + + + + + + + | 15 |
| Nervous System | | |
| Brain | + + + + + + + + + + + + + + + | 15 |
| Respiratory System | | |
| Larynx | + I + I + I + + + I + + + + + | 11 |
| Lung | + + + + + + + + + + + + + + + | 15 |
| Alveolar/bronchiolar adenoma | | 1 |
| Nose | + + + + + + + + + + + + + + + | 15 |
| Trachea | + + + + + + + + + + + + + + + | 15 |
| Special Senses System | | |
| Ear | | 2 |
| Eye | + + A + + + + + + + + + + + + + | 14 |
| Harderian gland | + + + + + + + + + + + + + + + | 15 |
| Zymbal's gland | + + + M + + + + M M I M + + + | 10 |
| Urinary System | | |
| Kidney | + + + + + + + + + + + + + + + | 15 |
| Urinary bladder | + + + + + + + + + + + + + + + | 15 |
| Transitional epithelium, papilloma | | 1 |
| Systemic Lesions | | |
| Multiple organs | + + + + + + + + + + + + + + + | 15 |
| Leukemia erythrocytic | | 1 |

TABLE B3
Statistical Analysis of Primary Neoplasms in Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|---|-------------|----------------|-------------|-------------|
| Lung: Alveolar/bronchiolar Adenoma | | | | |
| Overall rate ^a | 1/15 (7%) | 0/0 | 0/0 | 1/15 (7%) |
| Adjusted rate ^b | 7.2% | 0.0% | 0.0% | 7.4% |
| Terminal rate ^c | 1/12 (8%) | 0/0 | 0/0 | 1/12 (8%) |
| First incidence (days) ^d | 274 (T) | — ^e | — | 274 (T) |
| Poly-3 test ^f | — | — | — | P=0.757 |
| Skin: Squamous Cell Papilloma | | | | |
| Overall rate | 3/15 (20%) | 4/15 (27%) | 6/15 (40%) | 4/15 (27%) |
| Adjusted rate | 21.1% | 30.3% | 42.9% | 29.4% |
| Terminal rate | 2/12 (17%) | 4/12 (33%) | 6/12 (50%) | 4/12 (33%) |
| First incidence (days) | 236 | 274 (T) | 274 (T) | 274 (T) |
| Poly-3 test | P=0.489 | P=0.456 | P=0.201 | P=0.473 |
| Stomach (Forestomach): Squamous Cell Papilloma | | | | |
| Overall rate | 9/15 (60%) | 7/15 (47%) | 9/15 (60%) | 8/15 (53%) |
| Adjusted rate | 62.9% | 53.0% | 62.4% | 58.8% |
| Terminal rate | 8/12 (67%) | 7/12 (58%) | 7/12 (58%) | 8/12 (67%) |
| First incidence (days) | 224 | 274 (T) | 232 | 274 (T) |
| Poly-3 test | P=0.581N | P=0.446N | P=0.637N | P=0.567N |
| Tooth: Odontogenic Tumor | | | | |
| Overall rate | 2/15 (13%) | 0/15 (0%) | 6/15 (40%) | 3/15 (20%) |
| Adjusted rate | 14.1% | 0.0% | 40.1% | 20.3% |
| Terminal rate | 1/12 (8%) | 0/12 (0%) | 4/12 (33%) | 1/12 (8%) |
| First incidence (days) | 236 | — | 207 | 156 |
| Poly-3 test | P=0.319 | P=0.248N | P=0.120 | P=0.519 |
| All Organs: Erythrocytic Leukemia | | | | |
| Overall rate | 2/15 (13%) | 1/15 (7%) | 0/15 (0%) | 1/15 (7%) |
| Adjusted rate | 13.6% | 7.2% | 0.0% | 7.3% |
| Terminal rate | 0/12 (0%) | 0/12 (0%) | 0/12 (0%) | 0/12 (0%) |
| First incidence (days) | 224 | 196 | — | 255 |
| Poly-3 test | P=0.501N | P=0.521N | P=0.242N | P=0.522N |
| All Organs: Benign Neoplasms | | | | |
| Overall rate | 11/15 (73%) | 9/15 (60%) | 10/15 (67%) | 10/15 (67%) |
| Adjusted rate | 75.0% | 68.2% | 69.3% | 73.5% |
| Terminal rate | 9/12 (75%) | 9/12 (75%) | 8/12 (67%) | 10/12 (83%) |
| First incidence (days) | 224 | 274 (T) | 232 | 274 (T) |
| Poly-3 test | P=0.569 | P=0.509N | P=0.528N | P=0.635N |
| All Organs: Malignant Neoplasms | | | | |
| Overall rate | 3/15 (20%) | 1/15 (7%) | 0/15 (0%) | 2/15 (13%) |
| Adjusted rate | 20.0% | 7.2% | 0.0% | 14.5% |
| Terminal rate | 0/12 (0%) | 0/12 (0%) | 0/12 (0%) | 1/12 (8%) |
| First incidence (days) | 224 | 196 | — | 255 |
| Poly-3 test | P=0.623N | P=0.329N | P=0.119N | P=0.540N |

TABLE B3
Statistical Analysis of Primary Neoplasms in Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|-------------|-------------|-------------|-------------|
| All Organs: Benign or Malignant Neoplasms | | | | |
| Overall rate | 13/15 (87%) | 10/15 (67%) | 13/15 (87%) | 13/15 (87%) |
| Adjusted rate | 86.7% | 72.3% | 86.7% | 86.7% |
| Terminal rate | 10/12 (83%) | 9/12 (75%) | 10/12 (83%) | 10/12 (83%) |
| First incidence (days) | 224 | 196 | 207 | 156 |
| Poly-3 test | P=0.443 | P=0.311N | P=0.698 | P=0.698 |

(T) Terminal sacrifice

^a Number of neoplasm-bearing animals/number of animals examined. Denominator is number of animals examined microscopically for lung; for other tissues, denominator is number of animals necropsied.

^b Poly-3 estimated neoplasm incidence after adjustment for intercurrent mortality

^c Observed incidence at terminal kill

^d Beneath the control incidence is the P value associated with the trend test. Beneath the exposed group incidence are the P values corresponding to pairwise comparisons between the controls and that exposed group. The Poly-3 test accounts for the differential mortality in animals that do not reach terminal sacrifice. A negative trend or a lower incidence in an exposed group is indicated by N.

^e Not applicable; no neoplasms in animal group

^f Value of statistic cannot be computed

TABLE B4
Summary of the Incidence of Nonneoplastic Lesions in Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|---|----------|----------|----------|----------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 1 | | 2 | 2 |
| Natural deaths | 2 | 3 | 1 | 1 |
| Survivors | | | | |
| Terminal sacrifice | 12 | 12 | 12 | 12 |
| Animals examined microscopically | 15 | 14 | 14 | 15 |
| Alimentary System | | | | |
| Liver | (15) | (2) | | (15) |
| Inflammation, chronic | 2 (13%) | | | 3 (20%) |
| Necrosis, focal | 1 (7%) | | | |
| Tension lipidosis, focal | | | | 1 (7%) |
| Hepatocyte, inflammation, chronic | 1 (7%) | | | |
| Hepatocyte, necrosis, focal | 2 (13%) | | | 3 (20%) |
| Mesentery | (2) | | (1) | |
| Fat, inflammation, acute, focal | | | 1 (100%) | |
| Fat, necrosis, focal | 2 (100%) | | | |
| Tongue | (15) | (1) | | (15) |
| Ulcer | 1 (7%) | | | |
| Cardiovascular System | | | | |
| None | | | | |
| Endocrine System | | | | |
| Adrenal cortex | (14) | (1) | | (15) |
| Subcapsular, hyperplasia, focal | 9 (64%) | | | 6 (40%) |
| Zona reticularis, vacuolization cytoplasmic | 9 (64%) | | | 10 (67%) |
| General Body System | | | | |
| None | | | | |
| Genital System | | | | |
| Uterus | (13) | (2) | (1) | (14) |
| Endometrium, hyperplasia, cystic | 7 (54%) | 2 (100%) | 1 (100%) | 10 (71%) |
| Vagina | (13) | | | (15) |
| Hyperplasia | 1 (8%) | | | |

^a Number of animals examined microscopically at the site and the number of animals with lesion

TABLE B4
Summary of the Incidence of Nonneoplastic Lesions in Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|-----------|---------|----------|-----------|
| Hematopoietic System | | | | |
| Lymph node, mandibular | (12) | (1) | | (15) |
| Hyperplasia | 1 (8%) | | | 3 (20%) |
| Spleen | (15) | | (3) | (15) |
| Hematopoietic cell proliferation | 12 (80%) | | 3 (100%) | 13 (87%) |
| Pigmentation | 3 (20%) | | | 2 (13%) |
| Thymus | (11) | (1) | | (14) |
| Atrophy, diffuse | 1 (9%) | | | 1 (7%) |
| Integumentary System | | | | |
| None | | | | |
| Musculoskeletal System | | | | |
| Bone | (14) | (2) | (1) | (15) |
| Mandible, inflammation, chronic active | | 1 (50%) | | 1 (7%) |
| Nervous System | | | | |
| None | | | | |
| Respiratory System | | | | |
| Lung | (15) | | | (15) |
| Hemorrhage, focal | | | | 1 (7%) |
| Special Senses System | | | | |
| Eye | (13) | | | (14) |
| Retina, atrophy | 13 (100%) | | | 14 (100%) |
| Harderian gland | (14) | (1) | | (15) |
| Infiltration cellular, focal, lymphocyte | 1 (7%) | | | |
| Urinary System | | | | |
| Kidney | (15) | (1) | | (15) |
| Renal tubule, dilatation, focal | | | | 1 (7%) |

APPENDIX C
SUMMARY OF LESIONS
IN MALE p53 HAPLOINSUFFICIENT MICE
IN THE 9-MONTH FEED STUDY
OF ACESULFAME POTASSIUM

| | | |
|-----------------|---|-----------|
| TABLE C1 | Summary of the Incidence of Neoplasms in Male p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 74 |
| TABLE C2 | Individual Animal Tumor Pathology of Male p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 76 |
| TABLE C3 | Statistical Analysis of Primary Neoplasms in Male p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 81 |
| TABLE C4 | Summary of the Incidence of Nonneoplastic Lesions in Male p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 82 |

TABLE C1
Summary of the Incidence of Neoplasms in Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|-----------------------------------|------|------|----------|--------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 1 | | | 1 |
| Survivors | | | | |
| Terminal sacrifice | 14 | 15 | 15 | 14 |
| Animals examined microscopically | 15 | | 1 | 15 |
| Alimentary System | | | | |
| None | | | | |
| Cardiovascular System | | | | |
| None | | | | |
| Endocrine System | | | | |
| None | | | | |
| General Body System | | | | |
| None | | | | |
| Genital System | | | | |
| None | | | | |
| Hematopoietic System | | | | |
| None | | | | |
| Integumentary System | | | | |
| Skin | (15) | | (1) | (15) |
| Pinna, hemangiosarcoma | | | 1 (100%) | |
| Subcutaneous tissue, fibrosarcoma | | | | 1 (7%) |
| Musculoskeletal System | | | | |
| None | | | | |
| Nervous System | | | | |
| None | | | | |
| Respiratory System | | | | |
| None | | | | |

TABLE C1
Summary of the Incidence of Neoplasms in Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|---|----|------|----|----|
| Special Senses System | | | | |
| None | | | | |
| Urinary System | | | | |
| None | | | | |
| Neoplasm Summary | | | | |
| Total animals with primary neoplasms ^b | | | 1 | 1 |
| Total primary neoplasms | | | 1 | 1 |
| Total animals with malignant neoplasms | | | 1 | 1 |
| Total malignant neoplasms | | | 1 | 1 |

^a Number of animals examined microscopically at the site and the number of animals with neoplasm

^b Primary neoplasms: all neoplasms except metastatic neoplasms

TABLE C2
Individual Animal Tumor Pathology of Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium: 0%

| Number of Days on Study | 1 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 2 |
|------------------------------|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|----------------------|
| | 9 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| | 0 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| Carcass ID Number | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 6 | 6 | 6 | 6 | 6 | 6 | 6 |
| | 8 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 9 | 0 | 1 | 2 | 3 | 4 | 5 | |
| | | | | | | | | | | | | | | | | Total Tissues/Tumors |
| Alimentary System | | | | | | | | | | | | | | | | |
| Esophagus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Gallbladder | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Intestine large, colon | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Intestine large, rectum | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Intestine large, cecum | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Intestine small, duodenum | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Intestine small, jejunum | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Intestine small, ileum | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Liver | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Mesentery | | | | | | | | | + | | | | | | | 1 |
| Pancreas | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Salivary glands | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Stomach, forestomach | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Stomach, glandular | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Tongue | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Cardiovascular System | | | | | | | | | | | | | | | | |
| Blood vessel | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Heart | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Endocrine System | | | | | | | | | | | | | | | | |
| Adrenal cortex | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Adrenal medulla | + | + | + | + | + | + | + | + | + | M | + | + | + | + | + | 14 |
| Islets, pancreatic | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Parathyroid gland | + | + | + | + | M | M | + | + | M | M | M | M | M | M | M | 7 |
| Pituitary gland | + | I | + | + | I | + | + | + | + | + | I | + | I | I | I | 9 |
| Thyroid gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| General Body System | | | | | | | | | | | | | | | | |
| None | | | | | | | | | | | | | | | | |
| Genital System | | | | | | | | | | | | | | | | |
| Epididymis | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Preputial gland | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Prostate | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Seminal vesicle | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Testes | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Hematopoietic System | | | | | | | | | | | | | | | | |
| Bone marrow | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Lymph node, mandibular | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Lymph node, mesenteric | + | + | + | + | M | + | + | + | + | + | + | + | + | + | + | 14 |
| Lymph node, mediastinal | | | | | | | | | + | | | | | | | 1 |
| Spleen | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |
| Thymus | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | 15 |

+: Tissue examined microscopically
 A: Autolysis precludes examination

M: Missing tissue
 I: Insufficient tissue

X: Lesion present
 Blank: Not examined

TABLE C2
Individual Animal Tumor Pathology of Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium: 1%

| | | | |
|--------------------------------|-------------------------------|---|----------|
| Number of Days on Study | 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | | |
| | 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 | | |
| | 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 | | |
| Carcass ID Number | 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | | Total |
| | 8 8 8 8 8 8 8 8 8 8 9 9 9 9 9 | | Tissues/ |
| | 1 2 3 4 5 6 7 8 9 0 1 2 3 4 5 | | Tumors |
| Alimentary System | | | |
| None | | | |
| Cardiovascular System | | | |
| None | | | |
| Endocrine System | | | |
| None | | | |
| General Body System | | | |
| None | | | |
| Genital System | | | |
| None | | | |
| Hematopoietic System | | | |
| None | | | |
| Integumentary System | | | |
| Skin | | + | 1 |
| Pinna, hemangiosarcoma | | X | 1 |
| Musculoskeletal System | | | |
| None | | | |
| Nervous System | | | |
| None | | | |
| Respiratory System | | | |
| None | | | |
| Special Senses System | | | |
| None | | | |
| Urinary System | | | |
| None | | | |
| Systemic Lesions | | | |
| Multiple organs | | + | 1 |

TABLE C3
Statistical Analysis of Primary Neoplasms in Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|----------------|----------------|-----------|-----------|
| All Organs: Malignant Neoplasms | | | | |
| Overall rate ^a | 0/15 (0%) | 0/15 (0%) | 1/15 (7%) | 1/15 (7%) |
| Adjusted rate ^b | 0.0% | 0.0% | 6.7% | 6.7% |
| Terminal rate ^c | 0/14 (0%) | 0/15 (0%) | 1/15 (7%) | 0/14 (0%) |
| First incidence (days) | — ^e | — | 274 (T) | 225 |
| Poly-3 test ^d | P=0.312 | — ^f | P=0.509 | P=0.509 |
| All Organs: Benign or Malignant Neoplasms | | | | |
| Overall rate | 0/15 (0%) | 0/15 (0%) | 1/15 (7%) | 1/15 (7%) |
| Adjusted rate | 0.0% | 0.0% | 6.7% | 6.7% |
| Terminal rate | 0/14 (0%) | 0/15 (0%) | 1/15 (7%) | 0/14 (0%) |
| First incidence (days) | — | — | 274 (T) | 225 |
| Poly-3 test | P=0.312 | — | P=0.509 | P=0.509 |

(T) Terminal sacrifice

^a Number of neoplasm-bearing animals/number of animals necropsied.

^b Poly-3 estimated neoplasm incidence after adjustment for intercurrent mortality

^c Observed incidence at terminal kill

^d Beneath the control incidence is the P value associated with the trend test. Beneath the exposed group incidence are the P values corresponding to pairwise comparisons between the controls and that exposed group. The Poly-3 test accounts for the differential mortality in animals that do not reach terminal sacrifice.

^e Not applicable; no neoplasms in animal group

^f Value of statistic cannot be computed

TABLE C4
Summary of the Incidence of Nonneoplastic Lesions in Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|--|----------|------|----|----------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 1 | | | 1 |
| Survivors | | | | |
| Terminal sacrifice | 14 | 15 | 15 | 14 |
| Animals examined microscopically | 15 | | 1 | 15 |
| Alimentary System | | | | |
| Liver | (15) | | | (15) |
| Inflammation, chronic active | 1 (7%) | | | 5 (33%) |
| Hepatocyte, inflammation, chronic active | 1 (7%) | | | |
| Hepatocyte, necrosis, focal | 1 (7%) | | | |
| Hepatocyte, centrilobular, hypertrophy | | | | 1 (7%) |
| Mesentery | (1) | | | (1) |
| Fat, inflammation, chronic active, focal | 1 (100%) | | | |
| Fat, necrosis, focal | | | | 1 (100%) |
| Salivary glands | (15) | | | (15) |
| Infiltration cellular, focal, lymphocyte | 10 (67%) | | | 8 (53%) |
| Tongue | (15) | | | (15) |
| Inflammation | 1 (7%) | | | |
| Cardiovascular System | | | | |
| None | | | | |
| Endocrine System | | | | |
| Adrenal cortex | (15) | | | (15) |
| Hypertrophy, focal | | | | 1 (7%) |
| Capsule, inflammation, chronic active, diffuse | 1 (7%) | | | |
| Subcapsular, hyperplasia, focal | 1 (7%) | | | 1 (7%) |
| General Body System | | | | |
| None | | | | |
| Genital System | | | | |
| Epididymis | (15) | | | (15) |
| Infiltration cellular, lymphocyte | 1 (7%) | | | |
| Polyarteritis | | | | 1 (7%) |
| Epithelium, degeneration, focal | 1 (7%) | | | |
| Testes | (15) | | | (15) |
| Mineralization, focal | 1 (7%) | | | |
| Sertoli cell, germinal epithelium, degeneration, focal | 1 (7%) | | | |

^a Number of animals examined microscopically at the site and the number of animals with lesion

TABLE C4
Summary of the Incidence of Nonneoplastic Lesions in Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|----------|------|-----|----------|
| Hematopoietic System | | | | |
| Lymph node | | | | (1) |
| Inguinal, hyperplasia | | | | 1 (100%) |
| Lymph node, mandibular | (15) | | | (15) |
| Hyperplasia | | | | 2 (13%) |
| Lymph node, mediastinal | (1) | | | |
| Edema | 1 (100%) | | | |
| Inflammation, chronic active, diffuse | 1 (100%) | | | |
| Spleen | (15) | | | (15) |
| Hematopoietic cell proliferation | 1 (7%) | | | 1 (7%) |
| Thymus | (15) | | | (14) |
| Atrophy | 2 (13%) | | | 2 (14%) |
| Atrophy, focal | | | | 1 (7%) |
| Integumentary System | | | | |
| Skin | (15) | | (1) | (15) |
| Subcutaneous tissue, inflammation, chronic active, focal | 1 (7%) | | | |
| Musculoskeletal System | | | | |
| None | | | | |
| Nervous System | | | | |
| Brain | (15) | | | (15) |
| Inflammation, chronic active | 1 (7%) | | | |
| Respiratory System | | | | |
| Lung | (15) | | | (15) |
| Alveolar epithelium, inflammation, chronic active, focal | 1 (7%) | | | 1 (7%) |
| Nose | (15) | | | (15) |
| Inflammation, chronic active | 1 (7%) | | | |
| Inflammation, granulomatous | 1 (7%) | | | |
| Special Senses System | | | | |
| Harderian gland | (15) | | | (15) |
| Infiltration cellular, lymphocyte | | | | 1 (7%) |
| Inflammation, chronic active | 4 (27%) | | | 2 (13%) |
| Urinary System | | | | |
| Kidney | (15) | | | (15) |
| Inflammation, chronic active | 1 (7%) | | | |
| Renal tubule, dilatation, focal | 1 (7%) | | | 2 (13%) |
| Renal tubule, nephropathy | 1 (7%) | | | |

APPENDIX D
SUMMARY OF LESIONS
IN FEMALE p53 HAPLOINSUFFICIENT MICE
IN THE 9-MONTH FEED STUDY
OF ACESULFAME POTASSIUM

| | | |
|-----------------|---|-----------|
| TABLE D1 | Summary of the Incidence of Neoplasms in Female p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 86 |
| TABLE D2 | Individual Animal Tumor Pathology of Female p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 88 |
| TABLE D3 | Statistical Analysis of Primary Neoplasms in Female p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 94 |
| TABLE D4 | Summary of the Incidence of Nonneoplastic Lesions in Female p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 95 |

TABLE D1
Summary of the Incidence of Neoplasms in Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|--|------|----------|----------|----------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 1 | | 1 | |
| Natural deaths | | 1 | | 1 |
| Survivors | | | | |
| Terminal sacrifice | 14 | 14 | 14 | 14 |
| Animals examined microscopically | 15 | 1 | 2 | 15 |
| Alimentary System | | | | |
| Mesentery | | (1) | | |
| Sarcoma | | 1 (100%) | | |
| Cardiovascular System | | | | |
| Heart | (15) | | | (15) |
| Endocrine System | | | | |
| None | | | | |
| General Body System | | | | |
| Tissue NOS | | (1) | | |
| Pelvic, osteosarcoma, metastatic, bone | | 1 (100%) | | |
| Genital System | | | | |
| None | | | | |
| Hematopoietic System | | | | |
| Lymph node | | | | (1) |
| Pancreatic, histiocytic sarcoma | | | | 1 (100%) |
| Spleen | (15) | | | (14) |
| Histiocytic sarcoma | | | | 1 (7%) |
| Thymus | (15) | | | (14) |
| Histiocytic sarcoma | | | | 1 (7%) |
| Integumentary System | | | | |
| Mammary gland | (15) | | (1) | (13) |
| Osteosarcoma | | | 1 (100%) | |
| Musculoskeletal System | | | | |
| Bone | (15) | | | (14) |
| Femur, osteosarcoma | | 1 (7%) | | |

TABLE D1
Summary of the Incidence of Neoplasms in Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|---|--------|------|-----|--------|
| Nervous System | | | | |
| None | | | | |
| Respiratory System | | | | |
| Lung | (15) | | | (15) |
| Osteosarcoma, metastatic, bone | 1 (7%) | | | |
| Special Senses System | | | | |
| None | | | | |
| Urinary System | | | | |
| None | | | | |
| Systemic Lesions | | | | |
| Multiple organs ^b | (15) | (1) | (2) | (15) |
| Histiocytic sarcoma | | | | 1 (7%) |
| Lymphoma malignant | | | | 1 (7%) |
| Neoplasm Summary | | | | |
| Total animals with primary neoplasms ^c | 1 | 1 | 1 | 2 |
| Total primary neoplasms | 1 | 1 | 1 | 2 |
| Total animals with malignant neoplasms | 1 | 1 | 1 | 2 |
| Total malignant neoplasms | 1 | 1 | 1 | 2 |
| Total animals with metastatic neoplasms | 1 | | | |
| Total metastatic neoplasms | 2 | | | |

^a Number of animals examined microscopically at the site and the number of animals with neoplasm

^b Number of animals with any tissue examined microscopically

^c Primary neoplasms: all neoplasms except metastatic neoplasms

TABLE D2
Individual Animal Tumor Pathology of Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium: 0.3%

| | | |
|--------------------------------|-------------------------------|-----------------------------|
| Number of Days on Study | 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | |
| | 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 | |
| | 6 4 4 4 4 4 4 4 4 4 4 4 4 4 4 | |
| Carcass ID Number | 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | Total Tissues/ Tumors |
| | 3 2 2 2 2 3 3 3 3 3 3 3 3 3 4 | |
| | 7 6 7 8 9 0 1 2 3 4 5 6 8 9 0 | |
| Alimentary System | | |
| Mesentery | + | 1 |
| Sarcoma | X | 1 |
| Cardiovascular System | | |
| None | | |
| Endocrine System | | |
| None | | |
| General Body System | | |
| None | | |
| Genital System | | |
| None | | |
| Hematopoietic System | | |
| None | | |
| Integumentary System | | |
| None | | |
| Musculoskeletal System | | |
| None | | |
| Nervous System | | |
| None | | |
| Respiratory System | | |
| None | | |
| Special Senses System | | |
| None | | |
| Urinary System | | |
| None | | |
| Systemic Lesions | | |
| Multiple organs | + | 1 |

TABLE D2
Individual Animal Tumor Pathology of Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium: 3%

| | | |
|--------------------------------|---------------------------------|--------------------------|
| Number of Days on Study | 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | |
| | 4 7 7 7 7 7 7 7 7 7 7 7 7 7 7 | |
| | 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4 | |
| Carcass ID Number | 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | Total Tissues/ Tumors |
| | 5 5 5 5 6 6 6 6 6 6 6 6 6 6 7 | |
| | 7 6 8 9 0 1 2 3 4 5 6 7 8 9 0 | |
| Integumentary System | | |
| Mammary gland | A M + + + + + + + + + + + + + + | 13 |
| Skin | A + + + + + + + + + + + + + + | 14 |
| Musculoskeletal System | | |
| Bone | A + + + + + + + + + + + + + + | 14 |
| Skeletal muscle | A + + + + + + + + + + + + + + | 14 |
| Nervous System | | |
| Brain | A + + + + + + + + + + + + + + | 14 |
| Respiratory System | | |
| Larynx | A + + + + + + + + I + + I I + | 11 |
| Lung | + + + + + + + + + + + + + + | 15 |
| Nose | A + + + + + + + + + + + + + + | 14 |
| Trachea | A + + + + + + + + + + + + + + | 14 |
| Special Senses System | | |
| Eye | A + + + + + + + + + + + + + + | 14 |
| Harderian gland | A + + + + + + + + + + + + + + | 14 |
| Zymbal's gland | A + + + + + + + + + + + + M M | 12 |
| Urinary System | | |
| Kidney | A + + + + + + + + + + + + + + | 14 |
| Urinary bladder | A + + + + + + + + + + + + + + | 14 |
| Systemic Lesions | | |
| Multiple organs | + + + + + + + + + + + + + + | 15 |
| Histiocytic sarcoma | | 1 |
| Lymphoma malignant | X | 1 |

TABLE D3
Statistical Analysis of Primary Neoplasms in Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|-----------|-----------|-----------|------------|
| All Organs: Malignant Neoplasms | | | | |
| Overall rate ^a | 1/15 (7%) | 1/15 (7%) | 1/15 (7%) | 2/15 (13%) |
| Adjusted rate ^b | 6.7% | 6.7% | 6.7% | 13.3% |
| Terminal rate ^c | 0/14 (0%) | 0/14 (0%) | 0/14 (0%) | 1/14 (7%) |
| First incidence (days) | 166 | 176 | 127 | 243 |
| Poly-3 test ^d | P=0.360 | P=0.760 | P=0.760 | P=0.500 |
| All Organs: Benign or Malignant Neoplasms | | | | |
| Overall rate | 1/15 (7%) | 1/15 (7%) | 1/15 (7%) | 2/15 (13%) |
| Adjusted rate | 6.7% | 6.7% | 6.7% | 13.3% |
| Terminal rate | 0/14 (0%) | 0/14 (0%) | 0/14 (0%) | 1/14 (7%) |
| First incidence (days) | 166 | 176 | 127 | 243 |
| Poly-3 test | P=0.360 | P=0.760 | P=0.760 | P=0.500 |

^a Number of neoplasm-bearing animals/number of animals necropsied.

^b Poly-3 estimated neoplasm incidence after adjustment for intercurrent mortality

^c Observed incidence at terminal kill

^d Beneath the control incidence is the P value associated with the trend test. Beneath the exposed group incidence are the P values corresponding to pairwise comparisons between the controls and that exposed group. The Poly-3 test accounts for the differential mortality in animals that do not reach terminal sacrifice.

TABLE D4
Summary of the Incidence of Nonneoplastic Lesions in Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium^a

| | 0% | 0.3% | 1% | 3% |
|--|----------|------|----------|----------|
| Disposition Summary | | | | |
| Animals initially in study | 15 | 15 | 15 | 15 |
| Early deaths | | | | |
| Moribund | 1 | | 1 | |
| Natural deaths | | 1 | | 1 |
| Survivors | | | | |
| Terminal sacrifice | 14 | 14 | 14 | 14 |
| Animals examined microscopically | 15 | 1 | 2 | 15 |
| Alimentary System | | | | |
| Liver | (15) | | | (14) |
| Infiltration cellular, focal, lymphocyte | 2 (13%) | | | 1 (7%) |
| Inflammation, chronic active | 6 (40%) | | | 7 (50%) |
| Hepatocyte, necrosis, focal | 1 (7%) | | | |
| Salivary glands | (15) | | | (14) |
| Infiltration cellular, focal, lymphocyte | 10 (67%) | | | 9 (64%) |
| Cardiovascular System | | | | |
| None | | | | |
| Endocrine System | | | | |
| Adrenal cortex | (15) | | | (14) |
| Hyperplasia, focal | 2 (13%) | | | 2 (14%) |
| Subcapsular, hyperplasia, focal | 13 (87%) | | | 12 (86%) |
| Thyroid gland | (15) | | | (14) |
| Ectopic thymus | | | | 1 (7%) |
| General Body System | | | | |
| None | | | | |
| Genital System | | | | |
| Uterus | (15) | | (1) | (14) |
| Endometrium, hyperplasia, cystic | 13 (87%) | | 1 (100%) | 13 (93%) |
| Hematopoietic System | | | | |
| Lymph node, mandibular | (15) | | | (14) |
| Hyperplasia | 1 (7%) | | | |
| Spleen | (15) | | | (14) |
| Hematopoietic cell proliferation | 2 (13%) | | | 1 (7%) |
| Thymus | (15) | | | (14) |
| Atrophy | 6 (40%) | | | 2 (14%) |
| Atrophy, diffuse | 1 (7%) | | | |
| Atrophy, focal | | | | 4 (29%) |

^a Number of animals examined microscopically at the site and the number of animals with lesion

TABLE D4
Summary of the Incidence of Nonneoplastic Lesions in Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| | 0% | 0.3% | 1% | 3% |
|--|---------|------|-----|---------|
| Integumentary System | | | | |
| Mammary gland | (15) | | (1) | (13) |
| Inflammation, acute | 1 (7%) | | | |
| Musculoskeletal System | | | | |
| None | | | | |
| Nervous System | | | | |
| None | | | | |
| Respiratory System | | | | |
| Lung | (15) | | | (15) |
| Infiltration cellular, lymphocyte | 1 (7%) | | | 1 (7%) |
| Special Senses System | | | | |
| Harderian gland | (15) | | | (14) |
| Inflammation, chronic active | 1 (7%) | | | |
| Urinary System | | | | |
| Kidney | (15) | | | (14) |
| Inflammation, chronic active | | | | 1 (7%) |
| Renal tubule, dilatation, focal | 6 (40%) | | | 2 (14%) |
| Renal tubule, nephropathy | 1 (7%) | | | |
| Urinary bladder | (14) | | | (14) |
| Infiltration cellular, focal, lymphocyte | | | | 1 (7%) |

APPENDIX E

GENETIC TOXICOLOGY

| | |
|--|------------|
| MOUSE PERIPHERAL BLOOD MICRONUCLEUS TEST PROTOCOL | 98 |
| EVALUATION PROTOCOL | 98 |
| RESULTS | 98 |
| TABLE E1 Frequency of Micronuclei in Peripheral Blood Erythrocytes of Tg.AC Hemizygous Mice Following Administration of Acesulfame Potassium in Feed for 9 Months | 99 |
| TABLE E2 Frequency of Micronuclei in Peripheral Blood Erythrocytes of p53 Haploinsufficient Mice Following Administration of Acesulfame Potassium in Feed for 9 Months | 100 |

GENETIC TOXICOLOGY

MOUSE PERIPHERAL BLOOD MICRONUCLEUS TEST PROTOCOL

A detailed discussion of this assay is presented by MacGregor *et al.* (1990). At the end of the 9-month toxicity studies, peripheral blood samples were obtained from male and female mice. Smears were immediately prepared and fixed in absolute methanol. The methanol-fixed slides were stained with acridine orange and coded. Slides were scanned to determine the frequency of micronuclei in 2,000 normochromatic erythrocytes (NCEs) in up to 13 Tg.AC hemizygous and 15 p53 haploinsufficient mice per exposure group; 1,000 total erythrocytes were counted to determine the percent polychromatic erythrocytes (PCEs).

The results were tabulated as the mean of the pooled results from all animals within a treatment group plus or minus the standard error of the mean. The frequency of micronucleated cells among NCEs was analyzed by a statistical software package that tested for increasing trend over exposure groups with a one-tailed Cochran-Armitage trend test, followed by pairwise comparisons between each exposure group and the control group (ILS, 1990). In the presence of excess binomial variation, as detected by a binomial dispersion test, the binomial variance of the Cochran-Armitage test was adjusted upward in proportion to the excess variation. In the micronucleus test, an individual trial is considered positive if the trend test P value is less than or equal to 0.025 or if the P value for any single exposed group is less than or equal to 0.025 divided by the number of exposure groups. A final call of positive for micronucleus induction is preferably based on reproducibly positive trials (as noted above). Results of the 9-month studies were accepted without repeat tests, because additional test data could not be obtained. Ultimately, the final call is determined by the scientific staff after considering the results of statistical analyses, the reproducibility of any effects observed, and the magnitudes of those effects.

EVALUATION PROTOCOL

These are the basic guidelines for arriving at an overall assay result for assays performed by the National Toxicology Program. Statistical as well as biological factors are considered. For an individual assay, the statistical procedures for data analysis have been described in the preceding protocols. There have been instances, however, in which multiple aliquots of a chemical were tested in the same assay, and different results were obtained among aliquots and/or among laboratories. Results from more than one aliquot or from more than one laboratory are not simply combined into an overall result. Rather, all the data are critically evaluated, particularly with regard to pertinent protocol variations, in determining the weight of evidence for an overall conclusion of chemical activity in an assay. In addition to multiple aliquots, the *in vitro* assays have another variable that must be considered in arriving at an overall test result. *In vitro* assays are conducted with and without exogenous metabolic activation. Results obtained in the absence of activation are not combined with results obtained in the presence of activation; each testing condition is evaluated separately. The summary table in the Abstract of this Technical Report presents a result that represents a scientific judgement of the overall evidence for activity of the chemical in an assay.

RESULTS

Acesulfame potassium did not increase the frequency of micronucleated NCEs in peripheral blood of male or female Tg.AC hemizygous mice administered 0.3% to 3% of the chemical in feed for 9 months (Table E1). A similar study was conducted in p53 haploinsufficient mice, and in this study, a significant exposure-related increase in the frequency of micronucleated NCEs was noted in males but not females; micronucleus frequencies in the 1% and 3% male groups were significantly elevated over the control group (Table E2). There was no significant alteration in the percentage of PCEs among total erythrocytes in either Tg.AC hemizygous or p53 haploinsufficient mice.

TABLE E1
Frequency of Micronuclei in Peripheral Blood Erythrocytes of Tg.AC Hemizygous Mice
Following Administration of Acesulfame Potassium in Feed for 9 Months^a

| Dose (%) | Number of Mice with Erythrocytes Scored | Micronucleated NCEs/ 1,000 NCEs ^b | P Value ^c | % PCE |
|----------------------------|---|---|----------------------|-------|
| Male | | | | |
| NTP-2000 feed ^d | 13 | 1.42 ± 0.19 | | 3.4 |
| 0.3 | 11 | 1.59 ± 0.18 | 0.3180 | 3.4 |
| 1 | 13 | 1.62 ± 0.23 | 0.2867 | 2.9 |
| 3 | 13 | 1.58 ± 0.19 | 0.3252 | 3.4 |
| | | P=0.391 ^e | | |
| Female | | | | |
| NTP-2000 feed | 12 | 1.29 ± 0.16 | | 3.4 |
| 0.3 | 12 | 1.13 ± 0.22 | 0.7004 | 3.1 |
| 1 | 12 | 1.17 ± 0.27 | 0.6520 | 3.5 |
| 3 | 12 | 1.58 ± 0.23 | 0.1995 | 3.3 |
| | | P=0.098 | | |

^a Study was performed at SITEK Laboratories, Inc. The detailed protocol is presented by MacGregor *et al.* (1990); NCE=normochromatic erythrocyte; PCE=polychromatic erythrocyte.

^b Mean ± standard error

^c Pairwise comparison with the controls, significant at P≤0.008 (ILS, 1990)

^d Control

^e Significance of micronucleated NCEs/1,000 NCEs tested by the one-tailed trend test, significant at P≤0.025 (ILS, 1990)

TABLE E2
Frequency of Micronuclei in Peripheral Blood Erythrocytes of p53 Haploinsufficient Mice
Following Administration of Acesulfame Potassium in Feed for 9 Months^a

| Dose (%) | Number of Mice with Erythrocytes Scored | Micronucleated NCEs/1,000 NCEs ^b | P Value ^c | % PCE |
|----------------------------|---|---|----------------------|-------|
| Male | | | | |
| NTP-2000 feed ^d | 14 | 1.57 ± 0.22 | | 2.98 |
| 0.3 | 15 | 2.10 ± 0.24 | 0.0691 | 3.25 |
| 1 | 15 | 2.57 ± 0.19 | 0.0043 | 3.30 |
| 3 | 14 | 2.79 ± 0.33 | 0.0010 | 3.36 |
| | | P=0.003 ^e | | |
| Female | | | | |
| NTP-2000 feed | 14 | 1.29 ± 0.16 | | 3.44 |
| 0.3 | 14 | 1.71 ± 0.17 | 0.0950 | 3.29 |
| 1 | 14 | 1.75 ± 0.21 | 0.0791 | 3.27 |
| 3 | 14 | 1.50 ± 0.26 | 0.2483 | 3.24 |
| | | P=0.467 | | |

^a Study was performed at SITEK Laboratories, Inc. The detailed protocol is presented by MacGregor *et al.* (1990); NCE=normochromatic erythrocyte; PCE=polychromatic erythrocyte.

^b Mean ± standard error

^c Pairwise comparison with the controls, significant at P≤0.008 (ILS, 1990)

^d Control

^e Significance of micronucleated NCEs/1,000 NCEs tested by the one-tailed trend test, significant at P≤0.025 (ILS, 1990)

APPENDIX F

CHEMICAL CHARACTERIZATION AND DOSE FORMULATION STUDIES

| | |
|--|------------|
| PROCUREMENT AND CHARACTERIZATION OF ACESULFAME POTASSIUM | 102 |
| PREPARATION AND ANALYSIS OF DOSE FORMULATIONS | 102 |
| FIGURE F1 Infrared Absorption Spectrum of Acesulfame Potassium | 104 |
| FIGURE F2 Proton Nuclear Magnetic Resonance Spectrum of Acesulfame Potassium | 105 |
| TABLE F1 Preparation and Storage of Dose Formulations in the 9-Month Feed Studies of Acesulfame Potassium | 106 |
| TABLE F2 Results of Analyses of Dose Formulations Administered to Tg.AC Hemizygous Mice and p53 Haploinsufficient Mice in the 9-Month Feed Studies of Acesulfame Potassium .. | 107 |

CHEMICAL CHARACTERIZATION AND DOSE FORMULATION STUDIES

PROCUREMENT AND CHARACTERIZATION OF ACESULFAME POTASSIUM

Acesulfame potassium was obtained from Riedel-deHaen (St. Louis, MO) via Research Triangle Institute (RTI) (Research Triangle Park, NC) in one lot (8415-76-02 RTI), which was used in the 9-month studies in Tg.AC hemizygous mice and p53 haploinsufficient mice. Identity and purity analyses were conducted by the analytical chemistry laboratory, RTI, and the study laboratory (BioReliance Corporation, Rockville, MD); the study laboratory also conducted stability analyses. Reports on analyses performed in support of the acesulfame potassium studies are on file at the National Institute of Environmental Health Sciences.

Lot 8415-76-02 RTI of the chemical, a white, odorless, crystalline powder, was identified as acesulfame potassium by the analytical chemistry laboratory using infrared and proton nuclear magnetic resonance spectroscopy and by the study laboratory using infrared spectroscopy. All spectra were consistent with the structure of acesulfame potassium. The infrared and proton nuclear magnetic resonance spectra are presented in Figures F1 and F2.

The purity of lot 8415-76-02 RTI was determined by the analytical chemistry laboratory using high-performance liquid chromatography (HPLC), using Waters Chromatography (Milford, MA), an Alltima C₁₈ column, 25 cm × 3.2 mm (Alltech; Chicago, IL), a mobile phase of 0.0125 M potassium dihydrogen phosphate (pH3.5):acetonitrile (100:0 for 5 minutes; 100:0 to 90:10 in 25 minutes; 90:10 for 10 minutes), a flow rate of 1.0 mL/minute, with ultraviolet detection at 230 nm. The study laboratory confirmed purity by major peak comparisons versus a reference sample from the same lot (stored at -20° C) using HPLC with a variation in the mobile phase (potassium dihydrogen phosphate:acetonitrile; 95:5) and a Hewlett-Packard model 1100 (Palo Alto, CA) instrument.

For lot 8415-76-02 RTI, HPLC indicated one major peak and no impurities. The analytical chemistry laboratory determined the purity to be greater than 99%; the study laboratory determined the purity to be 101% relative to the reference standard. The overall purity of lot 8415-76-02 RTI was determined to be 99% or greater.

To ensure stability, the bulk chemical was stored at room temperature under a headspace of inert gas, protected from light, in polyethylene bags inside plastic pails. Stability of the bulk chemical was monitored during the study by the study laboratory using HPLC. No degradation of the bulk chemical was observed.

PREPARATION AND ANALYSIS OF DOSE FORMULATIONS

The dose formulations were prepared once every 2 to 4 weeks by mixing acesulfame potassium with feed (Table F1). A premix was prepared by hand and then blended with additional feed in a Patterson-Kelly twin-shell blender for 15 minutes using an intensifier bar for the initial 5 minutes. Formulations were stored in doubled polyethylene bags at room temperature for up to 35 days.

Homogeneity studies of the 0.1% and 5% acesulfame potassium formulations and stability studies of the 0.1% formulations were performed by the analytical chemistry laboratory using HPLC with a variation in the mobile phase ratio (95:5). Homogeneity studies of the 0.3% and 3% acesulfame potassium formulations were performed by the study laboratory using HPLC and the RTI extraction method modified to centrifuge samples before analysis. Homogeneity was confirmed, and stability was confirmed for at least 35 days for dose formulations stored in amber glass bottles at -20° C, -5° C, and 25° C, and when exposed to light and air for 7 days.

Periodic analyses of the dose formulations of acesulfame potassium were conducted by the study laboratory using HPLC. Samples of the formulation were extracted with 50:50 water/methanol, shaken for 10 minutes, and then centrifuged; theophylline (2 mg/mL) was used as an internal standard. During the 9-month studies, the dose formulations were analyzed five times; all 15 of the dose formulations analyzed were within 10 % of the target concentrations (Table F2). Animal room samples of these dose formulations were also analyzed, and 12 of 15 animal room samples were within 10% of the target concentrations (Table F2).




FIGURE F1
Infrared Absorption Spectrum of Acesulfame Potassium




FIGURE F2
Proton Nuclear Magnetic Resonance Spectrum of Acesulfame Potassium

TABLE F1**Preparation and Storage of Dose Formulations in the 9-Month Feed Studies of Acesulfame Potassium**

Preparation

A premix of feed and acesulfame potassium was prepared, then layered into the remaining feed and blended in a Patterson-Kelly twin-shell blender with the intensifier bar on for 5 minutes and off for 10 minutes. The dose formulations were prepared every two to four weeks.

Chemical Lot Number

8415-76-02 RTI

Maximum Storage Time

35 days

Storage Conditions

Stored in doubled polyethylene bags at room temperature, protected from light

Study Laboratory

BioReliance Corporation(Rockville, MD)

TABLE F2
Results of Analyses of Dose Formulations Administered to Tg.AC Hemizygous Mice
and p53 Haploinsufficient Mice in the 9-Month Feed Studies of Acesulfame Potassium

| Date Prepared | Date Analyzed | Target Concentration (%) | Determined Concentration ^a (%) | Difference from Target (%) |
|-------------------|--------------------------------|--------------------------|---|----------------------------|
| January 19, 2000 | January 19, 2000 | 0.3 | 0.283 | -6 |
| | | 1 | 1.06 | +6 |
| | | 3 | 2.89 | -4 |
| | March 7, 2000 ^b | 0.3 | 0.297 | -1 |
| | | 1 | 1.04 | +4 |
| | | 3 | 3.06 | +2 |
| February 29, 2000 | February 29, 2000 | 0.3 | 0.305 | +2 |
| | | 1 | 1.02 | +2 |
| | | 3 | 3.14 | +5 |
| | April 19, 2000 ^b | 0.3 | 0.301 | 0 |
| | | 1 | 0.857 | -14 |
| | | 3 | 3.01 | 0 |
| May 9, 2000 | May 9, 2000 | 0.3 | 0.298 | -1 |
| | | 1 | 0.985 | -2 |
| | | 3 | 3.01 | 0 |
| | June 8, 2000 ^b | 0.3 | 0.295 | -2 |
| | | 1. | 0.942 | -6 |
| | | 3. | 3.04 | +1 |
| July 25, 2000 | July 25, 2000 | 0.3 | 0.288 | -4 |
| | | 1 | 1.05 | +5 |
| | | 3 | 3.12 | +4 |
| | September 6, 2000 ^b | 0.3 | 0.293 | -2 |
| | | 1 | 0.847 | -15 |
| | | 3 | 2.76 | -8 |
| October 3, 2000 | October 3, 2000 | 0.3 | 0.301 | 0 |
| | | 1 | 1.01 | +1 |
| | | 3 | 3.08 | +3 |
| | November 8, 2000 ^b | 0.3 | 0.336 | +12 |
| | | 1 | 1.06 | +6 |
| | | 3 | 3.14 | +5 |

^a Results of duplicate analyses

^b Animal room samples

APPENDIX G
FEED AND COMPOUND CONSUMPTION
IN THE 9-MONTH FEED STUDIES
OF ACESULFAME POTASSIUM

| | | |
|-----------------|---|------------|
| TABLE G1 | Feed and Compound Consumption by Male Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 110 |
| TABLE G2 | Feed and Compound Consumption by Female Tg.AC Hemizygous Mice in the 9-Month Feed Study of Acesulfame Potassium | 111 |
| TABLE G3 | Feed and Compound Consumption by Male p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 112 |
| TABLE G4 | Feed and Compound Consumption by Female p53 Haploinsufficient Mice in the 9-Month Feed Study of Acesulfame Potassium | 113 |

TABLE G1
Feed and Compound Consumption by Male Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Week | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|-----------------------|-----------------|----------|-----------------|---------------------------|----------|-----------------|--------------|----------|-----------------|--------------|
| | Feed (g) ^a | Body Weight (g) | Feed (g) | Body Weight (g) | Dose (mg/kg) ^b | Feed (g) | Body Weight (g) | Dose (mg/kg) | Feed (g) | Body Weight (g) | Dose (mg/kg) |
| 2 | 3.9 | 22.6 | 4.1 | 22.0 | 554 | 4.5 | 22.7 | 1,978 | 4.6 | 22.5 | 6,093 |
| 3 | 4.2 | 23.3 | 4.4 | 23.6 | 564 | 4.5 | 23.7 | 1,901 | 4.6 | 23.7 | 5,838 |
| 4 | 4.3 | 24.2 | 4.3 | 24.9 | 524 | 4.5 | 24.8 | 1,817 | 4.6 | 24.7 | 5,615 |
| 5 | 4.2 | 24.9 | 4.4 | 25.7 | 509 | 4.3 | 25.3 | 1,686 | 4.4 | 25.1 | 5,309 |
| 6 | 4.2 | 25.5 | 4.1 | 26.3 | 470 | 4.4 | 26.4 | 1,659 | 4.5 | 25.6 | 5,290 |
| 7 | 4.1 | 26.3 | 4.1 | 26.5 | 460 | 4.3 | 26.8 | 1,607 | 4.5 | 26.9 | 5,021 |
| 8 | 3.8 | 25.9 | 4.2 | 27.2 | 467 | 4.3 | 27.5 | 1,567 | 4.5 | 27.4 | 4,918 |
| 9 | 4.2 | 27.4 | 4.1 | 27.6 | 445 | 4.1 | 27.7 | 1,464 | 4.3 | 27.7 | 4,656 |
| 10 | 4.1 | 27.5 | 4.1 | 27.6 | 450 | 4.2 | 28.0 | 1,515 | 4.4 | 27.7 | 4,739 |
| 11 | 4.1 | 27.0 | 4.2 | 28.2 | 445 | 4.3 | 28.4 | 1,528 | 4.4 | 27.6 | 4,790 |
| 12 | 4.2 | 28.0 | 4.3 | 29.0 | 444 | 4.4 | 29.2 | 1,502 | 4.6 | 28.8 | 4,832 |
| 13 | 4.0 | 28.2 | 4.2 | 28.9 | 436 | 4.3 | 29.4 | 1,463 | 4.5 | 29.0 | 4,683 |
| 14 | 3.9 | 28.8 | 4.1 | 29.8 | 413 | 4.2 | 29.7 | 1,418 | 4.5 | 29.3 | 4,589 |
| 15 | 4.1 | 29.3 | 4.3 | 29.9 | 434 | 4.4 | 30.0 | 1,458 | 4.6 | 29.9 | 4,575 |
| 16 | 4.0 | 29.5 | 4.0 | 29.8 | 406 | 4.2 | 30.3 | 1,402 | 4.5 | 30.1 | 4,485 |
| 17 | 4.2 | 29.6 | 4.2 | 29.8 | 420 | 4.4 | 30.5 | 1,454 | 4.7 | 30.4 | 4,669 |
| 18 | 4.1 | 29.2 | 4.4 | 30.5 | 429 | 4.4 | 30.7 | 1,440 | 4.5 | 30.6 | 4,399 |
| 19 | 4.2 | 29.4 | 4.0 | 30.4 | 396 | 4.1 | 30.5 | 1,331 | 4.5 | 30.7 | 4,399 |
| 20 | 3.9 | 29.5 | 4.0 | 29.7 | 409 | 4.2 | 30.5 | 1,391 | 4.4 | 30.6 | 4,342 |
| 21 | 4.0 | 29.5 | 4.2 | 30.1 | 417 | 4.3 | 31.2 | 1,367 | 4.4 | 31.1 | 4,232 |
| 22 | 3.8 | 29.5 | 4.1 | 30.9 | 393 | 4.3 | 31.6 | 1,359 | 4.6 | 31.5 | 4,379 |
| 23 | 4.1 | 30.4 | 4.1 | 32.0 | 387 | 4.1 | 31.4 | 1,298 | 4.4 | 31.8 | 4,130 |
| 24 | 4.1 | 31.5 | 3.9 | 31.1 | 374 | 4.0 | 31.8 | 1,242 | 4.4 | 31.9 | 4,156 |
| 25 | 4.0 | 31.6 | 4.4 | 32.2 | 408 | 4.2 | 32.2 | 1,299 | 4.4 | 32.1 | 4,073 |
| 26 | 4.0 | 31.8 | 4.2 | 32.2 | 393 | 4.1 | 32.2 | 1,264 | 4.5 | 32.8 | 4,109 |
| 27 | 4.0 | 32.0 | 4.1 | 32.2 | 386 | 4.2 | 32.2 | 1,318 | 4.4 | 32.3 | 4,065 |
| 28 | 3.8 | 32.0 | 4.2 | 32.6 | 386 | 4.2 | 32.9 | 1,276 | 4.4 | 32.8 | 4,037 |
| 29 | 4.3 | 32.4 | 4.3 | 32.8 | 394 | 4.2 | 33.1 | 1,280 | 4.3 | 32.5 | 3,977 |
| 30 | 3.9 | 32.5 | 4.2 | 33.3 | 378 | 4.1 | 33.4 | 1,231 | 4.3 | 33.0 | 3,948 |
| 31 | 4.1 | 32.5 | 4.2 | 33.3 | 377 | 4.2 | 33.1 | 1,266 | 4.2 | 33.0 | 3,778 |
| 32 | 4.1 | 32.5 | 4.2 | 33.2 | 379 | 4.3 | 33.6 | 1,287 | 4.7 | 33.7 | 4,187 |
| 33 | 4.1 | 32.3 | 4.1 | 32.5 | 379 | 4.2 | 33.2 | 1,280 | 4.3 | 33.0 | 3,888 |
| 34 | 4.1 | 32.2 | 4.4 | 32.7 | 402 | 4.3 | 33.4 | 1,281 | 4.4 | 33.4 | 3,935 |
| 35 | 4.3 | 32.6 | 4.2 | 32.8 | 383 | 4.5 | 33.4 | 1,346 | 4.3 | 33.4 | 3,818 |
| 36 | 4.3 | 33.0 | 4.3 | 32.7 | 390 | 4.5 | 33.7 | 1,325 | 4.5 | 33.4 | 4,071 |
| 37 | 4.4 | 32.7 | 4.3 | 33.5 | 387 | 4.6 | 34.1 | 1,344 | 4.6 | 33.3 | 4,171 |
| 38 | 4.3 | 32.9 | 4.6 | 33.5 | 412 | 4.4 | 33.4 | 1,318 | 4.6 | 33.2 | 4,162 |
| 39 | 4.2 | 33.2 | 4.0 | 32.9 | 361 | 4.4 | 34.4 | 1,275 | 4.6 | 33.9 | 4,070 |
| 40 | 4.2 | 33.2 | 4.5 | 33.6 | 400 | 4.2 | 34.1 | 1,245 | 4.5 | 33.5 | 4,059 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 4.1 | 25.9 | 4.2 | 26.5 | 481 | 4.3 | 26.7 | 1,641 | 4.5 | 26.4 | 5,149 |
| 14-40 | 4.1 | 31.3 | 4.2 | 31.8 | 396 | 4.3 | 32.2 | 1,326 | 4.5 | 32.1 | 4,174 |

^a Grams of feed consumed per animal per day

^b Milligrams of acesulfame potassium consumed per kilogram body weight per day

TABLE G2
Feed and Compound Consumption by Female Tg.AC Hemizygous Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Week | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|-----------------------|-----------------|----------|-----------------|---------------------------|----------|-----------------|--------------|----------|-----------------|--------------|
| | Feed (g) ^a | Body Weight (g) | Feed (g) | Body Weight (g) | Dose (mg/kg) ^b | Feed (g) | Body Weight (g) | Dose (mg/kg) | Feed (g) | Body Weight (g) | Dose (mg/kg) |
| 2 | 3.6 | 17.5 | 4.3 | 18.2 | 710 | 3.9 | 18.2 | 2,160 | 4.5 | 18.1 | 7,467 |
| 3 | 4.3 | 17.8 | 4.1 | 18.5 | 660 | 4.3 | 18.3 | 2,339 | 4.0 | 18.3 | 6,630 |
| 4 | 4.5 | 19.6 | 4.1 | 19.5 | 625 | 4.1 | 19.3 | 2,106 | 4.1 | 19.7 | 6,299 |
| 5 | 4.3 | 19.7 | 3.8 | 19.6 | 582 | 3.8 | 19.8 | 1,944 | 3.9 | 19.7 | 5,992 |
| 6 | 4.0 | 19.7 | 4.1 | 19.5 | 626 | 3.8 | 19.6 | 1,948 | 4.1 | 19.4 | 6,303 |
| 7 | 3.9 | 20.3 | 3.9 | 20.1 | 588 | 4.2 | 20.5 | 2,029 | 4.1 | 20.6 | 5,926 |
| 8 | 4.0 | 20.7 | 4.1 | 20.8 | 592 | 4.0 | 21.2 | 1,902 | 4.0 | 20.8 | 5,771 |
| 9 | 3.9 | 21.3 | 4.0 | 21.6 | 559 | 4.0 | 21.5 | 1,875 | 4.0 | 21.2 | 5,672 |
| 10 | 4.0 | 21.7 | 3.9 | 21.6 | 548 | 4.1 | 21.9 | 1,880 | 4.2 | 21.4 | 5,897 |
| 11 | 4.2 | 22.1 | 4.0 | 21.9 | 546 | 4.4 | 22.2 | 1,960 | 4.4 | 22.3 | 5,953 |
| 12 | 4.2 | 22.5 | 4.2 | 22.1 | 573 | 4.3 | 22.6 | 1,892 | 4.2 | 21.9 | 5,801 |
| 13 | 4.1 | 22.2 | 4.0 | 22.4 | 538 | 3.9 | 22.7 | 1,737 | 4.2 | 21.9 | 5,770 |
| 14 | 3.8 | 22.3 | 4.0 | 22.7 | 533 | 3.9 | 22.7 | 1,719 | 4.0 | 22.1 | 5,411 |
| 15 | 4.3 | 22.5 | 4.0 | 23.1 | 514 | 4.1 | 22.9 | 1,785 | 3.9 | 22.6 | 5,230 |
| 16 | 4.2 | 23.5 | 4.0 | 23.3 | 512 | 4.0 | 23.2 | 1,705 | 4.2 | 23.4 | 5,448 |
| 17 | 4.0 | 23.7 | 4.2 | 23.6 | 528 | 4.2 | 23.2 | 1,828 | 4.1 | 23.0 | 5,316 |
| 18 | 4.1 | 24.0 | 3.8 | 23.4 | 491 | 4.1 | 23.7 | 1,738 | 4.2 | 23.3 | 5,408 |
| 19 | 4.0 | 23.8 | 3.9 | 23.3 | 500 | 3.9 | 23.9 | 1,615 | 4.1 | 23.3 | 5,249 |
| 20 | 4.3 | 24.4 | 4.0 | 23.5 | 507 | 4.2 | 23.9 | 1,760 | 4.3 | 23.7 | 5,447 |
| 21 | 4.1 | 24.6 | 4.0 | 23.8 | 498 | 4.3 | 24.4 | 1,781 | 3.7 | 23.5 | 4,754 |
| 22 | 4.0 | 24.9 | 3.9 | 24.0 | 488 | 4.0 | 24.6 | 1,628 | 3.9 | 23.3 | 5,026 |
| 23 | 4.3 | 25.0 | 3.8 | 24.1 | 478 | 3.7 | 24.2 | 1,544 | 4.5 | 24.0 | 5,614 |
| 24 | 4.1 | 25.4 | 4.0 | 24.5 | 491 | 4.1 | 24.8 | 1,641 | 4.1 | 24.7 | 4,996 |
| 25 | 3.8 | 25.3 | 4.0 | 24.5 | 487 | 3.8 | 24.7 | 1,527 | 4.0 | 25.1 | 4,827 |
| 26 | 4.1 | 25.4 | 4.0 | 24.8 | 488 | 4.2 | 25.0 | 1,668 | 4.2 | 25.2 | 4,973 |
| 27 | 4.3 | 26.7 | 4.2 | 25.3 | 499 | 3.9 | 25.2 | 1,545 | 4.0 | 25.1 | 4,840 |
| 28 | 4.1 | 26.5 | 3.8 | 25.5 | 448 | 3.8 | 25.1 | 1,525 | 4.1 | 25.1 | 4,893 |
| 29 | 3.9 | 26.8 | 3.9 | 25.2 | 467 | 4.0 | 25.1 | 1,597 | 3.9 | 24.8 | 4,672 |
| 30 | 3.8 | 27.1 | 4.0 | 25.6 | 465 | 3.8 | 25.4 | 1,483 | 4.1 | 25.7 | 4,758 |
| 31 | 3.9 | 26.7 | 4.0 | 25.5 | 465 | 3.9 | 25.5 | 1,536 | 4.1 | 26.1 | 4,755 |
| 32 | 4.0 | 26.4 | 4.1 | 25.7 | 477 | 4.3 | 26.0 | 1,641 | 4.3 | 26.1 | 4,934 |
| 33 | 3.9 | 25.9 | 4.2 | 25.8 | 482 | 3.8 | 25.5 | 1,502 | 4.1 | 25.8 | 4,701 |
| 34 | 4.1 | 26.0 | 4.0 | 26.1 | 465 | 3.8 | 25.4 | 1,493 | 4.2 | 25.6 | 4,882 |
| 35 | 4.0 | 26.0 | 3.9 | 24.9 | 470 | 4.1 | 25.9 | 1,586 | 4.4 | 26.4 | 5,048 |
| 36 | 4.6 | 28.3 | 4.3 | 25.9 | 493 | 4.1 | 26.3 | 1,573 | 4.4 | 26.2 | 5,021 |
| 37 | 4.5 | 28.4 | 4.3 | 26.4 | 487 | 4.3 | 26.6 | 1,615 | 4.2 | 26.3 | 4,813 |
| 38 | 4.3 | 27.8 | 4.2 | 26.7 | 475 | 4.0 | 26.6 | 1,508 | 4.5 | 26.6 | 5,042 |
| 39 | 4.6 | 28.5 | 4.1 | 26.6 | 460 | 4.0 | 26.8 | 1,491 | 4.2 | 26.7 | 4,763 |
| 40 | 4.4 | 28.5 | 4.5 | 26.3 | 515 | 4.4 | 27.0 | 1,639 | 4.3 | 26.6 | 4,851 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 4.1 | 20.4 | 4.0 | 20.5 | 596 | 4.1 | 20.7 | 1,981 | 4.2 | 20.4 | 6,123 |
| 14-40 | 4.1 | 25.7 | 4.0 | 24.8 | 488 | 4.0 | 25.0 | 1,618 | 4.2 | 24.8 | 5,025 |

^a Grams of feed consumed per animal per day

^b Milligrams of acesulfame potassium consumed per kilogram body weight per day

TABLE G3
Feed and Compound Consumption by Male p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Week | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|-----------------------|-----------------|----------|-----------------|---------------------------|----------|-----------------|--------------|----------|-----------------|--------------|
| | Feed (g) ^a | Body Weight (g) | Feed (g) | Body Weight (g) | Dose (mg/kg) ^b | Feed (g) | Body Weight (g) | Dose (mg/kg) | Feed (g) | Body Weight (g) | Dose (mg/kg) |
| 2 | 4.2 | 21.9 | 4.1 | 21.7 | 571 | 4.2 | 22.0 | 1,922 | 4.2 | 22.0 | 5,731 |
| 3 | 4.4 | 22.1 | 3.7 | 21.9 | 509 | 4.1 | 22.3 | 1,854 | 3.8 | 22.1 | 5,170 |
| 4 | 4.3 | 22.9 | 4.2 | 22.5 | 560 | 3.8 | 22.8 | 1,690 | 4.1 | 22.7 | 5,416 |
| 5 | 4.4 | 23.6 | 4.7 | 23.6 | 599 | 4.2 | 23.9 | 1,755 | 4.6 | 24.1 | 5,696 |
| 6 | 4.1 | 24.1 | 4.1 | 23.9 | 509 | 4.1 | 24.4 | 1,673 | 4.1 | 24.4 | 5,091 |
| 7 | 3.9 | 24.5 | 4.0 | 24.2 | 491 | 4.2 | 25.0 | 1,674 | 4.1 | 24.7 | 5,011 |
| 8 | 3.9 | 25.3 | 4.1 | 24.8 | 491 | 4.2 | 25.5 | 1,638 | 4.3 | 25.2 | 5,085 |
| 9 | 3.9 | 25.4 | 3.9 | 24.9 | 468 | 4.0 | 25.6 | 1,549 | 4.1 | 25.4 | 4,789 |
| 10 | 4.0 | 25.6 | 4.1 | 25.1 | 486 | 4.1 | 26.1 | 1,559 | 4.2 | 25.6 | 4,922 |
| 11 | 4.0 | 25.9 | 4.1 | 25.4 | 483 | 4.1 | 26.8 | 1,520 | 4.3 | 25.7 | 5,042 |
| 12 | 4.2 | 26.5 | 4.1 | 25.6 | 483 | 4.1 | 27.1 | 1,520 | 4.3 | 26.3 | 4,937 |
| 13 | 4.0 | 26.6 | 4.3 | 25.9 | 499 | 4.1 | 26.9 | 1,514 | 4.1 | 26.5 | 4,601 |
| 14 | 4.1 | 27.0 | 4.3 | 26.7 | 482 | 4.0 | 27.2 | 1,473 | 4.2 | 27.0 | 4,631 |
| 15 | 4.2 | 27.5 | 4.2 | 27.2 | 464 | 4.4 | 28.3 | 1,561 | 4.3 | 27.9 | 4,614 |
| 16 | 4.3 | 28.0 | 4.3 | 27.9 | 463 | 4.1 | 28.4 | 1,459 | 4.3 | 28.1 | 4,566 |
| 17 | 4.3 | 28.5 | 4.5 | 27.7 | 484 | 4.5 | 28.9 | 1,544 | 4.5 | 28.2 | 4,802 |
| 19 | 4.2 | 29.2 | 4.1 | 28.3 | 434 | 4.2 | 29.8 | 1,406 | 4.2 | 28.9 | 4,407 |
| 20 | 4.0 | 29.4 | 4.2 | 28.6 | 436 | 4.0 | 29.2 | 1,359 | 4.4 | 29.1 | 4,539 |
| 21 | 4.3 | 29.7 | 4.3 | 28.8 | 449 | 4.3 | 30.0 | 1,443 | 4.4 | 29.4 | 4,510 |
| 22 | 4.1 | 29.5 | 4.1 | 29.0 | 426 | 4.1 | 29.9 | 1,370 | 4.2 | 29.5 | 4,303 |
| 23 | 3.9 | 29.2 | 4.0 | 28.5 | 425 | 4.0 | 29.6 | 1,357 | 4.5 | 29.5 | 4,565 |
| 24 | 4.3 | 29.6 | 4.8 | 29.2 | 489 | 4.3 | 30.3 | 1,410 | 4.5 | 30.3 | 4,422 |
| 25 | 4.2 | 29.6 | 4.2 | 28.9 | 441 | 4.3 | 30.5 | 1,392 | 4.3 | 29.9 | 4,281 |
| 26 | 4.2 | 29.6 | 4.5 | 29.4 | 462 | 4.3 | 30.5 | 1,405 | 4.4 | 30.3 | 4,351 |
| 27 | 3.9 | 29.0 | 4.3 | 29.0 | 443 | 4.4 | 30.9 | 1,417 | 4.4 | 30.4 | 4,325 |
| 28 | 4.2 | 29.3 | 4.3 | 29.8 | 432 | 4.5 | 31.9 | 1,422 | 4.5 | 31.0 | 4,402 |
| 29 | 4.3 | 30.3 | 4.4 | 30.2 | 438 | 4.3 | 32.1 | 1,327 | 4.1 | 30.5 | 4,001 |
| 30 | 4.3 | 30.4 | 4.5 | 30.5 | 441 | 4.3 | 32.4 | 1,322 | 4.5 | 31.1 | 4,357 |
| 31 | 4.3 | 30.1 | 4.4 | 30.0 | 438 | 4.5 | 32.2 | 1,405 | 4.4 | 31.1 | 4,210 |
| 32 | 4.2 | 29.9 | 4.4 | 29.3 | 447 | 4.4 | 32.0 | 1,377 | 4.6 | 30.9 | 4,454 |
| 33 | 4.4 | 29.6 | 4.6 | 29.5 | 469 | 4.3 | 31.1 | 1,383 | 4.7 | 30.4 | 4,648 |
| 34 | 4.5 | 29.2 | 4.5 | 29.2 | 466 | 4.7 | 31.3 | 1,501 | 4.5 | 30.2 | 4,442 |
| 35 | 4.6 | 29.2 | 4.7 | 29.3 | 484 | 4.5 | 30.8 | 1,458 | 4.5 | 29.6 | 4,558 |
| 36 | 4.3 | 29.1 | 4.7 | 29.3 | 484 | 4.6 | 31.0 | 1,473 | 4.7 | 29.4 | 4,764 |
| 37 | 4.4 | 29.1 | 4.7 | 29.7 | 473 | 4.7 | 31.1 | 1,517 | 4.7 | 29.7 | 4,773 |
| 38 | 4.5 | 29.1 | 4.7 | 29.5 | 477 | 4.6 | 31.3 | 1,477 | 4.7 | 29.9 | 4,722 |
| 39 | 4.6 | 29.2 | 4.7 | 29.9 | 469 | 4.8 | 31.7 | 1,531 | 4.7 | 30.1 | 4,728 |
| 40 | 4.5 | 29.5 | 4.6 | 30.1 | 458 | 4.4 | 31.6 | 1,407 | 4.5 | 29.9 | 4,520 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 4.1 | 24.5 | 4.1 | 24.1 | 513 | 4.1 | 24.9 | 1,656 | 4.2 | 24.6 | 5,124 |
| 14-40 | 4.3 | 29.2 | 4.4 | 29.1 | 457 | 4.4 | 30.5 | 1,431 | 4.4 | 29.7 | 4,496 |

^a Grams of feed consumed per animal per day

^b Milligrams of acesulfame potassium consumed per kilogram body weight per day

TABLE G4
Feed and Compound Consumption by Female p53 Haploinsufficient Mice
in the 9-Month Feed Study of Acesulfame Potassium

| Week | 0% | | 0.3% | | | 1% | | | 3% | | |
|-----------------------|-----------------------|-----------------|----------|-----------------|---------------------------|----------|-----------------|--------------|----------|-----------------|--------------|
| | Feed (g) ^a | Body Weight (g) | Feed (g) | Body Weight (g) | Dose (mg/kg) ^b | Feed (g) | Body Weight (g) | Dose (mg/kg) | Feed (g) | Body Weight (g) | Dose (mg/kg) |
| 2 | 4.2 | 18.5 | 4.2 | 17.9 | 708 | 3.7 | 18.4 | 2,027 | 4.0 | 17.8 | 6,756 |
| 3 | 4.4 | 18.4 | 4.5 | 18.7 | 719 | 4.2 | 19.2 | 2,207 | 4.3 | 18.5 | 6,987 |
| 4 | 4.3 | 19.6 | 4.4 | 19.5 | 674 | 4.3 | 19.9 | 2,160 | 4.0 | 19.1 | 6,327 |
| 5 | 5.0 | 20.2 | 5.1 | 20.7 | 737 | 4.7 | 20.6 | 2,282 | 4.8 | 20.1 | 7,184 |
| 6 | 4.5 | 20.9 | 4.8 | 21.1 | 685 | 4.5 | 21.2 | 2,124 | 4.7 | 21.0 | 6,771 |
| 7 | 4.2 | 21.0 | 4.6 | 21.2 | 645 | 4.0 | 21.5 | 1,871 | 4.0 | 21.1 | 5,757 |
| 8 | 4.7 | 21.4 | 4.3 | 21.5 | 595 | 3.8 | 21.5 | 1,789 | 3.9 | 20.8 | 5,654 |
| 9 | 4.5 | 21.4 | 4.2 | 21.5 | 587 | 3.9 | 21.7 | 1,801 | 4.1 | 21.3 | 5,844 |
| 10 | 4.4 | 21.8 | 4.4 | 21.6 | 608 | 3.9 | 21.9 | 1,797 | 4.2 | 21.6 | 5,798 |
| 11 | 4.4 | 22.4 | 4.6 | 22.4 | 616 | 4.2 | 22.4 | 1,888 | 4.3 | 21.8 | 5,890 |
| 12 | 4.7 | 22.6 | 4.8 | 22.3 | 644 | 4.3 | 22.6 | 1,895 | 4.7 | 22.3 | 6,271 |
| 13 | 4.4 | 23.0 | 4.9 | 23.1 | 633 | 4.6 | 23.2 | 1,995 | 4.5 | 22.8 | 5,940 |
| 14 | 4.5 | 23.2 | 4.5 | 23.6 | 574 | 4.6 | 23.5 | 1,937 | 4.5 | 22.9 | 5,873 |
| 15 | 4.4 | 23.6 | 4.9 | 24.2 | 605 | 4.4 | 23.7 | 1,862 | 4.2 | 22.9 | 5,543 |
| 16 | 4.9 | 24.1 | 4.9 | 25.2 | 578 | 4.6 | 24.2 | 1,906 | 4.8 | 23.6 | 6,146 |
| 17 | 4.7 | 24.3 | 4.8 | 25.1 | 578 | 4.6 | 24.2 | 1,919 | 4.7 | 23.7 | 5,980 |
| 18 | 5.1 | 24.7 | 5.1 | 25.6 | 601 | 4.8 | 25.0 | 1,912 | 4.7 | 24.1 | 5,874 |
| 19 | 4.9 | 24.8 | 4.6 | 25.8 | 531 | 4.7 | 24.9 | 1,888 | 4.8 | 24.6 | 5,855 |
| 20 | 4.9 | 25.1 | 4.6 | 26.1 | 526 | 4.5 | 25.5 | 1,784 | 4.8 | 24.9 | 5,816 |
| 21 | 5.0 | 25.4 | 4.8 | 26.4 | 549 | 4.4 | 25.7 | 1,720 | 4.7 | 25.3 | 5,575 |
| 22 | 4.6 | 25.6 | 4.6 | 26.8 | 517 | 4.5 | 26.1 | 1,708 | 4.7 | 25.5 | 5,483 |
| 23 | 4.8 | 26.1 | 5.0 | 27.0 | 550 | 4.4 | 26.3 | 1,682 | 5.0 | 25.6 | 5,878 |
| 24 | 4.3 | 26.1 | 4.7 | 26.9 | 522 | 4.5 | 26.6 | 1,679 | 4.6 | 26.0 | 5,317 |
| 25 | 4.6 | 26.4 | 4.6 | 26.9 | 516 | 4.4 | 26.7 | 1,647 | 4.5 | 26.3 | 5,190 |
| 26 | 4.6 | 26.5 | 4.8 | 27.1 | 535 | 4.6 | 26.5 | 1,721 | 4.9 | 26.3 | 5,568 |
| 27 | 4.6 | 26.7 | 4.4 | 27.3 | 479 | 4.5 | 27.2 | 1,657 | 4.8 | 26.9 | 5,391 |
| 28 | 4.7 | 26.9 | 4.6 | 27.8 | 495 | 4.4 | 27.4 | 1,596 | 4.7 | 27.4 | 5,167 |
| 29 | 4.5 | 27.0 | 4.4 | 28.0 | 475 | 4.5 | 28.0 | 1,618 | 4.7 | 27.1 | 5,150 |
| 30 | 4.7 | 28.2 | 4.8 | 28.6 | 501 | 4.4 | 28.5 | 1,548 | 5.0 | 28.0 | 5,357 |
| 31 | 4.7 | 27.8 | 4.9 | 28.6 | 512 | 4.7 | 28.5 | 1,634 | 4.7 | 27.9 | 5,020 |
| 32 | 4.7 | 28.1 | 4.9 | 27.9 | 531 | 4.6 | 28.5 | 1,622 | 4.7 | 27.5 | 5,127 |
| 33 | 4.7 | 27.5 | 5.1 | 28.0 | 546 | 4.9 | 28.4 | 1,718 | 5.3 | 27.9 | 5,736 |
| 34 | 4.6 | 27.7 | 5.1 | 27.8 | 552 | 4.6 | 28.3 | 1,641 | 4.8 | 27.5 | 5,274 |
| 35 | 4.8 | 27.6 | 5.3 | 27.3 | 581 | 4.7 | 27.7 | 1,690 | 4.7 | 26.8 | 5,284 |
| 36 | 4.4 | 27.3 | 4.8 | 28.1 | 510 | 4.7 | 28.2 | 1,668 | 4.6 | 27.1 | 5,049 |
| 37 | 4.8 | 27.5 | 5.0 | 28.0 | 539 | 4.7 | 28.0 | 1,676 | 4.7 | 27.6 | 5,054 |
| 38 | 4.6 | 27.7 | 4.5 | 28.1 | 483 | 4.8 | 27.9 | 1,708 | 4.8 | 27.6 | 5,267 |
| 39 | 4.5 | 27.5 | 4.9 | 28.3 | 521 | 4.6 | 28.2 | 1,617 | 4.9 | 28.0 | 5,213 |
| 40 | 4.7 | 28.3 | 4.9 | 28.8 | 514 | 4.8 | 28.8 | 1,677 | 4.7 | 28.3 | 4,968 |
| Mean for weeks | | | | | | | | | | | |
| 1-13 | 4.5 | 20.9 | 4.6 | 21.0 | 654 | 4.2 | 21.2 | 1,986 | 4.3 | 20.7 | 6,265 |
| 14-40 | 4.7 | 26.4 | 4.8 | 27.0 | 534 | 4.6 | 26.8 | 1,720 | 4.7 | 26.2 | 5,450 |

^a Grams of feed consumed per animal per day

^b Milligrams of acesulfame potassium consumed per kilogram body weight per day

