

Table I-26
AGE-ADJUSTED U.S. DEATH RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

Both Sexes

	All Races		White		Black	
	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004
All Sites	192.7	-1.2*	190.7	-1.1*	238.8	-1.8*
Lung and Bronchus	54.7	-1.0*	55.0	-0.8*	62.0	-1.6*
Colon and Rectum	19.4	-2.2*	18.9	-2.3*	26.7	-1.5*
Breast	14.5	-2.4*	14.1	-2.5*	21.9	-3.6*
Pancreas	10.6	0.1*	10.4	0.3*	20.1	-1.7*
Prostate ^f	10.5	-3.6*	9.6	-3.5*	13.8	-0.6*
Non-Hodgkin Lymphoma	7.6	-2.5*	7.9	-2.5*	8.2	-3.3*
Leukemia	7.5	-0.8*	7.7	-0.7*	7.1	-1.4*
Ovary ^f	5.0	-0.3	5.2	-0.3	6.7	-1.1*
Liver & IBD ^g	4.9	1.7*	4.8	-0.9*	6.5	1.3*
Brain and ONS ^g	4.4	-1.0*	4.5	0.0	6.0	-4.2*
Esophagus	4.4	0.3*	4.5	1.7*	5.2	-2.4*
Urinary Bladder	4.3	-0.2	4.3	1.2*	4.5	0.0
Stomach	4.2	-3.2*	4.3	-0.3	4.3	0.4
Kidney and Renal Pelvis	4.2	-0.4	3.7	-3.2*	4.1	-0.7*
Myeloma	3.7	-0.9*	3.5	-0.8*	3.9	-3.8*

	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004
All Sites	115.5	-1.8*	160.4	-0.3	129.1	-1.3*
Lung and Bronchus	26.9	-1.4*	39.9	-0.1	23.6	-1.6*
Colon and Rectum	12.3	-2.0*	17.0	-1.4	13.6	-0.9*
Liver & IBD ^g	10.6	-0.6	9.0	0.1	9.0	-2.6*
Stomach	8.0	-4.6*	8.4	2.2	8.3	-3.1*
Pancreas	7.4	-0.2	8.4	-2.2	8.3	0.2
Breast	7.0	-0.2	7.5	1.0	7.6	1.4*
Non-Hodgkin Lymphoma	4.8	-2.1*	7.2	-0.6	6.8	-2.1*
Prostate ^f	4.6	-5.6*	6.5	0.6	5.7	-3.1*
Leukemia	3.9	-1.9*	5.5	0.5	5.1	-1.2*
Ovary ^f	2.7	0.8	4.8	0.3	3.6	0.5
Oral Cavity and Pharynx	2.3	-2.6*	4.0	2.0	3.4	0.3
Brain and ONS ^g	2.0	0.5	3.8	-2.0	3.1	-0.7
Esophagus	1.8	-2.6	3.8	4.5*	2.8	-0.3
Urinary Bladder	1.8	-0.1	2.5	-0.3	2.4	0.2
Kidney and Renal Pelvis	1.7	-3.0*	2.2	-2.0	2.4	-2.2*

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2000-2004 age-adjusted rates for the race/ethnic group.

^b Mortality data used in calculating the rates are analyzed from a public use file provided by the National Center for Health Statistics (NCHS). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). The rates shown for sex-specific cancer sites are calculated using the population for both sexes combined.

^c The APC is the Annual Percent Change over the time interval. Mortality data used in calculating the trends are analyzed from a public use file provided by the National Center for Health Statistics (NCHS). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. The 2000-2004 Hispanic death rates do not include deaths from Minnesota, New Hampshire and North Dakota. The 1995-2004 Hispanic mortality trends do not include deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^f The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^g IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero ($p < .05$).