Surgical Operations in Short-Stay Hospitals

UNITED STATES-1973

Statistics are presented on the number and rate of inpatients with surgery discharged from non-Federal short-stay hospitals and the types of surgical operations performed. The data for this report are based on information abstracted by means of the Hospital Discharge Survey from a national sample of the hospital records for discharged inpatients. Estimates of the number of patients with surgery and of operations, grouped by surgical classes and categories, are shown by the demographic characteristics of the operated patients and by geographic region and size of hospital. The number and average length of stay is shown for first-listed operations by age and sex.

DHEW Publication No. (HRA) 76-1775

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
Health Resources Administration
National Center for Health Statistics
Rockville, Md. May 1976



Library of Congress Cataloging in Publication Data

Ranofsky, Abraham L

Surgical operations in short-stay hospitals, United States, 1973.

(Vital and health statistics: Series 13, Data from the National Health Survey; no. 24) (DHEW publication; no. (HRA) 76-1775)

Bibliography: p.

1. Operations, Surgical—United States—Statistics. 2. Hospital utilization—United States—Statistics. I. United States. National Center for Health Statistics. II. Title. III. Series: United States. National Center for Health Statistics. Vital and health statistics: Series 13, Data from the National Health Survey; no. 24. IV. Series: United States. Dept. of Health, Education, and Welfare. DHEW publication; no. (HRA) 76-1775.

RA407.3.A349

no. 24

[RA981.A2]

362.1'1'0973s

76-608026 [362.1'9'70973]

ISBN 0-8406-0069-0

NATIONAL CENTER FOR HEALTH STATISTICS

DOROTHY P. RICE, Director

ROBERT A. ISRAEL, Deputy Director
JACOB J. FELDMAN, Ph.D., Associate Director for Analysis
GAIL F. FISHER, Associate Director for the Cooperative Health Statistics System
ELIJAH L. WHITE, Associate Director for Data Systems
GEORGE P. FAILLA, Associate Director for Management
PETER L. HURLEY, Associate Director for Operations
JAMES M. ROBEY, Ph.D., Associate Director for Program Development
ALICE HAYWOOD, Information Officer

DIVISION OF HEALTH RESOURCES UTILIZATION STATISTICS

SIEGFRIED A. HOERMANN, Director
BEULAH K. CYPRESS, Ph.D., Chief, Family Planning Statistics Branch
W. EDWARD BACON, Ph.D., Chief, Hospital Care Statistics Branch
JAMES E. DeLOZIER, Chief, Ambulatory Care Statistics Branch
MANOOCHEHR K. NOZARY, Chief, Technical Services Branch
JOAN F. VAN NOSTRAND, Chief, Long-Term Care Statistics Branch

COOPERATION OF THE BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Center for Health Statistics, the Bureau of the Census, under a contractual arrangement, participated in planning the survey and collecting the data.

Vital and Health Statistics-Series 13-No. 24

DHEW Publication No. (HRA) 76-1775

Library of Congress Catalog Card Number 76-608026

CONTENTS

	Page
Introduction	1
Discharges with Surgery	2
Age and Sex	3
Color	4
Geographic Region	5
Size of Hospital	5
Surgical Operations by Patient and Hospital Characteristics	7
Surgery by Age and Sex	7
Surgery by Color	15
Surgery by Geographic Region	16
Surgery by Bed Size of Hospital	17
First-Listed Operations and Average Length of Stay	19
References	22
List of Detailed Tables	23
Appendix I. Technical Notes on Methods	46
Statistical Design of the Hospital Discharge Survey	46
Data Collection and Processing	46
Presentation of Estimates	50
Reliability of Estimates	50
Appendix II. Definitions of Certain Terms Used in This Report	53
Terms Relating to Hospitalization	53
Terms Relating to Surgery	53
Demographic Terms	54

SURGICAL OPERATIONS IN SHORT-STAY HOSPITALS

Abraham L. Ranofsky, Division of Health Resources Utilization Statistics

INTRODUCTION

This report, based on data collected by the Hospital Discharge Survey (HDS), presents estimates of the number of inpatients with surgery discharged from non-Federal short-stay hospitals during 1973 and measurements of the volume of surgical operations or procedures performed. This survey of short-stay hospital utilization has been conducted by the National Center for Health Statistics (NCHS) continuously since the beginning of 1965.

Data for the survey are abstracted from medical records of discharges selected from a sample of short-stay hospitals located in the 50 States and the District of Columbia. The sample for 1973 included about 225,000 medical records for inpatients discharged from 424 hospitals that participated in the survey. Data for newborn infants are excluded from this report. See appendix I for a description of the sample design, data collection procedures, and the estimation process. A detailed report on the design of the HDS has been published. 1

Demographic, diagnostic, and surgical data are abstracted from the medical record face sheet for each patient in the sample. A maximum of five diagnoses and three operations are coded for each medical record. Coding of the medical data is done according to the Eighth Revision International Classification of Diseases, Adapted for Use in the United States² (ICDA), with some modifications. Included with surgery are biopsies, exploratory laparotomies, and certain other diagnostic procedures. Excluded from the definition of surgery are certain obstetrical procedures,

diagnostic endoscopy and radiography, and certain other nonsurgical procedures.

Familiarity with the definition of surgery as used in this report is important for interpreting the data and for making comparisons with statistics on surgery which are available from other data collection sources. The HDS modifications of the ICDA section on Surgical Operations, Diagnostic and Other Therapeutic Procedures are described in appendix I under "Data Collection and Processing." Definitions of the terms used in this study relating to hospitalization and characteristics of inpatients and hospitals are presented in appendix II.

The data for patients with surgery and the volume of all-listed operations (a maximum of three operations coded for each medical record) are shown in this report by age, sex, and color of inpatients, and by the geographic regions and bed sizes of hospitals where the medical care was provided. The data presented on surgery are grouped by the broad surgical classes of the ICDA and by surgical categories which represent more detailed information on specific operations or groups of operations. The surgery performed in short-stay hospitals is measured by frequencies, percent distributions, and rates of surgery in the civilian noninstitutionalized population. In addition, for the first-listed operations on the medical record face sheets, estimates of the number of operations and average lengths of stay are provided for each surgical class and category by age and sex.

Data from the HDS on short-stay hospital utilization by surgical operations have been published in previous reports for 1965, 1968,

1971, and 1972. ³⁻⁶ Information on diseases and injuries diagnosed for patients discharged from short-stay hospitals has been published for each year from 1965 to 1972, with the exception of 1969 and 1970. ⁷⁻¹¹

Another program of NCHS, the Health Interview Survey (HIS), also collects information on hospitalization. The estimates provided by HIS are generally smaller for number of discharges and longer for average lengths of stay than HDS because of differences in collection procedures, population sampled, and definitions. Data from HIS are published by NCHS in Series 10 of Vital and Health Statistics reports.

DISCHARGES WITH SURGERY

One or more surgical procedures were performed for an estimated 13.3 million, or 41.3 percent, of the 32.1 million inpatients discharged from non-Federal short-stay hospitals during 1973 (table A). Patients with surgery included 5.2 million males and 8.1 million females. There were 18.4 million surgical operations performed for the 13.3 million patients with surgery, or an average of 1.4 operations per patient. Males accounted for 6.9 million operations and females 11.5 million. (Table 1.)

Table A. Number and rate of inpatients discharged from short-stay hospitals with and without surgery and percent with surgery, by age and sex: United States, 1973

		of disch thousand		Percent of dis- charges	- 1,000 population				
Sex and age	Total	Without surgery	With surgery	with surgery	Total	Without surgery	With surgery		
Total ¹							•		
A11 ages	32,125	18,859	13,266	41.3	156.1	91.6	64.4		
Under 15 years	3,933 13,482 7,772 6,937	2,073 7,445 4,483 4,858	1,861 .6,037 3,290 2,079	47.3 44.8 42.3 30.0	70.8 154.4 182.3 341.8	37.3 85.2 105.1 239.4	33.5 69.1 77.1 102.4		
<u>Male</u>									
All ages	12,835	7,667	5,168	40.3	129.2	77.2	52.0		
Under 15 years	2,231 3,873 3,637 3,094	1,182 2,122 2,247 2,116	1,048 1,751 1,390 978	47.0 45.2 38.2 31.6	78.8 91.7 179.1 367.0	41.8 50.2 110.6 251.0	37.0 41.4 68.5 116.0		
<u>Female</u>	}								
All ages	19,266	11,175	8,091	42.0	180.9	104.9	76.0		
Under 15 years	1,700 9,598 4,131 3,837	889 5,314 2,233 2,739	811 4,283 1,898 1,099	47.7 44.6 45.9 28.6	62.4 212.9 185.0 323.5	32.6 117.9 100.0 230.9	29.8 95.0 85.0 92.6		

¹Includes data for sex not stated.

Age and Sex

The estimated number of inpatients with surgery per 1,000 persons in the civilian noninstitutionalized population increased consistently with age from 33.5 for under 15 years of age to 102.4 for those 65 years and over (table A). This pattern of surgery by age was also exhibited for males, but for females the rate of discharges with surgery was highest for the age group 15-44 years.

There was less of an increase with age in the discharge rates for patients with surgery than without surgery. Rates of discharges with surgery were about three times higher for the oldest (65 years and over) than the youngest (under age

15) age groups for both males and females, but for inpatients without surgery the increases in the rates were sixfold for males and sevenfold for females. When compared with the young ages, older patients accounted for larger proportions of the population hospitalized for chronic illnesses such as heart disease, cerebrovascular disease, diabetes, emphysema, and osteoarthritis which are not generally treatable by surgery (data from tabulations).

Of the estimated 13.3 million inpatients with surgery, 9.2 million patients or 69.6 percent had one operation, 2.9 million patients or 22.0 percent had two operations, and 1.1 million patients or 8.5 percent had three operations (table B). The

Table B. Number and percent distribution of inpatients discharged from short-stay hospitals with surgery by number of operations, according to age and sex: United States, 1973

Sex and age	All dis- charges with	Number of operations			All dis- charges with		mber o	_
	surgery	One	Two	Three	surgery	One	Two	Three
Total ¹	Number	Number of discharges in thousands				nt distr	ibutio	n
All ages	13,266	9,230	2,912	1,124	100.0	69.6	22.0	8.5
Under 15 years 15-44 years 45-64 years 65 years and over	1,861 6,037 3,290 2,079	1,447 4,190 2,118 1,474	362 1,310 807 434	52 537 365 171	100.0 100.0 100.0 100.0	77.8 69.4 64.4 70.9	19.4 21.7 24.5 20.9	2.8 8.9 11.1 8.2
All ages	5,168	3,746	1,076	346	100.0	72.5	20.8	6.7
Under 15 years 15-44 years 45-64 years 65 years and over <u>Female</u>	1,048 1,751 1,390 978	792 1,282 987 684	224 342 292 218	32 127 111 76	100.0 100.0 100.0 100.0	75.6 73.2 71.0 69.9	21.4 19.5 21.0 22.3	3.1 7.3 8.0 7.7
All ages	8,091	5,479	1,835	777	100.0	67.7	22.7	9.6
Under 15 years 15-44 years 45-64 years 65 years and over	811 4,283 1,898 1,099	654 2,907 1,130 788	138 968 515 215	20 409 253 95	100.0 100.0 100.0 100.0	80.6 67.9 59.5 71.8	17.0 22.6 27.1 19.6	2.4 9.5 13.3 8.7

¹Includes data for sex not stated.

Table C. Number of inpatients discharged from short-stay hospitals with and without surgery and percent with surgery, according to color: United States, 1973

Color	Number ir	Percent of dis-		
	Total	Without surgery	With surgery	charges with surgery
Tota1	32,125	18,859	13,266	41.3
White	24,402 3,619 4,105	14,309 2,228 2,322	10,092 1,391 1,783	41.4 38.4 43.4

proportion of operated patients with multiple operations was smallest for under age 15 (22.2 percent) and largest for the age group 45-64 (35.6 percent). The percent of males with multiple operations ranged from 24.4 percent for under age 15 to 30.1 percent for age 65 years and over and for females the range was from 19.4 percent for under age 15 to 40.5 percent for the age group 45-64 years.

Color

Inpatients are presented in this report by the color groups "white" and "all other." Of the 32.1 million inpatients discharged from short-stay hospitals during 1973, an estimated 24.4 million

patients were identified on the medical records as white and 3.6 million as all other color groups (table C). Color was not stated for 4.1 million, or about 12.8 percent of the patients hospitalized. In view of the large number of patients for whom color was not stated, surgery rates were not computed and caution should be used in drawing conclusions from the data by color.

The proportions of the patients discharged with surgery by color were 41.4 percent for white and 38.4 percent for all other patients (table C). The percent distributions of discharges with surgery by the number of operations per patient were about the same for patients identified as white and all other (table D).

Table D. Number and percent distribution of inpatients discharged from short-stay hospitals with surgery by number of operations, according to color: United States, 1973

Color	All dis- charges		umber o		All dis- charges		umber o peratio		
	with		Two	Three	with surgery	One	Two	Three	
	Number of discharges in thousands				Percent distribution				
Tota1	13,266	9,230	2,912	1,124	100.0	69.6	22.0	8.5	
WhiteAll otherColor not stated	10,092 1,391 1,783	6,995 993 1,242	2,231 286 395	866 112 146	100.0 100.0 100.0	69.3 71.4 69.7	22.1 20.6 22.2	8.6 8.0 8.2	

Geographic Region

The number of inpatients with a surgical procedure discharged from short-stay hospitals during 1973 varied by geographic region from an estimated 2.3 million in the West Region to 4.2 million in the North Central Region (table E). These differences in number of operations are due principally to variations among the regions in the number of persons in the civilian noninstitutionalized population and partially to variations in the surgery rates. Populations ranged from 35.6 million in the West Region to 64.5 million in the South Region (appendix I, table II), and the number of patients discharged with surgery per 1,000 population ranged from 55.7 in the South Region to 73.1 in the North Central Region. The North Central Region with a higher rate of patients with surgery than the South ranked first and the South Region ranked second in number of operated patients, although the population size of the South Region was larger than that of the North Central Region.

The proportions of patients with or without surgery during episodes of hospitalization fluctuated by region. Operated patients as a percent of all patients discharged ranged from a low of 36.3 percent in the South Region to a high of 45.4 percent in the West Region.

Variations were small among the regions

in the percent distributions of the estimated number of discharges with surgery by the number of operations per patient (table F). In every region about 70 percent of the patients had one surgical operation or procedure and approximately 30 percent had multiple operations during an episode of hospitalization.

Size of Hospital

Surgery was reported for higher proportions of the patients discharged from the larger than the smaller hospitals (table G). The percent of patients with surgery increased from 29.0 percent in hospitals with 6-99 beds to 47.6 percent in hospitals with 500 beds or more. It is likely that larger hospitals account for higher proportions of patients with surgery because they are more apt to have the specialized personnel, equipment, and facilities required for performing complex surgery.

There were small differences in the percent distributions of patients with surgery by number of operations per patient in hospitals of various size groups (table F). As was shown previously for the other variables discussed, differences were relatively small in the distributions of surgical operations per patient by color and geographic region, but varied considerably by sex and age.

Table E. Number and rate of inpatients discharged from short-stay hospitals with and without surgery and percent with surgery, by geographic region: United States, 1973

Geographic region		of disch		Percent of dis-	Disch 1,00	arge rate O populat	per ion
Geographic region	Total	Without With surgery surger		charges with surgery	Total	Without surgery	With surgery
Total	32,125	18,859	13,266	41.3	156.1	91.6	64.4
Northeast North Central South West	7,208 9,975 9,905 5,036	3,972 5,824 6,311 2,752	3,237 4,151 3,594 2,284	44.9 41.6 36.3 45.4	147.3 175.7 153.6 141.4	81.2 102.6 97.8 77.3	66.1 73.1 55.7 64.1

Table F. Number and percent distribution of inpatients discharged from short-stay hospitals with surgery by number of operations, according to geographic region and bed size of hospital: United States, 1973

	All dis-		Number o	_ ,	All dis-	1	mber o	_
Geographic region and bed size of hospital	charges with surgery	One	Two	Three	charges with surgery	One	Two	Three
	Number	Number of discharges in thousands				t dist	ributi	.on
Total	13,266	9,230	2,912	1,124	100.0	69.6	22.0	8.5
Geographic region Northeast	3,237 4,151 3,594 2,284	2,250 2,874 2,499 1,607	712 924 777 500	275 354 319 177	100.0 100.0 100.0 100.0	69.2	22.0 22.3 21.6 21.9	8.5 8.5 8.9 7.7
6-99 beds	1,917 2,102 2,459 3,842 2,946	1,364 1,479 1,689 2,625 2,073	384 454 552 894 628	169 169 218 323 244	100.0 100.0 100.0 100.0 100.0	71.2 70.4 68.7 68.3 70.4	20.0 21.6 22.5 23.3 21.3	8.8 8.0 8.9 8.4 8.3

Table G. Number of inpatients discharged from short-stay hospitals with and without surgery and percent with surgery, according to bed size of hospital: United States, 1973

Bed size of hospital	Number in	Percent of dis- charges		
ned Size of Mospital	Total	Without surgery	With surgery	with surgery
Total	32,125	18,859	13,266	41.3
6-99 beds	6,604 5,485 5,625 8,228 6,183	4,687 3,383 3,166 4,386 3,237	1,917 2,102 2,459 3,842 2,946	29.0 38.3 43.7 46.7 47.6

SURGICAL OPERATIONS BY PATIENT AND HOSPITAL CHARACTERISTICS

Almost half (47.0 percent) of the surgical operations performed in short-stay hospitals during 1973 were related to the specialties of gynecological surgery (19.4 percent), abdominal surgery (14.9 percent), and orthopedic surgery (12.8 percent). About 37.6 percent of the surgical operations were for males and 62.3 percent for females. Gynecological surgery accounted for a major proportion of the difference in the volume of surgery by sex.

The estimates of surgical operations reported for inpatients discharged from short-stay hospitals are grouped in the detailed tables of this report by the surgical classes or specialties of the ICDA. The surgical operations and the ICDA codes comprising each class are presented in the detailed tables by categories which represent single surgical procedures which occur in large frequencies or groups of associated surgical operations. Unpublished data are available from HDS for most of the surgical codes listed in the ICDA for which the estimates are large enough to meet the standards of reliability.

The surgical categories of surgical operations and procedures with the largest frequencies are presented for the variables age, sex, color, geographic region, and bed size of hospitals in text tables H-O. Residual categories of operations are excluded. The text tables on surgery by age are cross-tabulated by sex. Similar cross-tabulations are not shown in the detailed tables.

Surgery by Age and Sex

An estimated 18.4 million operations were performed during 1973 for the 13.3 million inpatients with surgery (table 1). The corresponding rate was 89.5 operations per 1,000 persons in the civilian noninstitutionalized population (rates in tables are shown as per 100,000 population to accommodate small estimates). There were 6.9 million operations for males and 11.5 million for females. The rates of operations were 69.8 and 107.8 per 1,000 population, respectively, or about 54 percent higher for females than for males. Sex-specific surgical

procedures for females accounted for 4,830,000 operations compared with 682,000 sex-specific operations for males (data from tabulations). Exclusive of the sex-specific operations, the rates per 1,000 population were about the same for males (63.0) and for females (62.4).

As shown for all patients, surgical operations by sex were also clustered into a few of the 17 ICDA surgical classes. Over half (52 percent) of the operations for males were in the specialties abdominal, orthopedic, and urological surgery; for females over half (53 percent) of the operations were in the specialties gynecological, abdominal, and orthopedic surgery. The annual rates of surgery for the ICDA surgical classes by sex are shown in figure 1.

The number and rate for surgical operations reported most frequently during 1973 are shown in table H by surgical category and sex. Diagnostic dilation and curettage of uterus was the most frequently reported surgical procedure. This procedure accounted for 934,000 operations and an annual rate of 4.5 operations per 1,000 persons in the civilian noninstitutionalized population. The rate was 8.8 based on the female population. Other leading surgical procedures were biopsy, tonsillectomy with or without adenoidectomy, hysterectomy, and repair of inguinal hernia. The rates for these operations per 1,000 population were 4.5. 4.3. 3.4, and 2.6, respectively.

The operations for males with the largest frequencies were repair of inguinal hernia (464,000 operations), tonsillectomy with or without adenoidectomy (398,000 operations), and biopsy (300,000 operations). The corresponding rates per 1,000 males were 4.7, 4.0, and 3.0, respectively. For females the operations with the largest frequencies were diagnostic dilation and curettage of uterus (934,000 operations), hysterectomy (690,000 operations), and biopsy (618,000 operations). The corresponding rates per 1,000 females were 8.8, 6.5, and 5.8, respectively.

There were large variations by sex for many non-sex-specific surgical operations (table 1). The rates for males were larger than for females by about eight times for repair of inguinal hernia, by about three times for meatotomy, and by two times for lung lobectomy or pneumonectomy. Patients with breast surgery were predominantly females (94 percent). Other non-




Figure 1. Annual rate of all-listed operations for inpatients discharged from short-stay hospitals, excluding newborn infants, by surgical class and sex: United States, 1973

Table H. Number and rate of all-listed operations for inpatients of all ages discharged from short-stay hospitals, by selected surgical categories and sex: United States, 1973

Surgical category and ICDA codes	Number in	of opera thousand	itions ls		operatio 00 popula	
TODA Codes	Both sexes ¹	Male	Female	Both sexes ¹	Male	Female
All operations ²	18,426	6,936	11,480	8,951.8	6,984.7	10,776.5
Dilation and curettage of uterus, diagnostic70.3 Biopsy	934 919 884 690 525 455 425 411 339 321 307 299 284 279 273 249 246 227 218	300 398 464 205 175 174 169 249 82	934 618 486 690 61 250 425 318 163 147 138 299 164 162 273 246 145	453.8 446.6 429.6 335.2 255.3 221.2 206.5 199.8 164.5 156.0 149.1 145.5 137.9 135.6 132.6 120.8 119.6 110.4	302.3 400.9 467.1 206.3 94.4 176.7 174.8 169.9 120.0 116.8 250.4 82.4	876.8 580.0 455.9 647.7 57.5 234.7 399.1 298.1 153.0 138.1 129.8 281.1 154.4 152.3 256.2 231.1 136.0
Plastic repair of cystocele and/or rectocele71.4	204	111	107 204	106.0 99.3	111.8	100.5

Includes data for sex not stated.
Includes data for surgery not shown in table.
Itimited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery.

Table J. Number and rate of biopsies performed for inpatients discharged from shortstay hospitals, by body site and sex: United States, 1973

Biopsy and ICDA codes	Number of biopsies Rate of biopsies per in thousands 100,000 population					
	Both sexes ¹	Male	Female	Both sexes ¹	Male	Female
All biopsiesAl-A2	919	300	618	446.6	302.3	580.0
Biopsy of breastA2.3 Biopsy of internal female genital	186	5	181	90.4	4.7	169.8
organsA2.5 Biopsy of biliary tract, spleen,	177	•••	177	86.1	•••	166.4
and pancreasA1.9 Biopsy of boneA2.7 Other biopsy of circulatory and	94 90	45 42	48 48	45.6 43.6		45.2 45.0
lymphatic systemsA2.6 Biopsy of thoraxA1.6 Biopsy of stomach and intestinesA1.8	68 61 57	34 37 30	33 24 27	29.8 27.5		31.1 22.6 25.2
Biopsy of male genital organsA2.2 Biopsy of mouth and throatA1.4 Biopsy of skin and subcutaneous	33 31	33 20	11	15.8 15.0	32.8 20.0	10.3
tissueA2.9 Biopsy of urinary tractA2.1 Other biopsy of musculoskeletal	29 25	15 13	15 12	14.2 12.1	14.8 12.8	13.8 11.5
systemA2.8 Other biopsiesResidual	19 50	9 18	10 32	9.1 24.5	9.1 18.4	9.1 30.1

¹Includes data for sex not stated.

sex-specific operations with higher rates for females than for males were thyroidectomy by about five times, cholecystectomy by about three times, and excision and ligation of varicose veins by about three times. There were also many non-sex-specific surgical procedures such as operations on valves of heart, nephrectomy, and hemorrhoidectomy for which surgical rates were about the same for males and females.

More than twice as many biopsies were performed in short-stay hospitals for females (618,000 biopsies) than for males (300,000 biopsies) during 1973. The number and rate of biopsies reported are shown in table J by body site and sex. By body site, about two-fifths of all the biopsies were of the breast (20.2 percent) and of internal female genital organs (19.3 percent). For female patients, almost three-fifths (58.0 percent) of the biopsies were for these two body sites.

The age group 15-44 years accounted for 8.4 million surgical operations, or about 46 percent of all the operations performed in shortstay hospitals in 1973 (table 2). However, the number of operations per 1,000 persons in the civilian noninstitutionalized population increased consistently for each older age group from 41.9 for under 15 years of age to 140.7 for age 65 years and over (table 3). This pattern, however, varied considerably among the surgical categories. For example, the rates of operations increased with age for extraction of lens, cholecystectomy, and prostatectomy, but decreased with age for resection and recession of eve muscle, myringotomy, and tonsillectomy with adenoidectomy. As would be expected, almost all surgery concerned with childbearing such as the obstetrical procedures and bilateral ligation and division of fallopian tubes were reported for women in the age group 15-44 years. Surgical rates

for operations such as hemorrhoidectomy and diagnostic dilation and curettage of uterus were lower for the youngest and oldest age groups, but the rates were highest in these age groups for urethral meatotomy and closed reduction of fracture without fixation.

The numbers and rates for operations with large frequencies are presented by age and sex in tables K-N. The surgical categories shown in these tables were selected by frequency from detailed tables 2 and 3.

Age group under 15 years. - Tonsillectomy with or without adenoidectomy was the leading surgical category for patients under age 15. These procedures accounted for an estimated 647,000 operations, or over a fourth (27.8 percent) of all operations for patients under age 15 (table K). The corresponding rate was 11.6 per 1,000 population. Other leading surgical categories in this age group were myringotomy, repair of inguinal hernia, closed reduction of fracture without fixation, and appendectomy. The

Table K. Number and rate of all-listed operations for inpatients under 15 years of age discharged from short-stay hospitals, by selected surgical categories and sex: United States, 1973

Symptocal actoropy and TCDA codes	Number of operations Rate of operations per in thousands 100,000 population					
Surgical category and ICDA codes	Both sexes ¹	Male	Fe- male	Both sexes ¹	Male	Female
All operations, under 15 years of age ²	2,326	1,337	988	4,185.8	4,720.6	3,626.0
Tonsillectomy with or without adenoidectomy	647 198 117 103 103 61 58 58 53 52 46 34 30 28	317 116 101 67 59 35 27 8 53 29 29 34 20 24	330 82 15 36 44 26 31 50 23 17 	1,164.9 356.2 210.4 185.8 185.3 110.1 105.2 105.1 94.6 93.4 82.4 60.7 54.6 50.8	1,119.3 408.1 358.1 236.7 207.4 124.7 97.1 29.5 185.7 100.7 102.9 119.2 70.1 83.1	1,212.2 301.9 56.1 132.2 162.4 94.9 113.5 183.7 85.0 61.2
Incision of skin and subcutaneous tissue92.0	24	15	9	43.2	53.2	32.8

 $^{^1}$ Includes data for sex not stated. 2 Includes data for surgery not shown in table. 3 Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery.

rates for these operations were 3.6, 2.1, 1.9, and 1.9, respectively. The five leading surgical categories represented about half (50.2 percent) of all the operations reported for this age group.

The rate of operations for males under age 15 was 47.2 per 1,000 population, or about 30 percent higher than the rate of 36.2 for females. Of the leading operations for patients under 15 years of age the largest differences in surgical rates by sex were for repair of inguinal hernia with a rate of 3.6 for males compared with 0.6 for females and for dilation of urethra with a rate of 0.3 for males compared with 1.8 for females.

Age group 15-44 years. - The estimated surgical rate for persons aged 15-44 years was 96.4 operations per 1,000 population (table L). The rate for women was 134.6 compared with 55.6 for men, or almost two and a half times higher. Surgery in this age group was dominated by the sex-specific specialties of gynecological and obstetrical procedures (table 2). These spe-

Number and rate of all-listed operations for inpatients 15-44 years of age discharged from short-stay hospitals, by selected surgical categories and sex: United States, 1973

[Excludes newborn infants and Federal hospitals] Number of operations Rate of operations per in thousands 100,000 population Surgical category and ICDA codes Both sexes1 Both Male Female Male Female sexes1 All operations, 15-44 8,420 2,347 6,069 9,640.1 5,555.7 13,459.4 years of age²-----Dilation and currettage of 1,320.2 uterus, diagnostic ----- 70.3 595 595 681.5 Hysterectomy-----69.1-69.5 415 415 474.7 919.5 63 259 368.0 148.0 574.0 321 Biopsy----A1-A2 Ligation and division of fallopian tubes bilateral)-----68.5 295 295 337.9 654.5 Dilation and currettage after delivery or abortion-----78.1 269 269 308.2 597.0 . . . Cesarean section-----77 244 244 279.9 542.2 . . . Oophorectomy; salpingooophorectomy-----67.2-67.5 239 239 273.6 530.0 Tonsillectomy with or without 232 adenoidectomy-----21.1-21.2 80 152 266.1 188.4 337.7 219.2 223.3 Appendectomy 3 -----41.1 191 97 94 215.3 Excision of lesion of skin and subcutaneous tissue---92.1-92.2 182 74 108 208.2 174.9 239.0 Repair of obstetrical laceration-----78.2-78.3 180 180 205.8 398.6 Operations on muscles, tendons, fascia, and bursa-----88-89 Cholecystectomy-----43.5 214.2 156 90 178.3 144.6 154 18 136 176.4 43.0 301.5 Repair of inguinal 126 264.1 hernia-----38.2-38.3 112 14 143.8 31.0 Suture of skin and mucous 32 135.9 204.2 membrane----92.5 119 86 71.7

Includes data for sex not stated.

²Includes data for surgery not shown in table.

³Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery.

cialties accounted for two-fifths (41.1 percent) of all operations for this age group and for more than half (57.1 percent) of the surgery for women. By excluding the 3.5 million gynecological and obstetrical procedures from this comparison, the surgical rates were about the same for males and females.

Seven of the 15 surgical categories of operations with the largest frequencies listed in table L for patients aged 15-44 years were in the gynecological and obstetrical specialties. The leading operations for women and the rates per 1,000 women aged 15-44 years were diagnostic dilation and curettage of uterus (13.2), hysterectomy (9.2), and bilateral ligation and di-

vision of fallopian tubes (6.5). For men, the surgery rates were highest for repair of inguinal hernia (2.6); appendectomy (2.2); and operations on muscles, tendons, fascia, and bursa (2.1). The three leading non-sex-specific operations for this age group and their rates were biopsy (3.7), tonsillectomy with or without adenoidectomy (2.7), and appendectomy (2.2). For biopsy and tonsillectomy the surgical rates were substantially higher for females than for males, and for appendectomy the rates were approximately the same for both sexes.

Age group 45-64 years.—Surgery for patients aged 45-64 years was performed at a rate of 113.2 operations per 1,000 population (table M).

Table M. Number and rate of all-listed operations for inpatients 45-64 years of age discharged from short-stay hospitals, by selected surgical categories and sex: United States, 1973

[Excludes newborn infants and Federal hospitals]

[Excludes newborn infants and Federal hospitals]									
Surgical category and		of oper thousan		Rate o: 100,0	f operation 000 popula	ns per tion			
ICDA codes	Both sexes ¹			Both sexes ¹	Male	Female			
All operations, 45-64 years of age ²	4,826	1,904	2,920	11,317.1	9,377.1	13,074.8			
BiopsyA1-A2	338	115	222	793.0	567.3	995.5			
Dilation and curettage of uterus, diagnostic70.3 Hysterectomy69.1-69.5	296 243	• • • •	296 243	694.2 568.9		1,325.6 1,086.2			
Repair of inguinal hernia38.2-38.3	179	160	18	419.7	789.8	82.4			
Oophorectomy; salpingo- oophorectomy	165 159	44	165 114	386.0 371.8	218.6	737.0 511.2			
Excision of lesion of skin and subcutaneous tissue92.1-92.2	146	65	81	342.8	322.0	361.8			
Plastic repair of cystocele and/or rectocele71.4 Hemorrhoidectomy51.3 Operations on muscles,	93 90	47	93 43	217.0 210.8	229.8	414.3 192.8			
tendons, fuscia, and bursa88-89 Extraction of lens14.4-14.6 Prostatectomy58.1-58.3 Partial mastectomy65.2 Excision of intervertebral	82 78 71 61	40 39 71 *	42 40 60	192.7 184.1 166.1 143.9	197.7 190.8 348.7 *	188.1 177.4 268.0			
cartilage (prolapsed disk)86.4 Excision of bone, partial80.4	61 58	35 19	26 39	143.3 136.1	172.0 94.4	117.2 174.0			

Includes data for sex not stated.

²Includes data for surgery not shown in table.

Table N. Number and rate of all-listed operations for inpatients age 65 years and over discharged from short-stay hospitals, by selected surgical categories and sex: United States, 1973

Surgical category and	Number of operations in thousands			Rate of operations per 100,000 population				
ICDA codes	Both sexes ¹	Male	Female	Both sexes ¹	Male	Female		
All operations, 65 years of age and over ²	2,855	1,348	1,504	14,067.1	15,986.3	12,677.8		
BiopsyA1-A2 Extraction of lens14.4-14.6	229	106	122	1,127.1	1,254.3	1,031.9		
lens	189 175	71 175	117	929.8 862.3	843.3 2,075.5	985.9		
with fixation82.2 Repair of inguinal	140	28	112	687.7	326.5	943.4		
hernia38.2-38.3 Cholecystectomy43.5 Excision of lesion of	104 97	90 31	14 66	512.3 479.3	1,072.1 365.5	114.5 560.2		
skin and subcutaneous tissue92.1-92.2 Resection of small	75	37	38	371.7	439.1	323.0		
intestine or colon47.4-47.6 Local excision and	64	25	39	315.1	291.5	330.9		
destruction of lesion of bladder56.1-56.2 Closed reduction of	60	43	17	294.2	511.6	139.7		
fracture without fixation82.0 Dilation of urethra57.5	59 52	10 32	49 20	289.7 254.1	119.3 374.9	409.3 166.5		
Dilation and curettage of uterus, diagnostic70.3 Ileostomy, colostomy,	40	•••	40	195.5		334.5		
and other enterostomy47.7-47.9 Plastic repair of	34	16	19	168.1	185.2	156.0		
cystocele and/or rectocele71.4 Hysterectomy69.1-69.5	33 31	•••	33 31	163.7 152.3	•••	280.0 260.6		

By sex, the rates were 93.8 for men and 130.7 for women. Gynecological surgery, which accounted for 974,000 operations in this age group. contributed to the large differences in the rates by sex (table 2). Gynecological operations represented about a fifth (20.2 percent) of all

operations for all patients aged 45-64 and about a third (33.4 percent) of all operations for females.

Four of the 10 leading surgical operations for the 45-64 year age group were classified with gynecology. The leading surgical procedure

 $^{^1}_2$ Includes data for sex not stated. Includes data for surgery not shown in table.

was biopsy which accounted for 338,000 operations and a rate of 7.9 operations per 1,000 population. Other leading surgical operations and their rates were diagnostic dilation and curettage of uterus (6.9), hysterectomy (5.7), repair of inguinal hernia (4.2), and oophorectomy and salpingo-oophorectomy (3.9). As was observed for the other age groups, some non-sex-specific operations have a greater impact on one sex than the other. For females in the age group 45-64 years the surgery rate for biopsy was almost double the rate for males and for cholecystectomy 21/3 times higher. On the other hand, for males the rate for repair of inguinal hernia was over 9½ times higher than for females and the rate for excision of intervertebral cartilage was almost 1½ times higher than for females.

Age group 65 years and over.—An estimated 2,855,000 operations were performed for inpatients age 65 years and over discharged from short-stay hospitals during 1973 (table 2). The corresponding rate was 140.7 operations per 1,000 persons age 65 years and over in the civilian noninstitutionalized population (table 3). The rate for the oldest age group was more than three times that of the youngest age group.

The leading surgical categories for patients 65 years and over and the corresponding rates per 1,000 population by sex are presented in table N. The surgical rates were 159.9 for men and 126.8 for women. Surgical rates for males were higher than for females in the age groups under 15 and 65 years and over. The surgical categories with the highest rates were biopsy (11.3), extraction of lens (9.3), prostatectomy (8.6), reduction of fracture with fixation (6.9), and repair of inguinal hernia (5.1). Over two-thirds of all patients with prostatectomy and extraction of lens operations were age 65 years and over. The rates for 12 of the 15 leading surgical categories for the elderly in table N were substantially higher than for any other age group (table 3). The exceptions were the categories for diagnostic dilation and curettage of uterus, plastic repair of cystocele and/or rectocele, and hysterectomy.

Rates for the surgical categories varied considerably by sex. For persons 65 years and over, men had substantially higher surgical rates than women for repair of inguinal hernia, local excision and destruction of lesion of bladder,

and excision of lesion of skin and subcutaneous tissue. In contrast, for women 65 years and over the surgical rates were higher than for men for closed reduction of fracture without fixation, reduction of fracture with fixation, and cholecystectomy.

Surgery by Color

The number and percent distribution of all-listed operations for inpatients discharged from short-stay hospitals in 1973 are shown in table 4 according to color. Surgery rates were not computed because color was not identified on 12.8 percent of the medical records in the sample which accounted for 13.4 percent of the discharges with operations.

The surgical classes with the largest frequencies for both white and all other patients were gynecological, abdominal, orthopedic, and urological surgery, but not in the same order. In addition, otorhinolaryngology ranked fourth for white patients and obstetrical procedures ranked third for all other patients. Measured by the percentage of the number of operations in each class to all operations, the largest percent difference by color was for obstetrical procedures, 5.0 percent for white patients compared with 12.2 percent for all others. Other surgical specialties with large differences in these percentages by color were otorhinolaryngology which accounted for 10.0 percent of all operations for white patients and 6.3 percent for all other patients and vascular and cardiac surgery which accounted for 4.2 percent of the operations for white and 2.8 percent for all other patients.

The detailed surgical categories with the largest frequencies for both white and all other patients were biopsy, diagnostic dilation and curettage of uterus, tonsillectomy with or without adenoidectomy, and hysterectomy. These four surgical categories accounted for almost one out of five of all operations performed for white (18.5 percent) and all other (18.7 percent) patients. Ranked fifth in frequency were inguinal hernia for white patients and dilation and curettage after delivery or abortion for all other patients.

Variations by color are evident in the proportions which surgical categories represented

of all operations. For example, white patients had larger percentages of all operations for extraction of lens, myringotomy, and cholecystectomy than all other patients. However, for white patients the percentages were smaller than for all other patients operated for bilateral ligation and division of fallopian tubes, cesarean section, and dilation and curettage after delivery or abortion.

Surgery by Geographic Region

The surgical rates for all operations by region were lowest in the South and West Regions and were highest in the North Central and Northeast Regions. The estimated number of operations per 1,000 population ranged from 77.6 in the South Region to 101.9 in the North Central Region, or about 1.3 times larger (table 6). Patients with

Table O. Rate of all-listed operations for inpatients discharged from short-stay hospitals, by selected surgical categories and geographic region: United States, 1973

[Excludes newborn infants and Federal hospitals]

[Excludes newborn mants and rederal nospitals]								
Canadical astronomy and TODA and	United	Geographic region						
Surgical category and ICDA codes	States	North- east	North Central	South	West			
	Rate of	operatio	ons per 100	,000 popu	lation			
All operations ¹	8,951.8	9,191.3	10,185.6	7,764.0	8,807.3			
Dilation and curettage of uterus	453.8	586.1	546.5	335.9	337.5			
diagnostic70.3 BiopsyA1-A2 Tonsillectomy with or without	446.6	535.7	519.2	334.7	411.2			
adenoidectomy	429.6 335.2 255.3	346.0 275.9 302.2	530.2 333.5 276.6	375.4 370.9 198.1	482.4 354.8 260.2			
subcutaneous tissue92.1-92.2 Oophorectomy: salpingo-	221.2	220.5	233.1	226.6	193.6			
oophorectomy	206.5 199.8 164.5	210.4 229.7 145.1	207.2 219.5 183.9	208.0 178.9 163.8	197.5 165.4 161.5			
fixation82.0	156.0	141.3	182.6	140.5	161.9			
Operations on muscles, tendons, fascia, and bursa88-89 Ligation and division of fallopian	149.1	143.4	160.5	132.0	169.8			
Ligation and division of fallopian tubes (bilateral)68.5 Reduction of fracture with	145.5	173.9	137.6	150.5	110.1			
fixation82.2 Extraction of lens14.4-14.6 Dilation and curettage after delivery	137.9 135.6	115.3 143.7	159.8 154.1	127.8 112.3	152.6 137.1			
or abortion	132.6 120.8 119.6 110.4 106.0	149.7 146.0 129.3 97.4 98.3	142.2 142.6 108.2 161.6 116.4	133.9 90.6 128.9 100.9 111.8	91.5 106.2 107.6 63.6 89.5			
rectocele71.4	99.3	85.1	110.1	90.5	117.2			

Includes data for surgery not shown in table.

²Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery.

surgery in the United States and in each geographic region had an average of 1.4 surgical operations during their hospitalization. These statistics were obtained by comparing the number of surgical operations (table 5) with the number of discharges with surgery (table E).

The leading surgical classes in terms of frequency of operations were gynecological surgery, abdominal surgery, orthopedic surgery, otorhinolaryngology, and urological surgery. These were also the leading specialties in all the geographic regions. Gynecological surgery accounted for one in six operations in the West Region and one in five in the other regions.

Regional variations in surgical rates by surgical class were largest for dental surgery which ranged from 0.5 procedures per 1,000 population in the West Region to 2.6 in the Northeast Region, or about 5.8 times higher. For all other surgical classes, the highest rate for a region compared with the lowest rate was within the range of 1.2 to 1.8 times larger.

The five leading surgical categories in 1973 for inpatients of all short-stay hospitals in the United States were diagnostic dilation and curettage of uterus, biopsy, tonsillectomy with or without adenoidectomy, hysterectomy, and repair of inguinal hernia (table O). These were also the leading operations in each geographic region

except the South, but not in the same order. In the South Region excision of lesion of skin and subcutaneous tissue ranked fifth and repair of inguinal hernia ranked seventh.

There were variations among the regions in the estimated rates for the surgical categories. For example, of the leading surgical categories shown in table 0, the smallest difference in the rates per 1,000 population was for cophorectomy and salpingo-cophorectomy for which the range was from 2.0 in the West to 2.1 in the Northeast. The largest difference was for diagnostic dilation and curettage of the uterus for which the range was from 3.4 in the South to 5.9 in the Northeast, or 1.7 times larger.

Surgery by Bed Size of Hospital

The estimated number of all-listed operations in each surgical class and category is shown in table 7 by bed size of hospital where the surgery occurred. A percent distribution of these data is presented in table 8.

The smaller hospitals had smaller proportions of the total operations performed than of the total discharges and the larger hospitals had greater proportions of the operations than of the discharges (table P). This is also true when discharges are related to persons with surgery.

Table P. Number and percent distribution of inpatients discharged from short-stay hospitals, inpatients with surgery, and all-listed operations, according to bed size of hospital: United States, 1973

Bed size of hospital	Total dis- charges	Dis- charges with surgery	All- listed opera- tions	Total dis- charges	Dis- charges with surgery	All- listed opera- tions	
	Number	in thous	ands	Percent	distribution		
Tota1	32,125	13,266	18,426	100.0	100.0	100.0	
6-99 beds	6,604 5,485 5,625 8,228 6,183	1,917 2,102 2,459 3,842 2,946	2,640 2,893 3,447 5,383 4,062	20.6 17.1 17.5 25.6 19.2	14.5 15.8 18.5 29.0 22.2	14.3 15.7 18.7 29.2 22.0	
	,	'i '' '			1		

[Excludes newborn infants and Federal hospitals]

Table Q. Percent distribution of all-listed operations for inpatients discharged from short-stay hospitals by surgical class, according to bed size of hospital: United States, 1973

Surgical class and ICDA codes	All sizes	6-99 beds	100 - 199 beds	200 - 299 beds	300 - 499 beds	500 beds or more
				tributi		
All operations	100.0	100.0	100.0	100.0	100.0	100.0
Neurosurgery	0.4 3.9 1.3 14.9 3.1 7.9 1.8 19.4	1.3 0.7 17.4 3.0 5.8 1.7	1.7 0.8 16.4 3.1 8.9	0.4 3.6 1.3 14.8 3.6 9.0	1.4 13.9 3.2 7.9 1.9 19.1 5.4 12.6 5.2	4.2 8.1 0.5 7.2 1.8 13.6 2.4 7.6 1.55 17.2 7.5

Hospitals with fewer than 300 beds had 55.2 percent of the total discharges in 1973 and only 48.7 percent of the operations compared with hospitals of 300 beds or more which had only 44.8 percent of the discharges and 51.2 percent of the operations.

The leading surgical classes in hospitals of all bed size groups were gynecological surgery, abdominal surgery, orthopedic surgery, and otorhinolaryngology (table 7). Urological surgery ranked among the five leading surgical specialties in all hospitals except those with 6-99 beds, where plastic surgery ranked fifth and urological surgery sixth.

Surgical operations were highly clustered in the five leading surgical classes, which accounted for about two-thirds (65 percent) of all the operations reported. However, there was greater dispersion of the operations among the surgical classes in the larger than in the smaller hospitals.

Thus, the percentage which operations in the five leading specialties represented of all operations decreased from 73 percent in hospitals with 6-99 beds to 58 percent in hospitals with 500 beds or more. Offsetting these differences in the leading surgical classes by bed size of hospital. some specialties accounted for greater proportions of the total operations in the larger than the smaller hospitals. Ophthalmology increased consistently with bed size of hospital from 2.5 percent of all operations in hospitals with 6-99 beds to 4.2 percent in hospitals with 500 beds or more. vascular and cardiac surgery from 1.3 percent to 7.2 percent, thoracic surgery from 0.7 percent to 1.8 percent, oral and maxillofacial surgery from 0.5 percent to 1.2 percent, and biopsy from 3.7 percent to 6.0 percent (table Q). For other specialties such as otorhinolaryngology and proctological surgery definitive patterns of change were not evident by bed size of hospital.

Operations performed in 1973 were almost equally distributed between hospitals with fewer than 300 beds (48.7 percent) and hospitals with 300 beds or more (51.3 percent). However, the distributions of operations by surgical categories, the most detailed grouping of operations in this report, varied considerably between the smaller and larger hospitals.

Hospitals with fewer than 300 beds accounted for most of the following surgical operations: repair and plastic operations on joints of foot and toes (77.1 percent), suture of skin or mucous membrane (63.9 percent), appendectomy (63.7 percent), trachelectomy (62.2 percent), and closed reduction of fracture without fixation (60.3 percent). Hospitals with 300 beds or more accounted for most of the following surgical operations: reattachment of retina (87.9 percent), operations on valves of heart (87.8 percent), arthroplasty of hip (74.1 percent), nephrectomy (64.9 percent). and typanoplasty (64.9 percent). The proportions of total operations performed in small and large hospitals varied slightly for the surgical categories diagnostic dilation and curettage of uterus. hysterectomy, and oophorectomy and salpingooophorectomy.

FIRST-LISTED OPERATIONS AND AVERAGE LENGTH OF STAY

The number of first-listed operations (the only operation or the first-listed of multiple operations) and average lengths of stay are shown for 1973 by surgical category and sex in table 9 and by surgical category and age in tables 10 and 11. Average length of stay was computed for the operations and surgical procedures listed first on the face sheets of the medical records.

First-listed operations accounted for 13.3 million, or 72.0 percent, of the estimated 18.4 million operations performed during 1973. However, there were large variations in the percentages which first-listed operations represented of the all-listed operations (maximum of three operations coded for each medical record) by surgical category. These variations are evident when data in table 9 on number of first-

listed operations are compared with the data in table 1 on number of all-listed operations.

The proportion of first-listed operations to all-listed operations by surgical class ranged from 59.4 percent for gynecological surgery and 60.2 percent for biopsy to 93.7 percent for obstetrical procedures and 88.3 percent for operations on thyroid, parathyroid, thymus, and adrenals. For the more detailed surgical categories, some operations were almost always listed first and others were most frequently listed second or third on the face sheets of the medical records. First-listed operations accounted for 98 to 99 percent for the operations for extraction of lens, tonsillectomy with or without adenoidectomy, and cesarian section. In contrast, for oophorectomy and salpingo-oophorectomy the proportion first-listed was only 13.9 percent; for local excision of other lesions of uterus, cervix, and supporting structures 20,3 percent; and for colporrhaphy 27.9 percent.

Average length of stay was longer for patients discharged from short-stay hospitals with surgery than without surgery for all age groups and by sex, except for under 15 years of age (table R). For the youngest age group, average length of stay for all patients and by sex was shorter for discharges with surgery than without surgery, primarily because about two-fifths (43.0 percent) of the patients with surgery had a first-listed operation in the otorhinolaryngology specialty, for which the average length of stay was only 2.1 days.

Average length of stay for patients discharged with surgery was longer by age and sex for patients with multiple surgical operations performed than for patients with single operations. As shown in table B, 22.2 percent of the patients under age 15 had multiple operations compared with a high of 35.6 percent for age group 45-64.

The average lengths of stay for patients with surgery increased with age for all patients and by sex. For patients without surgery this pattern of longer average stays with advancing age was also evident for all patients and for males. However, for females without surgery average length of stay for under 15 years of age (5.2 days) was about the same as for ages 15-44 (5.0 days). Average length of stay was

Table R. Average length of stay for inpatients discharged from short-stay hospitals with and without surgery, by number of operations, sex, and age: United States, 1973

	Total	Total	Discharges with surgery					
Sex and age dis- charges wi		dis- charges without surgery	Total	One opera- tion	Two opera- tions	Three opera- tions		
<u>Total</u> ¹		Average 1	length of	stay ir	days			
All ages	7.8	7.6	7.9	7.0	9.1	12.8		
Under 15 years	4.5 5.7 9.1 12.1	5.0 5.4 8.7 11.2	4.0 6.0 9.7 14.3	5.2	4.1 7.1 10.7 16.6	9.3 9.8 13.8 21.4		
Male								
All ages	8.3	8.0	8.7	7.4	10.3	17.0		
Under 15 years	4.5 6.8 9.3 11.6	4.9 6.6 8.4 10.6	4.0 7.1 10.7 13.7	3.8 6.0 9.3 11.7	4.1 8.2 12.6 16.7	8.3 15.0 17.5 23.3		
<u>Female</u>						ji		
All ages	7.4	7.4	7.5	6.7	8.5	11.0		
Under 15 years	4.6 5.2 9.0 12.5	5.2 5.0 8.9 11.6	4.0 5.5 9.1 14.8	3.8 4.8 8.1 13.8	4.0 6.7 9.6 16.6	11.0 8.2 12.2 19.8		

¹Includes data for sex not stated.

shorter for women aged 15-44 without surgery than with surgery, primarily because of the large number of patients with first-listed diagnoses of uncomplicated deliveries which required short hospital stays but were not counted as operations in this report.

The average short-stay hospital stays by surgical class of first-listed operation ranged from 3.1 days for otorhinolaryngology to 15.0 days for neurosurgery. By the more detailed

surgical categories, average length of stay ranged from about 2 days to over 20 days. First-listed surgical categories which accounted for short average lengths of stay during 1973 include adenoidectomy without tonsillectomy (1.7 days), tonsillectomy with adenoidectomy (2.1 days), tonsillectomy without adenoidectomy (2.3 days), myringotomy (2.3 days), and dilation and curettage after delivery or abortion (2.4 days). Average lengths of stay were long for patients

whose first-listed operations were ileostomy, colostomy, and other enterostomy (25.1 days); emergency tracheotomy or tracheostomy (24.4 days); arthroplasty of hip (22.8 days); operations on valves of heart (22.2 days); and resection of

small intestine or colon (20.8 days). There were variations in average length of stay for the surgical categories by sex (table 9) and especially by age (table 11), which with few exceptions was longer for each older age group.

REFERENCES

¹National Center for Health Statistics: Development of the design of the NCHS Hospital Discharge Survey. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 2-No. 39. Public Health Service. Washington. U.S. Government Printing Office, Sept. 1970.

²National Center for Health Statistics: Eighth Revision International Classification of Diseases, Adapted for Use in the United States, 2 vols. PHS Pub. No. 1693. Public Health Service. Washington. U.S. Government Printing Office, 1967 and 1968.

³National Center for Health Statistics: Surgical operations in short-stay hospitals for discharged patients, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 7. Public Health Service. Washington. U.S. Government Printing Office, Apr. 1971.

⁴National Center for Health Statistics: Surgical operations in short-stay hospitals, United States, 1968. *Vital and Health Statistics*. Series 13-No. 11. DHEW Pub. No. (HSM) 73-1762. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Jan. 1973.

⁵National Center for Health Statistics: Surgical operations in short-stay hospitals, United States, 1971. Vital and Health Statistics. Series 13-No. 18. DHEW Pub. No. (HRA) 75-1769. Health Resources Administration. Washington. U.S. Government Printing Office, Nov. 1974.

⁶National Center for Health Statistics: Surgery in shortstay hospitals: United States, 1972. Monthly Vital Statistics Report. Vol. 23, No. 7, Supplement (3), (HRA) 75-1120. Rockville, Md. Health Resources Administration, Oct. 17, 1974.

⁷National Center for Health Statistics: Inpatient utilization of short-stay hospitals by diagnosis, United States,1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 6. Public Health Service. Washington. U.S. Government Printing Office, May 1970.

8 National Center for Health Statistics: Utilization of short-stay hospitals, by diagnosis: United States, 1966 and 1967. Monthly Vital Statistics Report. Vol. 20, No. 4, Supplement (HSM) 72-1124. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office. July 22, 1971.

⁹National Center for Health Statistics: Inpatient utilization of short-stay hospitals by diagnosis, United States, 1968. Vital and Health Statistics. Series 13-No. 12. DHEW Pub. No. (HSM) 73-1763. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Mar. 1973.

¹⁰National Center for Health Statistics: Inpatient utilization of short-stay hospitals by diagnosis, United States, 1971. Vital and Health Statistics. Series 13-No. 16. DHEW Pub. No. (HRA) 75-1767. Health Resources Administration. Washington. U.S. Government Printing Office, July 1974.

¹¹National Center for Health Statistics: Inpatient utilization of short-stay hospitals by diagnosis, United States, 1972. Vital and Health Statistics. Series 13-No. 20. DHEW Pub. No. (HRA) 76-1771. Health Resources Administration. Washington. U.S. Government Printing Office, Nov. 1975.

¹²National Center for Health Statistics: Development and maintenance of a national inventory of hospitals and institutions. Vital and Health Statistics. PHS Pub. No. 1000-Series 1-No. 3. Public Health Service. Washington. U.S. Government Printing Office, Feb. 1965.

13 National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics, United States, 1965. Vital and Health Statistics. PHS Pub. No. 1000-Series 13-No. 2. Public Health Service. Washington. U.S. Government Printing Office, Aug. 1967.

¹⁴National Center for Health Statistics: Utilization of short-stay hospitals by characteristics of discharged patients, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 3. Public Health Service. Washington. U.S. Government Printing Office, Dec. 1967.

LIST OF DETAILED TABLES

			Page
Table	1.	Number and rate of all-listed operations for inpatients discharged from short- stay hospitals, by surgical category and sex: United States, 1973	24
	2.	Number of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and age: United States, 1973	26
	3.	Rates of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and age: United States, 1973	28
	4.	Number and percent distribution of all-listed operations for inpatients discharged from short-stay hospitals by surgical category, according to color: United States, 1973	30
	5.	Number of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and geographic region: United States, 1973	32
	6.	Rates of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and geographic region: United States, 1973	34
	7.	Number of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and bed size of hospital: United States, 1973	36
	8.	Percent distribution of all-listed operations for inpatients discharged from short-stay hospitals by surgical category, according to bed size of hospital: United States, 1973	38
	9.	Number of first-listed operations and average length of stay for inpatients discharged from short-stay hospitals, by surgical category and sex: United States, 1973	40
:	10.	Number of first-listed operations for inpatients discharged from short-stay hospitals, by surgical category and age: United States, 1973	42
,	11.	Average length of stay for inpatients discharged from short-stay hospitals, by surgical category of first-listed operation and age: United States, 1973	44

Table 1. Number and rate of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and sex: United States, 1973

Surgical category and ICDA codes	Total ¹	Male	Female	Total ¹	Male	Female
		of all- ations ousands	in	ор	of all-li erations p 000 popula	er
All operations	18,426	6,936	11,480	8,951.8	6,984.7	10,776.5
Neurosurgery01-05	310	157	152	150.5	158.4	143.2
Ophtha1mology06-14	655	295	358	318.3	297.5	336.3
Resection and recession of eye muscle	82 33 279 260	37 21 116 122	46 13 162 138	40.0 16.2 135.6 126.5	37.0 20.7 116.8 123.0	42.9 12.0 152.3 129.2
Otorhinolaryngology16-21	1,835	903	930	891.4	909.8	873.3
Myringotomy	215 30 53 47 76 42 247 638 63 424	122 11 24 30 32 29 87 311 36 221	92 19 29 17 45 13 159 327 27 203	104.3 14.8 25.6 23.0 37.2 20.6 119.8 309.9 30.6 205.8	122.8 11.1 24.4 30.3 31.9 29.5 87.9 313.1 36.4 222.5	86.8 18.2 26.8 16.1 41.9 129.2 306.7 25.1 190.1
Operations on thyroid, parathyroid, thymus, and adrenals22-23	80	17	64	39.1	16.8	59.8
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	64 16	10 6	54 10	31.3 7.8	10.4 6.4	50.8 9.0
Vascular and cardiac surgery24-30	718	410	308	348.9	413.0	288.9
Incision of peripheral vessels	26 98 63 22 509	16 24 31 10 330	10 74 32 12 179	12.7 47.8 30.6 10.5 247.4	16.1 24.1 31.0 10.0 331.8	9.6 69.8 30.1 11.0 168.4
Thoracic surgery32-35	240	138	102	116.7	138.8	95.9
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	29 211	19 118	10 93	14.1 102.6	19.6 119.2	9.0 86.9
Abdominal surgery38-48	2,747	1,357	1,389	1,334.5	1,366.9	1,303.6
Repair of diaphragm and diaphragmatic hernia -38.0-38.1 Repair of inguinal hernia -38.2-38.3 Appendectomy -41.1 Cholecystectomy -45.1 Gastric resection, partial or complete -46.2-46.3 Vagotomy -46.8 Resection of small intestine or colon -47.4-47.6 Ileostomy, colostomy, and other enterostomy -47.7-47.9 Other abdominal surgery -Residual	30 525 339 411 31 64 60 146 71 1,070	12 464 175 94 16 39 40 65 37 415	18 61 163 318 15 25 20 81 34 655	14.4 255.3 164.5 199.8 15.0 31.0 29.1 71.0 34.5 519.8	12.1 467.1 176.7 94.4 16.3 39.4 40.3 65.5 37.2 418.0	16.6 57.5 153.0 298.1 13.8 23.2 18.7 76.1 32.1 614.5
Proctological surgery50-52	565	303	262	274.5	304.8	246.2
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	140 218 207	76 111 116	65 107 90	68.1 106.0 100.4	76.1 111.8 116.9	60.7 100.5 84.9
Urological surgery54-61	1,453	1,032	420	705.8	1,038.8	394.7
Nephrotomy and pyelotomy	28 32 74 101 25 54 227 249 72 26 60 99 382	15 15 43 67 18 40 7 82 249 72 26 60 99 240	12 18 31 7 14 18 145 	13.5 15.8 36.2 49.0 12.0 26.1 110.4 120.8 34.9 12.6 29.2 47.9 185.4	17.9 39.9 6.6 82.4 250.4 72.3 260.6 99.4	11.6 16.5 29.5 31.7 6.5 13.1 17.1 136.0

See footnotes at end of table.

Table 1. Number and rate of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and sex: United States, 1973—Con.

Surgical category and ICDA codes	Total ¹	Male	Female	Total ¹	Male	Female
	operations in oper		oper	of all-listed rations per 00 population		
Breast surgery65	336	19	317	163.2	18.8	297.8
Partial mastectomy65.2 Complete and radical mastectomy65.3-65.6 Other breast surgeryResidual	194 96 46	4 7 8	190 89 39	94.2 46.6 22.4	3.7 7.5 7.6	178.5 83.2 36.2
Gynecological surgery67-72	3,565	••••	3,565	1,732.2	•••	3,346.9
Local excision or destruction of lesion of ovary	74 425 299 690 116 934 127 41 204 655		74 425 299 690 116 934 127 41 204 655	36.1 206.5 145.5 335.2 56.2 453.8 61.5 20.0 99.3 318.2		69.7 399.1 281.1 647.7 108.7 876.8 118.7 38.6 191.8 614.7
Obstetrical procedures 374-78	1,077		1,077	523.2		1,011.0
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical proceduresResidual	246 273 182 376		246 273 182 376	119.6 132.6 88.2 182.9	:::	231.1 256.2 170.4 353.3
Orthopedic surgery80-90	2,351	1,217	1,133	1,142.3	1,225.1	1,064.0
Excision of bone, partial	1.57 321 74 284 147 53 120 49 29 32 307 779	64 174 45 119 86 22 17 28 14 19 169 461	93 147 29 164 61 31 102 21 15 13 138 318	76.3 156.0 35.8 137.9 71.4 25.7 58.1 24.0 14.0 15.3 149.1 378.7	64.0 174.8 45.6 120.0 86.3 21.9 17.4 28.6 14.0 18.8 169.9 463.8	87.7 138.1 26.8 154.4 57.5 29.2 96.1 19.6 12.1 1298.8
Plastic surgery92-94	1,041	543	497	505.8	546.7	466.5
Incision of skin and subcutaneous tissue	114 455 195 15 136 126	205 136 11	45 250 59 4 50 88	94.8 7.3 65.9	68.9 206.3 136.5 10.7 85.8 38.5	42.6 234.6 55.6 4.1 47.2 82.3
Oral and maxillofacial surgery95-98	174	92	82	84.5	92.6	76.8
Dental surgery99	358	153	204	174.1	154.5	191.
Extraction of tooth, forceps extraction	134 126 48 50	44 24	82	61.2	60.9 43.9 24.3 25.4	68. 77. 22. 23.
BiopsyA1-A2		300	618	446.6	302.3	580.

¹Includes data for inpatients discharged with sex not stated.

²Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery.

³Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 2. Number of all-listed operations for impatients discharged from short-stay hospitals, by surgical category and age:
United States, 1973

Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
		Number of all-listed operations in thousands			
All operations	18,426		8,420		2,855
Neurosugery01-05	310	28	111	117	54
Ophthalmology06-14	655	113	103	173	266
Resection and recession of eye muscle	82 33 279 260	58 * * 52	19 7 10 67	3 14 78 77	* 12 189 64
Otothinolaryngology16-21	1,835	995	596	174	69
Myringotomy	215 30 53 47 76 42 247 638 63 424	198 * 15 3 * 3 52 596 61 65	11 12 22 15 59 10 191 41 * 233	4 15 12 19 13 18 3 *-	* * * 4 10
Operations on thyroid, parathyroid, thymus, and adrenals22-23	80	4	36	31	10
Thyroidectomy	64 16	*	30 5	25 6	8 *
Vascular and cardiac surgery24-30	718	48	169	334	168
Incision of peripheral vessels	26 98 63 22 509	* 6 * 39	5 44 22 7 91	10 46 22 11 245	10 8 14 * 134
Thoracic surgery32-35	240	16	69	88	66
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	29 211	* 16	6 64	14 74	9 58
Abdominal surgery38-48	2,747	316	936	855	641
Repair of diaphragm and diaphragmatic hernia	30 525 339 411 31 64 60 146 71 1,070	* 117 103 * 4 * * * * 83	126 191 154 15 15 20 28 12 369	15 179 32 159 8 31 27 49 23 332	9 104 13 97 4 17 13 64 34 286
Proctological surgery50-52	565	10	298	193	65
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgeryResidual	140 218 207	* * 7	66 108 124	51 90 52	21 20 25
Urological surgery54-61	1,453	247	349	384	473
Nephrotomy and pyelotomy	28 32 74 101 25 54 25 227 249 72 26 60 99 382	* * 4 * * 28 * 58 * 34 21 * 53 43	9 10 28 11 4 10 11 64 * 12 5 20 33 129	11 11 29 28 7 7 8 53 71 16 * 17	7 9 14 60 11 9 6 522 175 10 * 23 93

See footnotes at end of table.

Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
		Number of operations			
Breast surgery65	336	5	166	118	47
Partial mastectomy65.2 Complete and radical mastectomy65.3-65.6 Other breast surgeryResidual	194 96 46	* *	116 17 33	61 47 10	14 31 *
Gynecological surgery67-72	3,565	22	2,402	974	168
Local excision or destruction of lesion of ovary	74 425 299 690	* * *	61 239 295 415	7 165 4 243	20 * 31
Hysterectomy	116 934 127 41 204 655	* 3 * * * 11	68 595 91 24 78 536	296 28 13 93 86	7 40 7 4 33 22
Obstetrical procedures ² 74-78	1,077	11	1,061	5	-
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical procedures	246 273 182 376	* * 7	244 269 180 368	* *	-
Orthopedic surgery80-90	2,351	270	1,020	646	416
Excision of bone, partial————————————————————————————————————	157 321 74 284 147 53 120 49 29 32 307 779	11 103 11 15 * * * * 46 67	71 103 34 79 80 44 26 11 15 156 398	58 19 12 5 82	16 59 12 140 6 28 15 * * 5 23 109
Plastic surgery92-94	1,041	150	470	279	142
Incision of skin and subcutaneous tissue92.0 Excision of lesion of skin and subcutaneous tissue92.1-92.2 Suture of skin or mucous membrane92.5 Plastic operations on lip and mouth93.1 Skin graft except lip and mouth93.2-93.6 Other plastic surgery	114 455 195 15 136 126	52 30 8 19	119 4	28 * 34	15 75 18 * 20 12
Oral and maxillofacial surgery95-98	174	24	84	45	22
Dental surgery99	358	36	229	73	20
Extraction of tooth, forceps extraction99.3 Surgical removal of tooth99.4 Alveoloplasty99.7 Other dental surgery	134 126 48 50	7 *	25	15 18	
BiopsyA1-A2	919	31	321	338	229

 $^{^1}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 3. Rates of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and age:
United States, 1973

Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
		Rate of al per 100	1-listed o		
All operations	8,951.8	4,185.8	9,640.1	11,317.1	14,067.1
Neurosurgery01-05	150.5	50.3	126.9	274.7	265.3
Ophthalmology06-14	318.3	203.3	118.5	405.9	1,309.0
Resection and recession of eye muscle	40.0 16.2 135.6 126.5	105.2 * * 93.4	22.2 7.7 12.0 76.6	8.0 32.4 184.1 181.4	56.8 929.8 316.6
Otorhinolaryngology16-21	891.4	1,791.5	682.8	408.7	339.4
Myringotomy -17.0 Stapedectomy with ossicular reconstruction -17.4 Tympanoplasty -17.6-17.7 Excision of lesion of nose -19.0 Section of nasal septum -19.1 Tracheostomy or tracheostomy, emergency -20.5 Tonsillectomy without adenoidectomy -21.1 Tonsillectomy with adenoidectomy -21.2 Adenoidectomy without tonsillectomy -21.3 Other operations on ears, nose, and throat -Residual	104.3 14.8 25.6 23.0 37.2 20.6 119.8 309.9 30.6 205.8	356.2 26.3 6.0 * 6.1 92.8 1,072.0 110.1 117.0	12.6 14.1 25.3 16.7 68.0 11.2 218.8 47.3 266.6	10.1 35.6 28.8 44.4 31.3 41.4 7.5 *	** 18.5 51.6 ** 57.0 ** **
Operations on thyroid, parathyroid, thymus, and adrenals22-23	39.1	6.7	40.9	73.2	48.2
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	31.3 7.8	5.5	34.8 6.1	59.6 13.6	39.5
Vascular and cardiac surgery24-30	348.9	86.8	193.2	782.2	826.4
Incision of peripheral vessels	12.7 47.8 30.6 10.5 247.4	* 11.1 * 69.9	5.7 50.2 24.7 8.2 104.4	23.4 108.0 50.6 24.9 575.4	50.3 37.8 66.9 * 659,9
Thoracic surgery32-35	116.7	29.3	79.4	207.0	327.2
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	14.1 102.6	28.6	6.4 73.0	33.1 173.9	43.7 283.4
Abdominal surgery38-48	1,334.5	569.1	1,071.1	2,003.9	3,157.2
Repair of diaphragm and diaphragmatic hernia -38.0-38.1 Repair of inguinal hernia -38.2-38.3 Appendectomy: -41.1 Cholecystectomy -45.5 Splenectomy: -45.1 Gastric resection, partial or complete -46.2-46.3 Vagotomy: -46.8 Resection of small intestine or colon: -47.4-47.6 Ileostomy, colostomy, and other enterostomy -47.7-47.9 Other abdominal surgery: -Residual	14.4 255.3 164.5 199.8 15.0 31.0 29.1 71.0 34.5 519.8	210.4 185.3 * 7.3 * 8.6 *	6.4 143.8 219.2 176.4 17.1 17.2 22.9 32.4 13.6 422.2	34.7 419.7 74.3 371.8 19.0 73.3 62.2 115.4 55.0 778.6	43.4 512.3 61.7 479.3 18.6 85.2 64.5 315.1 168.1 1,408.8
Proctological surgery50-52	274.5	17.2	340.9	451.8	320.4
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	68.1 106.0 100.4	* * 11.9	75.8 123.6 141.5	119.4 210.8 121.7	102.2 97.2 121.0
Urological surgery54-61	705.8	445.3	399.6	900.2	2,328.7
Nephrotomy and pyelotomy	13.5 15.8 36.2 49.0 12.0 26.1 10.4 120.8 34.9 12.6 29.2 47.9 185.4	6.4 50.8 105.1 60.7 37.4 94.6 76.9	10.7 11.6 32.6 12.8 4.8 11.0 12.1 73.3 * 13.9 5.3 23.0 37.8 148.2	25.1 26.5 67.2 66.4 17.4 16.6 18.2 124.8 166.1 37.8 40.7 19.8 272.7	33.5 43.6 68.0 294.2 55.8 43.7 30.5 254.1 862.3 48.5 23.1 459.2

See footnotes at end of table.

Table 3. Rates of all-listed operations for impatients discharged from short-stay hospitals, by surgical category and age:
United States, 1973—Con.

Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over			
	Rate of all-listed operations per 100,000 population							
	160.01	229.8						
Breast surgery65	163.2	9.4	190.3	276.5				
Partial mastectomy	94.2 46.6	*	132.6	143.9 109.4	69.9 150.6			
Complete and radical mastectomyResidual	22.4	*	37.7	23.1	*			
Other breast surgery	1,732.2	39.4	2,750.0	2,284.2	826.1			
Gynecological surgery67-72		1	-	16 5	17.8			
Local excision or destruction of lesion of ovary	36.1 206.5	*	70.3 273.6	16.5 386.0	98.4			
Oophorectomy; salpingo-oophorectomy	145.5 335.2	*	337.9 474.7	8.9 568.9	152.3			
Hysterectomy	56.2	*	78.0	93.6	34.7			
supporting tissues	453.8 61.5	5.6	681.5 104.2	694.2 66.2	195.5 35.2			
Trachelectomy71.3	20.0	*	27.1 89.2	30.5 217.0	19.4 163.7			
Plastic repair of cystocele and/or rectoceleResidual	99.3 318.2	19.2	613.5	202.4	108.5			
Other gynecological surgery	523.2	19.7	1,214.8	10.9	_			
Obstetrical procedures ² 74-78			,		1			
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical proceduresResidual	119.6 132.6	* *	279.9 308.2	*				
Dilation and curettage after delivery or abortion	88.2 182.9	12.4	205.8 421.0	*				
Other obstetrical proceduresResidual	102.7							
Orthopedic surgery80-90	1,142.3	486.1	1,167.6	1,513.9	2,049.7			
Excision of bone, partial	76.3 156.0	20.7 185.8	81.4 117.8	136.1 131.9	80.4 289.7			
Closed reduction of fracture without fixation82.1	35.8	19.5	39.4 90.4	38.2 119.0	60.5			
Reduction of fracture with fixation86.4	137.9 71.4	*	91.3	143.3	28.0 138.2			
Reduction of fracture with fixation	25.7 58.1	* 	4.5 50.1	46.7 135.6	74.9			
Repair and plastic operations on joints of loot and loots.	24.0 14.0	6.5	29.9 12.2 17.1	45.3 28.4	*			
Spinal fusion	15.3 149.1	11.7 82.4	17.1 178.3	12.4 192.7	114.7			
Arthrodesis and stabilization of joints (except spine)	378.7	120.0	455.3	484.3				
Plastic surgery92-94	505.8	269.3	538.1	654.6	701.0			
	l	43.2	56.8	59.2				
Incision of skin and subcutaneous tissue	221.2 94.8		208.2 135.9	342.8 65.9				
Suture of skin or mucous membrane93.1	7.3	14.9	72.0	79.1	100.9			
Incision of skin and subcutaneous tissue92.0 Excision of lesion of skin and subcutaneous tissue92.1 Suture of skin or mucous membrane92.5 Plastic operations on lip and mouth93.2 Skin graft except lip and mouth	65.9 61.2	29.7	60.5	104.3				
OF OF OR OTHER PLANTS SUIT SUIT SUIT SUIT SUIT SUIT SUIT SU	84.5	42.5	95.9	104.5	108.7			
Oral and maxillofacial surgery95-98	,	11			1			
Dental surgery99	174.1	64.0	262.1	172.1	101.0			
Extraction of tooth forceps extraction99.3	65.1							
Surgical removal of tooth99,	61.2	*	28.4	42.7	23.1			
Extraction of tooth, forceps extraction99. Surgical removal of tooth99. Alveoloplasty	24.3	29.0	25.9	22,8	`			
BiopsyAl-A	446.6	55.9	368.0	793.0	1,127.1			

 $^{^{1}}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 4. Number and percent distribution of all-listed operations for inpatients discharged from short-stay hospitals by surgical category, according to color: United States, 1973

			<i></i>			
Surgical category and ICDA codes	Total ¹	Color stated		1	Color stated	
		White	All other	Total ¹	White	All other
	Number operation	r of all-l	listed ousands	Percent distribution		
All operations	18,426	14,056	1,900	100.0	100.0	100.0
Neurosurgery01-05	310	240	27	1.7	1.7	1.4
Ophthalmology06-14	655	494	57	3.6	3,5	3.0
Resection and recession of eye muscle	82 33 279 260	61 28 217 188	7 * 17 32	0.4 0.2 1.5 1.4	0.4 0.2 1.5 1.3	0.4 * 0.9 1.7
Otorhinolaryngology16-21	1,835	1,409	121	10.0	10.0	6.3
Myringotomy -17.0 Stapedectomy with ossicular reconstruction -17.4 Tympanoplasty -17.6-17.7 Excision of lesion of nose -19.0 Section of nasal septum -19.1 Trachectomy or tracheostomy, emergency -20.5 Tonsillectomy without adenoidectomy -21.1 Tonsillectomy with adenoidectomy -21.2 Adenoidectomy without tonsillectomy -21.3 Other operations on ears, nose, and throat -Residual	215 30 53 47 76 42 247 638 63 424	160 25 41 40 68 32 178 493 45 327	8 * * * 6 26 43 3 30	1.2 0.2 0.3 0.3 0.4 0.2 1.3 3.5 0.3 2.3	1.1 0.2 0.3 0.5 0.2 1.3 3.5 2.3	0.4 * * 0.3 1.4 2.3 1.6
Operations on thyroid, parathyroid, thymus, and adrenals22-23	80	59	12	0.4	0.4	0.6
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	64 16	46 13	10 *	0.3 0.1	0.3	0.5 *
Vascular and cardiac surgery24-30	718	588	53	3.9	4.2	2.8
Incision of peripheral vessels	26 98 63 22 509	20 79 49 19 421	* 4 5 * 42	0,1 0,5 0,3 0,1 2,8	0.1 0.6 0.4 0.1 3.0	0.2 0.2 0.2 *
Thoracic surgery32-35	240	185	31	1.3	1.3	1.6
Lung lobectomy	29 211	23 162	3 27	0.2 1.1	0.2 1.2	0.2 1.4
Abdominal surgery38-48	2,747	2,126	277	14.9	15.1	14.6
Repair of diaphragm and diaphragmatic hernia -38.0-38.1 Repair of inguinal hernia -38.2-38.3 Appendectomy ² -41.1 Cholecystectomy -43.5 Splenectomy -45.1 Gastric resection, partial or complete -46.2-46.3 Vagotomy -46.8 Resection of small intestine or colon -47.4-47.6 Ileostomy, colostomy, and other enterostomy -47.7-47.9 Other abdominal surgery -Residual	1 1	25 414 264 329 27 49 45 113 55 805	* 47 29 26 * 6 7 13 8 140	0.2 2.9 1.8 2.2 0.3 0.3 0.4 5.8	0.2 2.9 1.9 2.3 0.2 0.3 0.3 0.4 5.7	2.5 1.5 1.4 0.3 0.4 0.7 0.4 7.4
Proctological surgery50-52	565	440	52	3.1	3.1	2.7
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	140 218 207	109 170 160	12 19 21	0.8 1.2 1.1	0.8 1.2 1.1	0.6 1.0 1.1
Urological surgery54-61	1,453	1,104	141	7.9	7.9	7.4
Nephrotomy and pyelotomy	28 32 74 101 25 54 25 227 249 72 26 60 99 382	22 24 58 82 19 41 19 178 193 53 20 51 54 292	*4 3 5 *6 ** 18 19 7 3 *34 33	0.1 1.2 1.3 0.4 0.1 0.3 0.5	0.6 0.1 0.3 0.1 1.3 1.4 0.1 0.4	0.2 0.2 0.2 0.3 * 1.0 0.3 0.2 * 1.8

See footnotes at end of table.

Table 4. Number and percent distribution of all-listed operations for impatients discharged from short-stay hospitals by surgical category, according to color: United States, 1973—Con.

		Color stated			Color stated	
Surgical category and ICDA codes	Total ¹	White	A11	Total ¹	White	A11
			other		WILLE	other
		er of all- ions in th		Percent distribution		
Breast surgery65	336	260	31	1.8	1.9	1.6
Partial mastectomy65.2 Complete and radical mastectomy65.3-65.6 Other breast surgeryResidual	194 96 46	147 76 37	20 6 5	1.1 0.5 0.3	1.0 0.5 0.3	1.0 0.3 0.3
Gynecological surgery67-72	3,565	2,672	410	19.4	19.0	21.6
Local excision or destruction of lesion of ovary	74 425 299 690	56 324 217 526	6 47 47 71	0,4 2,3 1,6 3,7	0.4 2.3 1.5 3.7	0.3 2.4 2.5 3.7
Local excision and destruction of other lesions of uterus, cervix, and supporting tissues	116 934 127 41 204 655	91 703 95 31 167 463	11 104 17 4 5 98	0.6 5.1 0.7 0.2 1.1 3.6	0.6 5.0 0.7 0.2 1.2 3.3	0.6 5.5 0.9 0.2 0.3 5.1
Obstetrical procedures ³ 74-78	1,077	708	232	5.8	5.0	12.2
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical procedures	246 273 182 376	172 187 120 230	46 48 41 97	1.3 1.5 1.0 2.0	1.2 1.3 0.9 1.6	2.4 2.5 2.1 5.1
Orthopedic surgery80-90	2,351	1,845	195	12.8	13.1	10.3
Excision of bone, partial————————————————————————————————————	157 321 74 284 147 53 120 49 29 32 307 779	126 257 57 230 115 44 97 39 23 25 231 603	12 24 8 19 10 * 10 4 3 * 36 63	0.9 1.7 0.4 1.5 0.8 0.3 0.6 0.3 0.2 0.2	0.9 1.8 0.4 1.6 0.3 0.7 0.3 0.2 1.6 4.3	0.6 1.3 0.4 1.0 0.5 0.5 0.2 0.2 0.2
Plastic surgery92-94	1,041	821	101	5.6	5.8	5.3
Incision of skin and subcutaneous tissue92.0 Excision of lesion of skin and subcutaneous tissue92.1-92.2 Suture of skin or mucous membrane92.5 Plastic operations on lip and mouth93.1 Skin graft except lip and mouth93.2-93.6 Other plastic surgery	114 455 195 15 136 126	84 365 150 12 104 106	19 34 20 * 18 8	0.6 2.5 1.1 0.1 0.7 0.7	0.6 2.6 1.1 0.1 0.7 0.8	1.0 1.8 1.1 * 0.9 0.4
Oral and maxillofacial surgery95-98	174	128	25	0.9	0.9	1.3
Dental surgery99	358	274	26	1.9	2.0	1.4
Extraction of tooth, forceps extraction99.3 Surgical removal of tooth99.4 Alveoloplasty	134 126 48 50	99 99 39 37	11 6 4 5	0.7 0.7 0.3 0.3	0.7 0.7 0.3 0.3	0.6 0.3 0.2 0.3
Biopsy	919	703	111	5.0	5.0	5,8

¹Includes data for inpatients discharged with color not stated.
²Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery.
³Codes 75,0-75,6 and 75.9 are not used by HDS.

Table 5. Number of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and geographic region: United States, 1973

		7			
	United	Geo	graphic :	region	
Surgical category and ICDA codes	States	Northeast	North Central	South	West
		Number of operations			
All operations	18,426	4,498	5,783	5,008	3,138
Neurosurgery01-05	310	64	84	88	74
Ophthalmology06-14	655	152	218	156	129
Resection and recession of eye muscle	82 33 279 260	19 8 70 55	28 14 87 88	16 6 72 62	19 6 49 55
Otorhino1aryngo1ogy16-21	1,835	393	666	427	349
Myringotomy	215 30 53 47 76 42 247 638 63 424	44 10 11 11 26 10 45 124 14 97	83 12 19 17 32 11 82 219 28 165	43 * 12 9 8 12 66 176 12 85	45 6 11 10 10 9 54 118 10 77
Operations on thyroid, parathyroid, thymus, and adrenals22-23	80	21	23	24	13
Thyroidectomy22,1-22,2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	64 16	17 4	19 5	19 5	11 *
Vascular and cardiac surgery24-30	718	165	242	177	134
Incision of peripheral vessels	26 98 63 22 509	6 33 14 5 108	9 31 24 6 173	7 18 15 5 132	5 16 10 5 97
Thoracic surgery32-35	240	57	64	72	48
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	29 211	6 51	. 57	8 63	8 41
Abdominal surgery38-48	2,747	684	831	777	455
Repair of diaphragm and diaphragmatic hernia 38.0-38.1 Repair of inguinal hernia 38.2-38.3 Appendectomy - 41.1 Cholecystectomy - 43.5 Splenectomy - 45.1 Gastric resection, partial or complete 46.2-46.3 Vagotomy - 46.8 Resection of small intestine or colon - 47.4-47.6 Ileostomy, colostomy, and other enterostomy 47.7-47.9 Other abdominal surgery -	30 525 339 411 31 64 60 146 71 1,070	5 148 71 112 7 19 12 41 21 248	8 157 104 125 10 17 16 47 24 323	10 128 106 115 8 16 17 32 15 330	7 93 58 59 6 12 14 26 11 169
Proctological surgery50-52	565	147	170	163	86
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgeryResidual	140 218 207	36 48 63	46 66 58	39 72 52	20 32 34
Urological surgery54-61	1,453	357	478	409	209
Nephrotomy and pyelotomy	28 32 74 101 25 54 227 249 72 26 60 99	7 8 22 29 6 8 48 48 71 19 8	8 11 25 36 9 19 9 92 81 24 7 17 23 117	7 10 19 23 6 21 8 65 58 18 7 19	6 4 8 13 4 6 * 23 31 4 9 15 66

Table 5. Number of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and geographic region: United States, 1973—Con.

		Geo	egion		
Surgical category and ICDA codes	United States	Northeast	North Central	South	West
	1	Number of operations			l,
Breast surgery65	336	95	101	90	50
Partial mastectomy	194 96 46	55 29 11	56 30 16	55 23 11	28 14 8
Gynecological surgery67-72	3,565	910	1,088	1,033	535
Local excision or destruction of lesion of ovary	74 425 299 690	19 103 85 135	23 118 78 189	20 134 97 239	11 70 39 126
Local excision and destruction of other lesions of uterus, cervix, and supporting tissues70.2 Dilation and curettage of uterus, diagnostic70.3 Trachelectomy	116 934 127 41 204 655	42 287 22 7 42 168	35 310 44 11 63 217	29 217 39 16 58 183	10 120 22 8 42 87
Obstetrical procedures ² 74-78	1,077	311	278	300	187
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical proceduresResidual	246 273 182 376	63 73 32 143	61 81 54 83	83 86 60 71	38 33 36 81
Orthopedic surgery80-90	2,351	481	741	630	499
Excision of bone, partial	157 321 74 284 147 53 120 49 29 32 307 779	27 69 18 56 30 10 12 7 6 7 70 167	38 104 27 91 44 20 29 16 10 91 262	47 91 18 82 43 10 36 14 8 9 85 189	45 58 12 54 31 13 43 12 5 60 162
Plastic surgery92-94	1,041	231	313	316	180
Incision of skin and subcutaneous tissue92.0 Excision of lesion of skin and subcutaneous tissue92.1-92.2 Suture of skin or mucous membrane92.5 Institute operations on lip and mouth93.1 Skin graft except lip and mouth93.2-93.6 Other plastic surgery	114 455 195 15 136 126	26 108 40 4 30 22	32 132 60 4 40 45	33 146 58 4 41 34	22 69 37 * 25 24
Oral and maxillofacial surgery95-98	174	40	57	51	26
Dental surgery99	358	129	132	81	16
Extraction of tooth, forceps extraction99.3 Surgical removal of tooth99.4 Alveoloplasty99.7 Other dental surgery	134 126 48 50	50 49 18 12	43 49 19 21	36 24 9 12	5 4 * 5
BiopsyA1-A2	919	262	295	216	147

 $^{^{1}}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 6. Rates of all-listed operations for impatients discharged from short-stay hospitals, by surgical category and geographic region: United States, 1973

,	United		Geographic	region		
Surgical category and ICDA codes	States	Northeast	North Central	South	West	
			Rate of all-listed operations per 100,000 population			
All operations	8,951.8	9,191.3	10,185.6	7,764.0	8,807.3	
Neurosurgery01-05	150.5	130.7	147.3	136.1	208.9	
Ophthalmology06-14	318.3	309.9	384.2	242.3	362.2	
Resection and recession of eye muscle	40.0 16.2 135.6	38.7 15.4 143.7	49.8 25.0 154.1	24.3 9.0 112.3	54.7 16.0 137.1	
		112.0	155.4	96.7	154.4	
Otorhinolaryngology16-21	891.4	802.6	1,173.9	661.6	979.5	
Myringotomy	104.3 14.8 25.6 23.0 37.2 20.6 119.8 309.9 30.6 205.8	89.5 20.3 22.3 53.8 20.7 92.7 253.4 28.9 197.7	145.4 20.9 33.7 29.9 56.1 19.1 143.6 386.6 48.6 290.1	67.1 * 18.6 13.9 19.1 102.4 273.0 17.9 132.0	126.2 15.5 30.1 28.3 27.9 25.4 150.5 331.9 27.4 216.4	
Operations on thyroid, parathyroid, thymus, and adrenals22-23	39.1	43.1	40.8	36.5	35.4	
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	31.3 7.8	34.3 8.9	32.6 8.2	28.9 7.6	29.5	
Vascular and cardiac surgery24-30	348.9	338.1	426.9	273.9	375.4	
Incision of peripheral vessels	12.7 47.8 30.6 10.5 247.4	11.4 67.2 28.1 10.6 220.8	15.2 54.6 41.4 11.4 304.2	10.4 28.5 24.0 7.1 203.9	14.5 45.0 28.5 15.1 272.3	
Thoracic surgery32-35	116.7	115.9	112.2	111.1	135.4	
Lung lobectomy34.2-34.4 Other thoracic surgery	14.1 102.6	11.7 104.2	12.6 99.6	13.1 98.0	21.6 113.7	
Abdominal surgery38-48	1,334.5	1,397.4	1,464.0	1,204.0	1,278.2	
Repair of diaphragm and diaphragmatic hernia -38.0-38.1 Repair of inguinal hernia -38.2-38.3 Appendectomy¹ -41.1 Cholecystectomy -45.1 Gastric resection, partial or complete -46.2-46.3 Vagotomy* -46.8 Resection of small intestine or colon -47.4-47.6 Ileostomy, colostomy, and other enterostomy -47.7-47.9 Other abdominal surgery*	14.4 255.3 164.5 199.8 15.0 31.0 29.1 71.0 34.5 519.8	9.8 302.2 145.1 229.7 14.4 38.6 25.4 83.7 42.3 506.3	14.3 276.6 183.9 219.5 16.8 29.8 28.2 82.6 42.4 569.7	15.0 198.1 163.8 178.9 12.4 24.2 26.6 49.5 23.8 511.7	20.0 260.2 161.5 165.4 17.5 35.1 40.2 74.2 30.8 473.3	
Proctological surgery50-52	274.5	299.4	299.1	252.8	240.4	
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	68.1 106.0 100.4	72.9 98.3 128.2	80.3 116.4 102.3	60.2 111.8 80.9	56.6 89.5 94.3	
Urological surgery54-61	705.8	729.4	842.3	633.5	587.0	
Nephrotomy and pyelotomy	13.5 15.8 36.2 49.0 12.0 26.1 110.4 120.8 34.9 12.6 29.2 47.9 185.4	14.0 16.8 45.6 59.4 12.1 15.7 8.2 97.4 146.0 39.8 16.5 32.1 42.7 183.2	14.7 19.4 44.7 63.2 15.6 33.3 16.6 142.6 42.5 11.8 29.5 40.1 206.7	11.0 14.8 29.0 35.9 9.4 32.1 12.5 100.9 90.6 27.2 10.3 29.4 62.3 168.2	15.5 10.4 22.6 35.9 10.8 18.1 63.6 106.2 29.9 12.5 24.6 185.8	

Table 6. Rates of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and geographic region: United States, 1973—Con.

		T						
	United		Geographic	region				
Surgical category and ICDA codes	States	Northeast	North Central	South	West			
	,	Rate of all per 100,	-listed or 000 popula	erations tion				
Breast surgery65	163.2	194.1	178.5	139.0	140.4			
Partial mastectomy65.2 Complete and radical mastectomy65.3-65.6 Other breast surgery	94.2 46.6 22.4	112.8 58.5 22.8	97.9 52.5 28.2	85.6 36.0 17.3	78.2 40.3 21.9			
Gynecological surgery67-72	1,732.2	1,859.0	1,915.6	1,601.5	1,502.2			
Local excision or destruction of lesion of ovary	36.1 206.5 145.5 335.2	39.3 210.4 173.9 275.9	41.0 207.2 137.6 333.5	31.5 208.0 150.5 370.9	32.0 197.5 110.1 354.8			
Local excision or destruction of lesion of overy	56.2 453.8 61.5 20.0 99.3 318.2	85.3 586.1 45.8 13.5 85.1 343.7	62.0 546.5 77.1 18.9 110.1 381.7	44.6 335.9 60.1 25.1 90.5 284.3	28.3 337.5 60.5 21.2 117.2 243.1			
Obstetrical procedures ² 74-78	523.2	635.3	490.5	465.5	526.0			
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical procedures	119.6 132.6 88.2 182.9	129.3 149.7 65.0 291.2	108.2 142.2 94.7 145.5	128.9 133.9 93.3 109.4	107.6 91.5 100.3 226.5			
Orthopedic surgery80-90	1,142.3	982.5	1,305.6	977.4	1,400.3			
Excision of bone, partial	76.3 156.0 35.8 137.9 71.4 25.7 58.1 24.0 14.0 15.3 149.1 378.7	55.7 141.3 36.3 115.3 60.5 20.7 25.3 15.0 12.4 15.0 143.4 341.6	66.4 182.6 47.0 159.8 77.7 35.7 51.2 28.3 17.0 18.3 160.5	73.3 140.5 27.7 127.8 65.9 15.4 55.1 21.8 11.9 13.5 132.0 292.5	125.5 161.9 32.4 152.6 86.0 35.3 119.7 133.3 14.9 169.8 454.4			
Plastic surgery92-94	505.8	472.7	552.2	489.6	506.5			
Incision of skin and subcutaneous tissue92.0 Excision of lesion of skin and subcutaneous tissue92.1-92.2 Suture of skin or mucous membrane92.5 Plastic operations on lip and mouth93.1 Skin graft except lip and mouth93.2-93.6 Other plastic surgery	55.3 221.2 94.8 7.3 65.9 61.2	53.9 220.5 81.9 9.1 61.5 45.8	56.8 233.1 105.0 7.5 70.4 79.3	50.7 226.6 89.9 6.1 63.1 53.2	63.1 193.6 104.9 * 70.1 68.0			
Oral and maxillofacial surgery95-98	84.5	82.7	100.5	78.5	72.6			
Dental surgery99	174.1	262.8	232.8	126.0	45.7			
Extraction of tooth, forceps extraction99.3 Surgical removal of tooth99.4 Alveoloplasty99.7 Other dental surgery	65.1 61.2 23.5 24.3	101.5 100.7 36.2 24.5	76.2 86.6 33.4 36.6	55.9 37.1 14.1 18.9	13.9 10.3 * 14.0			
BiopsyAl-A2	446.6	535.7	519.2	334.7	411.2			

 $^{^1}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 7. Number of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and bed size of hospital: United States, 1973

	-	1					
	All		Bed	size of h	ospital		
Surgical category and ICDA codes	sizes	6-99 beds	100-199 beds	200-299 beds	300-499 beds	500 beds or more	
		Number of all-listed operations in thousands					
All operations	18,426	2,640	2,893	3,447	5,383	4,062	
Neurosurgery01-05	310	40	24	47	99	100	
Ophthalmology06-14	655	67	91	121	203	173	
Resection and recession of eye muscle	82 33 279 260	9 * 21 37	13 * 40 36	17 * 54 48	24 10 95 74	20 19 69 64	
Otorhinolaryngology16-21	1,835	283	268	388	567	328	
Myringotomy	215 30 53 47 76 42 247 638 63 424	16 * 7 15 * 52 132 * 54	28 7 5 14 * 40 107 9	49 6 12 9 11 52 145 12 84	79 8 19 19 21 13 68 180 23 138	. 43 9 15 8 17 16 34 73 15	
Operations on thyroid, parathyroid, thymus, and adrenals22-23	80	8	13	13	26	21	
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	64 16	7 *	11 *	12 *	20 6	15 6	
Vascular and cardiac surgery24-30	718	35	51	124	215	294	
Incision of peripheral vessels	26 98 63 22 509	13 7 - 12	16 7 - 25	5 15 13 * 89	9 36 17 6 147	7 19 19 13 235	
Thoracic surgery32-35	240	19	23	45	78	75	
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	29 211	* 18	* 21	6 39	10 68	10 65	
Abdominal surgery38-48	2,747	460	475	510	748	554	
Repair of diaphragm and diaphragmatic hernia	30 525 339 411 31 64 60 146 71 1,070	6 97 82 73 * 7 10 16 7	86 66 75 * 11 13 21 10 186	5 99 68 77 5 12 9 26 13 198	6 155 75 114 11 19 14 48 23 284	8 89 48 73 9 15 15 35 20 243	
Proctological surgery50-52	565	79	91	124	172	99	
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	140 218 207	18 37 24	20 40 31	32 44 48	43 62 68	27 36 37	
Urological surgery54-61	1,453	152	257	309	427	307	
Nephrectomy and pyelotomy	28 32 74 101 25 54 25 227 249 72 26 60 99 382	* 6 7 * 5 * 19 20 10 4 10 20 43	4 12 16 4 11 42 42 42 14 *	4 6 17 22 6 10 61 52 13 6 14 16 74	10 10 23 33 7 19 7 65 75 20 7 16 27 109	7 11 17 23 5 8 4 41 60 14 10 16 16 18 18	

Table 7. Number of all-listed operations for inpatients discharged from short-stay hospitals, by surgical category and bed size of hospital: United States, 1973—Con.

			Bed	size of h	ospital	pital	
Surgical category and ICDA codes	All sizes	6-99 beds	100-199 beds	200-299 beds	300-499 beds	500 beds or more	
		Number of all-listed operations in thousands					
Breast surgery65	336	44	58	71	102	62	
Partial mastectomy	194 96 46	25 12 7	36 15 7	44 19 8	60 29 13	29 20 12	
Gynecological surgery67-72	3,565	512	684	640	1,030	700	
Local excision or destruction of lesion of ovary	74 425 299 690	13 60 44 100	14 82 64 142	16 73 42 112	20 127 75 186	12 84 73 150	
Supporting tissues	116 934 127 41 204 655	19 131 23 * 33 85	23 166 25 10 38 121	23 172 31 5 36 130	37 291 31 11 58 194	14 173 17 12 40 125	
Obstetrical procedures?74-78	1,077	121	192	169	292	303	
Cessream section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical procedures	246 273 182 376	29 40 26 26	46 55 26 65	41 48 29 50	64 74 45 108	66 55 55 127	
Orthopedic surgery80-90	2,351	453	303	438	678	479	
Excision of bone, partial	157 321 74 284 147 53 120 49 29 32 307 779	55 75 11 37 12 * 72 * 8 56 121	157 12 45 155 7 6 * 6 42 92	21 62 17 57 28 7 14 13 7 7 7 57	40 81 20 85 50 19 17 15 9 7 91 244	26 47 14 58 42 21 11 13 10 * 61	
Plastic surgery92-94	1,041	232	148	179	279	203	
Incision of skin and subcutaneous tissue92.0 Excision of lesion of skin and subcutaneous tissue92.1-92.2 Suture of skin or mucous membrane92.5 Plastic operations on lip and mouth93.1 Skin graft except 1tp and mouth93.0 Other plastic surgery	114 455 195 15 136 126	23 93 57 * 17 41	15 76 30 * 14 13	21 73 38 * 26 18	29 131 42 5 43 30	27 83 28 4 36 24	
Oral and maxillofacial surgery95-98	174	13	21	31	62	47	
Dental surgery99	358	25	70	71	121	72	
Extraction of tooth, forceps extraction99.3 Surgical removal of tooth99.4 Alveoloplasty99.7 Other dental surgery	134 126 48 50	13 * * 6	30 22 9 9	25 26 13 6	40 46 16 19	26 29 7 10	
BiopsyA1-A2	919	98	126	166	284	246	

 $^{^1}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 8. Percent distribution of all-listed operations for inpatients discharged from short-stay hospitals by surgical category, according to bed size of hospital: United States, 1973

			Bed	size of l	nospital	
Surgical category and ICDA codes	All sizes	6-99 beds	100-199 beds	200-299 beds	300-499 beds	500 beds or more
		L				
All operations	100.0	14.3	15.7	distribut	29.2	22.0
Neurosurgery01-05	100.0	12.0	7.6	75.0	91.0	
Ophthalmology06-14	1 1	12.9	7.6 13.9	15.2	31.9	32.4
• •				18.5	31.0	26.3
Resection and recession of eye muscle	100.0	10.4	16.2	20.2	29.1 30.4	24.1 57.5
Other eye surgery	100.0	7.5 14.3	14.4 14.0	19.3 18.4	34.1 28.5	24.8 24.8
Otorhinolaryngology16-21	100.0	15.4	14.6	21.2	30.9	17.9
Myringotomy	100.0	7.4	13.0 22.3	22.9 20.0	36.6 26.4	20.0 29.5
Tympanoplasty	100.0	*	10.3	22.0	36.4	28.5
Tympanoplasty	100.0	15.7 19.4	9.9 18.1	18.6 13.8	39.5 27.1	16.3 21.7
Tracheotomy or tracheostomy, emergency	100.0	*	*	20.7	30.71	36.6
Tonsillectomy with adenoidectomy	100.0	21.2 20.8	16.3 16.8	21.0	27.6 28.2	13.8
Adenoidectomy without tonsillectomy21.3	100.0	20.8	14.3	22.8 19.5	36.0	11.4 24.2
Other operations on ears, nose, and throatResidual	100.0	12.6	11.5	19.5 19.7	32.6	23.5
Operations on thyroid, parathyroid, thymus, and adrenals22-23	100.0	9.6	16.1	16.3	31.7	26.3
Thyroidectomy	100.0 100.0	10.6	16.7	18.4	30.3 37.7	24.0 35.3
Vascular and cardiac surgery24-30	100.0	4.8	7.0	17.3	29.9	40.9
Incision of peripheral vessels	100.0 100.0 100.0 100.0 100.0	13.0 11.1 2.4	16.4 11.8 5.0	20.2 15.1 20.5 *	34.6 36.2 26.9 27.2 29.0	28.7 19.3 29.7 60.6 46.2
Thoracic surgery32-35	100.0	8,0	9.6	18.8	32.4	31.1
Lung lobectomy34,2-34,4 Other thoracic surgeryResidual	100.0 100.0	* 8.7	10.1	22.3 18.3	35.0 32.0	33.0 30.9
Abdominal surgery38-48	100.0	16.7	17.3	18.6	27.2	20.2
Repair of diaphragm and diaphragmatic hernia -38.0-38.1 Repair of inguinal hernia -38.2-38.3 Appendectomy! -41.1 Cholecystectomy- -45.5 Splenectomy- -45.1 Gastric resection, partial or complete- -46.2-46.3 Vagotomy- -46.8 Resection of small intestine or colon- -47.4-47.6 Ileostomy, colostomy, and other enterostomy- -47.7-47.9 Other abdominal surgery- -Residual	100.0 100.0 100.0 100.0 100.0 100.0 100.0 100.0 100.0	20.2 18.4 24.1 17.8 11.5 17.0 11.0 9.3 15.0	14.6 16.4 19.6 18.1 17.7 20.9 14.3 13.6 17.3	16.5 18.9 20.0 18.6 17.0 18.2 14.5 17.8 17.7	21.7 29.4 22.2 27.6 34.5 29.1 23.1 33.0 26.5	27.1 16.9 14.1 17.7 30.8 23.4 24.6 23.8 27.5
Proctological surgery50-52	100.0	13.9	16.0	22.0	30.5	17.5
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	100.0 100.0 100.0	13.1 16.8 11.6	14.2 18.3 14.9	22.9 20.3 23.1	30.8 28.3 32.7	19.0 16.4 17.7
Urological surgery54-61	100.0	10.5	17.7	21.2	29.4	21.2
Nephrotomy and pyelotomy	100.0	* * 7.8 6.5 9.9 * 8.3 7.9 14.0 15.6 3 20.2 11.3	14.6 12.6 16.1 16.3 17.3 20.8 * 18.3 16.9 19.7 * 17.9 20.1 18.2	15.8 17.2 22.9 22.1 23.3 18.5 26.8 27.0 21.0 23.5 15.8 19.5	35.2 29.6 30.4 32.3 29.7 35.0 29.9 28.5 30.0 28.3 27.9 25.8 27.5	26.3 35.4 22.9 22.8 19.9 15.8 16.7 17.8 24.2 19.3 16.5 16.5

Table 8. Percent distribution of all-listed operations for inpatients discharged from short-stay hospitals by surgical category, according to bed size of hospital: United States, 1973—Con.

			Bed	sizes of	hospital	
Surgical category and ICDA codes	All sizes	6-99 beds	100-199 beds	200-299 beds	300-499 beds	500 beds or more
	J		Percent	distribut	ion	L
Breast surgery65	100.0	13,1	17.3	21.0	30.3	18.3
Partial mastectomy65.2 Complete and radical mastectomy65.3-65.6 Other breast surgeryResidual	100.0 100.0 100.0	12.8 13.0 14.3	18.8 15.7 14.1	22.6 19.9 17.0	30.9 30.5 27.5	15.0 20.8 27.1
Gynecological surgery67-72	100.0	14.4	19.2	17.9	28.9	19.6
Local excision or destruction of lesion of ovary	100.0 100.0 100.0 100.0	17.7 14.0 14.8 14.5	18.5 19.2 21.5 20.6	21.2 17.2 14.2 16.2	26.7 29.8 25.1 27.0	15.9 19.8 24.4 21.7
Hysterectomy	100.0 100.0 100.0 100.0 100.0 100.0	16.3 14.1 18.2 * 16.2 12.9	20.1 17.7 19.4 24.4 18.5 18.4	19.5 18.5 24.7 11.3 17.6 19.8	31.6 31.2 24.7 26.9 28.2 29.7	12.5 18.6 13.0 28.5 19.4 19.1
Obstetrical procedures ² 74-78	100.0	11.2	17.8	15.7	27.1	28.2
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical proceduresResidual	100.0 100.0 100.0 100.0	11.6 14.8 14.3 7.0	18.9 20.1 14.2 17.2	16.7 17.7 16.2 13.4	26.1 27.3 25.0 28.6	26.6 20.1 30.4 33.9
Orthopedic surgery80-90	100.0	19.2	12,9	18.6	28,8	20.4
Excision of bone, partial	100.0 100.0 100.0 100.0 100.0 100.0 100.0 100.0 100.0 100.0	34.8 23.3 14.6 13.2 8.1 60.3 * 24.3 18.2 15.5	9.4 17.8 16.4 15.9 10.1 9.2 5.5 11.8 * 18.7 13.7	13.6 19.2 22.8 20.2 18.9 13.2 21.3 25.4 22.8 22.1 18.5	25.5.1 27.0 30.1 34.3 35.3 30.3 31.1 22.8 31.3	16.8 14.5 19.2 20.6 28.6 38.8 9.0 26.9 33.1 *
Plastic surgery92-94	100.0	22.3	14.3	17.1	26.8	19.5
Incision of skin and subcutaneous tissue92.0 Excision of lesion of skin and subcutaneous tissue92.1-92.2 Suture of skin or mucous membrane93.1 Skin graft except lip and mouth93.1 Skin graft except lip and mouth93.2-93.6 Other plastic surgery	100.0 100.0 100.0 100.0 100.0 100.0	19.8 20.4 29.4 * 12.5 32.3	13.2 16.6 15.2 * 10.2 10.2	18.1 16.1 19.3 * 18.8 14.6	25.1 28.8 21.5 30.5 31.7 23.8	23.8 18.2 14.6 27.1 26.8 19.1
Oral and maxillofacial surgery95-98	100.0	7.3	11.9	18.3	35.5	27.0
Dental surgery99	100.0	6.9	19.5	19.8	33.8	20.0
Extraction of tooth, forceps extraction99.3 Surgical removal of tooth99.4 Alveoloplasty99.7 Other dental surgery	100.0 100.0 100.0 100.0	9.9 * 11.4	22.4 17.1 18.4 18.8	18.7 20.7 27.4 12.9	29.8 36.7 33.3 37.6	19.2 23.0 15.0 19.4
BiopsyA1-A2	100.0	10.6	13.7	18.1	30.9	26.7

 $^{^{1}}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 9. Number of first-listed operations and average length of stay for inpatients discharged from short-stay hospitals, by surgical category and sex: United States, 1973

	,				,	
Surgical category and ICDA codes	Total ¹	Male	Female	Total ¹	Male	Female
	оре	of first rations housands			ge leng y in da	
All operations	13,266	5,168	8,091	7.9	8.7	7.5
Neurosurgery01-05	220	111	108	15.0	15.7	14.2
Ophthalmology06-14	550	249	300	5.6	5.4	5.8
Resection and recession of eye muscle	65 32 275 178	29 20 114 86	37 12 160 91	3.7 7.3 6.6 4.5	3.6 6.9 6.3 4.4	3.7 7.9 6.8 4.6
Otorhinolaryngology16-21		693	740	3.1	3.4	2.8
Myringotomy -17.0 Stapedectomy with ossicular reconstruction -17.4 Tympanoplasty -17.6-17.7 Excision of lesion of nose -19.0 Section of nasal septum -19.1 Tracheotomy or tracheostomy, emergency -20.5 Tonsillectomy without adenoidectomy -21.1 Tonsillectomy with adenoidectomy -21.2 Adenoidectomy without tonsillectomy -21.3 Other operations on ears, nose, and throat -Residual	67 27 39 29 51 27 244 629 50 271	39 10 17 19 25 18 87 906 29 146	27 17 22 10 27 9 157 323 22 126	2.3 2.9 2.8 3.5 4.4 22.3 2.1 1.7	2.3 2.7 3.2 3.5 23.1 2.3 2.2 4.6	2.2 2.8 2.8 3.9 5.1 2.1 1.9 1.6 3.9
Operations on thyroid, parathyroid, thymus, and adrenals22-23	71	14	57	7.1	7.2	7.1
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	60 11	9 5	51 6	6.8 8.7	7.2 7.2	6.7 9.9
Vascular and cardiac surgery24-30	568	319	248	11.8	12.1	11.4
Incision of peripheral vessels	19 93 38 16 401	11 23 19 7 259	8 70 19 10 141	17.4 7.0 9.4 22.2 12.5	12.6 8.5 10.9 21.0 12.3	23.9 6.5 7.8 23.0 12.8
Thoracic surgery32-35	173	99	74	13.9	14.7	12.8
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	19 154	12 87	6 67	19.2 13.2	19.2 14.0	19.3 12.2
Abdominal surgery38-48	1,970	1,033	937	10.9	9.9	12.0
Repair of diaphragm and diaphragmatic hernia	19 490 292 371 17 42 36 99 23 583	8 436 166 82 9 24 22 46 12 227	10 54 127 289 7 17 13 53 11 355	13.3 6.2 6.1 12.2 17.9 18.5 16.2 20.8 25.1	11.7 6.2 6.1 14.2 17.1 18.3 16.9 20.1 24.7	14.6 6.3 6.1 11.7 19.1 18.7 14.9 21.5 25.6 12.9
Proctological surgery50-52	386	216	171	7.4	7.1	7.8
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	79 169 139	44 90 82	35 79 57	6.4 7.3 8.1	6.0 7.2 7.7	6.9 7.5 8.6
Urological surgery54-61	1,007	707	300	9.4	9.7	8.5
Nephrotomy and pyelotomy	16 196 203 32 18	11 12 36 49 11 29 5 69 203 32 18 22 84	9 14 26 24 4 10 11 127 	14.7 17.7 8.8 10.4 17.0 3.5 6.1 6.5 13.9 5.3 5.1 6.3	14.6 18.9 7.7 10.5 18.3 3.1 6.1 8.3 13.9 5.3 5.3 3.3 10.4	14.8 16.6 10.4 10.2 13.6 4.7 6.0 5.6

Table 9. Number of first-listed operations and average length of stay for inpatients discharged from short-stay hospitals, by surgical category and sex: United States, 1973—Con.

Surgical category and ICDA codes	Total ¹	Male	Female	Total ¹	Male	Female
	ope	of first rations chousands	in	Avera sta	gth of ays	
Breast surgery65	277	17	260	5.1	3.7	5.2
Partial mastectomy65 2	178	11 ,	174	2 -	11	
Partial mastectomy	66 34	3 7 7	59 27	3.5 9.7 4.5	3.0 3.7 4.0	10.4
Gynecological surgery67-72	2,117		2,117	6.2	 	6.2
Local excision or destruction of lesion of ovary	30 59 190 556	•••	30 59 190 556	7.8 9.4 4.3 9.5		7.8 9.4 4.3 9.5
Supporting tissues 70.2 Dilation and curettage of uterus, diagnostic 70.3 Trachelectomy 70.4 Colporrhaphy 71.4 Plastic repair of cystocele and/or rectocele 71.4 Other gynecological surgery Residual	24 812 39 11 70 326		24 812 39 11 70 326	6.6 4.1 5.6 7.4 9.4 5.4	•••	6.6 4.1 5.6 7.4 9.4 5.4
Obstetrical procedures374-78	1,009		1,009	3.9		3.9
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical procedures	244 263 162 341		244 263 162 341	7.2 2.4 3.7 2.7		7.2 2.4 3.7 2.7
Orthopedic surgery80-90	1,828	939	888	10.7	9.6	12.0
Excision of bone, partial	90 290 66 267 143 48 98 21 20 28 217 540	40 153 40 107 83 20 12 12 12 17 120 322	50 137 26 159 60 28 86 9 9 11 96 218	6.1 8.8 12.2 18.2 15.3 22.8 5.6 16.6 9.0 6.7 4.2	6.5 7.8 11.3 14.5 20.5 5.3 16.8 9.1 5.5 4.5 9.1	5.8 10.0 13.5 20.9 16.4 24.5 5.6 16.3 8.8 8.5 3.8
Plastic surgery92-94	721	391	329	7.8	7.5	8.0
Incision of skin and subcutaneous tissue	96 319 144 10 71 81	58 153 101 7 45 27	38 165 43 * 26 54	8.2 6.7 5.6 5.3 20.5 4.5	7.5 6.4 5.4 5.2 18.6 3.9	9.3 6.9 6.0 * 23.9 4.8
Oral and maxillofacial surgery95-98	119	62	57	6.5	6.8	6.1
Dental surgery99	262	104	157	3.2	3.2	3.2
Extraction of tooth, forceps extraction	111 120 3 28	49 40 * 14	62 80 * 14	3.7 3.0 3.4 2.2	4.2 2.5 * 1.7	3.4 3.2 * 2.6
BiopsyA1-A2	553	213	339	10.8	12.1	10.0

¹Includes data for inpatients discharged with sex not stated.

²Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery.

³Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 10. Number of first-listed operations for inpatients discharged from short-stay hospitals, by surgical category and age: United States, 1973

WATER TO THE PARTY OF THE PARTY					
Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Numi	ber of firs	t-liste		tions
All operations	13,266	1,861	6,037	3,290	2,079
Neurosurgery01-05	220	22	76	83	39
Ophthalmology06-14	550	87	76	143	244
Resection and recession of eye muscle	65 32 275	47 * *	15 6 10	* 14 77 49	11 186
Other eye surgery	178 1,434	38 801	45 466	121	46 47
17.0	67	54	8	4	*
Stapedectomy with ossicular reconstruction	27 39 29 51 27 244	** 11 ** 3 517 587 49	10 16 7 39	14 10 12 10 11 3 *	* 8 * 6 *
Adenoidectomy without tonsillectomy21.3 Other operations on ears, nose, and throat	272	42	148	57	25
Operations on thyroid, parathyroid, thymus, and adrenals22-23	71	4	32	27	9
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	60 11	*	29 3	23 4	7 *
Vascular and cardiac surgery24-30	568	40	129	265	134
Incision of peripheral vessels	19 93 38 16 401	* * 5 * 33		7 44 12 8 194	8 8 9 * 108
Thoracic surgery32-35	173	13	46	64	50
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	19 154	* 13	4 42	9 55	6 45
Abdominal surgery38-48	1,970	272	678	606	414
Repair of diaphragm and diaphragmatic hernia -38.0-38.1 Repair of inguinal hernia -38.2-38.3 Appendectomy¹ -41.1 Cholecystectomy -43.5 Splenectomy -45.1 Gastric resection, partial or complete -46.2-46.3 Vagotomy -46.8 Resection of small intestine or colon -47.4-47.6 Ileostomy, colostomy, and other enterostomy -47.7-47.9 Other abdominal surgery -Residual	19 490 292 371 17 42 36 99 23 583	** 104 98 * * * * * 63	120 162 143 10 9 13 15 *	9 169 24 144 * 21 16 35 7 178	6 97 9 83 * 11 7 47 12 142
Proctological surgery50-52	386	7	213	121	45
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	79 169 139	* * 5	36 84 93	26 70 25	15 14 16
Urological surgery54-61	1,007	178	242	264	323
Nephrotomy and pyelotomy	20 26 62 73 15 15 16 196 203 32 18 22 84 200	* * * 3 3 * * 23 3 * 53 * 7 15 - 444	6 8 25 7 * 8 7 55 * 8 3 12 29 70	8 10 24 20 5 5 4 46 59 11 6 7	6 10 45 7 3 4 4 42 142 6 4 4 43

Table 10. Number of first-listed operations for inpatients discharged from short-stay hospitals, by surgical category and age: United States, 1973—Con.

			,		,=====
Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Number of first-listed operat in thousands				ations
Breast surgery65	277	5	147	88	37
Partial mastectomy65.2 Complete and radical mastectomy65.3-65.6	178	*	107	55	1.3
Complete and radical mastectomy65.3-65.6 Other breast surgery	66 34	*	14 26	28 5	23 *
Gynecological surgery67-72	2,117	14	1,441	564	97
Local excision or destruction of lesion of ovary67.1	30	*	25	*	*
Ophorectomy; salpingo-ophorectomy	59 190	*	45	10	3
Hysterectomy	556	*	189 340	192	23
Local excision and destruction of other lesions of uterus, cervix, and supporting tissues70.2	330		340	1 -72	1
supporting tissues70.2	24	*	12	8	3 34
Dilation and curettage of uterus, diagnostic	812	*	510	265	34
Trache Lectomy	39 11	*	28	7 3	3 *
Plastic repair of cystocele and/or rectocele71.4	70]	6 16	38	16
Supporting tissues	326	7	271	37	îi
Obstetrical procedures ² 74-78	1,009	9	995	4	*
Gesarean section77	244	*	242	*	*
Dilation and curettage after delivery or abortion78.1	263	*	260	*	-
Repair of laceration	162 341	* 6	160 333	*	-
Negratal	341	ا	333	"	•
Orthopedic surgery80-90	1,828	223	765	485	355
Excision of bone, partial	90	ا و	47	28	6
Closed reduction of fracture without fixation82.0	290	100	86	51	54
Open reduction of fracture without fixation82.1	66	10	30	1.5	11
Reduction of iracture with fixation	267	13	69	48	137
Arthroniasty of hip	143 48	*	78 3	60 19	6 26
Repair and plastic operations on joints of foot and toes87.2	98	*	34	48	13
Spinal fusion87.4	21	*	11	8	*
Arthrodesis and stabilization of joints (except spine)87.5-87.6	20	*	- 9	7	*
Glosed reduction of dislocation of joint8,7,7	28	5	14	-4	.5
Arthrodesis and stabilization of joints (except spine)	217 540	33 45	113 273	56 142	15 80
Plastic surgery92-94					
Incision of skin and subcutaneous tissue92.0	721	122	323	175	100
Excision of legion of skin and subsystematic tiesus	96 319	22 45	42 124	19 94	11 55
Suture of skin or mucous membrane92.5	144	23	87	20	14
Suture of skin or mucous membrane	10	7	*	*	*
Skin graft except lip and mouth93.2-93.6	71	12	33	16	10
Í	81	13	34	25	9
Oral and maxillofacial surgery95-98	119	17	57	28	17
Dental surgery99	262	26	176	47	13
Extraction of tooth, forceps extraction	111	7	69	27	8
Surgical removal of tooth99.4	120	7	97	13	4
Alveoloplasty99.7	3		*	*	*
Ocner dental surgeryResidual	28	13	9	5	*
BiopsyA1-A2	553	21	173	204	155

 $^{^{1}}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

Table 11. Average length of stay for inpatients discharged from short-stay hospitals, by surgical category of first-listed operation and age: United States, 1973

					
Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Average length of stay in de			lays	
All operations	7.9	4.0	6.0	9.7	14.3
Neurosurgery01-05	15.0	13.7	12.2	16.2	18.4
Ophthalmology06-14	5.6	3.2	4.7	5.8	6.6
Resection and recession of eye muscle	3.7 7.3 6.6 4.5	3.1 * * 3.2	5.2 7.3 5.7 4.0	7.1 6.4 4.5	7.5 6.7 6.0
Otorhinolaryngology16-21	3,1	2.1	3.3	6.6	8.8
Myringotomy- 17.0 Stapedectomy with ossicular reconstruction 17.4 Tympanoplasty- 17.6-17.7 Excision of lesion of nose- 19.0 Section of nasal septum- 19.1 Tracheotomy or tracheostomy, emergency- 20.5 Tonsillectomy without adenoidectomy- 21.1 Tonsillectomy without adenoidectomy- 21.2 Adenoidectomy without tonsillectomy- 21.3 Other operations on ears, nose, and throat- Residual	2.3 2.9 2.8 3.5 4.4 24.4 2.3 2.1	2.0 * 2.3 * 25.4 2.0		5.5 3.2 3.4 3.3 4.1 25.3 2.8	* * * 4.7 * 24.1 *
Adenoidectomy without tonsillectomy21.3 Other operations on ears, nose, and throatResidual	1.7 4.3	1.7 2.6	3.6	5.8	8.1
Operations on thyroid, parathyroid, thymus, and adrenals22-23		3.3	6.4	7.8	9.2
Thyroidectomy22.1-22.2 Other operations on thyroid, parathyroid, thymus, and adrenalsResidual	6.8 8.7	* 3.0	6.3 7.1	7.1 12.0	7.9
Vascular and cardiac surgery24-30	11.8	8.5	8.5	11.8	15.9
Incision of peripheral vessels	17.4 7.0 9.4 22.2 12.5	* 5.0 * 8.8	5.5	15.1 7.7 11.0 24.9 12.2	22.9 11.6 14.1 * 15.7
Thoracic surgery32-35	13.9	8.7	10.4	15.2	16.7
Lung lobectomy34.2-34.4 Other thoracic surgeryResidual	19.2 13.2	8.7	14.6 10.0	20.9 14.2	19.7 16.3
Abdominal surgery38-48	10.9	4.9	8.7	12.2	16.7
Repair of diaphragm and diaphragmatic hernia		2.6 5.8 * * * * * * 6.1	16.7	12.7 7.2 8.4 12.1 * 16.6 15.1 18.6 26.3	15.8 9.1 10.7 16.8 22.5 22.5 22.4 25.0 19.0
Proctological surgery50-52	7.4	6.6	5.8	8.2	12.8
Local excision and destruction of lesion of rectum and anus50.2,51.2 Hemorrhoidectomy51.3 Other proctological surgery	6.4 7.3 8.1	* * 8.1	5.5 6.5 5.4	6.1 7.5 12.3	9.6 11.4 16.8
Urological surgery54-61	9.4	3.8	7.1	10.2	13.4
Nephrotomy and pyelotomy	10.4 17.0 3.5 6.1 6.5 13.9 5.3 5.1	* * * 6.0	3.2 4.7 5.9 * 4.8 8.8 2.1 3.2	15.2 4.9 6.2 7.8 12.6 5.9 9.5 4.2	19.3 20.2 10.1 10.9 18.5 14.3 8.5 11.3 7.2 13.3 6.4 14.4

Table 11. Average length of stay for inpatients discharged from short-stay hospitals, by surgical category of first-listed operation and age: United States, 1973—Con.

. Surgical category and ICDA codes	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Ave	erage lengt	h of st	ay in d	ays
Breast surgery65	5.1	2.9	3.6	5.8	9.7
Partial mastectomy	3.5 9.7 4.5	* *	3.0 6.5 4.4	3.9 9.8 4.7	6.4 11.9 *
Gynecological surgery67-72	6.2	4.8	5.6	6.7	11.0
Local excision or destruction of lesion of ovary	7.8 9.4 4.3 9.5	* * *	7.1 8.7 4.3 9.2	10.8 * 9.7	15.6 12.7
Supporting tissues	6.6 4.1 5.6 7.4	* * - *	4.7 3.8 4.9 6.1	4.9 4.1 6.6 8.9	19.4 7.3 9.2
Plastic repair of cystocele and/or rectocele71.4 Other gynecological surgery	9.4 5.4	3.6	9.2 4.9	9.1 6.4	10.4 16.2
Obstetrical procedures ² 74-78	3.9	2.7	3.9	4.2	*
Cesarean section77 Dilation and curettage after delivery or abortion78.1 Repair of laceration78.2-78.3 Other obstetrical proceduresResidual	7.2 2.4 3.7 2.7	* * * 1.8	7.2 2.4 3.7 2.7	* * *	* - - *
Orthopedic surgery80-90	10.7	6.1	7.9	10.8	19.6
Excision of bone, partial	6.7 4.2 10.0	5.2 5.3 5.9 11.5 ** ** 5.7 4.3 6.9	10.0 11.5 14.1 22.9 5.0 15.7 8.3 5.9	6.6 9.3 13.2 16.4 16.1 19.7 5.7 14.7 9.0 3.6 5.0 10.9	8.5 14.3 21.9 22.9 22.5 25.3 6.5 * 12.9 4.6 21.4
Plastic surgery92-94		5.6	6.7	9.0	11.6
Incision of skin and subcutaneous tissue	8.2 6.7 5.6 5.3 20.5 4.5	5.2 3.9 3.6 6.1 18.1 4.0	6.0 5.1 *	11.2 7.1 8.1 23.2 5.7	14.0 9.5 7.7 * 29.8 6.6
Oral and maxillofacial surgery95-98	6.5	4.4	5.7	7.2	9.7
Dental surgery99		1.9	2.9	4.5	6.1
Extraction of tooth, forceps extraction	3.7 3.0 3.4 2.2	2.5 2.1 1.4	*	*	*
BiopsyAl-A2	1	19.3	6.4	10.4	15.0

 $^{^1}$ Limited to estimated number of appendectomies excluding those performed incidental to other abdominal surgery. 2 Codes 75.0-75.6 and 75.9 are not used by HDS.

APPENDIX I

TECHNICAL NOTES ON METHODS

Statistical Design of the Hospital Discharge Survey

Scope of the survey.—The scope of the HDS encompasses patients discharged from noninstitutional hospitals, exclusive of military and Veterans Administration (VA) hospitals, that have six beds or more for inpatient use, are located in the 50 States and the District of Columbia, and in which the average length of stay for all patients is less than 30 days. Although all discharges of inpatients from these hospitals are within the scope of this survey, all newborn infants and discharges from Federal hospitals are excluded from this report.

Sampling frame and size of sample.—The sampling frame (universe) for hospitals in the HDS is the Master Facility Inventory of Hospitals and Institutions (MFI). A detailed description of how the MFI was developed, its contents, plans for maintaining it, and procedures for assessing the completeness of its coverage has been published. 12

The universe for the survey consisted of 6,965 short-stay hospitals, excluding military and Veterans Administration hospitals, contained in the MFI in 1963. The universe was expanded in 1969 by the addition of 442 hospitals which were added to the MFI. The distribution of the hospitals by size and geographic region in the MFI and the HDS sample for 1973 is shown in table I.

The sample of hospitals for 1973 consisted of 497 hospitals. Of these hospitals, 42 refused to participate and 31 were out of scope either because the hospital had gone out of business or because they failed to meet the definition of a short-stay hospital. Thus 424 hospitals participated in the survey during 1973. Approximately 225,000 abstracts of medical records were received from the 424 hospitals.

Sample design.—All hospitals with 1,000 beds or more in the universe of short-stay hospitals were selected with certainty in the sample. All hospitals with fewer than 1,000 beds were stratified, the primary strata being the 24 size-by-region classes shown in table I. Within each of these 24 primary strata, the allocation of the hospitals was made through a con-

trolled selection technique so that hospitals in the sample would be properly distributed with regard to ownership and geographic division. Sample hospitals were drawn with probabilities ranging from certainty for the largest hospitals to 1 in 40 for the smallest hospitals.

The within-hospital sampling ratio for selecting sample discharges varied inversely with the probability of selection of the hospital. The smallest sampling fraction of discharged patients was taken in the largest hospitals, and the largest fraction was taken in the smallest hospitals. This was done to compensate for the fact that hospitals were selected with probabilities proportionate to their size class and to assure that the overall probability of selecting a discharge would be approximately the same in all hospitals.

In nearly all the sample hospitals, the daily listing sheet of discharges was the frame from which the subsamples of discharges were selected. The sample discharges were selected by a random technique, usually on the basis of the terminal digit(s) of the patient's medical record number—a number assigned when the patient was admitted to the hospital. If the hospital's daily discharge listing did not show the medical record numbers, the sample was selected by starting with a randomly selected discharge and taking every kth discharge thereafter.

Data Collection and Processing

Data collection.—Depending on the study procedure agreed on with the hospital administrator, the sample selection and the transcription of information from the hospital records to abstract forms were performed either by the hospital staff or by representatives of NCHS, or by both. In approximately 70 percent of the hospitals that participated in the HDS during 1973, this work was performed by the medical records department of the hospital. In all the remaining hospitals, the work was performed by personnel of the U.S. Bureau of the Census acting for NCHS.

Survey hospitals used an abstract form to transcribe data from the hospital records (figure I). The abstract form provides for recording demographic

Table I. Distribution of short-stay hospitals in the universe (MFI) and in the Hospital Discharge Survey sample, and the number of hospitals that participated in the survey, by size of hospital and geographic region: United States, 1973

Size of hospital	All regions	North- east	North Central	South	West	
All sizes		Number of hospitals				
Universe Total sample Number participating	7,407 407 424	1,146 129 115	2,064 146 128	2,832 148 118	1,365 74 63	
<u>6-49 beds</u>					1	
Universe Total sample Number participating	3,304 64 43	209 7 5	865 18 15	1,549 26 14	681 13 9	
<u>50-99 beds</u>						
Universe	1,746 72 61	293 13 10	467 19 16	642 27 24	344 13 11	
100-199 beds						
Universe Total sample Number participating	1,224 103 91	288 26 26	392 31 26	365 32 25	179 14 14	
200-299 beds						
Universe Total sample Number participating	583 89 76	191 30 27	158 26 23	140 19 14	94 14 12	
300-499 beds						
Universe Total sample Number participating	397 93 84	111 25 23	131 30 28	102 26 24	53 12 9	
500-999 beds						
Universe Total sample Number participating	135 58 51	45 19 15	48 19 17	20 18 12	13 7 7	
1,000 beds or more						
Universe Total sample Number participating	18 18 18	9 9 9	3 3 3	5 5 5	1 1 1	

data, admission and discharge dates, discharge status, and information on discharge diagnoses and surgical operations or procedures. All discharge diagnoses and operations were listed on the abstract form in the order in which they were entered on the face sheet of the hospital medical records.

Shipments of completed abstract forms for each sample hospital were transmitted, along with sample selection control sheets, to NCHS for processing. Every shipment of abstracts was reviewed and each abstract form was checked for completeness.

Medical coding and edit.—The medical information recorded on the sample patient abstracts for data

year 1973 were coded centrally by NCHS staff. A maximum of five diagnostic codes and three codes for surgical operations and procedures were assigned for each sample abstract. Following the conversion of the data on the medical abstract to computer tape, a final medical edit was accomplished by computer inspection runs and a review of rejected abstracts. If sex and/or age of patient was incompatible with the recorded medical information, priority was given to the latter in the editing decision.

The basic system used for coding the medical terminology on HDS sample patient abstracts is the ICDA, However, some modifications were made to ac-

CONFIDENTIAL - All information which would permit identification of an individual or of an establishment will be held confidential, will be used only by persons engaged in and for the purposes of the survey and will not be disclosed or released to other persons or used for any other purpose.

DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE Public Health Service Health Services and Mental Health Administration

National Center for Health Statistics MEDICAL ABSTRACT — HOSPITAL DISCHARGE SURVEY

I. Patier	it Identification				
	Hospital number	4. Date of admission	·		
2.	HDS number		Month	Day	Year
3.	Medical record number	5. Date of discharge	Month	Day	Year
II. Patier	t Characteristics				
1.	Date of birth;	2. Age (complete ONL) if date of birth not g		{s}_2[years months days
3.	Sex: 1 Male 2 Female				
4.	Race or color: 1 White 2 Negro	3 Other nonwhite 4 🗆 "	Nonwhite"	5 🗌 Not s	stated
5.	Marital status: 1 ☐ Married 2 ☐ Single	3 Widowed 4 Divorced	5 Separated	6 🔲 או	ot stated
6.	Discharge status: 1 ☐ Alive 2 ☐ De	ead	********		
	Final diagnoses:				
				see	reverse side
2.	Operations:				
				see	reverse side
Complete	d by	Date			
FOR NCH	S USE ONLY				
Diagnose Operation	s				

Figure I. Medical Abstract Form

commodate incomplete or ill-defined terminology on the source documents. $% \left\{ \mathbf{r}_{i}^{\mathbf{r}_{i}}\right\} =\mathbf{r}_{i}^{\mathbf{r}_{i}}$

The Hospital Discharge Survey modifications of the ICDA diagnostic coding system exclude the sections Accidents, poisonings, and violence (external cause) (E800-E999) and Fetal death (Y30). The ICDA class XV, Certain Causes of Perinatal Morbidity (760-779) was modified to exclude diseases, difficult labor, and conditions of mothers of newborn infants (760-771), termination of pregnancy (773), and fetal death of unknown cause (779). Birth injury without mention of

cause (772) was changed to birth injury. Codes retained in this ICDA class are 772 and 774-778. Code 793 for observation, without need for further medical care was restated as observation and tests with negative or unspecified findings and was moved from ICDA class XVI, Symptoms and Ill-Defined Conditions to the supplementary classification section on Special Conditions Without Sickness (Y00-Y13).

The Hospital Discharge Survey modifications of the ICDA system for coding surgical operations and procedures are shown on page 49.

HDS modifications of the ICDA section Surgical Operations, Diagnostic and Other Therapeutic Procedures

ICDA Code and Title	HDS Modification
01.1-Transsphenoid surgical approach to brain stem	01.1 not used; included in 01.7-Hypophysectomy
14.4-Extraction of lens, extracapsular	
14.5-Extraction of lens, intracapsular	
***************************************	Add 14.6-Extraction of lens or cataract, not otherwise specified
22.6-Excision of branchial cleft cyst	22.6 not used; included in 21.6-Pharyngectomy and destruction of lesion of pharynx
30.6-Open heart technique	Redefined 30.6-Open heart surgery (with cardiopulmonary bypass) not otherwise specified
30.7-Open heart technique with cardiopulmonary bypass	30.7 not used; included in 30.6
56.7-Urethrovesical suspension	56.7 not used; included in 57.4-Repair and plastic operations on urethra
65.7-Repair or plastic operations on breast	
	Add 65.8-Augmentation mammoplasty
65.9-Other operations on breast	Redefined 65.9-Other operations on breast (includes reduction or amputative mammoplasty)
74-78-Obstetrical procedures	75.0-75.6, 75.9 not used
77.0-Cesarean section, classical	Redefined 77.0-Cesarean section, all types
	77.1-77.2, 77.8-77.9 not used
82-84-Reduction of fracture and fracture dislocation of bones	82.3-82.9, 83-84 not used
82.0-Closed reduction of separated upper femoral epiphysis	Redefined 82.0-Reduction (closed or not otherwise specified) of fracture in 82-84 without mention of fixation
82.1-Open reduction of separated upper femoral epiphysis with or without internal fixation	Redefined 82.1-Reduction (open) of fracture in 82-84 without mention of fixation
82.2-Closed reduction of intertrochanteric fracture	Redefined 82.2-Reduction (closed or open) of fracture in 82-84 with mention of fixation
87.0-Arthroplasty of hip without mechanical device	Redefined 87.0-Arthroplasty of hip with or without mechanical device
87.1-Arthroplasty of hip with mechanical (prosthetic) device	87.1 not used; included in 87.0
94.2-Plastic operation of nose	94.2 not used; included in 19.3-Rhinoplasty and repair of nose
94.4-Augmentation mammoplasty	94.4 not used; see code 65.8
98-Reduction of fracture and fracture-dislocation of jawbone	98.2-98.6 not used
98.0-Closed reduction, malar, zygoma and zygomatic arch	Redefined 98.0-Reduction (closed), malar, zygoma, zygomatic arch, maxilla, mandible, alveolus
98.1-Open reduction, malar, zygoma and zygomatic arch	Redefined 98.1-Reduction (open), malar, zygoma, zygomatic arch, maxilla, mandible, alveolus
A4-A5-Diagnostic endoscopy	A4-A5 not used
A8-A9-Diagnostic radiography	A8-A9 not used
R1-Radiotherapy and related therapies	R1 not used
R4-Physical medicine and rehabilitation	R4 not used
R9-Other surgical procedures	R9 not used

Presentation of Estimates

Grouping of operations.—Estimates of the number, rate, and distribution of operations by surgical class are based on the classification of operations reported on sample patient abstracts in the 3-digit detail provided by the ICDA. The groupings that are used in this report are specialties numbered 1-17 of the ICDA section Surgical Operations, Diagnostic and Other Therapeutic Procedures. The surgical categories, the most detailed groupings of surgical operations or procedures shown in this report, are subsets of the major groups or classes. In developing the tabular list of operations, an effort was made to maximize specificity of the operations consistent with clarity of characterization and with the frequency of their occurrence within the context of the HDS for 1973.

Patient characteristics "not stated."-Age and sex of patient were not stated on the hospital records of sample hospitals (the face sheet of patient's medical record) for less than one-fourth of 1 percent of the discharges. If age was not stated, it was imputed by assigning the patient an age consistent with the ages of other patients with the same diagnostic code. If the dates of admission or discharge were not given, and if they could not be obtained from the monthly sample listing sheet transmitted by the sample hospital, a length of stay was imputed by assigning the patient a stay consistent with the stays of other patients of the same age. Other missing demographic items were coded and tabulated as "not stated," Color was not stated for 12.8 percent of all discharges. For this reason, rates by color were not computed and caution should be used in drawing conclusions from the data by color. In the detailed tables presenting frequencies and rates, the totals include the not stated cases.

Rounding of numbers.—Estimates of the number of discharges, discharges with surgery, and operations have been rounded to the nearest thousand for tabular presentation. For this reason, detailed figures within the tables do not always add to totals. Rates, percents, and average lengths of stay presented in the report were calculated on the basis of unrounded figures and will not necessarily agree with rates and other calculations that may be calculated from the rounded data.

Population estimates.— The population estimates used in computing rates for 1973 HDS data are unpublished estimates for the U.S. civilian noninstitutionalized population as of July 1, 1973, provided by the U.S. Bureau of the Census.

The population estimates for the United States by age and sex and by geographic region, presented in table II, are consistent with the population estimates published by the U.S. Bureau of the Census in *Current Population Reports*, Series P-25. However, they are not official population estimates of the U.S. Bureau of the Census.

Table II. Civilian noninstituitionalized population used to compute rates shown in this publication, by age, sex, and geographic region: United States, July 1, 1973

Characteristic	Both sexes	Male	Female
	Populati	on in the	usands ¹
Total	205,836	99,307	106,529
Age			
Under 15 years 15-44 years 45-64 years 65 years and over	55,559 87,342 42,641 20,294	28,313 42,253 20,310 8,431	27,246 45,088 22,331 11,862
Geographic region			
Northeast North Central South West	48,940 56,772 64,499 35,625	•••	•••

¹These estimates of the U.S. civilian noninstitutionalized population are consistent with the population estimates published by the U.S. Bureau of the Census in <u>Current Population</u> <u>Reports</u>, Series P-25.

Reliability of Estimates

Estimation.—Statistics produced by HDS are derived by a complex estimating procedure. The basic unit of estimation is the sample inpatient discharge abstract. The estimating procedure used to produce essentially unbiased national estimates in the HDS has three principal components: inflation by reciprocals of the probabilities of sample selection, adjustment for nonresponse, and ratio adjustment to fixed totals. These components of estimation are described in appendix I of two earlier publications. 13,14

Measurement errors.—As in any survey, the results are subject to nonsampling or measurement errors, which include errors due to hospital nonresponse, missing abstracts, information incompletely or inaccurately recorded on abstract forms, and processing errors. Some of these errors were discussed earlier in this report. Quality control programs have been instituted to alleviate these types of errors.

Sampling errors.— The standard error is primarily a measure of variability that occurs by chance because a sample rather than the entire universe is surveyed. In this report, the standard error also reflects part of the measurement error, but does not measure any systematic biases in the data. The relative standard error of the estimate is obtained by dividing the standard error of the estimate by the estimate itself and is expressed as a percentage of the estimate.

Table III. Approximate standard error of percentages shown in this report for discharges or operations: Patient characteristics crossclassified by geographic region and bed size of hospital and for all hospitals

Number of discharges or	Estimated percent						
operations (base of per- cent in thou- sands)	2 or 98	4 or 96	10 or 90	20 or 80	30 or 70	50	
	Standard error expressed in percentage points						
1.00	0.8 0.6 0.3 0.2 0.1 0.1 0.1	1.2 0.8 0.5 0.4 0.3 0.2 0.1 0.1		2.4 1.7 1.0 0.8 0.5 0.3 0.2 0.1	2.8 2.0 1.1 0.9 0.6 0.4 0.3 0.2	3.0 2.1 1.2 1.0 0.7 0.4 0.3 0.2 0.2	

NOTE: Illustration of use of table III: Table 8 shows that 18.6 percent of the 411,000 cholecystectomies reported in table 7 were performed in hospitals of 200-299 beds. Linear interpolation between the values shown in table III yields an approximate standard error of 1.3 percent for an estimate of 18.6 percent with a base of 411,000.

The chances are about 68 out of 100 that the value obtained in a complete enumeration is contained in the interval represented by the estimate plus or minus one standard error of the estimate; 95 out of 100 for two standard errors; and 99 out of 100 for 2½ standard errors. Applying the illustration at the bottom of figure II, the chances are about 68 out of 100 that the value that would be obtained in a complete enumeration is contained in the interval 125,000 + 10.7 percent of 125,000 (between 111,625 and 138,375); 95 out of 100 for the interval 125,000 + 10.7 percent of 125,000 multiplied by 2; 99 out of 100 for the interval 125,000 multiplied by 2.5.

The standard error of one statistic is generally different from that of another, even when the two come from the same survey. In order to derive standard errors that would be applicable to a wide variety of statistics that could be prepared at a moderate cost, a number of approximations are required. As a result, figure II and tables III and IV shown in this section provide general standard errors for a wide variety of estimates rather than the specific error for a particular statistic.

The relative standard errors (figure II) and approximate standard errors of percentages (table III) and of average lengths of stay (table IV) that have been prepared for this report are applicable to estimates of discharges and number of operations for all hospitals and by type of surgery or specific procedure

Table IV. Approximate standard errors of average lengths of stay shown in this report:
Patient characteristics cross-classified by all hospitals

Number of discharges (base of	Average length of stay in days							
average in thousands)	2	6	10	14	¹⁸ .	22	26	30
	Standard error in days							
4	0.9 0.6 0.3 0.2 0.1 0.1 0.1 0.1	1.4 0.9 0.5 0.4 0.3 0.2 0.2 0.2	2.1 1.4 0.7 0.6 0.4 0.4 0.4 0.4	2.9 1.8 0.9 0.6 0.6 0.5 0.5	3.7 2.4 1.2 1.0 0.7 0.7 0.7 0.7 0.7	4.6 2.9 1.5 1.2 0.9 0.8 0.8 0.8	5.5 3.5 1.7 1.4 1.0 1.0 1.0	6.4 4.1 2.0 1.6 1.2 1.2 1.2

NOTE: Illustration of use of table IV: Table 11 shows that the average length of stay was 14.1 days for the estimated 78,000 discharged patients age 15-44 years with surgery for excision of intervertebral cartilage (table 10). Linear interpolation between the values shown in table IV will yield an approximate standard error of 0.8 days for an estimated average length of stay of 14.1 days with a base of 78,000.

Figure II. Approximate relative standard errors of the estimated numbers of discharges or operations for inpatients discharged from all short-stay hospitals, and for discharges by geographic region and bed size of hospital




Illustration for use of figure II: As shown in table 5, an estimated 125,000 cholecystectomies were performed during 1973 for inpatient discharges from short-stay hospitals within the North Central Region. The relative standard error of the estimate as read from the curve "Region groups" is approximately 10.7 percent: the standard error of 125,000 is 13,375 (10.7 percent of 125,000).

cross-tabulated by age, sex, color, geographic region, and size of hospital. The curve in figure II to which one refers to obtain a sampling error is contingent on whether the type of estimate (for example, operations) relates to all hospitals, geographic region, or a hospital size group. The approximate standard errors of

estimated percentages in table III, when the characteristic(s) used to form the numerator of the percentage is a subclass of the denominator, and the approximate standard errors of average lengths of stay in table IV, are applicable to all percentages and all average lengths of stay presented in this report.



APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

Terms Relating to Hospitalization

Short-stay hospitals.—General and short-term special hospitals having six beds or more for inpatient use and an average (mean) length of stay of less than 30 days. Federal hospitals and hospital units of institutions are not included. The terms "hospitals" and "short-stay hospitals" are used synonymously.

Inpatient.—A person who is formally admitted to the inpatient service of a short-stay hospital for observation, care, diagnosis, or treatment. In this report, the number of inpatients refers to the number of discharges during 1973, including multiple discharges (if any) of the same individual from one short-stay hospital or more. Newborn infants admitted by birth to the hospital from which they are discharged are excluded in this report. The terms "inpatient" and "patient" are used synonymously.

Discharge.—The formal release of an inpatient by a hospital, that is, the termination of a period of hospitalization by death or by disposition to place of residence, nursing home, or another hospital. In this report, the number of discharges from short-stay hospitals (alive or dead) is exclusive of newborn infants. The number of discharges by death is limited to hospital deaths that occurred following formal admission of the patient to the inpatient service of the short-stay hospital. The terms "discharges," "inpatient discharges," and "patients (or inpatients) discharged" are used synonymously.

Average length of stay.—The total number of inpatient days accumulated at the time of discharge by patients with surgery discharged during 1973 divided by the number of patients with surgery. A stay of less than 1 day (admission and discharge on the same calendar day) is counted as 1 day in the summation of total inpatient days. The terms "average length of stay" and "average hospital stay" are used synonymously.

Bed size of hospital.—Measured by the number of beds, cribs, and pediatric bassinets regularly maintained (set up and staffed for use) for inpatients; bassinets for newborn infants are not included. In this report the classification of hospitals by bed size is

based on the number of beds at or near midyear reported by the hospitals.

Terms Relating to Surgery

Discharges with surgery. —The estimated number of surgically treated patients discharged from non-Federal short-stay hospitals during 1973.

Operation.—One or more surgical operations, procedures, or special treatments that are assigned by the physician to the medical record of patients discharged from the inpatient service of short-stay hospitals. In this survey, all terms listed on the face sheet (summary sheet) of the medical record under the captions "operation," "operative procedures," "operations and/or special treatments," and the like are transcribed in the order listed. A maximum of three 3-digit codes are assigned per sample discharge according to the ICDA and Hospital Discharge Survey directives. (See "Medical coding and edit" in appendix I for further details.) "Operations," "surgical operations," and "surgical procedures" are used synonymously.

All-listed operations.—In terms of the classification system used and the coding of not more than three operations per sample discharge, the estimated number of surgical operations and procedures performed for inpatient discharges from non-Federal short-stay hospitals during 1973. Refers to the aggregate of individually coded operations, procedures, and special treatments in code positions 1-3 exclusive of certain obstetrical procedures, diagnostic endoscopy and radiography, radiotherapy, and certain other treatments not generally considered as surgery.

First-listed operations.—The aggregate of individually coded surgical operations and procedures listed first, including single (only) and first of multiple operations listed. The number of first-listed operations is equivalent to the number of discharges with surgery.

Surgically treated patients.—Inpatients for whom at least one operation or procedure is performed during one period of hospitalization exclusive of certain obstetrical procedures, diagnostic endoscopy and

radiography, radiotherapy, and certain other treatments not generally considered as surgery.

Volume of operations.—The estimated number of all-listed operations performed for inpatient discharges from non-Federal short-stay hospitals during 1973.

Surgery rate.—The ratio of the number of alllisted operations performed during 1973 to the number of persons in the civilian noninstitutionalized population as of July 1, 1973.

Obstetrical procedures.—Certain procedures in ICDA surgery class 12. The HDS includes antepartum obstetrical operations, operations inducing or assisting delivery, cesarean section, and operations after delivery or abortion, but excludes certain routine procedures. (See "Medical coding and edit" in appendix I for further details.)

Demographic Terms

Age.—Refers to age at last birthday prior to admission to the hospital inpatient service.

The elderly.—Persons 65 years of age and over. Color.—In this report, patients are classified into two groups, "white" and "all other," based on information available on the hospital records (face sheet of the inpatient's medical record) of sample hospitals. "White" includes Mexican and Puerto Rican unless patient is specifically identified as other than white.

 ${\it United States.}{
m - The 50 \ States}$ and the District of Columbia.

Geographic region.—In this report, hospitals are classified by location according to the four geographic regions of the United States which correspond to those of the U.S. Bureau of the Census. The States included in each region are as follows:

ized population	Region	States Included
procedures in es antepartum ing or assist- erations after n routine pro-	Northeast	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania
' in appendix I	North Central	Ohio, Indiana, Illinois, Michigan, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, Kansas
age and over. are classified er," based on records (face rd) of sample	South	Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, Texas
as other than	West	Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, California, Alaska, Hawaii
	0 —	

★U.S. GOVERNMENT PRINTING OFFICE: 1976-210-981:38

VITAL AND HEALTH STATISTICS PUBLICATIONS SERIES

Formerly Public Health Service Publication No. 1000

- Series 1. Programs and Collection Procedures.—Reports which describe the general programs of the National Center for Health Statistics and its offices and divisions, data collection methods used, definitions, and other material necessary for understanding the data.
- Series 2. Data Evaluation and Methods Research.—Studies of new statistical methodology including experimental tests of new survey methods, studies of vital statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, contributions to statistical theory.
- Series 3. Analytical Studies.—Reports presenting analytical or interpretive studies based on vital and health statistics, carrying the analysis further than the expository types of reports in the other series.
- Series 4. Documents and Committee Reports.—Final reports of major committees concerned with vital and health statistics, and documents such as recommended model vital registration laws and revised birth and death certificates.
- Series 10. Data from the Health Interview Survey.—Statistics on illness; accidental injuries; disability; use of hospital, medical, dental, and other services; and other health-related topics, based on data collected in a continuing national household interview survey.
- Series 11. Data from the Health Examination Survey.—Data from direct examination, testing, and measurement of national samples of the civilian, noninstitutionalized population provide the basis for two types of reports: (1) estimates of the medically defined prevalence of specific diseases in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics; and (2) analysis of relationships among the various measurements without reference to an explicit finite universe of persons.
- Series 12. Data from the Institutionalized Population Surveys.—Discontinued effective 1975. Future reports from these surveys will be in Series 13.
- Series 13. Data on Health Resources Utilization.—Statistics on the utilization of health manpower and facilities providing long-term care, ambulatory care, hospital care, and family planning services.
- Series 14. Data on Health Resources: Manpower and Facilities.—Statistics on the numbers, geographic distribution, and characteristics of health resources including physicians, dentists, nurses, other health occupations, hospitals, nursing homes, and outpatient facilities.
- Series 20. Data on Mortality.—Various statistics on mortality other than as included in regular annual or monthly reports. Special analyses by cause of death, age, and other demographic variables; geographic and time series analyses; and statistics on characteristics of deaths not available from the vital records, based on sample surveys of those records.
- Series 21. Data on Natality, Marriage, and Divorce.—Various statistics on natality, marriage, and divorce other than as included in regular annual or monthly reports. Special analyses by demographic variables; geographic and time series analyses; studies of fertility; and statistics on characteristics of births not available from the vital records, based on sample surveys of those records.
- Series 22. Data from the National Mortality and Natality Surveys.—Discontinued effective 1975. Future reports from these sample surveys based on vital records will be included in Series 20 and 21, respectively.
- Series 23. Data from the National Survey of Family Growth.—Statistics on fertility, family formation and dissolution, family planning, and related maternal and infant health topics derived from a biennial survey of a nationwide probability sample of ever-married women 15-44 years of age.
- For a list of titles of reports published in these series, write to:
 Scientific and Technical Information Branch
 National Center for Health Statistics
 Public Health Service, HRA
 Rockville, Md. 20852

DHEW Publication No. (HRA) 76-1775 Series 13-No. 24

NCHS

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

Public Health Service
HEALTH RESOURCES ADMINISTRATION
5600 Fishers Lane
Rockville, Md. 20852

OFFICIAL BUSINESS
Penalty Private Use, \$300

For publications in the Vital and Health Statistics Series call 301-443-NCHS. POSTAGE AND FEES PAID U.S. DEPARTMENT OF H.E.W

HEW 390

THIRD CLASS BLK. RATE

