

Effects of Cattle Manure Handling and Management on Fate and Transport of Hormones in the Feedlot and the Field

EPA-G2006-STAR-M1: Fate and Effects of Hormones in Waste From Concentrated Animal Feeding Operations (CAFOS)



Research Team

- Dan Snow, SNR Water Sciences Laboratory
- Shannon Bartelt-Hunt, Civil Engineering
- Bill Kranz, Biological Systems Engineering
- Terry Mader, Animal Science
- Charles Shapiro, Soil Scientist - Crop Nutrition
- David Shelton, Biological Systems Engineering
- Tian Zhang, Civil Engineering
- Simon von Donk, Water Resources Engineer
- David Tarkalson, USDA-ARS NW Irrigation and Soils Research Lab, Kimberly, ID
- Steve Ensley, College of Veterinary Medicine, Iowa State University, Ames. IA

UNIVERSITY OF
Nebraska
Lincoln

Pioneering new frontiers.



EPA Request for Proposals

Purpose of Program

- “Characterize the occurrence, magnitude, and extent of the impact of natural and synthetic steroid hormones in liquid and solid animal waste from concentrated animal feeding operations (CAFOs) on the environment and human health”
- “Determine the impact of current CAFO waste management strategies (i.e. storage and disposal) on the transport, fate, and effects of steroid hormones originating from CAFOs”

Background


- Estimated 1.3 million livestock operations in the U.S
- Roughly 20% confined generate ~500 million tons of waste annually
- Large facilities have limited land available for effective use of the generated waste as fertilizer
- Potential for natural and synthetic steroid hormones in livestock waste to reach groundwater and surface waters through many pathways



?



Cattle and Calves - Inventory: 2002



2000 Census
People 1.7 million head
Hogs 3.0 million head
Cattle 6.2 million head


1 Dot = 10,000 Cattle and Calves

United States Total
95,497,994




Steroid Hormones

- Steroids (terpinoid lipids) that act as hormones
- Hormone is a chemical messenger from one cell or group of cells to another (3 classes)
- Natural steroid hormones are generally synthesized from cholesterol




General
chemical
structure




Steroid Hormones in Cattle Waste

- Endogenous (natural)


- Estrodiol
- Estrone
- Estriol
- Testosterone
- Androstenedione
- Progesterone




17β-estradiol



estrone




Testosterone




Progesterone

- Exogenous (synthetic)


- Trenbolone
- Zearalonol
- Melengestrol Acetate



17α-trenbolone



Melengestrol acetate




α-zearalonol

Steroid hormones excreted in livestock manure

- Levels are highly variable
 - up to 1 $\mu\text{g/g}$
 - <100 ng/g
 - depends on compound, sex, diet, reproductive status
- Conjugated forms can be converted back to free hormones



Figure 2. Biliary 3-ketotrienic metabolites in the bile of a heifer



Chemical Properties

<i>Common Name</i>	<i>Aqueous solubility ($\mu\text{g/L}$)@ 25°C</i>	<i>Log P_{ow}</i>
<i>Estrogens</i>		
17 β -estradiol	12,960	4.01
Estrone	12,420	3.13
Estriol	13,250	2.45
<i>Androgens</i>		
Testosterone	5,570	3.23
Androstenedione	27,000	2.9
Androsterone	8,750	3.69
<i>Progestins</i>		
Progesterone	6,600	4.0
<i>Synthetic Steroids/progestins</i>		
Trenbolone	20,000	--
Zearalanol	--	3.9
Melengestrol Acetate	1,060	3.67
Ethinylestradiol	483	3.67

Atrazine solubility = 30,000 $\mu\text{g/L}$ Log Pow=2.3

Persistence




Figure 4. Degradation of (A) TbOH-17 α , (B) TbOH-17 β , and (C) TbO in liquid manure during storage.


“The Fate of Trenbolone Acetate and Melengestrol Acetate after Application as Growth Promoters in Cattle: Environmental Studies” *Bettina Schiffer, Andreas Daxenberger, Karsten Meyer, and Heinrich H.D. Meyer.* Environmental Health Perspectives • VOLUME 109 | NUMBER 11 | November 2001.

Project Objectives

- 1) *Quantify hormones in various stages of the manure pathway in cattle feedlots.*
- 2) *Determine the effects of different handling practices of cattle feedlot wastes on the stability and availability of hormones.*
- 3) *Determine the effects of different land application strategies on the fate and transport of hormones in vadose zone soils.*
- 4) *Determine if grasses grown in conservation buffers assimilate hormones.*

Environmental Fate and Transport of Steroid Hormones

145 day feeding period



Research Hypotheses

- 1) **Hormone levels** in manure are greater for cattle treated with hormones compared to untreated cattle.
- 2) **Composting manure** will facilitate the degradation of hormones in manure compared to stockpiling.
- 3) **Hormone losses** in runoff will be greater when manure is surface applied compared to when it is incorporated into the soil
- 4) **Hormone losses** in runoff will be related to the duration and timing of rainfall events with respect to land application of manure
- 5) **Grasses** commonly used in conservation buffer strips will assimilate hormones
- 6) **Hormones will persist** in soils and will leach through soils toward groundwater



Project Tasks

- Task 1: *Survey* existing feedlots and agricultural fields to determine the fate of hormones in the manure handling pathway over a climatic gradient.
- Task 2: *Quantify* the fate of hormones as influenced by manure handling practices such as stockpiling, composting, and runoff retention basins.
- Task 3: *Conduct* runoff studies with controlled rainfall simulators to determine the effect of manure application strategies on hormone transport.
- Task 4: *Quantify* the uptake of hormones applied in manures to selected grass species commonly used in grass buffer strips.
- Task 5: *Determine* the fate and transport of hormones in vadose zone soil after land application of manure.

Task 1. Survey of hormones in pens, run-off basins, stockpiles and fertilized cropland

- Where: Four University of Nebraska feedlots and manure-fertilized crop land
 - Collect manure from 3 pens and 2 stockpiles
 - Composite liquid from 2 run-off basins
 - Collect 12 composite samples at 3 depths from fields having received manure application within 1 month, 1 year, 2 year, and 3 years
- Collect data on feeding strategies, pen cleaning history, implant history, and manure handling




Figure 2. Locations of University of Nebraska Feedlots that will be used for Task 1.

Task 2. Measure differences in cattle waste as influence by handling practices

- Where: Haskell Ag Lab
- 96 heifers split between six pens
- 48 will receive implants (TBA, Z, E) and MGA supplement
- Composite pen surface samples and fresh manure collected at 0, 7, 45, and 125 days after implanting
- Waste removed after 150 days, stockpiled and composted
- Samples of stockpiled and composted manure will be collected to determine changes in hormone levels





Task 2. Measure differences in cattle waste as influence by handling practices

- **Two feedlot studies completed as of September 2008**
 - Feedlot surface samples collected at 7, 45 and 125 days after cattle placed in pens
 - ~30 runoff events sampled
 - Cattle treated with implants gained 3.23 lb/day, untreated cattle gained 2.65 lb/day
- **Waste handling study**
 - Manure from treated and untreated cattle placed in compost piles and anaerobic stockpiles
 - Piles stored and sampled over 6-8 month period



Aerobic versus anaerobic degradation

- Bioreactor tests started in August to help characterize transformation products under aerobic and anaerobic conditions
- Using ^{14}C -labelled testosterone and estradiol to track degradation rates and products



Task 3. Effect of tillage on hormones levels in field run-off

- Where: Haskell Ag Lab
- Five manure sources
 - treated compost and stockpile manure
 - untreated compost and stockpile manure
 - control with no manure applied
- Compare surface application (no-till), single disk incorporation, mold-board plow
- Simulated rainfall



Task 3. Effect of tillage on hormones levels in field run-off

- Run-off studies began in August 2008
 - 540 samples collected so far




Task 4. Uptake of hormones applied in manures to selected grass species commonly used in grass buffer strips

- Where: Haskell Ag Lab
- Hormone concentrations measured in buffer strip grasses receiving manure from steroid treated cattle
- Grass harvested after 30, 60, 90, 180, and 365 days after manure application
- Potential for exposure through grazing



Task 5. Fate and transport of hormones in vadose zone soil

- Where: WREC North Platte
- Lysimeter-instrumented test plots fertilized with stockpiled and composted manure from treated cattle in May
- Soil water will be collected from lysimeters over a 18-month period
- Determine potential for impacting ground water



Method Development: Extraction and analysis of steroid hormones

Solid Samples



Microwave-assisted solvent extraction

Liquid Samples




Automated solid-phase extraction



Analyze by liquid chromatography-tandem mass spectrometry

LC/MS/MS of Steroid Hormones




LC/MS/MS Detection Limits

Compound	Direct Injection	on-line SPE
	ppb	ppb
17 β -estradiol	1.07	0.0036
Estrone	0.77	0.0034
17 α -ethynyl estradiol	0.92	0.0056
Estriol	1.00	0.0064
α -Zearalonol	0.64	0.0032
Testosterone	1.46	0.0011
11-Ketotestosterone	0.58	0.0040
4-Androstenedione	1.71	0.0005
Progesterone	1.00	0.0008
Melengestrel Acetate	1.03	0.0016
17 β -trenbolone	0.93	0.0018
Androsterone	3.16	0.0038



LC/MS/MS of steroid hormones

- Steroids
 - difficult to ionize by electrospray (ESI)
 - matrix effects severe
- Atmospheric pressure photoionization (APPI)
 - more selective
 - improved ionization efficiency
 - better for steroid analysis



Preliminary On-Column Detection Limit (pg)

Compound	ESI	Recovery	APPI	Recovery
Testosterone	36.5	116 %	30.5	97 %
11-Ketotestosterone	14.5	91 %	13.8	103 %
4-Androstenedione	42.8	118 %	33.5	156 %
Progesterone	25.0	99 %	23.3	104 %
Melengestrol Acetate	25.8	96 %	20.5	125 %
17 β -Trenbolone	23.3	154 %	18.0	142 %

Preliminary Results

- Feed lot runoff samples
 - Few samples contain detectable levels of synthetic growth promoters
 - Concentrations of natural steroids low
 - Androsterone, estrone, 17β -estradiol, progesterone
 - Averages \ll 0.500 ppb
 - Many samples remaining to be analyzed



Contact Information

Daniel D. Snow
Director and Associate Professor
202 Water Sciences Laboratory
University of Nebraska – Lincoln
Lincoln, NE 68583-0844
dsnow1@unl.edu
402-472-7539

UNIVERSITY OF
Nebraska
Lincoln

Pioneering new frontiers.