

# Mangosteen

Robert E. Paull<sup>1</sup> and Saichol Ketsa<sup>2</sup>

<sup>1</sup>Department of Tropical Plant and Soil Sciences  
University of Hawaii at Manoa, Honolulu, HI

<sup>2</sup>Department of Horticulture, Kasetsart University, Bangkok, Thailand

**Scientific Name and Introduction:** Mangosteen (*Garcinia mangostana* L.) is one of the most praised of tropical fruit. It is also known as mangostanier, mangoustanier, mangouste, mangostier (French), mangostan (Spanish), manggis, mesetor, semetah, sementah (Malaysian), manggustan, mangis, mangostan (Philippine), mongkhut (Cambodian), mangkhut (Thai), cay mang cut (Vietnamese), manggis, manggistan (Dutch), and mangostao, mangosta, mangusta (Portuguese) (Nakasone and Paull, 1998). The globe-shaped, smooth berry is 4 to 7 cm (1.6 to 2.8 in) across, with a persistent calyx. The pericarp is 6 to 10 mm (0.24 to 0.4 in) thick and turns purple during ripening. It contains a bitter, yellowish latex and purple-staining juice. The edible, white aril has 4 to 8 segments with 1 or 2 larger segments containing apomictic seeds; there is no true seed.

**Quality Characteristics and Criteria:** Fruit have pinkish-red skin when mature, turning to a dark purple skin and white flesh. The skin should be thick and soft, but firm, when ripe. Fruit have a soft, sweet, slightly-acid flesh with a pleasant flavor. Misshapen and damaged fruit should be removed. Burst latex vessels leave a yellow dried latex on the fruit skin that should be scraped off and the fruit then washed with a soft brush. Fruit are graded to remove damaged fruit and for size. Some Thai growers/exporters coat cleaned fruit with lacquer, giving fruit skin damaged by thrips prior to harvest a more attractive appearance.

**Horticultural Maturity Indices:** Skin color is the major criteria used to judge maturity. Immature fruit that have a light, greenish-yellow skin with scattered pinkish spots do not ripen to full flavor if harvested. The minimum harvest stage for high quality fruit is when the skin has distinct irregular, pink-red spots over the whole fruit surface. Fruit are at the edible, ripe stage when the skin has darkened to a reddish-purple, no latex remains in the skin, and the flesh segments separate easily from the skin (Tongdee and Suwanagal, 1989). Careful handling is essential in order to avoid mechanical injury.

**Grades, Sizes and Packaging:** There are no U.S. or international standards. Fruit are graded by size and color. They are normally sold in single-layer, fiberboard cartons of 2.25 kg (5 lb) with padding, or sometimes in trays with fruit individually wrapped to prevent injury (20 to 24 fruit per tray). In Southeast Asia, fruit are sold in baskets or strung in bundles of from 10 to 25 fruit.

**Pre-cooling Conditions:** Room-cooling is normally used (Augustin and Azudin, 1986).

**Optimum Storage Conditions:** Recommendations vary from 3.9 to 5.6 °C (39 to 42 °F) with 85 to 90% RH for 7 weeks (Pantastico, 1975) to 13.3 °C (56 °F) with 85 to 90% RH for 14 to 25 days. Storage at 4 °C (39.2 °F) or 8 °C (46.4 °F), can lead to significant hardening of the skin (Augustin and Azudin, 1986), although the flesh may still be acceptable after 44 days. Current practice is to store fruit at 12 to 14 °C (54 to 57 °F) giving storage-life of about 20 days without chilling injury. Application of surface coatings reduces weight loss and prevents calyx wilting (Choehom, 1997).

**Controlled Atmospheres (CA) Considerations:** An atmosphere of 5% O<sub>2</sub> + 5% CO<sub>2</sub> has been used for 1 mo (Yahia, 1998) and resulted in best overall retention of peel appearance and internal quality (Rattanachinnakorn et al., 1996). Holding fruit in polyethylene film bags reduces weight loss and disease (Daryono and Sabari, 1986). However, it is not clear if the effects are due to the prevention of water loss

or to the modified atmospheres in the bags.

**Retail Outlet Display Considerations:** Display in over-wrapped trays or in closed, styrene, clam-shell containers with no perforations at 10 to 14 °C (50 to 57 °F). Do not mist.

**Chilling Sensitivity:** Storage at < 10 °C (50 °F) leads to rapid hardening and darkening of pericarp when fruit are returned to ambient temperature (Uthairatanakij and Ketsa, 1996; Choehom, 1997).

**Ethylene Production and Sensitivity:** Mangosteen is a climacteric fruit. Ethylene production is about 29 nL kg<sup>-1</sup> h<sup>-1</sup>. The respiratory peak occurs sooner when fruit are treated with ethylene. Ethylene treatment triggers autocatalytic ethylene production (Noichinda, 1992).

**Respiration Rate:** 21 mg (12 µL) CO<sub>2</sub> kg<sup>-1</sup> h<sup>-1</sup> at 25 °C. Heat production is 4,620 BTU ton<sup>-1</sup> day<sup>-1</sup> (1,281 kcal tonne<sup>-1</sup> day<sup>-1</sup>).

**Physiological Disorders:** Fruit damage during harvesting and marketing can affect > 20% of fruit. The “gamboges” disorder occurs where latex seeps into the flesh (aril) turning it yellow and giving it a bitter taste. The “gamboges” also moves onto the outer surface of the fruit. This is a pre-harvest disorder of unknown cause that makes it difficult to separate the aril from the surrounding tissue, even in ripe fruit; it causes hardening of the pericarp.

This should not be confused with impact injury that leads to hardening of the pericarp at the point of impact and aril collapse, dehydration, pink color development, or browning (Tongdee and Suwanagul, 1989). A drop of 10 cm can cause slight pericarp damage, indicated as hardening at the point of impact within 24 h. Higher drops causing significantly greater damage and often lead to downgrading of the fruit (Tongdee and Suwanagul, 1989; Ketsa and Atantee, 1998).

Another disorder of mangosteen fruit is translucent aril (*nue-kaew*) believed to be induced by heavy rain during fruit growth and development, even if just before harvest (Laywisakul, 1994). The specific gravity of fruit with translucent aril is > 1.0, while that of normal aril is < 1.0. This allows separation of fruit by floating them in water (Poddee, 1998). Fruit with translucent aril have lower SSC and TA than normal fruit (Pankasemsuk et al., 1996).

**Postharvest Pathology:** *Botryodiplodia theobromae*, *Diplodia* spp., *Pestalotia flagisetula*, *Phomopsis* spp., and *Rhizopus* spp. have been reported; they harden the skin and decay the aril.

**Quarantine Issues:** Mangosteen is a fruit fly host. Irradiation at 300 Grays has potential for disinfestation. Alternatively, harvested fruit are carefully cut open and the aril inspected; fruit are then frozen whole and shipped.

**Suitability as Fresh-cut Product:** Some potential.

**Special Considerations:** None.

## References:

- Augustin, M.A. and M.N. Azudin. 1986. Storage of mangosteen (*Garcinia mangostana*, L.). ASEAN Food J. 2:78-80.
- Choehom, R. 1997. Effect of waxing and plant growth regulators on quality and storage-life of mangosteen (*Garcinia mangostana* L.) fruit during cold storage. Graduate Special Problem, Department of Horticulture, Kasetsart University, Bangkok. 61 pp. (In Thai).
- Daryono, M. and S. Sabari. 1986. The practical method of harvest time on mangosteen fruit and its characteristics in storage. Bull. Penelitian Hort. Indonesia 14(2):38-44.

- Ketsa, S. and S. Atantee. 1998. Phenolics, lignin, peroxidase activity and increased firmness of damaged pericarp of mangosteen fruit after impact. *Postharv. Biol. Technol.* 14:117-124.
- Laywisakul, S. 1994. Factors influencing the development of translucent disorder in mangosteen. M.S. Thesis. Kasetsart University, Bangkok. 59 pp. (In Thai)
- Nakasone, H.Y. and R.E. Paull. 1998. *Tropical fruits*. CAB Intl., Wallingford, U.K. 445 pp.
- Noichinda, S. 1992. Effect of modified atmosphere on quality and storage-life of mangosteen (*Garcinia mangostana* L.) fruit. M.S. Thesis. Kasetsart Univ., Bangkok. 74 p. (In Thai).
- Pankasemsuk, T., J.O. Garner Jr., F.B. Matta and J.L. Silver. 1996. Translucent flesh disorder of mangosteen fruit (*Garcinia mangostana* L.). *HortScience* 31:112-113.
- Pantastico, E.B. 1975. *Postharvest physiology, handling and utilization of tropical and subtropical fruits and vegetables*. AVI Pub., Westport, Conn.
- Podee, R. 1998. Nondestructive methods for separation of translucent pulp in mangosteen by specific gravity and x-ray computed tomography. M.S. Thesis. Chiang Mai Univ., Chiang Mai, Thailand, 73 pp. (In Thai).
- Rattanachinnakorn, B., J. Phumhiran and S.S. Nanthachai. 1996. Controlled atmosphere storage of mangosteen. *Ann. Tech. Conf., Proc. Hort. Res. Inst., Dept. Agric., Bangkok, Thailand.* (In Thai).
- Tongdee, S.C. and A. Suwanagul. 1989. Postharvest mechanical damage in mangosteen. *ASEAN Food J.* 4:151-155.
- Uthairatanakij, A. and S. Ketsa. 1995. Physico-chemical changes of pericarp of mangosteen fruits after low temperature storage. In: S. Vijaysegaran, M. Pauziah, M.S. Mohamed and S. Ahmad Tarmizi (eds) *Proc. Intl. Conf. Trop. Fruits*. Vol. 1, Malaysian Agric. Res. Dev. Inst. (MARDI), Serdang, Salangor, pp. 411-422.
- Yahia, E.M. 1998. Modified and controlled atmospheres for tropical fruits. *Hort. Rev.* 22:123-183.