

# Orchid

Yin-Tung Wang

Department of Horticultural Sciences

Texas A&M University, Agricultural Experiment Station, Weslaco, TX

**Introduction:** For years, potted orchids were produced mainly for hobbyists. They are usually shipped in containers and arrive in buyers' hands by overnight delivery or in a few days. Many orchid growers do not ship orchids during Winter months to avoid chilling or freezing injury to the plants. However, during the past decade, there has been a tremendous worldwide booming of orchid production for sale as potted blooming plants for the public, particularly the *Phalaenopsis* and *Dendrobium* orchids. Potted, blooming orchids represent a \$100 million industry in the U.S. alone (USDA, 2001). However, for the most part, research in proper postharvest handling of orchids has lagged behind the increase in production and demand.

**Ethylene Sensitivity:** Orchid flowers are extremely sensitive to ethylene, even at very low levels (Davison, 1949; Beyer, 1976; Goh et al., 1985). Pollination and emasculation trigger ethylene production by flowers and result in rapid wilting. Blooming orchids should be handled carefully so that accidental pollination of flowers and removal of pollen caps (emasculation) do not occur. Accelerated flower wilting as a result of ethylene evolution following pollination in *Cymbidium*, *Dortitaenopsis*, *Dendrobium*, and *Phalaenopsis* orchids is induced by a loss of water from cells of the upper layer of the petals, leading to their in-folding and water-soaked appearance (Lee and Lin, 1992; Porat et. al., 1994). Treating orchid flowers with the ethylene biosynthesis inhibitor aminooxyacetic acid (AOA) did not alter their sensitivity to ethylene. Recent research also showed that pulsing excised *Dendrobium* flowers for 24 h with 0.5 mM AOA before exposing them to 0.2  $\mu\text{L L}^{-1}$  ethylene did not prolong their life (Wang and Wang, 2000); all treated flowers wilted within 3 days after exposure. Thus, AOA has no effect on longevity when wilting is triggered by exogenous ethylene.

**Treatments for Extending Shelf-life:** It was found that shelf-life of *Oncidium* Goldiana cut sprays increased after pulsing with silver nitrate for 30 min (Ong and Lim, 1983). Pulsing *Arranda* cut inflorescences for a short 10 min with a 4 mM solution of STS prolonged their display-life (Hew et al., 1987). Exposing excised *Dendrobium* and *Phalaenopsis* flowers to 0.1  $\mu\text{L L}^{-1}$  1-methylcyclopropene (1-MCP, an ethylene action inhibitor) for 12 h, or pulsing them with 0.5 mM silver thiosulfate (STS; ethylene inhibitor) for 24 h completely blocked the deleterious effect of ethylene (Wang and Wang, 2000). *Phalaenopsis* flowers that were treated with 0.1 to 0.4  $\mu\text{L L}^{-1}$  of 1-MCP are protected against ethylene as high as 10  $\mu\text{L L}^{-1}$ . However, the protective effect of 1-MCP lasts for no longer than 7 days at room temperature. Commercially, cut *Phalaenopsis* inflorescences are pulsed with 0.5 mM STS immediately following cutting.

When used at 50 to 100  $\text{mg L}^{-1}$ , 8-hydroxyquinoline sulfate (8-HQS) extended shelf-life of *Dendrobium* Pompadour cut flowers (Kesta and Amutiratana, 1986). Conflicting results were reported on the effect of sucrose in keeping solutions as a carbon source for extending vase-life of cut orchid flowers. Sprays of tropical orchids should be stored/transported at 12 to 18 °C (54 to 64 °F) (Akamine, 1976), whereas flowers of temperate orchids, ex., *Cymbidium*, may be stored at temperatures as low as 5 °C (41 °F) (Sheehan, 1954).

**Optimum Storage Conditions:** Minimal research has been conducted on optimum orchid storage conditions. In one study (Wang, 1997a), bare-root *Phalaenopsis* plants were packed in boxes and subjected to temperatures of 15, 20, 25, or 30 °C (59, 68, 77, or 86 °F) for 4, 7 or 14 days. Weight loss from plants increased with increasing temperature and storage duration. Symptoms of chilling injury were inversely

related with 15 and 20 °C storage temperatures. Chilling injury became more severe as storage duration increased. Plants had little or no chilling injury at 25 or 30 °C, regardless of the duration. Leaf loss was most severe on plants stored at 15 °C for 7 or 14 days or 30 °C for 14 days. Plants that had severe leaf loss after storing at 30 °C had delayed spiking and flowering.

Pre-conditioning greenhouse-grown potted *Phalaenopsis* plants at 25 °C (77 °F) for 10 days, followed by another 10 days at 20 °C (68 °F) reduced severity of chilling injury after storage at 15 °C (59 °F) for 2 weeks. Therefore, *Phalaenopsis* orchids harvested during the warm period of the year should be held above 20 °C (68 °F), and preferably closer to 25 °C (77 °F), during shipping. Under such conditions, plants can lose as much as 20% of their fresh weight at harvest without adversely affecting subsequent performance. While most *Phalaenopsis* species and hybrids need relatively cool air to initiate the flowering process, plants do not respond to cool air while being shipped in the absence of light (Wang, 1995).

Although prolonged duration in complete darkness is not recommended for keeping plants in bloom (Wang and Blessington, 1989), *Phalaenopsis* orchids tolerate low light levels once they have started flowering. One study show that exposing them to light levels between 10 and 50  $\mu\text{mol m}^{-2} \text{sec}^{-1}$  for 12 h daily produced orchids with similar longevity (individual flowers lasting over 120 days) and flower size (Wang, 1997b). Therefore, most retail stores have adequate light levels to maintain blooming *Phalaenopsis* while they are being marketed.

**Shipping Information:** Due to quarantine regulations, when shipped internationally, orchids are often removed of their potting medium. The bare-root plants are allowed to dehydrate slightly before packing. This is done for several reasons. Slight dehydration allows for packing more plants per box, avoids breaking leaves, and reduces the chances of moisture buildup to weaken shipping boxes. Drier plants also reduce incidence of rotting before boxes are opened. Plants are usually packed in layers alternating with shredded paper or other material. Often, plants are individually wrapped in newspaper, particularly those that have developing inflorescences.

Millions of bare-root *Phalaenopsis* orchids are shipped domestically and internationally each year. After planting in pots and placing on greenhouses benches, occasionally, yellow spots develop on either side of the mid-rib near the base of leaves due to chilling injury. Irregular yellow blotches also occur on other parts of leaf blades. Other times, leaves turn yellow and fall off, resulting in weak plants that do not produce flowers.

### Literature Cited

- Akamine, E.K. 1976. Postharvest handling of tropical ornamental cut crops in Hawaii. HortScience 11:125-126.
- Beyer, E.M. 1976. A potent inhibitor of ethylene action in plants. Plant Physiol. 58:268-271.
- Davison, O.W. 1949. Effect of ethylene on orchid flowers. Proc. Amer Soc Hort Sci. 53:440-446.
- Hew, C.S., K.H. Wee, and F.Y. Lee. 1987. Factors affecting the longevity of cut *Aranda* flowers. Acta Hort. 205:195-202.
- Goh, C.J., A.H. Halevy, R. Engel and A.M. Kofranek. 1985. Ethylene evolution and sensitivity in cut orchid flowers. Sci. Hort. 26:57-67.
- Kesta, S. and D. Amutriratana. 1986. Effect of sucrose, silver nitrate, and 8-hydroxyquinoline sulfate on postharvest behavior of *Dendrobium* Pompadour flowers. In: Proc. 6th ASEAN Orchid Congr., Bangkok, Thailand, pp. 124-129.
- Lee, N. and Y.S. Lin. 1992. Respiration of *Phalaenopsis* flowers. J. Chinese Soc. Hort. Sci. 38:228-240.
- Ong, H.T. and L.L. Lim. 1983. Use of silver nitrate and citric acid to improve shelf-life of *Oncidium* Golden Showers flowers. Orchid Rev. 90:264-266.
- Porat, P., A. Borochoy, .H. Halevy, and S.D. O'Neill. 1994. Pollination-induced senescence of *Phalaenopsis* petals. Plant Growth Regul. 15:129-136.
- Sheehan, T.J. 1954. Orchid flower storage. Amer. Orchid Soc. Bul. 23:579-584. USDA. 2001. Floriculture Crops, 2000 Summary. Natl. Agric. Stat. Serv., Wash DC.

- Wang, Y.T. 1995. Phalaenopsis orchid light requirement during the induction of spiking. HortScience 30:59-61.
- Wang, Y.T. 1997a. Effect of postharvest temperature and storage duration on growth and flowering of the *Phalaenopsis* orchids. HortScience 32:517.
- Wang, Y.T. 1997b. Phalaenopsis light requirements and scheduling of flowering. Orchids 66:934-939.
- Wang, Y.T. and T.M. Blessington. 1989. Dark-storage temperature and duration influences flowering and quality retention of hibiscus. HortScience 24:646-647.
- Wang, M.K. and Y.T. Wang. 2000. 1-MCP and STS counteract ethylene effects on cut orchid flowers. HortScience 35(3):349.