



United States
Department of
Agriculture

Marketing and
Regulatory
Programs

Animal and
Plant Health
Inspection
Service



Mexican Fruit Fly Cooperative Eradication Program

Monterey Park, California

Environmental Assessment October 2002

Mexican Fruit Fly Cooperative Eradication Program

Monterey Park, California

Environmental Assessment
October 2002

Agency Contact:

Stephen A. Knight
Senior Staff Officer
Invasive Species and Pest Management
Plant Protection and Quarantine
Animal and Plant Health Inspection Service
U.S. Department of Agriculture
4700 River Road, Unit 134
Riverdale, MD 20737-1236
Telephone: 301-734-8039

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, or marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director of Civil Rights, Room 326-W, Whitten Building, 1400 Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD).
USDA is an equal opportunity provided and employer.

Mention of companies or commercial products in this report does not imply recommendation or endorsement by the U.S. Department of Agriculture over others not mentioned. USDA neither guarantees nor warrants the standard of any product mentioned. Product names are mentioned solely to report factually on available data and to provide specific information.

This publication reports research involving pesticides. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.

Table of Contents

I.	Need for the Proposal	1
II.	Alternatives.....	2
	A. No Action	2
	B. Nonchemical Control	2
	C. Integrated Control (Preferred Alternative) ...	3
III.	Potential Environmental Consequences	4
	A. No Action.....	4
	B. Nonchemical Control.....	5
	C. Integrated Control (Preferred Alternative)	6
IV.	Agencies, Organizations, and Individuals Consulted.....	22
	Appendix A. References Cited	A-1

Tables

1.	Acute Oral LD ₅₀ s for Selected Species Dosed with Malathion (mg/kg)	8
2.	Malathion 96-hour LC ₅₀ s for Selected Aquatic Species (µg/L).....	8
3.	Acute Oral LD ₅₀ s for Selected Species Dosed with Diazinon (mg/kg)	12

I. Need for the Proposal

The Mexican fruit fly, *Anastrepha ludens* (Loew), is native to central Mexico and is a major pest of agriculture throughout many parts of the world. Commercial and home grown produce that is attacked by the pest is unfit to eat because the larvae tunnel through the fleshy part of the fruit, damaging the fruit and subjecting it to decay from bacteria and fungi. Because of its wide host range (over 40 species of fruits) and its potential for damage, a permanent infestation of Mexican fruit fly would be disastrous to agricultural production in the United States. In the past, eradication programs have been implemented successfully to prevent the pest from becoming permanently established on the U.S. mainland.

On October 9 and 12, 2002, an unmated wild male and an unmated wild female Mexican fruit fly were trapped in the Monterey Park area of Los Angeles County, California. Since those detections, two immature females were trapped on October 23 and a third instar larva was found in a sapote fruit on some adjacent property. The infestation is presently found in a residential area of Los Angeles County, although that situation may change in the future. This Mexican fruit fly infestation detected in southern California represents a major threat to the agriculture and environment of California and other U.S. mainland States. The U.S. Department of Agriculture's (USDA) Animal and Plant Health Inspection Service (APHIS) and the California Department of Food and Agriculture (CDFA) are proposing a cooperative program to eradicate the infestation and eliminate that threat.

APHIS' authority for cooperation in the program is based upon the Organic Act (7 United States Code (U.S.C.) 147a), which authorizes the Secretary of Agriculture to carry out operations to eradicate insect pests, and the Plant Protection Act (Title 4 of the Agricultural Risk Protection Act of 2000), which authorizes the Secretary of Agriculture to use emergency measures to prevent the dissemination of plant pests new to or not widely distributed throughout the United States.

This site-specific environmental assessment (EA) analyzes the environmental consequences of alternatives which have been considered for Mexican fruit fly control and considers, from a site-specific perspective, environmental issues that are relevant to this particular program. The control measures being considered for this program have been discussed and analyzed comprehensively within the fruit fly chemical risk assessments (USDA,

APHIS, 1998a, b). Those documents are incorporated by reference and summarized within this environmental assessment.

II. Alternatives

Alternatives considered for this proposed program include (1) no action, (2) nonchemical control, and (3) integrated control. APHIS' preferred alternative for the program is Integrated control (using chemicals) to facilitate timely eradication of the current Mexican fruit fly infestation.

A. No Action

Under this alternative, APHIS would not participate in efforts to eradicate the current infestation of the Mexican fruit fly in Los Angeles County, California. An eradication could proceed under the direction of the State and/or County government, but the lack of Federal/State coordination would likely jeopardize timely and efficient implementation of this program. This could result in delays in achieving eradication, expansion of the infested area, and permanent establishment of the Mexican fruit fly. Potential adverse environmental effects of this alternative would be at least as severe as those under the proposed integrated program alternative, and would be more severe if the infestation expanded substantially or could not be eradicated. Establishment of Mexican fruit flies would lead to increased damage to crops and backyard produce, uncoordinated use of insecticides by commercial and backyard growers, and increased environmental risk from the insecticide applications. Such adverse effects would be of an indirect, but continuing and escalating nature.

B. Nonchemical Control

Under this alternative, APHIS would participate in a cooperative program to eradicate the existing infestation of Mexican fruit fly in Los Angeles County, California, with solely nonchemical methods. Examples of such methods include: sterile insect technique, physical control, cultural control, male annihilation, and regulatory control. Biological control and biotechnological control are other nonchemical methods that were considered, but have not yet been proven efficacious or technologically feasible. Federal/State approval of such a nonchemical program is unlikely because nonchemical technologies cannot respond quickly enough to the infestation to contain and eliminate it before it has had the opportunity to spread. Nonchemical

methods such as sterile insect technique have greater effectiveness when used as components of integrated programs, or in preventive programs designed to eliminate pest introductions before they become infestations. Cold treatments, vapor heat treatments, and irradiation treatments are regulatory treatments that are applicable to some commodities. The potential adverse environmental impacts of a nonchemical program would be expected to be as severe as under the no action alternative, because of the anticipated inability of such a program to quickly and effectively eradicate the infestation. The infestation would grow, resulting in increased damage to crops and backyard produce, uncoordinated use of insecticides by commercial and backyard growers, and increased environmental risk from insecticide applications. Such adverse impacts would be of an indirect, but continuing and escalating nature.

C. Integrated Control (Preferred Alternative)

The proposed integrated program would use any of a combination of control methods, based upon site-specific requirements that take into account program efficacy and environmental considerations. As a form of integrated pest management, integrated control may include the use of both chemical and nonchemical methods in a timely manner to achieve the program goal of eradication and minimize potential environmental consequences that could arise from program activities. This is the preferred alternative, from both program and environmental perspectives.

Specifically, this integrated program could use any or a combination of the following methods: chemical control, sterile insect technique, physical control, cultural control, male annihilation, and regulatory control. Biological control and biotechnological control also were considered, but have not yet been proven to be efficacious or technologically feasible for this species of pest. The eradication program is likely to consist of three ground applications of malathion bait, applied at 14-day intervals, prior to the release of sterile Mexican fruit flies. Diazinon will be drenched with water into the soil within the drip line of plants known or suspected to contain Mexican fruit fly larvae. Other control options include the use of mass trapping, host removal, and regulatory control. Regulatory control involves quarantine of fresh produce and commodities from host plants of Mexican fruit fly. Specific regulatory treatments are required for transport of produce grown within the designated quarantine area to destinations outside this regulated area. The treatment of produce and nursery stock may involve malathion

bait spray applications, diazinon soil treatments, or methyl bromide fumigations.

There are potential adverse environmental impacts from the use of chemicals in the integrated program, but those impacts are fewer and less severe than in the other alternatives. In general, the integrated program would have direct adverse impacts of a non-continuing nature.

III. Potential Environmental Consequences

The analysis of potential environmental consequences will consider the alternatives of no action, nonchemical control, and integrated control. Because the principal environmental concern over an integrated control program relates to its use of chemical pesticides, this assessment will focus on the potential environmental consequences of the pesticides on human health, nontarget species, and endangered and threatened species.

A. No Action

Under the no action (no APHIS effort) alternative, Mexican fruit fly control would be left to the State, grower groups, or individuals. Without a coordinated efforts between APHIS and cooperators, it is likely that the infestation would spread to other areas of California and the U.S. mainland. Any response to control such an expanded infestation by individuals or organizations would probably result in a greater magnitude of environmental impact than would be associated with a coordinated APHIS/State eradication program. Under those conditions, any available controls (including more hazardous chemical pesticides) could be used, resulting in greater environmental impact than is associated with the action alternatives analyzed within this assessment.

1. Human Health

Under the no action alternative, private homeowners and commercial growers would have few options other than pesticides to reduce the Mexican fruit fly damage to their crops. Any pesticides registered for use could be applied in an unsupervised and uncoordinated manner. Accordingly, greater pesticide amounts and higher frequencies of application are likely to be used than would be expected with a coordinated, cooperative government program. In addition to the direct toxic effects of those pesticides, humans

could also be affected by cumulative impacts resulting from synergistic effects of combining various pesticides for use against Mexican fruit fly. Human exposure to pesticides and resulting adverse consequences probably would be greater than if pesticides were applied in a cooperative government program. The spread of the infestation will reduce the amount of locally available produce and may restrict the fruit consumption of some members of the public. Some members of the public may depend upon this source of fruit as a substantial portion of their diet.

2. Nontarget Species

Broader pesticide use resulting from lack of APHIS effort to combat Mexican fruit fly would increase the pesticide load to the environment and, therefore, increase the probability of effects to nontarget species. The potential expansion and establishment of the pest also would have unknown effects on insect community structure and on predators in those systems.

3. Endangered and Threatened Species

Section 7 of the Endangered Species Act of 1973 (ESA) requires Federal agencies to consult with the U.S. Department of the Interior, Fish and Wildlife Service (FWS), if species listed or proposed for listing are likely to be adversely affected. The CDFG advised that the California Natural Diversity Database indicates that Least Bell's vireo and southwestern pond turtle reside within the current eradication zone. The no action alternative involves no activity by APHIS that would affect these species or the riparian, marsh, or sand habitats of these species. Further expansion of the Mexican fruit fly's range would be likely to include endangered and threatened species habitats, with unquantified risk to those species from uncoordinated pesticide use under the no action alternative. No adverse impacts to endangered or threatened species would result directly from APHIS' implementation of the no action alternative.

B. Nonchemical Control

The nonchemical control methods proposed for use under this alternative include sterile insect technique, physical control, cultural control, cold treatment, irradiation treatment, and vapor heat treatment. Although biological control and biotechnological control are being researched for development, these methods have not yet been proven efficacious or technologically feasible, so their potential environmental consequences are not analyzed here.

1. Human Health

Under the nonchemical control alternative, human health is not expected to be adversely affected. The sterile insect technique, physical control, and

cultural control do not pose a risk to human health. The control program includes some regulatory treatments (cold treatments, irradiation, and vapor heat treatments) that occur in restricted access facilities and that are strictly supervised to ensure no effects occur to human health. In general, the use of nonchemical methods reduces substantially the need for chemical applications, thereby decreasing the magnitude of impact from chemical usage. The nonchemical control alternative may not be successful for larger infestations and the human health consequences of inadequate control could be comparable to the no action alternative if the infestation of Mexican fruit fly expanded due to insufficient containment of the pest.

2. Nontarget Species

The nonchemical techniques that may be employed could disturb nontarget species, due to noise or mere human presence. In general, little risk is associated with these disturbances. Use of a sterile insect technique could have a positive effect, that of providing a food source to some insectivores. Nonchemical methods have the potential for less pesticide use than the other alternatives, but control and containment of Mexican fruit fly under this alternative depend upon low pest populations. If nonchemical methods were insufficient to eradicate the pest population, the ultimate expansion of the infestation could result in pesticide usage comparable to that of no action.

3. Endangered and Threatened Species

Nonchemical methods should not directly impact endangered or threatened species. The FWS has determined that the sterile insect technique is compatible with endangered or threatened species. The nonchemical regulatory treatments are not made to these species or their critical habitats.

Cultural and physical control methods can affect some species through habitat disturbance, but consultation with the FWS about the use of these methods within program areas is made prior to program action to ensure that no program actions will affect the endangered or threatened species or their critical habitat. None of the nonchemical control methods are known to affect endangered or threatened species within the program area.

C. Integrated Control (Preferred Alternative)

The environmental consequences of nonchemical methods were discussed under the nonchemical control alternative and this information will not be repeated in this section. The components of the proposed program which potentially have the greatest impact on the environment are the chemical pesticides. Special registration procedures are required for pesticides used against exotic pests, such as the Mexican fruit fly which is not native to this

country. A section 18 (emergency) or section 24c (special local needs) exemption under the Federal Insecticide, Fungicide, and Rodenticide Act allows their use. The environmental consequences from the use of these pesticides (malathion, diazinon, and methyl bromide) are discussed below. Because of the limited and restricted nature of these chemical control actions necessary in this integrated control, it has been analyzed within the framework of an environmental assessment.

Three major factors influence the risk associated with pesticide use: fate of the pesticide in the environment, its toxicity to humans and nontarget species, and the exposure of humans and nontarget species to the pesticide. These factors will be evaluated for each of the chemicals analyzed.

1. Malathion Bait Spray

a. Fate

Malathion is an amber-colored liquid that is combined with a protein bait to form a sticky spray. The formulation used in the program is 0.175 pounds of active ingredient per acre mixed with 9.6 fluid ounces of protein hydrolysate bait per acre, for both aerial and ground applications. The half-life of malathion in soil or on foliage ranges from 1 to 6 days; in water, from 6 to 18 days. Malathion bait spray is applied from the ground, generally as a spot treatment to individual trees, or from the air. Trees, shrubs, and other surfaces such as soil, roads, and ponds are likely to receive spray from aerial applications, although efforts, including the use of buffers, are made to avoid directly spraying water bodies. Malathion is generally of more concern in aquatic areas because of its high toxicity to aquatic organisms.

b. Toxicity

Malathion is an organophosphate that acts by inhibiting acetylcholinesterase. Mildly acutely toxic, malathion is classified by EPA as category III (Caution) based on oral, dermal, and inhalation exposure routes. At high doses, toxic effects from malathion may include headache, nausea, vomiting, blurred vision, weakness, and muscular twitching. In humans and other mammals, metabolism by one degradation pathway leads to the formation of malaoxon, a more potent cholinesterase inhibitor than malathion. The more common degradation pathways yield nontoxic intermediates.

The U.S. Environmental Protection Agency (EPA) has recently evaluated the carcinogenic potential of malathion. Their new classification describes malathion as having “suggestive evidence of carcinogenicity, but not sufficient

to assess human carcinogenic potential.” This indicates that any carcinogenic potential of malathion is so low that it cannot be quantified based upon the weight of evidence. The low exposures to malathion from program applications would not be expected to pose any carcinogenic risks to workers or the general public. Malathion may have synergistic effects when used with other organophosphate or carbamate pesticides.

Oral doses of malathion are slightly to moderately acutely toxic to mammals and birds (table 1). Signs of poisoning are similar to the reactions of humans. Malathion is highly toxic to some forms of aquatic life, including invertebrates, amphibians, and fish (table 2). The EPA has established a chronic water quality criteria of 0.1 µg/L (micrograms per liter) for protection of freshwater and marine aquatic life. Fish kills that may have been associated with aerial malathion bait spray applications have been documented.

Table 1. Acute Oral LD₅₀s¹ for Selected Species Dosed with Malathion (mg/kg)

Mouse	720 - 4,060
Female rat	1,000
Male rat	1,375
Mallard	1,485
Pheasant	167

¹LD₅₀ = Lethal dose for 50% of animals treated

Table 2. Malathion 96-hour LC₅₀s¹ for Selected Aquatic Species (µg/L)

Tadpole	200
Rainbow trout	4.1 - 200
Bluegill	20 - 110
Daphnia	1 - 1.8
Stone flies	1.1 - 8.8

¹LC₅₀ = Lethal concentration for 50% of animals treated

c. Exposure and Risk

(1) Human Health

Potential exposure to humans is by dermal absorption, inhalation, or ingestion of residues. Due to the potential for aerial application of malathion bait spray, dermal absorption from direct application or contact with treated surfaces is the primary exposure route for the public. Public exposure from a ground malathion bait spray application will be lower than exposure from an aerial application because less area is treated and less pesticide is used. Workers, such as ground applicators and the ground crew for aerial applications, may have inhalation exposure as well as dermal exposure.

Results of the quantitative risk assessment of malathion bait suggest that exposures to pesticides from comparable program operations are not likely to result in substantial adverse human health effects. Residues on commodities or backyard fruits resulting from the malathion bait spray application are unlikely to greatly increase exposure to the consuming public. Malathion concentrations on vegetation estimated by the California Department of Health Services (Kizer, 1991) indicate that levels of malathion on vegetation are not likely to exceed the residue tolerance levels set by EPA. Residue tolerances for malathion on many food items are established (40 CFR 180.11) and most are 8 ppm (parts per million). The provisional acceptable daily intake is 0.02 mg/kg per day.

The human health risks of comparable treatments are evaluated quantitatively in the Human Health Risk Assessment for Fruit Fly Cooperative Control Programs (USDA, APHIS, 1998a). Results suggest that exposure from normal program operations will not present a human health risk either to workers or the public. In addition, risks to humans have been analyzed qualitatively, with reliance on information from past fruit fly eradication programs in California. The exposure scenarios from previous fruit fly eradication efforts will not differ substantially from the current program.

(2) Nontarget Species

Malathion bait spray will kill insects other than the Mexican fruit fly. Malathion is highly toxic to bees, and direct application to areas of blooming plants can be expected to result in a high bee kill. Although malathion is not phytotoxic, there could be potential indirect effects on plant populations due to lower pollination rates if bee or other pollinator populations are reduced.

This is a concern of aerial application. Secondary pest outbreaks have occurred concurrently with the use of aerial applications of malathion bait spray, but have not been determined conclusively to be associated with the applications. In 1981, fish kills also occurred from a similar treatment method. Since then, the State of California has instituted procedures to reduce the likelihood of fish kills. None have been known to occur from aerial applications of malathion bait spray since the procedures were implemented.

Terrestrial animals are exposed to malathion primarily through dermal and oral routes. Ingesting prey containing residues, rubbing against treated vegetation, and grooming contribute to total dose. Aquatic species can be exposed to direct application and runoff. Exposure of malathion bait spray by aerial application poses high risk to nontarget invertebrates and some aquatic species. Some insectivores may be affected. Ground application of malathion bait spray has far fewer environmental consequences because the treated area is smaller and delivery is more accurate. Fewer species would be exposed and thus the treatment poses less total risk to nontarget species than does aerial application.

(3) Endangered and Threatened Species

It was determined that Least Bell's vireo and southwestern pond turtle reside within the current program eradication zone. Several other endangered or threatened species are found in nearby areas of Los Angeles County and adjacent counties. If the program were to expand and if the range of Federally listed species and the treatment area overlapped, protective measures may be required to protect species from adverse environmental consequences of the program. There are no plans to treat any riparian, marsh, or sand dune habitats or areas not adjacent to paved roads. The proposed program methods were determined to pose no effect to the endangered and threatened species found in the area based upon previous programmatic consultation. The species that may be affected by control efforts are dependent upon the control methods used (i.e., not all control methods affect all species equally). Thus, protective measures will vary depending on the control method being used and the species found within the treatment area.

Malathion bait spray is not selective for Mexican fruit fly alone. Ingestion of bait/malathion and cuticular exposure to malathion by insects other than Mexican fruit fly could result in their deaths. If their habitats overlapped with

the program treatments, those species could be adversely affected by aerial application of malathion bait. Repeated aerial sprays of malathion bait generally would reduce insect numbers. Reduction of insect populations could reduce pollinator species for threatened and endangered plants, and would reduce potential food resources for endangered and threatened insectivores. Malathion is highly toxic to many aquatic species, both vertebrate and invertebrate, and spray drift could result in aquatic system disruption. The ecosystem is resilient enough to absorb some reduction in nontarget populations and the resultant food web effects, but the severity of the reductions would increase with increased applications of malathion.

Many of the endangered and threatened species are dependent upon aquatic habitats. Loss of a single individual of a listed species from program activities would be a violation of the ESA. Thus, aerial application of malathion bait spray should be controlled both within the range of endangered and threatened insect-pollinated plants (especially annuals) and in aquatic habitats.

2. Diazinon Soil Treatments

a. Fate

Technical grade diazinon is a sweet, aromatic, amber-brown liquid. The program formulation is applied at a rate of 5 pounds active ingredient per acre. Its half-life in soil ranges from 1.5 to 10 weeks and in water at neutral pH ranges from 8 to 9 days. Small amounts of diazinon are used to treat soil within the drip line of trees that have fruit infested with Mexican fruit fly larvae. Surface vegetation may retain residues and, depending on soil type, local hydrology, and topography, diazinon may occur in runoff water.

b. Toxicity

Although diazinon is widely used and generally is not considered a hazard to human health under its registered uses, it can be toxic to humans. The EPA has classified the formulation of diazinon as category II (Warning) for program use in soil treatment. Although not a primary dermal or eye irritant, it can be absorbed through these routes and, at high concentrations or prolonged exposure, causes severe irritation.

The mode of toxic action of diazinon occurs through inhibition of the enzyme, acetylcholinesterase. Symptoms of poisoning in humans, who are much less susceptible to the effects of diazinon than insects, include blurred vision, nausea, vomiting, slurred speech, and mental confusion. Death, which can

occur from high doses, results from respiratory arrest caused by muscle paralysis and bronchoconstriction. Accidental oral poisonings have resulted in death from doses between 50 and 500 mg/kg.

Diazinon has many metabolites, but toxicity data on most are not currently available. Although the metabolite diazoxon is more toxic than diazinon, it is also more easily metabolized and excreted. Diazinon may exhibit synergistic effects with other commercial pesticide formulations currently in use. Diazinon is not considered to be a carcinogen and is nonmutagenic.

Animals differ in their sensitivity to diazinon, both within and between species. Toxicity varies widely and depends on sex and life stage (table 3). Diazinon is toxic to vertebrate laboratory animals and very toxic to livestock. Diazinon is extremely toxic to birds, which are sensitive because their blood has no enzymes to hydrolyze diazoxon (a toxic metabolite), as does mammalian blood (Eisler, 1986). Signs of intoxication include salivation, stiff-legged gaits, wing spasms, and wing-beat convulsions (Hudson *et al.*, 1984). Many incidents of avian (particularly geese and other waterfowl) mortality on golf courses have occurred because of the use of granular formulations of diazinon. These incidents led the EPA to cancel use of diazinon on golf courses and sod farms in 1986. Some terrestrial invertebrates (such as bees) are extremely sensitive to diazinon. Diazinon causes high earthworm mortality but does not have a similar effect on nematodes.

Freshwater cladocerans (water fleas, common to aquatic areas) are among the aquatic species most sensitive to diazinon; *Gammarus fasciatus* has a 96-hour LC₅₀ of 0.20 grams per liter. There is some evidence that juvenile fish are more sensitive than eggs. Sublethal effects include reduced growth and reproduction in both marine and freshwater invertebrates, including reduced emergence of insects (Eisler, 1986). Algae are unaffected by concentrations fatal to aquatic invertebrates.

Table 3. Acute Oral LD₅₀s¹ for Selected Species Dosed with Diazinon (mg/kg)

Rabbit	130
Mouse	80 -135
Female rat	76 - 250
Male rat	108 - 285
Guinea pig	280

Calf	0.5
Starling	110
Mallard (3 to 4 months old)	3.5
Pheasant (3 to 4 months old)	4.3
Bobwhite quail	3.4 - 10
Chicken (5 days old)	8.4
Redwinged blackbird	2.0
Butterfly	8.8
Honey bee	0.372/bee

¹LD₅₀ = Lethal dose for 50% of animals treated

c. Exposure and Risk

(1) Humans

Potential exposure to humans is by ingestion or dermal absorption. The soil drenching application (rate of 52 mg per square foot of treated area) techniques prevent inhalation exposure. Because the diazinon is watered into the soil and the drenched area is small, public exposure will be limited. Program use of the pesticide precludes exposure to residues from produce on host plants because any fruit will be stripped from the plants before treatment. Occupational exposure will be reduced by wearing gloves when handling or applying diazinon. The only human health risk associated with diazinon is the consumption of soil from the drenched area by toddlers. The public will be notified when a drench has occurred and will be advised of the necessary precautions.

(2) Nontarget Species

Diazinon exposure to nontarget organisms is restricted to those organisms that traverse or visit the treated area as well as relatively immobile species that inhabit the area directly treated. The treatments are limited (generally less than 10 gallons per year) and occur only within the drip line of host trees. However, due to diazinon's high toxicity, organisms that are directly exposed are at high risk. Limiting exposure will reduce this risk.

(3) Endangered and Threatened Species

Because birds are highly mobile and are among the most sensitive vertebrates to diazinon, endangered and threatened avian species are of special concern. Least Bell's vireo is known to reside within the current eradication zone.

However, those areas being treated are not habitat to this species. If the program were to expand, the limited nature of the soil treatments and implementation of appropriate protective measures would combine to protect any Federally listed endangered and threatened bird species.

Diazinon is used only to treat soil under hosts that are infested with Mexican fruit fly larvae. Very little is used in a program (usually less than 10 gallons annually, for a combined area of under 2 acres). Therefore, it is unlikely that endangered and threatened birds would even encounter any treatments.

Endangered and threatened birds may be protected from exposure to diazinon by the presence of program personnel, who remain in the area until the pesticide has soaked into the soil. Program monitoring may include carcass searches to ensure that no endangered and threatened species are affected by the program. If there is any confirmation that the program has adversely affected an endangered and threatened species, immediate action would be taken to determine an appropriate program response that would be required to protect those species.

It is anticipated that swift initiation of eradication activities upon detection of a Mexican fruit fly infestation will minimize the area requiring treatment and make it unlikely that treatments will occur where endangered and threatened species are present. Recent Mexican fruit fly infestations have occurred in urban and suburban areas where natural areas are small and endangered and threatened species are few or absent. Additionally, the incorporation of protective measures should further protect endangered and threatened species from potential adverse effects attributable to program eradication activities.

3. Methyl Bromide Fumigation

a. Fate

Methyl bromide is an odorless, colorless, volatile gas which is three times as heavy as air. Its half-life is 3 to 7 days. Methyl bromide is released when a fumigation chamber is aerated. Because methyl bromide is heavier than air, the gas can collect in isolated pockets, which could create hazardous conditions when there is little air circulation or mixing, such as during thermal inversions or periods of low wind.

b. Toxicity

Methyl bromide gas and liquid are acutely toxic to humans. Contact with liquid or vapors can cause serious skin or eye injury. Inhalation can cause acute illness, including pulmonary edema (fluid buildup in the lungs), gastrointestinal distress, and convulsions which can be fatal. The LD₅₀ (lethal dose for 50% of animals treated) of rats to methyl bromide is 2,700 ppm for a 30-minute exposure. In humans, 1,583 ppm (6.2 mg/L (milligrams per liter)) methyl bromide is lethal after 10 to 20 hours of exposure and 7,890 ppm (30.9 mg/L) is lethal after 1½ hours of exposure (EPA, 1986). The EPA has derived an RfC (reference concentration) of 0.48 mg/m³ (milligrams per cubic meter) for general population exposure to methyl bromide (EPA, 1992).

Methyl bromide is rapidly absorbed by the lungs and affects both the lungs and kidneys. Increased exposure to methyl bromide results in elevation of bromine levels in the blood; poisoning symptoms occur at a level of 2.8 mg/100 ml of blood (Curley, 1984). Symptoms of acute exposure typically are headache, dizziness, visual problems, gastrointestinal disturbances, and respiratory problems. In more extreme cases, muscular pain, numbness, or twitching precede convulsions, unconsciousness, and possibly death.

Chronic exposure can result in behavioral changes, loss of ability to walk, neurological damage, and renal and liver function disturbances (Verberk *et al.*, 1979). Because there are a number of toxicity data gaps, the chronic and subchronic toxicity of methyl bromide is not well characterized. For this reason, and the implication of its contribution to ozone depletion, the EPA has issued a call-in notice to provide this information for reregistration. Manufacturers must supply more information.

Based on laboratory studies of the effects of methyl bromide inhalation and ingestion, nontarget species of mammals and birds exhibit symptoms similar to humans: weakness, lack of muscular coordination, neurological and behavioral abnormalities, and death from high doses. Due to its restricted use as a fumigant, wild animals are rarely exposed to methyl bromide and toxicity data is limited to farm animals. Residues in hay ranging from 6,800 ppm to 8,400 ppm caused symptoms of intoxication in cattle, horses, and goats (Knight and Costner, 1977).

c. Exposure and Risk

(1) Humans

Inhalation is the primary exposure route for methyl bromide. Concentrations of methyl bromide are electronically monitored during the fumigation. Because the gas is odorless and nonirritating during exposure and the onset of symptoms is delayed, leaks and spills causing extreme exposure can occur without persons being aware of its presence. Protective clothing and self-contained breathing apparatus are worn whenever concentrations of methyl bromide are anticipated to reach or exceed 5 ppm. The American Conference of Governmental Industrial Hygienists (ACGIH) has established exposure standards (Threshold Limit Value) of 5 ppm (20 mg/m³) to protect against adverse neurotoxic and pulmonary effects (ACGIH, 1990). Dermal exposure to workers could occur in the unlikely event of a spill of liquid methyl bromide.

Ingestion of methyl bromide residues and its degradation products is a third exposure route. Following aeration of the commodity, the small amount of methyl bromide that remains dissipates and degrades, leaving only inorganic bromide residues. However, residues from the methyl bromide fumigation will remain on the commodity. EPA tolerances for residues of methyl bromide, measured as inorganic bromides (40 CFR 180.123), range from 5 ppm (for apples, pears, and quinces) to 240 ppm (for popcorn), with most commodities at 50 ppm or less. Ingestion of these small amounts of residues is considered to have no adverse toxicological effect.

EPA has classified methyl bromide as a class I ozone depleting chemical in a manner consistent with the Montreal Protocol. The EPA is expected to require the phase-out of most uses of methyl bromide by 2005. There is, however, an exemption to this phase-out for Quarantine and Pre-Shipments uses. The relative importance of methyl bromide to ozone depletion, however, is subject to fundamental uncertainties.

Workers will have little exposure to methyl bromide because fumigations are contained. The public will be restricted from access to the fumigation chamber by a 30-foot wide barrier zone. Residues in fumigated commodities will be within tolerance limits. There is very little risk to human health from a methyl bromide fumigation.

(2) Nontarget Species

Few nontarget species will be exposed to methyl bromide directly. The aeration duct will deliver a plume which will disperse quickly. Species within this plume, such as insects which inadvertently fly in, might die. However, these effects are restricted to areas within the 30-foot wide barrier zone (Bergsten, personal communication). In addition, ground-dwelling organisms immediately outside the fumigation chamber vent are not anticipated to survive.

(3) Endangered and Threatened Species

Fumigation chambers are generally located in high traffic areas; tarped fumigations occur in agricultural areas. These areas are highly disturbed and are very unlikely to harbor endangered and threatened species. Therefore, it is not expected that any endangered or threatened species will be exposed to methyl bromide fumigation.

4. Cumulative Impacts

Cumulative impacts are those impacts, either direct or indirect, that result from incremental impact of the program action when added to other past, present, and reasonably foreseeable future actions. It is difficult to quantitatively predict the cumulative impacts for a potential emergency program in an environmental assessment such as this. The impacts can be considered from a subjective perspective.

Some chemicals, when used together, have been shown to act in a manner that produces greater toxicity than would be expected from the addition of both. This effect is known as potentiation or synergism. Malathion bait spray and diazinon could be applied during the same treatment regimen. Because malathion has frequently been observed as one constituent of a potentiating pair of organophosphorus insecticides (Murphy, 1980), synergistic effects from the combination of malathion and diazinon (both organophosphorus insecticides) could occur. However, malathion bait spray is applied to the tree canopy and diazinon to the soil within the drip line of the canopy, so synergistic effects are limited to animals that are active on both foliage and soil. In addition, the restriction of diazinon treatments to plants with infested fruits make it unlikely that any animals would get concurrent exposure to both insecticides.

Impacts from implementation of the program are expected to be temporary with potential adverse effects ending shortly after the infestation is eradicated.

No bioaccumulation or environmental accumulation of malathion or diazinon is foreseen due to the rapid degradation rates. In contrast, the ongoing applications expected from the no action alternative would be expected to have cumulative effects. Therefore, any cumulative impacts of the program are expected to be less than those that might occur under the no action alternative, an alternative which most likely would result in escalating use of pesticides by the public.

Because the eradication may require the simultaneous use of malathion bait spray and diazinon, both of which are organophosphate cholinesterase inhibitors, there could be cumulative effects of using two pesticides. The history of the eradication efforts for the Mexican fruit fly shows that this use pattern does not result in adverse effects to the general resident population nor the workers. Because most nontarget species are mobile, it is unlikely that an individual will be exposed to more than one treatment. In addition, diazinon treatments are restricted to locations where Mexican fruit fly larvae are detected. Domestic animals and less mobile organisms, such as those dwelling near the soil surface, could be exposed.

The cumulative impacts from fumigations with methyl bromide have been discussed in detail in an environmental impact statement designed to address these issues (USDA, APHIS, 2002). This document included analysis of the limited use of methyl bromide in local eradication programs such as this one and determined that their potential cumulative contribution to ozone depletion is not significant.

In terms of the cumulative effects of pesticide use from the proposed action with pesticide use from other fruit fly programs, the small area requiring treatment for this program should not substantially increase exposure to workers, public, or nontarget species.

5. Methods To Reduce Risk

Human pesticide exposure would be primarily to workers, especially in the case of the soil drench pesticide, diazinon, or methyl bromide which is used only in certified fumigation chambers or under tarpaulins (enclosures). Residents within the eradication area will be exposed to malathion bait spray and diazinon to an extent depending on where the pesticides are applied. The public could be exposed to residues on any treated material moved out of the eradication area.

Current worker safety measures protect fumigators and other pesticide applicators from excessive exposure to methyl bromide, diazinon, and

malathion, during routine operations. To minimize worker exposure to methyl bromide, the fumigation chamber is opened only after concentrations are reduced below 5 ppm. Proper sealing of fumigation enclosures and proper aeration facilitate dispersal of the fumigant. Diazinon Worker exposure to diazinon can be prevented by gloves and safety goggles, which are indicated as protective clothing requirements on the label (Meister, 1990). Studies on exposure to diazinon during yard applications reveal that 85% of the exposure to workers is to their hands. Dermal exposure of workers to malathion can also be substantially reduced by the use of protective clothing. Written public notification will provide information about the schedule for pesticide treatments and applications and about specific precautions that residents should take to avoid excessive exposure, such as remaining indoors during malathion bait spray applications or diazinon soil treatments or not harvesting malathion-treated produce for 3 days after application. However, individuals with greater sensitivity to cholinesterase inhibitors or the protein bait may need to take extra precautions to avoid even minimal exposure.

The program, properly implemented, represents a relatively low risk to human health except for extremely sensitive individuals who have had problems with similar programs in the past. However, this assessment does contain uncertainties associated with toxicity data gaps and estimations of exposure. Furthermore, synergistic interactions between the pesticides which could be used in this program as well as other pesticides not associated with the program and possibly used in the same area could increase toxicity and the associated risk. Potential risk will be substantially diminished due to the localized nature and short duration of the program.

Risks to nontarget organisms can be reduced by limiting exposure. If aerial applications are conducted, beekeepers and backyard pond owners should be notified. A survey of water bodies within the treatment area should be conducted and mapped so they will be avoided by establishing “no treatment” zones during aerial operations. Ground application of malathion bait spray poses little direct risk. Pet owners should be notified to limit animals’ exposure to treated trees. Soil treatments pose more risk due to higher toxicities and a barrier or other safeguards should be used. Timing of the treatment should be considered to reduce exposure. Standard operating procedures for methyl bromide fumigations include fencing or other barriers to limit access to the fumigation and aeration area and preclude exposure of many vertebrates.

The FWS or the California Natural Diversity Database will be consulted if the program area is expanded to ensure that endangered or threatened species are not adversely impacted.

The potential environmental impacts of the program's alternatives and component treatment methods have been discussed and analyzed in detail within the risk assessments. In addition, potential cumulative impacts were analyzed as well. Refer to the risk assessments for greater detail.

6. Site-specific Issues

This environmental analysis focuses on site-specific issues and conditions, especially with respect to any effects they might have on potential environmental impacts. Issues of concern associated with this proposed action include (1) potential effect on human health from chemical pesticide applications, (2) potential effect on wildlife (including endangered and threatened species) from program activities and treatments, and (3) potential effect on environmental quality.

The area of the proposed program has urban and suburban characteristics. The current eradication zone (where eradication treatments will occur) is the area including and immediately surrounding the Mexican fruit fly detections – an area approximately 14 square miles. The eradication zone is predominantly residential, but includes some plant nurseries and fruit sellers. Parts of Alhambra, Montebello, Monterey Park, and Rosemead are within the eradication area. Outside of the previously described considerations for endangered and threatened species habitat, there are no sensitive sites within the treatment zone. However, expansion of the program area could place some sensitive sites within the quarantine zone. The Angeles National Forest is further to the north. The Los Angeles River is on the eastern edge of the eradication zone. The program has adjusted treatments in the spray areas to minimize human exposures through the use of ground applications rather than aerial applications. If the treatment zone should expand in the future to include the national forest, appropriate protection measures will be employed to avoid adverse impacts to these areas.

Consistent with Executive Order 12898, "Federal Actions To Address Environmental Justice in Minority Populations and Low-Income Populations," APHIS considered the potential for disproportionately high and adverse human health or environmental effects on minority populations and low-income populations. The population of this area is diverse and lacks any special characteristics that differ from that of the general population. There are, however, some areas that have minority communities within the

county. The potential for continuing expansion of the treatment area could result in potential exposure of many different communities to treatment chemicals, but there is no evidence that any one population is likely to have disproportionate effects from these program activities.

APHIS also recognizes that a proportion of the population may have unusual sensitivity to certain chemicals or environmental pollutants and that program treatments pose higher dangers for these individuals. Consistent with Executive Order 13045, "Protection of Children From Environmental Health Risks and Safety Risks," APHIS considered the potential for disproportionately high and adverse health or safety effects to children. The potential risks from potential program actions were determined to pose no excess risk to children. Special notification procedures and precautions are required and serve to minimize the risks to individuals and groups considered to have potential for increased sensitivity.

The alternatives were compared with respect to their potential to affect local nontarget species. Paralleling the findings for human health, we have determined that a well-coordinated eradication program using integrated pest management (IPM) technologies would result in the least use of chemical pesticides overall with minimal adverse impact to nontarget species. The no action alternative and nonchemical control alternative would be expected to result in broader and more widespread use of pesticides by homeowners and commercial growers, with correspondingly greater potential for adverse impact.

APHIS has consulted with the U.S. Department of the Interior, Fish and Wildlife Service (FWS), under the provisions of section 7 of the Endangered Species Act of 1973. California Natural Diversity Database indicates that Least Bell's vireo and southwestern pond turtle reside within the current eradication zone. There are no plans to treat any riparian, marsh, or sand dune habitats or areas not adjacent to paved roads. This precludes any effects on the species found within the eradication zone. If the program expands into other areas of those counties, and if there is a potential for affecting federally listed or proposed endangered and threatened species, APHIS will consult with FWS over protective measures that may be required. No adverse impacts to endangered or threatened species, or their habitats, are foreseen.

The area was considered with respect to any special characteristics that would tend to influence the effects of program operations. Potentially

sensitive areas have been identified, considered, and accommodated through special selection of control methods and use of specific mitigative measures. The present program area contains no special characteristics that would require a departure from the standard operating procedures and mitigative measures.

In conclusion, the majority of the risk in the program is associated with pesticide use. Pesticide exposure and subsequent risk to humans and nontarget species is not expected to be substantial in this program because of the localized nature of the infestation, the limited use of pesticides, the precise targeting of pesticides, and the safety procedures employed. Although minimal exposure could pose higher risk to some sensitive individuals and some nontarget organisms, pesticide exposure is generally expected to be minimal and program standard operating procedures and mitigations (especially notifications) serve to minimize that risk. Risk to environmental quality is considered minimal. No significant cumulative impacts are expected as a consequence of the proposed program or its component treatment methods.

IV. Agencies, Organizations, and Individuals Consulted

California Department of Food and Agriculture
Department of Plant Industry
Sacramento, California

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
Invasive Species and Pest Management
4700 River Road, Unit 134
Riverdale, MD 20737-1236

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Policy and Program Development
Environmental Services
4700 River Road, Unit 149
Riverdale, Maryland 20737-1238

Appendix A. References Cited

ACGIH. American Conference of Governmental Industrial Hygienists, 1990. Threshold limit values for chemical substances and physical agents and biological exposure indices. Pub. No. 0205. Cincinnati, OH.

Bergsten, D., 1992. Observations as a PPQ Officer, New Jersey.

Curley, W.H., 1984. Methyl bromide: A profile of its fate and effect in the environment. Submitted by Roy F. Weston, Inc., West Chester, PA, to U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine. Contract No. 53-3294-3-39.

Eisler, R., 1986. Diazinon Hazards to Fish, Wildlife, and Invertebrates: A synoptic review. U.S. Department of the Interior, Fish and Wildlife Service, Patuxent Wildlife Research Center. Biological Rpt. No. 85 (1.9).

EPA. See U.S. Environmental Protection Agency.

Hudson, R.H., Tucker, R.K., and Haegele, M.A., 1984. Handbook of toxicity of pesticides to wildlife (2nd ed.). U.S. Department of the Interior, Fish and Wildlife Service. Resource Pub. 153. Washington, DC.

Kizer, K.W., 1991. Health risk assessment of aerial application of malathion-bait. California Department of Health Services, Berkeley, CA.

Knight, H.D., and Costner, G.C., 1977. Bromide intoxication of horses, goats, and cattle. J. Am. Vet. Med. Assoc. 171(5):446-448.

Meister, R.T. (ed.), 1990. Farm Chemicals Handbook. Meister Publishing Co., Willoughby, OH.

Murphy, S.D., 1980. Pesticides. In J. Doull, C.D. Klassen, and M.O. Amdur (eds.), Toxicology: The basic science of poisons (2d ed.), p. 357-408. Macmillan Publishing Co., Inc., New York.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2002. Rule for the importation of unmanufactured wood articles from Mexico, with consideration for cumulative impact of methyl bromide use. Final environmental impact statement - September 2002. USDA, APHIS, Riverdale, MD.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1998a. Human health risk assessment for fruit fly cooperative control programs. USDA, APHIS, Riverdale, MD.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1998b. Nontarget species risk assessment for fruit fly cooperative control programs. USDA, APHIS, Riverdale, MD.

U.S. Environmental Protection Agency, 1992. Office of Research and Development. Integrated Risk Information System (IRIS). Cincinnati, OH.

U.S. Environmental Protection Agency, 1986. Pesticide Fact Sheet No. 98: Methyl bromide (August 22). Office of Pesticides and Toxic Substances, Office of Pesticide Programs, Washington, DC.

Verberk, M.M., Rooyackers-Beemster, T., De Vlieger, M., and Van Vliet, A.G.M., 1979. Bromine in blood, EEG and transaminases in methyl bromide workers. *British J. Ind. Med.* 36:59-62.

**Finding of No Significant Impact
for
Mexican Fruit Fly Cooperative Eradication Program
Monterey Park (Los Angeles County), California
Environmental Assessment,
October 2002**

The U.S. Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), has prepared an environmental assessment (EA) that analyzes potential environmental consequences of alternatives for eradication of the Mexican fruit fly, an exotic agricultural pest that has been found in the Monterey Park area of Los Angeles County, California. The EA, incorporated by reference in this document, is available from:

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
9580 Micron Avenue, Suite 1
Sacramento, CA 95827

or

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
Invasive Species and Pest Management
4700 River Road, Unit 134
Riverdale, MD 20737-1236

The EA analyzed alternatives of (1) no action, (2) nonchemical control, and (5) integrated control (including chemicals). Each alternative was determined to have potential environmental consequences. APHIS selected eradication of Mexican fruit fly using integrated control for the proposed program because of its capability to achieve eradication in a way that also reduces the magnitude of those potential environmental consequences. Program standard operational procedures and mitigative measures serve to negate or reduce the potential environmental consequences of this program.

The California Natural Diversity Database indicates that two endangered or threatened species reside within the current eradication zone, but no effects to these species or their habitats are anticipated. APHIS will adhere to those protective measures designed specifically for this program and mutually agreed upon with U.S. Fish and Wildlife Service.

I find that implementation of the proposed program will not significantly impact the quality of the human environment. I have considered and based my finding of no significant impact on the quantitative and qualitative risk assessments of the proposed pesticides and on my review of the program's operational characteristics. In addition, I find that the environmental process undertaken for this program is entirely consistent with the principles of "environmental justice," as expressed in Executive Order No. 12898 and protection of children as expressed in Executive Order No. 13045. It is expected that the program will pose no disproportionate adverse impacts to minority populations, low-income populations, or children. Lastly, because I have not found evidence of significant environmental impact associated with this proposed program, I further find that an environmental impact statement does not need to be prepared and that the program may proceed.

/s/

Helene Wright
State Plant Health Director - California
Plant Protection and Quarantine
Animal and Plant Health Inspection Service

11/04/02

Date