

TABLE 1
 THE 20 MOST FREQUENTLY REPORTED SALMONELLA SEROTYPES
 FROM HUMAN SOURCES REPORTED TO CDC IN 1996 AND FROM
 NONHUMAN SOURCES REPORTED TO CDC AND USDA IN 1996

HUMAN 1996				NONHUMAN 1996			
RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT
1	ENTERITIDIS	9570	24.5	1	TYPHIMURIUM *	2958	18.0
2	TYPHIMURIUM *	9501	24.3	2	HEIDELBERG	1641	10.0
3	HEIDELBERG	1998	5.1	3	KENTUCKY	991	6.0
4	NEWPORT	1985	5.1	4	MONTEVIDEO	953	5.8
5	MONTEVIDEO	1227	3.1	5	DERBY	802	4.9
6	JAVIANA	749	1.9	6	AGONA	647	3.9
7	ORANIENBURG	690	1.8	7	CHOLERAESUIS **	606	3.7
8	HADAR	658	1.7	8	ENTERITIDIS	522	3.2
9	AGONA	606	1.6	9	ANATUM	480	2.9
10	MUENCHEN	595	1.5	10	HADAR	440	2.7
11	THOMPSON	586	1.5	11	BRANDENBURG	413	2.5
12	SAINTPAUL	562	1.4	12	MUENSTER	409	2.5
13	BRAENDERUP	531	1.4	13	SENFENBERG	409	2.5
14	INFANTIS	503	1.3	14	INFANTIS	357	2.2
15	TYPHI	440	1.1	15	WORTHINGTON	342	2.1
16	POONA	415	1.1	16	CERRO	305	1.9
17	PARATYPHI B	298	0.8	17	MBANDAKA	286	1.7
18	JAVA	289	0.7	18	SCHWARZENGRUND	268	1.6
19	ANATUM	271	0.7	19	DUBLIN	221	1.3
20	MBANDAKA	223	0.6	20	MELEAGRIDIS	195	1.2
	SUB TOTAL	31697	81.2			13245	80.8
	TOTAL	39035				16399	

* TYPHIMURIUM INCLUDES VAR. COPENHAGEN

** CHOLERAESUIS INCLUDES VAR. KUNZENDORF