

Illustrations by Kelly Campbell



# K-8 Tidepool Math

U.S. Department of the Interior  
Minerals Management Service  
Pacific OCS Region



# Tidepool Math

## K-8

### Lesson Summary

Students use a plot of a mussel bed to better understand the scientific applications of counting and estimating. They will also learn that their environment is constantly changing, both physically and biologically. Through classroom exercises, they will also learn ways to monitor change in the environment.


### Learning Objectives

This curriculum ties in with the CA Math Standards for Grade K-8. Upon completion of this lesson, students will be able to use a mussel bed to:

- Estimate
- Count
- Determine the mean
- Identify Outliers
- Sample

This curriculum also helps students:

- Describe a specific environment
- Define variables in an experiment



### Process Skills Used in this Lesson

1. Be able to sample animals by counting and multiplying.
2. Describe their classroom environment.
3. Observe changes in their local environment.

Material developed by:  
Mary Elaine Dunaway and  
Kelly Campbell  
Minerals Management  
Service  
770 Paseo Camarillo  
Camarillo, CA 93010  
Illustrations by  
Kelly Campbell

Portions adapted from  
materials developed  
by Deby Everton for  
Minerals Management  
Service

# Background for Teachers



## 1. Introduction

**Math is a skill used by many scientists to study changes in the environment. Simple counts and measurements are commonly used by biologists to monitor changes in habitats, populations and ecosystems. By allowing students to glimpse the application of the math skills they're learning, they may find math "more fun." For example, it is much easier to understand why one must estimate the number of mussels in a mussel bed when confronted with a photo of a rocky bench (a large flat rock) covered with them.**

**Scientific methods taught in their classroom also take on new meaning when students see how this is done by field biologists to monitor the environment. Observation, recording, considering and eliminating variables in the experimental design, and critical thinking are all natural parts of monitoring a tidepool, a beach or a classroom.**

## 2. Concepts

**Several terms and concepts are introduced, including *sampling*, *estimating*, *monitoring*, *variables* and *physical and biological environment*.**

**To understand what changes in the environment may be caused by man's actions, we have to know how the environment changes naturally. Because the world is so large, we break it up into smaller components for study; the *physical environment* (non-living things such as rocks, air, and water) and the *biological environment* (living things such as plants and animals).**

**Physical environment is often overlooked, yet in many cases, it overwhelms natural systems. In the marine environment, physical forces such as the tides, storm waves, wind, heat caused by the sun, etc. dominate. Living things have been forced to adapt to the many physical changes in the environment.**

Illustration by  
Kelly Campbell

*Sampling:* selecting and analyzing a subset of elements in order to evaluate the whole. For example, a small group of animals may be selected to evaluate characteristics of the entire population.

*Estimating:* taking a guess about the size of a group. Guessing the number of beans in a jar would be an estimate. Better methods (i.e. sampling) can be used to improve the accuracy of the estimate. Counting the number of beans in one row and column in the jar (a sample) would improve the estimated bean count.

*Monitoring:* taking repeated samples in the same way at the same locations over time. Monitoring increases the sample number and allows the scientists to detect change over time.

*Variables:* Parameters in space and time which are not fixed. Time of year is a variable. Counts of ducks in an area need to be collected at different times of the year to account for seasonal migration.



### 3. Mussel Beds and Monitoring

**Mussel beds are one of the most important biological habitats because of the diversity and density of organisms found in them. Barnacles, snails, limpets, worms, and insects are found in large numbers inside mussel beds.**

Scientists monitor the health of mussel beds along the coast as part of MARINE (Multi-Agency Rocky Intertidal Network). The size of the bed and the changes in density of the mussels over time provide scientists with information to assess changes to a wide variety of plants and animals which depend on the mussel beds. Lobsters often feed in mussel beds; many birds and invertebrates, such as sea stars, depend on mussels for food.


Because mussel beds are so large, it is not possible to count the exact number of mussels; scientists rely on estimates based on collecting repeated samples. Photographs are taken in rectangular plots. The plots are initially laid out in a random pattern and are then marked with permanent bolts on the rock. Each time the scientists return,

they are able to locate the bolt and take photos in the exact same location.

These fixed plots are established in numerous locations. Every site is continuously monitored so measurements of change over time can be made.

Scientists take the digital photos back to the lab and reproduce them on a computer screen. A series of points is overlaid and the species under each point is identified. This measurement will provide the scientist with the percent cover of mussels for that plot.

Often measures of density are also recorded for mussels. To do this, mussels in each plot are counted and recorded either directly in the field, or using a photo of the plot. An estimate of the total number of mussels at a site can be made by multiplying the number of mussels counted in the plots required to cover the mussel beds at an entire site.



(background con't)

# California Mussel

**Latin name:** *Mytilus californianus*

**Range:** Alaska to Baja

**Occurs:** In the rocky intertidal. on piers, oil platform legs and occasionally subtidal to depths of 24m



Illustration by  
Kelly Campbell

## Other Information:

Juvenile mussels settle on hard surfaces, often near adult mussels. We study “recruitment” of mussels, the number of baby mussels which settle in an area, by nailing household “tuffy pads” at the site and later counting the tiny mussels that attach to them.

Mussels growing on the offshore platforms can reach sizes exceeding 12-14”. This fast growth is enabled by currents that race past the platforms carrying nutrients. Testing of mussels harvested regularly off the platforms for food show that they are several magnitudes cleaner than mussels harvested in mariculture areas along the coast. This is because the water offshore is much cleaner than the water near the coast, where the water is polluted by runoff, sewage and other discharges.



# EXERCISE 1:

Using Tidepool plots to  
Count, Estimate, Determine  
a Mean, and Look for Outliers

## Summary

Students use a plot of a mussel bed to better understand the scientific applications of counting and estimating.

## Process Skills

- Sample animals by counting and estimating
- Provide reasons for changes in their local environment

## Instructional Strategy:

**Start first by asking the students to count the number of students in their class. Get students to think about what inferences can be made about environmental conditions by counting the number of individuals in an area.**

**Ask** what their conclusions would be if they came to this same classroom and only 5 students were present.  
-Their reasons may include vacation, chicken pox, and flu. A critical thinker may suggest that you are counting at the wrong time, say before school.

**Show slide of mussel bed**  
**Introduce the importance of mussel beds.**

**Discuss** which intertidal animals depend on them for shelter and food.


-Examples include barnacles, snails, limpets, worms, insects, birds, lobsters, sea stars, and other invertebrates

**Discuss** why it would be important to know how many mussels are found here. **Discuss** why knowing about changes in the mussel bed would be useful.

-The condition of a mussel bed can be a good indicator of the overall condition of the ecosystem in the sampled area. Changes in the mussel bed may reflect changes in the surrounding environment.

**Ask** what a decrease in numbers might mean.

-They were eaten, they got sick, a big storm pulled them off the rocks, etc. A bright student may suggest that you are counting in a season when mussels were fewer (this doesn't apply to mussels but it is true for other animals).




# Classroom Activity 1:

## Counting Mussels

**Materials:** Enlarged photo of mussel plot, laminate (2' by 3'), "vis-a-vis", masking tape

**Preparation:** Enlarge mussel photo to 2' by 3 and mount on foam board



A. Place photo on floor as if it were a plot of mussels at the shoreline.

B. Overlay Laminate

C. Place masking tape over laminate to divide plot into fourths

D. Ask students to estimate the number of mussels in the total plot and in each square.

E. Have four students each count one square. Mark counted mussels with "vis-à-vis" or similar dry erase marker.

F. Have each student write the number they counted on the board. Have one student add them up to find the total and have another divide them by four to find a mean.

G. Compare these numbers to the estimates. Compare the mean to the individual counts. Discuss any outliers.

H. Multiply each square total to get four estimates of the entire plot. Compare to total, Discuss how much easier it is to count one square rather than the entire plot

**Note:** Younger students may want to count each section of the plot with you. Consider making the photo an overhead and counting as a class.

**Challenge:** Older students could discuss the fact that plots may not be homogenous- there may be an uneven distribution distribution of mussels within the plot. Some areas may be tightly packed while others may be barren rock. Discuss how you would accurately estimate the mussels in plots that are highly variable. Discuss why counting more than one plot in an area would increase the accuracy of the estimate. Discuss the importance of keeping *outliers* in the data set if the objective is to determine the density of mussels at a highly variable site.

In this case all densities found need to be included in determining an overall estimate.  
**Extra Practice:** Have students complete the "Estimating with Sea Stars" worksheet.



# EXERCISE 2:

Understanding the Environment through monitoring

## Summary

Students will identify the biological and physical environment inside and outside their classroom.

## Process Skills

- Describe the biological and physical environments
- Understand sampling and monitoring
- Identify controlled variables

## Instructional Strategy:

This exercise provides students with increased awareness about the *physical environment*. Very few people take note of the impact the physical environment has on the changing world. Except for the factors that directly impact the comforts of everyday life, many physical changes in the environment go unnoticed.

**Ask** your students to define environment.

- The environment is everything. The two components of the environment are physical and biological. Everything on earth falls into these two categories.

**Ask** what some physical parts of the environment are?

-Examples include air, sunlight, chairs, windows.

**Ask** what some parts of the *biological environment* are?

-Examples include humans, animals, plants

**Ask** the students what scientists have to do before they can tell if something, like installing an oil platform, will impact the environment?

-They must determine the present conditions and perform continuous *monitoring* in order to assess the impact.



## Monitoring Discussion:

**Introduce the term monitoring using the example of ducks in a pond.**

**Ask** if the environment changes? If a biologist wanted to know about the ducks in a pond, what would be her approach?  
-She would go to the pond and watch them


**Ask** them: Suppose you went to this pond in December and discovered five ducks swimming in the pond. Would this tell you how many ducks lived there?  
-Students may suggest that there could have been a lot more birds but they all flew south in winter.

**Ask** would the same five ducks be there tomorrow? Is it possible that other ducks live there but weren't there when the biologist was? How can the biologist get a truer picture of how many ducks live at that pond?  
-The biologist has to go back to the same pond over and over again to see what ducks are there. This is called *monitoring*.

**Introduce the term "variable."**  
**Variables in this case could be time of day, time of year, temperature outside.**

**Ask** what other variables might effect the duck population while monitoring at a particular time  
-Human presence, water or air temperature, maybe ducks forage in a different place when it is very cold or very hot.

**Discuss** what the biologist might find at different times of the day, or different months of the year.



## Monitoring Plan:

Have your students write down a possible monitoring plan that would allow them to monitor the ducks at this pond adequately. Have them specify which variables need to be recorded during each monitoring period.



## Classroom Activity 2:

### How does your environment change?

**Materials:** Journals, pencils

A. Ask two students to step outside the classroom for a few minutes. Help remaining students change one aspect of the physical environment (open a door or window, move a chair, etc.)

B. Ask the returning students to identify changes in the physical environment.

C. With journals in hand, ask students to go outside their classroom and record as many items as they can find in their biological and physical environment.

D. Have them compare lists and discuss the importance of being a detailed observer.

**Tell the students:** After a biologist has made a detailed description of the existing environment, they need to predict what they think can or will happen if a certain project is carried out. They have to look at impacts on the environment -what is going to happen and what activity is planned.

**Ask** how they affected the classroom environment today.


-Stepped on an insect, introduced new germs, changed the number of people in the room, changed the oxygen content of the air

These human impacts affect the classroom environment.

When determining the environmental impact in the real world, **the big question is:** Is the impact something to worry about? Science and judgement are combined to reach the answer to this question.

Will the effect last a very long time, such as 100 years, or a short time, such as a minute? Will the effect be felt over a large area, or a very small area? Will it affect the entire community or just one particular species?





**Ask** how they would rate stepping on some ants? The effect is probably felt over a very small area, and given the high number of ants, the population of ants will probably not be affected at all. How about losing several endangered birds? Loss of several endangered birds could negatively affect the entire population, an effect that would be felt over a large area and could require many years to recover.





# Estimating with Sea Stars

Name \_\_\_\_\_

<p>1</p> 	<p>Total arms _____ Total stars _____</p> <p>Average arms per star _____</p>
<p>2</p> 	<p>Total arms _____ Total stars _____</p> <p>Average arms per star _____</p>
<p>3</p> 	<p>Total arms _____ Total stars _____</p> <p>Average arms per star _____</p>
<p>4</p> 	<p>Total arms _____ Total stars _____</p> <p>Average arms per star _____</p>

Illustrations by Kelly Campbell

Sea stars have the ability to regenerate broken arms. This adaptation is both a defense mechanism and a mode of reproduction. If a large enough portion of the star is broken off it can develop into a an entire new star.

Count the total number of arms in each sea star community. Find the average number of arms per star by dividing the total arms by the total stars in each community.

What is the average number of arms per star for all four communities? \_\_\_\_\_





How many whole arms would the average star in community 3 have? \_\_\_\_\_

Are there any outlier sea stars? \_\_\_\_\_ How many arms do they have? \_\_\_\_\_

Why is estimating a useful and important tool for scientists?

# Estimating with Starfish

Name \_\_\_\_\_

<p>1</p> 	<p>Total arms <u>56</u> Total stars <u>16</u></p> <p>Average arms per star <u>3.5</u></p>
<p>2</p> 	<p>Total arms <u>60</u> Total stars <u>21</u></p> <p>Average arms per star <u>2.9</u></p>
<p>3</p> 	<p>Total arms <u>66</u> Total stars <u>18</u></p> <p>Average arms per star <u>3.7</u></p>
<p>4</p> 	<p>Total arms <u>64</u> Total stars <u>19</u></p> <p>Average arms per star <u>3.3</u></p>

Illustrations by Kelly Campbell

Sea stars have the ability to regenerate broken arms. This adaptation is both a defense mechanism and a mode of reproduction. If a large enough portion of the star is broken off it can develop into an entire new star.

Count the total number of arms in each sea star community. Find the average number of arms per star by dividing the total arms by the total stars in each community. Depending on math skills, answers can be averages as fractions (arms/stars), mixed numbers or decimals (key has decimals with two significant digits\*)

What is the average number of arms per star for all four communities? 3.4

How many whole arms would the average star in community 3 have? 3

The average is 3.7, but arms are counted here in whole numbers. The answer must be rounded down to 3.

Are there any outlier sea stars? yes How many rays do they have? 1

Why is estimating a useful and important tool for scientists?

Estimating allows scientist to look at a small portion of an area of interest and make assumptions about the entire area. It also enables them to make generalizations about an area, focusing on the condition of the majority. It can be assumed that stars in population 2 (where average arms is 2.9) may be more likely to lose limbs than stars in population 3 (where average arms is 3.7).