

INITIAL REVIEW DRAFT

**ENVIRONMENTAL ASSESSMENT / REGULATORY IMPACT REVIEW /
INITIAL REGULATORY FLEXIBILITY ANALYSIS**

for

**Revised Management Authority of Pelagic Shelf Rockfish Complex in the GOA and the
Other rockfish complex in the BSAI (Dark Rockfish)**

Proposed amendments to the
Fishery Management Plans for Gulf of Alaska Groundfish and Bering Sea Aleutian Islands
Groundfish

Prepared by staff of the
North Pacific Fishery Management Council



For Further Information Contact:
Diana Stram
North Pacific Fishery Management Council
605 West 4th Avenue, #306
Anchorage, Alaska 99501-2252
(907) 271-2809

January 19, 2007

[THIS PAGE INTENTIONALLY LEFT BLANK]

EXECUTIVE SUMMARY

This Environmental Assessment, Regulatory Impact Review and Initial Regulatory Flexibility Analysis describes the proposed amendment to the Gulf of Alaska Groundfish and Bering Sea Aleutian Islands Groundfish Fishery Management Plans (FMPs). This amendment proposes to remove dark rockfish (*Sebastes ciliatus*) from the GOA and BSAI groundfish FMPs. This species is currently contained in the pelagic shelf rockfish (PSR) assemblage in the GOA and in the other rockfish complex in the BSAI. It makes up a small proportion of the total biomass in each complex, is more often found in nearshore waters, and is caught in State fisheries. Removing this species from these FMPs would turn management for this species in both State and Federal waters over to the State of Alaska.

The following problem statement is proposed for this analysis:

Dark rockfish are a nearshore, shallow water species which are rarely caught in offshore, Federal waters. For management purposes they are contained within the pelagic shelf rockfish complex in the GOA, whose OFL and ABC are based primarily on the stock assessment for dusky rockfish which makes up the majority of the total exploitable biomass estimate for the PSR complex. In the BSAI dark rockfish are contained within the other rockfish complex whose biomass is largely comprised of dusky rockfish and thornyhead rockfish. As dark rockfish have now been identified as a separate species, are found in nearshore, shallow waters, and could potentially be locally overfished within the larger PSR complex TAC in the GOA, the Council should consider removing this species from the GOA groundfish FMP thereby transferring their management to the State of Alaska. For consistency in management the Council should also consider removing this species from the BSAI FMP.

Two actions are analyzed in this document with two alternatives for each action: Action 1 refers to the GOA groundfish FMP. Under this action there are two alternatives: alternative 1, to continue managing dark rockfish within the larger pelagic shelf rockfish complex; and alternative 2, to remove dark rockfish from the GOA FMP and turn over to the State of Alaska for management. Action 2 refers to the BSAI groundfish FMP. Under this action there are two alternatives: alternative 1, to continue managing dark rockfish within the other rockfish complex; and alternative 2, to remove dark rockfish from the BSAI FMP and turn over to the State of Alaska for management.

Environmental Assessment

There is limited impact in the Federal fishery of removing this species from either FMP. Dark rockfish comprise a small proportion of the total biomass in the PSR assemblage, which is dominated by the target species, dusky rockfish. Impacts to other PSR stocks as well as other groundfish stocks are minimal due to the relatively minor contribution to the overall exploitable biomass from the dark rockfish stock. Dark rockfish makes up a very minor component of the total biomass in the other rockfish complex in the BSAI. This is not a target fishery, and retained catch is dominated by shortspine thornyhead rockfish and dusky rockfish. These two species make up the majority of the biomass in the complex.

Management of dark rockfish by the State is anticipated to be an improvement over Federal management within the PSR complex due to the State's ability to manage this stock as a single stock and on smaller management areas to protect against the potential for localized depletion. There are no anticipated impacts to marine mammals, seabirds, threatened or endangered species, habitat or the ecosystem.

Regulatory Impact Review

Removal of dark rockfish from the pelagic shelf rockfish complex in the GOA could result in minor decreases in the pelagic shelf rockfish TAC, but since dark rockfish are such a small part of the stock of

the complex any decline in the TAC is likely to be nominal. Removal of dark rockfish from the other rockfish complex in the BSAI will result in a minimal decrease in the TAC for this complex.

Initial Regulatory Flexibility Analysis

Transfer of management of dark rockfish to the State is likely to result in some changes in regulation of catch. The State could develop a directed fishery for dark rockfish, most likely for fixed gear vessels. Since fixed gear vessels tend to be small, it is possible that the development of such a directed fishery would have a positive impact on small entities, by increasing fishing opportunities. The IRFA in this document is preliminary until the Council selects a preferred alternative. At that point, the potential impact on affected small entities of the action will be developed further in the analysis.

TABLE OF CONTENTS

Environmental Assessment.....	i
Regulatory Impact Review.....	i
Initial Regulatory Flexibility Analysis.....	ii
1.0 PURPOSE AND NEED FOR ACTION	1
1.1 Purpose and Need	1
1.1.1 Problem Statement.....	1
1.2 Next Steps in the Process	2
2.0 DESCRIPTION OF ALTERNATIVES	3
2.1 Action 1: GOA groundfish FMP	3
2.1.1 Alternative 1: Status quo.....	3
2.1.2 Alternative 2: Remove dark rockfish from the Gulf of Alaska FMP.....	3
2.2 Action 2: BSAI groundfish FMP	4
2.2.1 Alternative 1: Status Quo.....	4
2.2.2 Alternative 2: Remove dark rockfish from the BSAI FMP	4
2.3 Alternatives Considered but not Carried Forward.....	5
3.0 AFFECTED ENVIRONMENT	6
3.1 General distribution and habitat requirements of dark rockfish	6
3.1.1 Life history characteristics of <i>Sebastes</i> rockfish species	8
3.1.2 Biomass by species	10
3.1.3 Stock Assessment	30
3.2 Pelagic Shelf Rockfish Fishery (GOA)	31
3.3 BSAI other rockfish fishery.....	36
3.4 Other Groundfish Stocks	40
3.4.1 GOA black rockfish fishery	41
3.4.2 BSAI black rockfish fishery	44
3.5 Threatened and Endangered Species	45
3.6 Marine Mammals.....	47
3.7 Seabirds	47
3.8 Habitat and Essential Fish Habitat.....	47
3.9 Ecosystem Considerations.....	49
4.0 ENVIRONMENTAL IMPACTS	52
4.1 Action 1 GOA FMP.....	52
4.1.1 Alternative 1: Status quo.....	52
4.1.2 Impacts on Pelagic Shelf Rockfish Stocks.....	52
4.1.3 Impacts on Other Groundfish Stocks.....	52
4.1.4 Impacts on Threatened or Endangered Species	53
4.1.5 Impacts on Marine Mammals	53
4.1.6 Impacts on Seabirds.....	53
4.1.7 Impacts on Habitat and EFH.....	53
4.1.8 Impacts on the Ecosystem.....	53
4.1.9 Socioeconomic Impacts	53
4.2 Alternative 2	53
4.2.1 Impacts on Pelagic Shelf Rockfish Stocks.....	53
4.2.2 Impacts on other groundfish stocks	55
4.2.3 Impacts on threatened or endangered species	55
4.2.4 Impacts marine mammals	55
4.2.5 Impacts on seabirds.....	55
4.2.6 Impacts on habitat and EFH.....	55
4.2.7 Impacts on the ecosystem	56

4.2.8	Socio-economic impacts	56
4.3	Action 2 BSAI FMP	56
4.3.1	Alternative 1: Status quo.....	56
4.3.2	Impacts on Other Rockfish Stocks.....	56
4.3.3	Impacts on Other Groundfish Stocks.....	56
4.3.4	Impacts on Threatened or Endangered Species	56
4.3.5	Impacts on Marine Mammals	57
4.3.6	Impacts on Seabirds	57
4.3.7	Impacts on Habitat and EFH.....	57
4.3.8	Impacts on the Ecosystem.....	57
4.3.9	Socioeconomic Impacts	57
4.4	Alternative 2	57
4.4.1	Impacts on Other Rockfish Stocks.....	57
4.4.2	Impacts on other groundfish stocks	58
4.4.3	Impacts on threatened or endangered species	58
4.4.4	Impacts marine mammals	59
4.4.5	Impacts on seabirds.....	59
4.4.6	Impacts on habitat and EFH.....	59
4.4.7	Impacts on the ecosystem	59
4.4.8	Socio-economic impacts	59
4.4.9	Cumulative Impacts	59
5.0	REGULATORY IMPACT REVIEW	61
5.1	Introduction	61
5.2	What is a Regulatory Impact Review?	61
5.3	Statutory Authority	61
5.4	Purpose and Need for Action.....	61
5.5	Alternatives Considered	62
5.6	Action 1: GOA groundfish FMP	62
5.6.1	Alternative 1: Status quo.....	62
5.6.2	Alternative 2: Remove dark rockfish from the Gulf of Alaska FMP.....	62
5.7	Action 2: BSAI groundfish FMP.....	63
5.7.1	Alternative 1: Status Quo.....	63
5.7.2	Alternative 2: Remove dark rockfish from the BSAI FMP	64
5.8	Background.....	64
5.9	Analysis of the Alternatives	68
6.0	INITIAL REGULATORY FLEXIBILITY ANALYSIS	70
6.1	Introduction	70
6.2	The Purpose of an IRFA.....	70
6.3	What is required in an IRFA?.....	71
6.4	What is a small entity?	71
6.5	Reason for considering the action	72
6.6	Objectives of, and legal basis for, the proposed action	73
6.7	Number and description of small entities regulated by the proposed action	73
6.8	Impacts on Regulated Small Entities.....	73
6.9	Recordkeeping and Reporting Requirements	73
6.10	Federal rules that may duplicate, overlap, or conflict with proposed action	73
7.0	CONSISTENCY WITH APPLICABLE LAW AND POLICY	73
7.1	Magnuson-Stevens Act.....	73
7.1.1	National Standards.....	73
7.1.2	Section 303(a)(9) – Fisheries Impact Statement.....	74
7.2	Marine Mammal Protection Act.....	74

7.3 Coastal Zone Management Act 74

8.0 CONSULTATION AND PREPARERS..... 75

8.1 List of Persons and Agencies Consulted 75

8.2 List of Preparers 75

9.0 REFERENCES 75

APPENDIX 1. MRA Tables..... 77

LIST OF FIGURES AND TABLES

Figure 3-1	Locations where dark rockfish were captured during ADF&G surveys, 2001-2006. Survey locations were not systematically distributed but targeted known fish concentrations.....	7
Figure 3-2	Pie charts of black (black portion) and dark (gray portion) rockfish catches in: A. Spruce Island near the city of Kodiak, B. Mountain Point on Nagai Island in the Shumagin Island group, C. the north side of Akutan Island in the eastern Aleutian Islands.....	7
Figure 3-3	Percent Index of Relative Importance (a composite index based on frequency of occurrence, numbers consumed, and weight of prey items, Cortés 1997) for dark and black rockfish from the same area of the Shumagin Islands, August 2005.....	8
Figure 3-4	Age at length of male dark rockfish from Kodiak Island waters with a fitted von Bertalanffy growth curve ($\rho = 0.8363$, $k = 0.1787$, $L_{\infty} = 41.69$	10
Figure 3-5	Dark rockfish CPUE from survey 1999-2005.....	13
Figure 3-6	Dusky rockfish survey catch per unit effort (CPUE), survey years 1996-2003.....	15
Figure 3-7	Widow rockfish survey catch per unit effort (CPUE), survey years 1984-2005.....	17
Figure 3-8	Yellowtail rockfish survey catch per unit effort (CPUE) trawl surveys 1984-2005.....	20
Figure 3-9	Dark rockfish catch in survey hauls by weight 1996-2005.....	23
Figure 3-10	dark rockfish CPUE from the AI survey 1997-2006.....	26
Figure 3-11	Biomass of dark rockfish in survey data by depth strata and region.....	28
Figure 3-12	Occurrence of dark and dusky rockfish in surveys as a percentage of tow biomass in the BS, AI and GOA in conjunction with other State managed species.....	29
Figure 3-13	Percent species composition landed in the 2002 Black rockfish jig fishery (Source ADF&G).....	43
Figure 3-14	Percent species composition landed in the Black rockfish jig fishery 2003, 2005 (Source ADF&G).....	43
Figure 3-15	Observed slope rockfish pelagic trawl effort (hauls/25 square kilometers), 1990 to 2002.....	48
Figure 3-16	Observed slope rockfish non-pelagic (bottom) trawl effort (hauls/25 square kilometers), 1990 to 2002.....	49
Table 1	Instantaneous rate of natural mortality and maximum age for pelagic shelf rockfish, based on the break-and-burn method of aging otoliths. Area indicates location of study: Gulf of Alaska (GOA) or British Columbia (BC).....	9
Table 2	Biomass estimates (mt) for species in the pelagic shelf rockfish assemblage in the Gulf of Alaska, based on results of bottom trawl surveys from 1984 through 2005 (Lunsford et al. 2005).....	11
Table 3	Contribution of dark rockfish survey biomass to overall PSR survey biomass estimate.....	12
Table 4	Dark rockfish in surveyed tows 1996-2005.....	23
Table 5	Dusky rockfish in surveyed tows 1996-2005.....	24
Table 6	Black rockfish in surveyed tows 1996-2005.....	24
Table 7	Biomass from the Aleutian Islands surveys.....	24
Table 8	Biomass from the SE EBS surveys.....	25
Table 9	Biomass totals (by year) AI.....	25
Table 10	Biomass totals (by year) EBS.....	25
Table 11	dark rockfish biomass from survey data by depth strata and region.....	27
Table 12	Relative contribution of dark rockfish survey biomass to the total survey biomass in the AI 1997-2006.....	28
Table 13	Dark rockfish data from surveyed tows in the AI.....	28
Table 14	Dusky rockfish data from surveyed tows in the AI.....	29

Table 15	2007 OFL and ABC, calculated by species.....	30
Table 16	Overfishing levels (OFL), acceptable biological catch (ABC) and total allowable catch (TAC) levels for the GOA pelagic shelf rockfish complex 1998-2006.....	31
Table 17	Commercial catch ^a (mt) of fish in the pelagic shelf rockfish assemblage in the Gulf of Alaska, with Gulfwide values of acceptable biological catch (ABC) and total allowable catch (TAC), 1988-2005. Updated through October 18, 2005. (Lunsford et al. 2005).....	32
Table 18	Percentage of assemblage catch (from observer data).....	34
Table 19	Percent shore-based trawl catch in Central GOA area 1996-2004 (Lunsford et al 2005).....	34
Table 20	Retained catch (mt) of PSR species by gear type 1998-2005 (screened for confidentiality). Source: NMFS Catch Accounting.....	35
Table 21	OFL, ABC and catch for the other rockfish complex in the BSAI 2004-2007.....	36
Table 22	Summary of catches (mt) of other rockfish in the eastern Bering Sea and Aleutian Islands regions. (from Reuter and Spencer, 2006) data from NMFS/AK regional website.	38
Table 23	The common and scientific names of rockfish in the “other rockfish” reporting category identified, 1990- 2001, by AFSC research surveys (at least one observation) and U.S. fishery observers (greater than 1% of hauls) in the eastern Bering Sea and Aleutian Islands. Source: Reuter and Spencer, 2006.....	39
Table 24	Total fishery catch (mt) of top species in other rockfish group in the Aleutian Islands and eastern Bering Sea from 2003-2006. Source: Reuter and Spencer, 2006. data from Catch Accounting System, NMFS AK Regional Office.....	39
Table 25	Catch and effort, excluding discards, for the Kodiak Area black rockfish fishery 1998-2004 (from Sagalkin and Spalinger 2005).....	41
Table 26	Species composition of pelagic shelf rockfish sampled in the Cook Inlet Area jig fishery and surveys 2001-2004.....	44
Table 27	Black rockfish landings (in pounds) in the State Aleutian Islands fishery 1997-2006.....	45
Table 28	Species listed as endangered and threatened under the ESA that may be present in the Federal waters off Alaska include:.....	46
Table 29	Analysis of ecosystem considerations for pelagic shelf rockfish and the dusky rockfish fishery.	50
Table 30	ABC for the pelagic shelf rockfish complex 2002-2006 and the relative contribution from the dark rockfish stock to the overall complex ABC.....	54
Table 31	Ex-vessel value of rockfish catch in the GOA by vessel category and year (\$ millions) from Hiatt et al. 2006.....	64
Table 32	Ex-vessel value of rockfish catch in the BSAI by vessel category and year (\$ millions).....	64
Table 33	Number of vessels and retained catch of pelagic shelf rockfish by vessel category in the GOA.....	65
Table 34	Number of vessels by gear that caught rock fish by vessel category in the BSAI.....	65
Table 35	Total catch of rockfish by gear by vessel category in the BSAI (1,000 metric tons, round weight).....	65
Table 36	Number of catcher vessels, landings, ex-vessel revenues and average ex vessel prices in the Central GOA pelagic shelf rockfish fishery (1998-2002).....	66
Table 37	Number of vessels operating by NMFS management area and gear types for pelagic rockfish (primarily dusky, dark, yellowtail, and widow rockfish).....	67
Table 38	Catch and effort, excluding discards, for the Kodiak Area black rockfish fishery 1998-2004.....	68

1.0 PURPOSE AND NEED FOR ACTION

The groundfish fisheries in the Exclusive Economic Zone (EEZ) (3 to 200 miles offshore) in the Gulf of Alaska (GOA) and the Bering Sea Aleutian Islands (BSAI) are managed under the Fishery Management Plan (FMP) for the Gulf of Alaska Groundfish and the FMP for the Bering Sea Aleutian Islands Groundfish. These FMPs were developed by the North Pacific Fishery Management Council (Council) under the Magnuson Fishery Conservation and Management Act (Magnuson Act).

Actions taken to amend FMPs or implement other regulations governing the groundfish fisheries must meet the requirements of Federal laws and regulations. In addition to the Magnuson Act, the most important of these are the National Environmental Policy Act (NEPA), the Endangered Species Act (ESA), the Marine Mammal Protection Act (MMPA), Executive Order (E.O.) 12866, and the Regulatory Flexibility Act (RFA).

NEPA, E.O. 12866 and the RFA require a description of the purpose and need for the proposed action as well as a description of alternative actions which may address the problem. This information is included in Chapters 1 and 2 of this document. Chapter 3 contains information on the biological and environmental impacts of the alternatives as required by NEPA. Impacts on endangered species and marine mammals are addressed in Chapter 4. Chapter 5 contains a Regulatory Impact Review (RIR) which addresses the requirements of both E.O. 12866 and the RFA that economic impacts of the alternatives be considered. Chapter 6 discusses the potential impacts on small entities per the Regulatory Flexibility Act.

1.1 Purpose and Need

Dark rockfish are part of the pelagic shelf rockfish (PSR) assemblage in the Gulf of Alaska Groundfish Fishery Management Plan (FMP). Members of this assemblage include the following four species: dusky rockfish (*Sebastes variabilis*), dark rockfish (*S. ciliatus*), yellowtail rockfish (*S. flavidus*), and widow rockfish (*S. entomelas*). In the Bering Sea Aleutian Islands FMP dark rockfish are contained within the “other rockfish” complex which contains the following eight species: red banded rockfish (*Sebastes babcocki*), dark rockfish, dusky rockfish, redstripe rockfish (*S. proriger*), yelloweye rockfish (*S. ruberrimus*), harlequin rockfish (*S. variegatus*), sharpchin rockfish (*S. zacentrus*), shortspine thornyhead (*Sebatolobus alascanus*).

The forms of dusky rockfish commonly recognized as “light dusky rockfish” and “dark dusky rockfish” are now officially recognized as two species (Orr and Blackburn 2004). *S. ciliatus* applies to the dark shallow-water species with a common name dark rockfish, and *S. variabilis* applies to variably colored deeper-water species with a common name dusky rockfish.

Dark rockfish are found predominantly in nearshore, shallow waters. Assessment authors have suggested for years that dark rockfish be turned over the State of Alaska for management in the GOA as data in the stock assessment for PSR are predominantly from dusky rockfish (the offshore variety) not dark rockfish (the nearshore, shallow water variety). Most of the available information is from the offshore trawl surveys and offshore commercial fishery and dusky rockfish makes up the majority of the exploitable biomass and catch from the assemblage. A similar concern has been raised by the BSAI plan team for dark rockfish in the overall other rockfish assemblage.

1.1.1 Problem Statement

Given that dark rockfish are located predominantly in nearshore, shallow waters, if specifically targeted the potential exists for them to be locally overfished under the relatively high TAC for the entire pelagic shelf rockfish complex. Amendment 46 to the GOA groundfish FMP addressed a similar situation in the

PSR complex by removing black and blue rockfish, nearshore rockfish populations which were not thought to be well-assessed by the trawl survey, from the GOA groundfish FMP and turned management over to the State of Alaska (NPFMC 1998). A similar situation exists for dark rockfish, and management by the State of Alaska would better address localized harvest requirements for this nearshore species than is currently provided by Federal management under the larger PSR complex in the GOA and the other rockfish complex in the BSAI.

Since official recognition as a separate species, the GOA Plan Team has also endorsed removing dark rockfish from the FMP based on the following rationale: (1) separation at species level, (2) distribution of dark rockfish to nearshore habitats that are not specifically assessed by the GOA trawl survey, and (3) the risk of overfishing dark rockfish in local areas given the relatively high TAC for the pelagic shelf rockfish assemblage as a whole. In 2004, the SSC endorsed the rationale and agreed with the Plan Team's recommendation of removing dark rockfish from the FMP. The Council initiated this in 2005 but action was delayed until the 2005 GOA trawl survey data became available for analysis. An initial review draft of a GOA only amendment was presented to the Council in April 2006. At that time the Council chose to add an alternative to evaluate a similar action for the BSAI FMP due to suggestions made by the BSAI groundfish plan team and the SSC to that effect. The current analysis now evaluates removing dark rockfish from both the GOA FMP and the BSAI FMP.

The following problem statement is put forward to address the analysis for both BSAI and GOA FMPs:

Dark rockfish are a nearshore, shallow water species which are rarely caught in offshore, Federal waters. For management purposes they are contained within the pelagic shelf rockfish complex in the GOA, whose OFL and ABC are based primarily on the stock assessment for dusky rockfish which makes up the majority of the total exploitable biomass estimate for the PSR complex. In the BSAI dark rockfish are contained within the other rockfish complex whose biomass is largely comprised of dusky rockfish and thornyhead rockfish. As dark rockfish have now been identified as a separate species, are found in nearshore, shallow waters, and could potentially be locally overfished within the larger PSR complex TAC in the GOA, the Council should consider removing this species from the GOA groundfish FMP thereby transferring their management to the State of Alaska. For consistency in management the Council should also consider removing this species from the BSAI FMP.

1.2 Next Steps in the Process

This analysis is scheduled for initial review at the February Council meeting. Pending the review process by the Council, the analysis will be revised and released for public review following the February Council meeting. Final action on this amendment is scheduled for April 2007.

2.0 DESCRIPTION OF ALTERNATIVES

Two actions are analyzed in this document with two alternatives for each action: Action 1 refers to the GOA groundfish FMP. Under this action there are two alternatives: Alternative 1, to continue managing dark rockfish within the larger pelagic shelf rockfish complex; and alternative 2, to remove dark rockfish from the GOA FMP and turn over to the State of Alaska for management. Action 2 refers to the BSAI groundfish FMP. Under this action there are two alternatives: Alternative 1, to continue managing dark rockfish within the other rockfish complex; and Alternative 2, to remove dark rockfish from the BSAI FMP and turn over to the State of Alaska for management.

2.1 Action 1: GOA groundfish FMP

2.1.1 Alternative 1: Status quo

Under this alternative, dark rockfish would continue to be managed within the pelagic shelf rockfish assemblage. The Council and the National Marine Fisheries Service would retain management authority for dark rockfish within the PSR complex in the EEZ. Overfishing limits (OFLs), acceptable biological catch (ABC) limits and total allowable catch (TAC) limits are established for the complex as a whole and managed accordingly. In season, catch is managed through monitoring directed fishing, with the fishery closed when directed fishing is estimated to leave only the portion of the TAC necessary to support incidental catch in other directed fisheries. Once the directed fishery is closed, incidental catch is managed under the aggregate rockfish MRA, which limits catch of all rockfish of the genera *Sebastes* and *Sebastolobus* (which includes Pacific Ocean perch, northern rockfish, pelagic shelf rockfish, demersal shelf rockfish, and “other rockfish”) to 15 percent of directed fishing harvests.

2.1.2 Alternative 2: Remove dark rockfish from the Gulf of Alaska FMP

Under this alternative, management authority for dark rockfish is redefined by withdrawing dark rockfish from the Federal GOA groundfish FMP. Under the Magnuson-Stevens Act, State management authority may be extended into Federal waters off Alaska in the absence of Federal management of the species in question. Under this alternative, the State of Alaska could assume management authority for dark rockfish. Management plans for this species would be prepared by ADF&G staff for the Gulf of Alaska state management regions and reviewed by the Board of Fisheries.

OFLs, ABCs and TACs would continue to be specified for the PSR complex, but this complex would no longer include dark rockfish. The State would take on the responsibility for assessment and management of the dark rockfish stock.

In managing dark rockfish, the State of Alaska would develop a fishery management plan for the species under which gear type, season and guideline harvest level (GHL) for the species would be specified. The State may impose on State-registered vessels fishing in Federal fisheries only additional State measures such as bycatch retention limits for dark rockfish, as are consistent with the applicable Federal fishing regulations for the fishery in which the vessel is operating. It is not the intention of the Council or NMFS to give the State authority to indirectly regulate other Federal fisheries through State implementation of gear restrictions, area closures or other bycatch control measures. Most likely, State management of dark rockfish would include regulation of any directed fishing for dark rockfish. Dark rockfish catch in Federal fisheries would be limited by the current MRA for aggregate rockfish or a separate bycatch limit as established by the State.

While specific management plans have not yet been formulated by the State, it is likely that measures used currently (e.g., in management of black rockfish) would be among those considered for dark rockfish

management by the State (D. Carlile, pers. comm.).

These candidate measures would include, but not necessarily be limited to the following:

- Guideline harvest limits (GHLs, or quotas)
- Gear-, area- and directed-fishery-specific bycatch limits, wherein catch in excess of bycatch limits would be reported as bycatch overage on an ADF&G fish ticket, the excess bycatch would be required to be landed, with all proceeds from the sale of excess dark rockfish bycatch surrendered to the State.
- Full retention of all rockfish caught, with proceeds of the sale of any bycatch overage paid to the State of Alaska.
- Directed fisheries for dark rockfish in some areas of the State; in others perhaps bycatch only.
- No-take zones, wherein dark rockfish might not be allowed to be taken in a directed fishery and proceeds from any bycatch would be surrendered to the State.
- Gear restrictions (e.g. jig only) for directed fisheries.
- Trip limits.
- Reporting requirements such as submission of ADF&G fish tickets and/or logbooks.
- Vessel registrations for specific directed dark rockfish fishery areas.

2.2 Action 2: BSAI groundfish FMP

2.2.1 Alternative 1: Status Quo

Under this alternative, dark rockfish would continue to be managed within the other rockfish assemblage in the BSAI. The Council and the National Marine Fisheries Service would retain management authority for dark rockfish within the other rockfish complex in the EEZ. Overfishing limits (OFLs), acceptable biological catch (ABC) limits and total allowable catch (TAC) limits are established for the complex as a whole and managed accordingly. In season, catch is managed through monitoring directed fishing, with the fishery closed when directed fishing is estimated to leave only the portion of the TAC necessary to support incidental catch in other directed fisheries. Once the directed fishery is closed, incidental catch is managed under the aggregate rockfish MRA, which limits catch of all rockfish of the genera *Sebastes* and *Sebastolobus* (which includes Pacific ocean perch, northern rockfish, pelagic shelf rockfish, demersal shelf rockfish, and “other rockfish”) to 15 percent of directed fishing harvests.

2.2.2 Alternative 2: Remove dark rockfish from the BSAI FMP

Under this alternative, management authority for dark rockfish is redefined by withdrawing dark rockfish from the Federal BSAI groundfish FMP. Under the Magnuson-Stevens Act, State management authority may be extended into Federal waters off Alaska in the absence of Federal management of the species in question. Under this alternative, the State of Alaska could assume management authority for dark rockfish. Management plans for this species would be prepared by ADF&G staff for the Aleutian Island and Bering Sea state management regions and reviewed by the Board of Fisheries.

OFLs, ABCs and TACs would continue to be specified for the other rockfish complex, but this complex would no longer include dark rockfish. The State would take on the responsibility for assessment and management of the dark rockfish stock.

In managing dark rockfish, the State of Alaska would develop a fishery management plan for the species under which gear type, season and guideline harvest level (GHL) for the species would be specified.

Candidate measures to be included in any State management plan would be similar to those listed for the GOA FMP (see section 2.1.2).

2.3 Alternatives Considered but not Carried Forward

One alternative which was considered but not carried forward for analysis involves transferring management authority of dark rockfish to the State of Alaska while retaining the species under the Federal FMP. Demersal shelf rockfish in Southeast Alaska is under a similarly delegated management program to the State of Alaska. This alternative was not carried forward for dark rockfish for many reasons. A similar alternative was considered and rejected for black and blue rockfish under amendment 46 to the GOA groundfish FMP. Reasons for rejecting this for that amendment are the following: 1) State personnel would be required to comply with additional management processes; 2) the State would need to meet both state and federal requirements which are often on different time-frames for management (e.g., public meetings and reports); and 3) the State did not believe it could meet the costly assessment requirements for managing a nearshore species under a federal management plan (NPFMC 1998). Instead conservative management of the species under a state management jurisdiction only would be less costly and more conservative.

These reasons are also valid for the delegating state management of dark rockfish. Furthermore the State has indicated that it is not interested in delegated management authority for this species and would only be willing to take on management of dark rockfish if it was removed from the Federal FMP. Given this indication, this alternative was not carried forward for analysis in this document.

3.0 AFFECTED ENVIRONMENT

3.1 General distribution and habitat requirements of dark rockfish

In the GOA FMP, dark rockfish are managed as part of the shelf rockfish (PSR) assemblage. Four species comprise this assemblage: dusky rockfish (*Sebastes variabilis*), dark rockfish (*S. ciliatus*), yellowtail rockfish (*S. flavidus*), and widow rockfish (*S. entomelas*). In the Bering Sea Aleutian Islands FMP dark rockfish are contained within the “other rockfish” complex which contains the following eight species: red banded rockfish (*Sebastes babcocki*), dark rockfish, dusky rockfish, redstripe rockfish (*S. proriger*), yelloweye rockfish (*S. ruberrimus*), harlequin rockfish (*S. variegatus*), sharpchin rockfish (*S. zacentrus*), shortspine thornyhead (*Sebatolobus alascanus*).

The forms of dusky rockfish commonly recognized as “light dusky rockfish” and “dark dusky rockfish” are now officially recognized as two species (Orr and Blackburn 2004). *S. ciliatus* applies to the dark shallow-water species with a common name dark rockfish, and *S. variabilis* applies to variably colored deeper-water species with a common name dusky rockfish. Dusky rockfish are often found in large aggregations over the outer continental shelf and upper slope to depths of 675m (Orr and Blackburn, 2004). Dark rockfish are found in more shallow habitats from nearshore rocky reefs to depths no greater than 160m (Orr and Blackburn 2004).

The range of dark rockfish extends from the western Aleutian Islands and eastern Bering Sea, through the Gulf of Alaska to southeast Alaska (Orr and Blackburn 2004). Throughout its range it is common in depths ranging from 5m to 160m (Orr and Blackburn 2004). Dark rockfish are commonly collected with black rockfish (*S. melanops*) by trawl and hook-and-line gear in shallow waters and are often mis-identified as black rockfish (Orr and Blackburn 2004). In deeper trawls in the Aleutian Islands and Gulf of Alaska dark rockfish are found in association with Pacific Ocean perch (*S. alutus*), northern rockfish (*S. polyspinus*) and dusky rockfish (Blackburn and Orr 2004). Dark rockfish are occasionally found in association with other rockfishes such as harlequin rockfish, sharpchin rockfish, and redstripe rockfish (Orr and Blackburn 2004).

Habitat use changes with ontogeny. The smallest fish sampled, 10-30 cm and less than 10 years old, were collected in 1-5 m of water using herring jigs and gillnets and were found very near shore in boulder fields, commonly in harbor breakwaters. With increasing age, dark rockfish move offshore to deeper water and were captured with jig gear in 6-50 m. Video observations by ADF&G have shown that adult dark rockfish are semi-demersal, occur in rocky areas, and sometimes utilize boulder interstitial areas. Preliminary results of reproductive studies conducted in the Kodiak area by ADF&G indicate copulation in dark rockfish occurs between January and February, with fertilization in April and parturition peaking between May and June. Age and size of maturity for dark rockfish are currently under investigation (D. Urban, ADF&G, pers. comm.).

Dark and black rockfish often occur in the same locations. Of 1,133 sampling locations by ADF&G in Gulf of Alaska and Eastern Aleutian Islands from 2001 to 2006, 26% captured both dark and black rockfish (Figure 3-1). Co-occurrence was seen across the central and western Gulf of Alaska as well as the eastern Aleutian Islands (Figure 3-2). Because the sampling was done with jig gear which is subject to fishing bias, these results may not document relative population densities, but do reflect at minimum presence of the two species.




Figure 3-1 Locations where dark rockfish were captured during ADF&G surveys, 2001-2006. Survey locations were not systematically distributed but targeted known fish concentrations.




Figure 3-2 Pie charts of black (black portion) and dark (gray portion) rockfish catches in: A. Spruce Island near the city of Kodiak, B. Mountain Point on Nagai Island in the Shumagin Island group, C. the north side of Akutan Island in the eastern Aleutian Islands and D. the NE side of Unalaska is near Unalaska Bay.

The ecological separation of these two morphologically similar congeners is not well understood although underwater video reveals the darks to be more solitary and demersal while the blacks typically are a schooling fish well up in the water column (Dan Urban, ADF&G, personal observations). A food habits

study of 142 black and 84 dark rockfish was conducted by ADF&G in the Shumagin Islands. Stomachs were collected over a 10 day period in August 2005. It showed that these two species had a 29% diet overlap (Renkonen Index) with similar niche breadth (standardized Levin's measure, dark RF = 0.25, black RF 0.29). Black rockfish generally ate more fish (mostly sand lance and Pacific cod) while dark rockfish relied more on invertebrates, largely pteropods, decapod larvae, and jellyfish (Figure 3-3, ADF&G unpublished data).




Figure 3-3 Percent Index of Relative Importance (a composite index based on frequency of occurrence, numbers consumed, and weight of prey items, Cortés 1997) for dark and black rockfish from the same area of the Shumagin Islands, August 2005.

3.1.1 Life history characteristics of *Sebastes* rockfish species

Life history characteristics for all *Sebastes* species include an egg stage completed within the female and a pelagic larval stage (Lunsford et al. 2005). Larval studies for dusky rockfish (the best studied of the species in the PSR assemblage) are hampered by a lack of genetic analyses thus post-larval dusky rockfish have not been identified but are assumed to be similar to other *Sebastes* species and hence to be pelagic. Information for dark rockfish is presumed to be similar to known information for dusky rockfish. The habitat of young juveniles is unknown but a demersal stage follows the pelagic stage as evidenced by the appearance of juveniles less than 25 cm fork length in bottom trawl surveys (Clausen et al. 2002). Older juveniles have been taken only infrequently in trawl surveys and then in inshore more shallow waters than the adults (Lunsford et al. 2005). Limited food information for this species indicates that euphausiids are an important prey item for adult dusky rockfish (Yang 1993).

The size of dusky rockfish taken in the fishery generally appears to have increased after 1992; in particular, the mode increased from 42 cm in 1991-92 to 44-47 cm in 1993-97. The mode then decreased to 42 cm in 1998, and rose back to 45 cm in 1999-2002 (Lunsford et al. 2005). Age data from the fishery indicates a range of ages from 4-76 years (Lunsford et al. 2005). Age and length data from the Federal fishery data are only available for dusky rockfish.

Mortality rates and maximum age for pelagic shelf rockfish species are presented in Table 1. The estimates range from 0.06–0.09 and were based on dusky rockfish samples (Lunsford et al. 2005). A value of 0.09 has typically been used in stock assessments for pelagic shelf rockfish species because these species were typically younger than other long-lived rockfish (Lunsford et al. 2005). A value of 0.07 was recently computed for dark rockfish based upon a study completed in the GOA (Chilton. *In Review*). This study indicated a higher maximum age than had been previously assumed for dark rockfish. This value of 0.07 was utilized to compute ABCs and OFLs for dark, widow and yellowtail rockfish in the recent stock assessment for pelagic shelf rockfish (Lunsford et al. 2005).

Table 1 Instantaneous rate of natural mortality and maximum age for pelagic shelf rockfish, based on the break-and-burn method of aging otoliths. Area indicates location of study: Gulf of Alaska (GOA) or British Columbia (BC).

Species	Mortality Rate	Maximum Age	Area	Reference
Dusky Rockfish	0.09	59	GOA	1
	0.09	51 ^b	GOA	7
	0.08	59 ^c	GOA	5
	0.06	76	GOA	6
Dark Rockfish	0.07	75	GOA	2
Yellowtail Rockfish	0.07	53	BC	3
Widow Rockfish	0.05a	59	BC	4

^a Instantaneous rate of total mortality (Z).

^b Maximum survey age.

^c Maximum survey age.

References: (1) Clausen and Heifetz (1991); (2) Chilton, L. *In Review*. Growth and natural mortality of dark rockfish (*Sebastes ciliatus*) in the western Gulf of Alaska. 23rd. Lowell Wakefield Fisheries Symposium on Biology, Assessment, and Management of North Pacific Rockfishes; (3) Leaman and Nagtegaal (1987); (4) Chilton and Beamish (1982); (5) Malecha et al. (2004); (6) Calculated for this document using Hoenig (1983) $(-\ln(0.001)/t_m)$; (7) back calculated maximum age using Hoenig (1983) $(-\ln(0.001)/M)$.

Limited age and length data are available from ADF&G for dark rockfish from dockside sampling efforts from the 2002-2004 black rockfish commercial jig fishery from 1993-2006. and from black and dark rockfish surveys completed off Kodiak, Chignik, South Peninsula, and Eastern Aleutians from 2001 – 2006. Preliminary 2002 length data for dark rockfish ranged from 25–50 cm in the Kodiak region while ages ranged from 7-52 years (N. Sagalkin, unpublished data). Lengths of dark rockfish sampled range from 10 – 52 cm and 1 – 81 years old (ADF&G, unpublished data, Figure 3-4).




Figure 3-4 Age at length of male dark rockfish from Kodiak Island waters with a fitted von Bertalanffy growth curve ($\rho = 0.8363$, $k = 0.1787$, $L^\infty = 41.69$)

3.1.2 Biomass by species

3.1.2.1 GOA Pelagic Shelf rockfish complex

Dusky rockfish are the most abundant species in the pelagic shelf rockfish assemblage gulfwide. The remaining three species make up a small proportion of the assemblage. Biomass estimates from GOA trawl surveys are shown in Table 2. GOA trawl surveys were triennial until 1999 and biennial since that time. Starting in 1996 a distinction was made between “light” and “dark” dusky rockfish (and since 2005 they have been referred to by their now official names of dusky rockfish and dark rockfish). Data are presented through the most recent GOA trawl survey in 2005.

Biomass in all years is dominated by dusky rockfish. Biomass of dark, widow and yellowtail rockfish is patchy from one year to the next, with occasional single tows during the survey dominating the biomass estimate for that species. In 1999, dusky rockfish predominated, but a relatively large biomass of yellowtail rockfish was also seen in the Southeastern area. This yellowtail rockfish biomass can be mostly attributed to one relatively large catch in Dixon Entrance near the U.S./Canada boundary. In 2005, the dusky and dark rockfish biomass estimates were the highest ever recorded. The dark rockfish biomass was influenced by a large catch of 1,154 kg in the Shumagin area. The next largest catch of dark rockfish was 167 kg (Lunsford et al. 2005). With the exception of 2005 the relative contribution to the overall survey biomass from dark rockfish has been low (Table 3).

Table 2 Biomass estimates (mt) for species in the pelagic shelf rockfish assemblage in the Gulf of Alaska, based on results of bottom trawl surveys from 1984 through 2005 (Lunsford et al. 2005)

Species	Statistical Area					Total
	Shumagin	Chirikof	Kodiak	Yakutat	Southeastern	
1984						
Dusky rockfish	3,843	7,462	4,329	15,126	307	31,068
Yellowtail rockfish	0	0	0	17	454	471
Total, all species	3,843	7,462	4,329	15,143	761	31,539
1987						
Dusky rockfish	12,011	4,036	46,005	18,346	1,097	81,494
Widow rockfish	0	0	0	51	96	147
Total, all species	12,011	4,036	46,005	18,397	1,193	81,641
1990						
Dusky rockfish	2,963	1,233	16,779	5,808	953	27,735
Widow rockfish	0	0	0	285	0	285
Total, all species	2,963	1,233	16,779	6,093	953	28,020
1993						
Dusky rockfish	11,450	12,880	23,780	7,481	1,626	57,217
Total, all species	11,450	12,880	23,780	7,481	1,626	57,217
1996						
Light dusky rockfish	3,553	19,217	36,037	14,193	1,480	74,480
Dark dusky rockfish	152	139	59	0	0	350
Widow rockfish	0	10	0	0	919	929
Yellowtail rockfish	0	0	20	0	65	85
Total, all species	3,704	19,366	36,116	14,193	2,464	75,843
1999						
Light dusky rockfish	2,538	9,157	33,729	2,097	2,108	49,628
Dark dusky rockfish	2,130	31	49	0	0	2,211
Widow rockfish	0	0	69	0	115	184
Yellowtail rockfish	0	0	0	162	12,509	12,671
Total, all species	4,668	9,188	33,847	2,259	14,732	64,694
2001						
Light dusky rockfish	5,352	2,062	23,590	7,924 ^a	1,738 ^a	40,667 ^a
Dark dusky rockfish	362	15	36	0 ^a	0 ^a	413 ^a
Widow rockfish	0	0	0	0 ^a	345 ^a	345 ^a
Yellowtail rockfish	0	0	0	54 ^a	4,192 ^a	4,245 ^a
Total, all species	5,714	2,077	23,626	7,978 ^a	6,275 ^a	45,670 ^a
2003						
Light dusky rockfish	4,039	46,729	7,198	11,519	1,377	70,862
Dark dusky rockfish	235	49	16	0	0	300
Widow rockfish	0	0	0	0	32	32
Yellowtail rockfish	0	0	0	71	635	705
Total, all species	4,274	46,778	7,214	11,590	2,044	71,899
2005						
Dusky rockfish	69,295	38,216	60,097	2,488	389	170,484
Dark rockfish	21,454	389	2,348	0	0	24,191
Widow rockfish	0	0	51	0	77	128
Yellowtail rockfish	0	0	0	0	1,121	1,121
Total, all species	90,749	38,605	62,445	2,448	1,587	195,924

^aNote: The Yakutat and Southeastern areas were not sampled in the 2001 survey. Estimates of biomass for these two areas in 2001 were obtained by averaging the corresponding area biomasses in the 1993, 1996, and 1999 surveys.

Table 3 Contribution of dark rockfish survey biomass to overall PSR survey biomass estimate

Year	% Survey Biomass
2001*	0.90
2003	0.42
2005	12.35


*Note the 2001 survey did not cover the eastern GOA

Trawl survey data shows locations by species in the pelagic shelf rockfish assemblage observed in the Gulf of Alaska since 1996. Dark rockfish shows high biomass in selected tows in the Shumagin area in 1999 (Figure 3-5a) and 2005 (Figure 3-5e). Trawl survey data also shows selected high tows east and southeast of Kodiak (Figure 3-5e).

Dusky rockfish trawl survey data shows consistent high tows albeit patchily distributed from one survey to the next (Figure 3-5a-e). The 2005 survey showed the highest biomass of dusky rockfish since the survey has been conducted (Lunsford et al. 2005).

Survey Biomass data for widow and yellowtail rockfish are shown for the 1984-2005 survey years (Figure 3-7a-i and Figure 3-8a-i). Widow rockfish data showed only one high biomass tow in 1996 in the southeast leading to a biomass estimate in that area of >900 mt. Yellowtail rockfish showed higher biomass tows in southeast in 1984, 1996, and 2005 (Figure 3-8a-i). The high survey biomass estimate for yellowtail rockfish in 1999 was attributed to one relatively large catch in the Dixon entrance area (Figure 3-8f).

3-5a.



3-5b.





Figure 3-5 Dark rockfish CPUE from survey 1999-2005.

3-5c.



3-5d.




Figure 3-5 continued.






Figure 3-6 Dusky rockfish survey catch per unit effort (CPUE), survey years 1996-2003








Figure 3-6 continued.



3-7a.




3-7b.




3-7c.


Figure 3-7 Widow rockfish survey catch per unit effort (CPUE), survey years 1984-2005



3-7d.




3-7e.




3-7f.


Figure 3-7 continued.



3-7g.




3-7h.




3-7i.


Figure 3-7 continued.



3-8a.




3-8b.




3-8c.


Figure 3-8 Yellowtail rockfish survey catch per unit effort (CPUE) trawl surveys 1984-2005



3-8d.




3-8e.




3-8f.


Figure 3-8 continued.



3-8g.



3-8h.



3-8i.

Figure 3-8 continued.

Further analysis of trawl survey data for the GOA is included in order to investigate the relative prevalence of dark rockfish amongst rockfish species sampled as well as their habitat preference. Figure 3-9 shows the weight of dark rockfish found in survey hauls by bins. Large hauls of dark rockfish are extremely uncommon, with more than half of the hauls which catch dark rockfish containing less than 5 kg.




Figure 3-9 Dark rockfish catch in survey hauls by weight 1996-2005

Table 4 shows the relative weight (minimum and maximum) of dark rockfish in surveyed tows as well as the average bottom depth of the tow. Other than the single tow in 2005 with a maximum weight of 1154 kg, the maximum weight over the time period was 94 kg in 1999 (Table 4). Average bottom depth is relatively shallow and the number of hauls in which dark rockfish were identified is quite low (Table 4).

Table 4 Dark rockfish in surveyed tows 1996-2005

Year:	1996	1999	2001	2003	2005
Survey Data					
Minimum weight	0.35	1.30	1.01	0.52	0.21
Maximum weight	8.80	93.80	13.90	11.04	1153.98
Average weight	3.16	20.64	5.37	4.13	115.87
Average bottom depth	111.44	109.20	65.75	93.83	94.77
Number of hauls	9	5	4	6	13

In contrast, Table 5 shows similar survey information for dusky rockfish. Maximum weights are much higher, average bottom depth is much deeper and number of tows in which dusky rockfish are identified is much higher (Table 5). Data for black rockfish were also compiled for comparison with the depth strata for dusky and dark rockfish. Black rockfish are found in shallow waters and infrequently encountered in the bottom trawl survey due to their habitat preference (Table 6). Minimum weight, maximum weight, encounter rate in the survey and depth are all more similar to dark rockfish than to dusky.

Table 5 Dusky rockfish in surveyed tows 1996-2005

Year:	1996	1999	2001	2003	2005
Survey Data					
Minimum weight	0.15	0.14	0.09	0.09	0.32
Maximum weight	2403.55	874.00	926.31	2605.66	2239.44
Average weight	63.85	28.02	35.22	50.67	86.35
Average bottom depth	157.06	157.16	130.29	150.44	153.01
Number of hauls	109	89	70	115	140

Table 6 Black rockfish in surveyed tows 1996-2005

Year:	1996	1999	2001	2003	2005
Survey Data					
Minimum weight	0.50	1.27	0.66	1.73	0.99
Maximum weight	107.00	4.80	1.41	32.48	363.15
Average weight	36.05	2.32	1.08	14.00	89.59
Average bottom depth	54.00	46.50	55.33	97.25	82.89
Number of hauls	3	8	3	4	9

3.1.2.2 BSAI other rockfish complex

Biomass of species in the other rockfish complex is generally dominated by shortspine thornyhead rockfish and dusky rockfish. Dark rockfish are encountered infrequently in the Aleutian Island survey. Biomass total within each year as well as summary information across years for all other rockfish species are presented in tables 7-10. When encountered in the BSAI region, dark rockfish were nearly always in the AI survey. In the Bering Sea dark rockfish were rarely encountered (Table 10). Figure 3-10 shows locations by haul of dark rockfish in the Aleutian Islands region, while Table 11 shows the breakdown of biomass in the survey for dark rockfish by Aleutian Island region and depth strata. The majority of dark rockfish when encountered were found in the Western Aleutian region in the depth strata from 0-100m (Table 11, Figure 3-10). Coefficients of variation on these biomass estimates are very high given the patchy nature of surveying these species (Table 11). Dark rockfish make up a small percentage of the overall survey biomass in the Aleutian Islands in any year, ranging from 0.8 to 4.5 % since 1997 (Table 12)

Table 7 Biomass from the Aleutian Islands surveys

	1980	1983	1986	1991	1994	1997	2000	2002	2004	2006
Aleutian Islands										
dark dusky rockfish						524	99	315	320	982
harlequin rockfish	0	6	18	22	20	68	25	24	4,663	48
light dusky rockfish						712	1,306	612	2,089	6,687
redbanded rockfish	0	5	0	1	0	2	0	1	5	5
sharpchin rockfish	1	0	0	3	2	0	0	0	3	0
shortspine thornyhead	695	3,627	6,860	6,341	7,311	10,441	11,700	15,255	18,280	18,844
dusky rockfish	35	1,135	2,925	525	291					
Grand Total	730	4,774	9,803	6,891	7,624	11,747	13,130	16,208	25,359	26,567

Table 8 Biomass from the SE EBS surveys

SE EBS	YEAR									
	1980	1983	1986	1991	1994	1997	2000	2002	2004	2006
dark dusky rockfish						0	0	5	8	2
harlequin rockfish	0	0	18	0	2	0	0	24,167	7	
light dusky rockfish						138	55	971,359	731	
redbanded rockfish	0	0	0	1	0	0	0	0	0	0
sharpchin rockfish	0	0	0	3	1	0	0	0	3	0
shortspine thornyhead	23	566	423	1871	1,071	1,545	1,051	1,012	945	968
dusky rockfish	13	2362	812	58	99					
Grand Total	36	8023	2,253	2481	1,721	2,683	1,107	1,176	4,811	708

Table 9 Biomass totals (by year) AI**Aleutian Islands**

	1997-2006		1980-2006	
dark dusky rockfish	2,240	2%		
harlequin rockfish	4,828	5%	4,894	4%
light dusky rockfish	11,406	12%		
redbanded rockfish	13	0%	19	0%
sharpchin rockfish	3	0%	8	0%
shortspine thornyhead	74,521	80%	99,354	81%
dusky rockfish	0	0%	18,557	15%
Grand Total	93,011		122,831	

Table 10 Biomass totals (by year) EBS**EBS (SE portion)**

	1997-2006		1980-2006	
dark dusky rockfish	16	0%		
harlequin rockfish	4,176	35%	4,196	24%
light dusky rockfish	2,380	20%		
redbanded rockfish	0	0%	1	0%
sharpchin rockfish	3	0%	6	0%
shortspine thornyhead	5,522	46%	7,791	44%
dusky rockfish	0	0%	5,613	32%
Grand Total	12,096		17,608	




Figure 3-10 dark rockfish CPUE from the AI survey 1997-2006




Figure 3-10 continued.

Table 11 dark rockfish biomass from survey data by depth strata and region

	1997	2000	2002	2004	2006
Southern Bering Sea, 1 - 100m	0	0	5.4	8	1.3
Southern Bering Sea, 101 - 200m	0	0	0	0	0.8
Eastern Aleutian, 1 - 100m	32.4	0	0	0	0
Eastern Aleutian, 101 - 200m	0	0	0	0	8.4
Central Aleutian, 1 - 100m	0	0	0	0	72.9
Central Aleutian, 101 - 200m	9.9	0	0	2.3	0
Western Aleutian, 1 - 100m	481.6	98.6	310	308	898.4
Western Aleutian, 101 - 200m	0	0	0	1.9	0
Total	523.9	98.6	315.4	320.2	981.8
CV for total	61%	96%	57%	58%	47%




Figure 3-11 Biomass of dark rockfish in survey data by depth strata and region

Table 12 Relative contribution of dark rockfish survey biomass to the total survey biomass in the AI 1997-2006

Year	Survey biomass of darks	Total survey biomass of other rockfish complex	Percent contribution of dark rockfish to survey total
1997	524	11753	4.5
2000	99	13130	0.8
2002	315	16207	2.0
2004	320	25360	1.3
2006	982	26566	3.7

Table 13 shows the relative weight (minimum and maximum) of dark rockfish in surveyed tows as well as the average bottom depth of the tow. Similar data for dusky rockfish are presented in Table 14. Maximum weight of dark rockfish is lower than for dusky rockfish in two of the four years. Average bottom depth of haul is relatively shallow and number of hauls is relatively low in which dark rockfish were identified (Table 13).

Table 13 Dark rockfish data from surveyed tows in the AI

Year	1997	2000	2002	2004
Survey Data				
Minimum weight	0.18	0.15	1.32	0.79
Maximum weight	33.00	16.35	27.86	49.95
Average weight	6.66	5.67	11.63	7.81

Year	1997	2000	2002	2004
Average bottom depth	111.71	139.00	80.60	90.40
Number of hauls	7	3	5	10

Table 14 Dusky rockfish data from surveyed tows in the AI

Year	1997	2000	2002	2004
Survey Data				
Minimum weight	0.22	0.07	0.29	0.46
Maximum weight	15.45	121.50	27.50	161.58
Average weight	3.68	7.90	4.41	13.40
Average bottom depth	150.70	166.66	154.48	163.21
Number of hauls	20	41	29	33

Tentative biomass estimates are available from State surveys for black rockfish species but are not available for dark rockfish at this time in State waters. Hydroacoustic survey experience in State waters indicates that as dark rockfish tend toward the bottom they are likely found in the hydroacoustic dead zone and can't be easily detected via this method (Dan Urban, pers. comm.). These species may be difficult species to survey other than with submersibles or ROV transects (Dan Urban, pers. comm.).




Figure 3-12 Occurrence of dark and dusky rockfish in surveys as a percentage of tow biomass in the BS, AI and GOA in conjunction with other nearshore species.

Figure 3-12 shows the occurrence of dark and dusky rockfish in surveys in the BS, AI and GOA in conjunction with other State managed species in these areas. Dark rockfish are caught in approximately

1% of all survey tows in these areas, which is considerably less than many State managed species (Figure 3-11).

3.1.3 Stock Assessment

3.1.3.1 GOA pelagic shelf rockfish assessment

A single ABC is estimated for the pelagic shelf complex as a whole. An age-structured model is used to estimate the ABC and OFL for the dusky rockfish stock. This stock is currently in Tier 3. Under Amendment 56, Tier 3, the maximum permissible fishing mortality for ABC is $F_{40\%}$ and fishing mortality for OFL is $F_{35\%}$. These fishing rates are applied to the model estimated biomass to generate the ABC and OFL for the stock. The ABC is then apportioned over the three GOA management areas. For widow, yellowtail and dark rockfish, the average of exploitable biomass from the three most recent trawl surveys is used to determine the ABC (Tier 5). In Tier 5, F_{ABC} is defined to be $\leq 0.75 \times M$. For M of 0.07 for the three species, F_{ABC} is then $0.75 \times M$, which equals 0.0525. Multiplying this value of F by the current exploitable biomass for dark, widow, and yellowtail rockfish (10,493 mt) yields an ABC of 551 mt for 2007. The ABC is then apportioned over the GOA management areas. Table 15 provides the 2007 OFL and ABC calculated by species based on the 2006 stock assessment. There was no 2006 GOA trawl survey thus estimates for Tier 5 species (e.g. all but dusky rockfish) are based upon the 2005 stock assessment results. Changes to the ABC and OFL for the PSR complex in 2007 from the previous year's assessment are due to updated catch information included in the projection model for dusky rockfish.

Table 15 2007 OFL and ABC, calculated by species.

Species	OFL	ABC
Dusky	5,723	4,991
Dark	735 (combines all three species)	436
Widow		9
Yellowtail		106
Total PSR	6,458	5,542

The 2007 complex OFL is 6,458mt and the ABC is 5,542mt. This is apportioned over the three GOA areas as the following for 2007 WGOA=1,466mt, CGOA = 3,325mt, WYAK =307mt and EYAK/SEO = 444 mt.

3.1.3.2 BSAI other rockfish assessment

A single OFL is estimated for the other rockfish complex. ABCs are specified by individual area for the EBS and the AI. The complex is assessed at the Tier 5 level. In previous assessments Reuter and Spencer (2003; 2004) have recommended that shortspine thornyhead be split out of the other rockfish complex given that this species biomass makes up over 90% of the other rockfish biomass. The authors have also noted that the species is demographically different from other species in the complex with biomass estimates that have lower uncertainty than those of the remaining members of the complex (Reuter and Spencer, 2006). The Plan Team and SSC have continued to recommend that shortspine thornyhead remain in the complex. The Plan Team and SSC agree with the authors approach however in calculating OFL and ABC using different natural mortality rates for shortspine thornyhead ($M=0.03$) and for the remaining other rockfish biomass ($M=0.09$).

The BSAI OFL represents the sum of the individually calculated shortspine thornyhead (SST) OFL with the OFL from the remaining species in the complex (calculated as a group). The ABC is calculated

separately by area (for EBS and AI). Each area-specific ABC represents the sum of the individually calculated ABC for shortspine thornyhead together with the group ABC for the remaining species in the complex. The respective BSAI biomass estimates are calculated by adding the average biomass (1997-2006 surveys) of the AI (SST = 14,905 mt; Other rockfish = 3,698 mt) with the average EBS slope survey (2002-2004) (SST = 17,906 mt, Other rockfish 19 mt) estimate and the EBS shelf survey (Other rockfish 142 mt). BSAI OFL equals ((SST BSAI biomass (32,811) x 0.03 = 984) + (Other rockfish BSAI biomass (3,859 mt) x 0.09 = 347)) = 1,331. For calculation of the respective ABCs each of the biomass estimates were multiplied by 0.75 of M (SST 0.75 x 0.03 = 0.0225 and Other rockfish 0.75 x 0.09 = 0.0675). The resulting OFLs and ABCs for 2007 are shown below:

Other rockfish complex Tier 5 for 2007 (from Reuter and Spencer, 2006):

Region	M	Exploitable biomass (mt)	F _{ABC}	ABC (mt)	F _{OFL}	OFL (mt)
BSAI_{SST}	0.03	32,811			0.03	984
BSAI_{Orock}	0.09	3,859			0.09	347
BSAI_{Total}						1,331
EBS_{SST}	0.03	17,906	0.0225	403		
EBS_{Orock}	0.09	161	0.0675	11		
EBS_{Total}				414		
AI_{SST}	0.03	14,905	0.0225	335		
AI_{Orock}	0.09	3,698	0.0675	250		
AI_{Total}				585		

3.2 Pelagic Shelf Rockfish Fishery (GOA)

Pelagic shelf rockfish (GOA) have been caught almost exclusively with bottom trawls although some contribution from observed longline vessels has occurred. OFLs are specified gulfwide while ABCs and TACs are apportioned by area in the GOA. Overfishing levels in recent years are lower than in the period from 1998-2003 while ABCs have remained fairly constant (Table 16). Generally, in the PSR fishery in the GOA, the TAC has been established as equal to the ABC (Table 17).

Table 16 Overfishing levels (OFL), acceptable biological catch (ABC) and total allowable catch (TAC) levels for the GOA pelagic shelf rockfish complex 1998-2006

Year	OFL	ABC (total all areas)	TAC (total all areas)
1998*	9,420	4,880	4,880
1999	9,420	4,880	4,880
2000	9,040	5,980	5,980
2001	8,220	5,980	5,490
2002	8,220	5,490	5,490
2003	8,220	5,490	5,490
2004	5,570	4,470	4,470
2005	5,680	4,553	4,553
2006	6,662	5,436	5,436

*includes black and blue rockfish which were removed from the GOA FMP in 1998

The majority of the catch occurs in the Central GOA management area (Table 17).

Table 17 Commercial catch^a (mt) of fish in the pelagic shelf rockfish assemblage in the Gulf of Alaska, with Gulfwide values of acceptable biological catch (ABC) and total allowable catch (TAC), 1988-2005. Updated through October 18, 2005. (Lunsford et al. 2005)

Year	Category	Regulatory Area ^b					Gulfwide		
		Western	Central	Eastern	West Yakutat ^c	Southeast Outside ^d	Total	ABC	TAC
1988	Foreign	0	0	0	-	-	0		
	U.S.	400	517	168	-	-	1,085		
	JV	Tr	1	0	-	-	1		
	Total	400	518	168	-	-	1,086	3,300	3,300
1989	U.S.	113	888	737	-	-	1,738	6,600	3,300
1990	U.S.	165	955	527	-	-	1,647	8,200	8,200
1991	U.S.	215	1,191	936	-	-	2,342	4,800	4,800
1992	U.S.	105	2,622	887	-	-	3,605	6,886	6,886
1993	U.S.	238	2,061	894	-	-	3,193	6,740	6,740
1994	U.S.	290	1,702	997	-	-	2,989	6,890	6,890
1995	U.S.	108	2,247	536	471	64	2,891	5,190	5,190
1996	U.S.	182	1,849	265	190	75	2,296	5,190	5,190
1997	U.S.	96	1,959	574	536	38	2,629	5,140	5,140
1998	U.S.	60	2,477	576	553	22	3,113	4,880	4,880
1999	U.S.	130	3,835	694	672	22	4,659	4,880	4,880
2000	U.S.	190	3,074	467	445	22	3,731	5,980	5,980
2001	U.S.	121	2,436	451	439	12	3,008	5,980	5,980
2002	U.S.	185	2,680	457	448	9	3,322	5,490	5,490
2003	U.S.	164	2,194	617	607	10	2,975	5,490	5,490
2004	U.S.	281	2,182	211	199	12	2,885	4,470	4,470
2005	U.S.	118	1,843	218	215	3	2,397	4,553	4,553

^aCatches for 1988-97 include black rockfish and blue rockfish, which were members of the assemblage during those years.

^bCatches for West Yakutat and Southeast Outside areas are not available for years before 1996. Eastern area is comprised of the West Yakutat and Southeast Outside areas combined.

^cWest Yakutat area is comprised of statistical areas 640 and 649.

^dSoutheast Outside area is comprised of statistical areas 650 and 659.

Catches have been below TACs. Annual catches have generally increased from 1988 to 1992 and have fluctuated since that time. The pattern can largely be explained by management actions affecting rockfish during this time period. Prior to 1991 TACs for more desirable rockfish species such as Pacific ocean perch were relatively large thus the incentive to target lower valued rockfish (such as dusky rockfish in the PSR complex) was low. As TACs for slope rockfish became more restrictive in the 1990's the incentive to target other rockfish increased, resulting in higher catches for PSR species and a high in 1992 of 3605mt gulfwide. In-season management measures have largely prevented further increases in the dusky rockfish fishery. In some years (e.g., 1997-1998 and 2000-2005) the PSR trawl fishery in the Central GOA was closed prior to reaching the TAC. The fishery was closed either to ensure that catch did not exceed TAC or to prevent excessive bycatch of species such as Pacific Ocean perch and Pacific halibut (Lunsford et al. 2005).

Under the current management the Gulf of Alaska rockfish fisheries open on January 1st for non-trawl gear participants. The opening for trawl gear is near July 1st, but varies year-to-year. The trawl opening is

generally timed to coincide with the availability of the quarterly halibut PSC allocation. The fishery is also timed to accommodate the sablefish longline survey that occurs later in the summer. The rockfish fisheries, which also take some sablefish, must be completed early enough to allow the redistribution of sablefish stocks to avoid possible survey bias. The opening is also scheduled to accommodate in-season management so that managers have adequate catch and effort information to make Federal Register closure announcements, if needed, avoiding the 4th of July holiday weekend. The opening typically coincides with the openings of the Aleutian Islands Pacific ocean perch and Bering Sea flathead sole fisheries to distribute effort among the fisheries.

Both the trawl and non-trawl fisheries are prosecuted from a single TAC, with the harvest from the trawl fishery limited to the remaining available TAC after the non-trawl fleet has prosecuted the fishery from its January 1st opening. Since the non-trawl fleet has shown little interest in the fisheries historically, most of the TAC has been harvested by the trawl fleet.

Most participants target Pacific ocean perch first, until the TAC of that species is fully harvested. Pacific ocean perch are a larger biomass and typically are easier to target than the other two species. The season for Pacific ocean perch usually lasts between one and two weeks. Once the Pacific ocean perch fishery is closed, vessels will usually move on to the northern rockfish or pelagic shelf rockfish directed fisheries. The directed fisheries for northern rockfish and pelagic shelf rockfish typically last less than one month, closing before the end of July. Managers have exercised some caution in managing the fishery, occasionally closing the fisheries to ensure that the TAC is not exceeded. When sufficient TAC has remained available, managers have reopened the fisheries later to allow participants to complete the harvest.

Typically, harvests of the rockfish TACs have resulted in closure of the fisheries, although at times halibut PSC in the deep-water complex has closed the fisheries. In 2000, halibut PSC closed the pelagic shelf rockfish fishery. In 2001, halibut PSC closed both the northern rockfish and pelagic shelf rockfish fisheries in July. The fisheries were reopened on October 1st, when the fourth quarter halibut allocation became available. The fisheries closed again near the end of October, after harvest of the deep-water halibut PSC allocation.

From 1991-2005, dark rockfish have not made up more than 2.6 percent of the assemblage catch for pelagic shelf rockfish (Table 5). In most of these years dark rockfish made up only trace amounts of the catch with more than 99% of the catch made up of dusky rockfish. In 1999, dark rockfish made up 2.6% with dusky rockfish making up 97.4% of the catch. In 2004, widow rockfish made up a larger relative percentage of the total catch than in previous years with dusky rockfish making up 95.5% and dark rockfish 0.4%. In both of these years the high observed catch for dark rockfish (2.6% in 1999) and widow rockfish (4.5% in 2004) respectively were due to abnormally large individual tows recorded by observers (C. Lunsford, pers. comm.). In most years large tows of dark rockfish are not recorded by observers, indicating large catches of dark rockfish are uncommon in the trawl fishery. In 2005, the catch composition was 98.8% dusky rockfish and 1.1% dark rockfish (Table 18).

Table 18 Percentage of assemblage catch (from observer data)

Year	Dusky	Dark	Yellowtail	Widow
1991	93.5	0.2	5.1	1.2
1992	98.9	0.3	trace	0.8
1993	98.1	trace	0.5	1.4
1994	98.3	1.2	0.1	0.4
1995	99.2	trace	trace	0.8
1996	99.7	trace	trace	0.3
1997	99.9	trace	trace	0.1
1998	99.9	trace	trace	trace
1999	97.4	2.6	trace	trace
2000	99.2	0.6	0.1	0.2
2001	99.7	0.3	trace	trace
2002	99.4	0.5	trace	0.1
2003	98.8	0.8	trace	0.3
2004	95.1	0.4	trace	4.5
2005	98.8	1.1	0.2	trace

Source: C. Lunsford, NMFS

Catches for dusky rockfish are concentrated on several relatively shallow, offshore banks on the outer continental shelf particularly the “W” grounds west of Yakutat, Portlock Bank (northeast of Kodiak Island) and around Albatross Bank south of Kodiak Island (Lunsford et al. 2005). Highest CPUE in the commercial fishery is generally at depths of 100-149 m (Reuter 1999).

From 1988-1995 nearly all of the catch of dusky rockfish was taken by large factory trawlers that processed the fish at sea. Since 1999 a larger proportion of the catch has been taken by smaller shore-based trawlers in the Central GOA and the catch has been delivered to Kodiak-based processing plants. These shore-based trawlers have accounted for the following percentages of trawl catch in the CGOA from 1996-2004 (Table 19).

Table 19 Percent shore-based trawl catch in Central GOA area 1996-2004 (Lunsford et al 2005)

Year	Percent shore-based trawl catch in Central GOA area
1996	27.1
1997	18.1
1998	25.0
1999	45.2
2000	74.4
2001	58.0
2002	49.7
2003	n/a
2004	64.6

Overall catch by gear type from 1998-2005 is shown in Table 9. Some fish are not identified to species and end up in an aggregate PSR catch category. Here dusky rockfish contains both dark and dusky rockfish. Trawl catch accounts for the majority of all catch in the pelagic shelf rockfish fishery. Dark rockfish are caught by jig gear and the jig catch listed in Table 20 could be primarily dark rockfish. The highest jig catch in recent years was 53 mt in 2004. Trawl catch of dusky rockfish dominates all catch by year and gear type in this assemblage. Separate species codes are being developed to identify dusky rockfish and dark rockfish in future catch accounting given the differentiation to species level. In order to identify dark rockfish as a separate species in the Federal catch accounting system the federal reporting

requirements need revision and fairly complex data processing revisions are also required (A. Smoker, pers. comm.). New reporting requirements will be necessary whether or not the Council chooses to move dark rockfish for State management.

**Table 20 Retained catch (mt) of PSR species by gear type 1998-2005 (screened for confidentiality).
Source: NMFS Catch Accounting**

Species and year	Trawl	Fixed gear*	Jig Gear
1998			
Dusky rockfish	1,288	84	4
PSR**	1,510	0	0
Widow rockfish	18	0	0
Yellowtail rockfish	0	0	2
1999			
Dusky rockfish	2,364	19	3
PSR**	2,136	0	0
Widow rockfish	0	0	0
Yellowtail rockfish	0	1	3
2000			
Dusky rockfish	2,395	15	5
PSR**	1,092	0	0
Widow rockfish	0	0	0
Yellowtail rockfish	0	1	2
2001			
Dusky rockfish	1,932	9	9
PSR**	892	0	0
Widow rockfish	0	0	0
Yellowtail rockfish	24	0	1
2002			
Dusky rockfish	1,807	3	15
PSR**	1,195	0	0
Widow rockfish	0	0	0
Yellowtail rockfish	0	0	1
2003			
Dusky rockfish	2,946	9	8
Widow rockfish***	n/a	n/a	n/a
Yellowtail rockfish	0	0	3
2004			
Dusky rockfish	2,410	8	53
Widow rockfish	n/a	n/a	n/a
Yellowtail rockfish	0	1	1
2005			
Dusky rockfish	2,023	18	17
Widow rockfish	n/a	n/a	n/a
Yellowtail rockfish	0	n/a	1

*fixed gear includes hook and line and pot gear. Jig gear is not included as it is broken out separately.

**PSR aggregate were not identified to species

***total only available in 2003 (7mt)

Dark rockfish are also caught in the state jig fishery. Dark rockfish have often been misidentified as black rockfish and caught in the black rockfish commercial fishery (Orr and Blackburn 2004). Dark

rockfish have not been separately identified in the black rockfish fishery, although recent dockside sampling efforts by ADF&G have identified dark rockfish and other pelagic shelf rockfish species during the state jig fishery (see section 3.3 for additional information).

Major bycatch species for hauls targeting pelagic shelf rockfish include primarily northern rockfish and fish in the “other slope” rockfish category, followed by Pacific ocean perch (Ackley and Heifetz 2001). The “other slope” rockfish category includes 15 rockfish species with the primarily caught species in the category being sharpchin, redstripe, harlequin, silvergrey, yellowmouth and redbanded rockfish. Dusky rockfish was the primary bycatch species for hauls targeting northern rockfish (Ackley and Heifetz 2001). Bycatch of pelagic shelf rockfish species in the non-rockfish fisheries is presumed to be small (Lunsford et al 2005).

Discard rates of pelagic shelf rockfish have been lower than the rates for other slope rockfish species and in recent years (200-2004) have ranged from 2.4% to 4.7% (Lunsford et al 2005). Dark rockfish are included in the MRA for aggregate rockfish in the GOA. MRAs for aggregate rockfish range from 5-15% by fishery except for arrowtooth flounder which remains at 0 (Appendix 1). The Council is considering management measures to adjust the arrowtooth flounder MRA in the GOA.

3.3 BSAI other rockfish fishery

Dark rockfish are managed as part of the “other rockfish” complex in the Aleutian Islands/Eastern Bering Sea. Dusky rockfish and shortspine thornyheads are the two most abundant species in this complex. The distributions of other species in this complex including dark rockfish are not well documented (Reuter and Spencer, 2006). There is no targeted fishery for “other rockfish” in the AI or EBS. In the Aleutians, “other rockfish” are primarily caught by the atka mackerel trawl fishery (dusky rockfish) and to a lesser extent the sablefish longline fishery (shortspine thornyheads). In the Bering Sea “other rockfish are taken in small amounts by several fisheries, primarily the pacific cod trawl and longline fishery. From 1990-2001 dark rockfish comprised <1% of the “other rockfish” catch in the EBS and 3% in the AI catch (Table 20). For catch accounting purposes dark rockfish are grouped with redbanded, redstripe, yelloweye, and shapchin rockfish. In 2006 the catch of these four species was 61 mt in the AI and 6 mt in the BS (Table 21).

OFLs for the other rockfish complex are set for the entire BSAI area, while ABCs and TACs are set by area for the EBS and AI (Table 20). The TAC in the EBS has been set below ABC in recent years while the AI TAC is set equal to ABC. TACs are set to meet incidental catch needs.

Table 21 OFL, ABC and catch for the other rockfish complex in the BSAI 2004-2007

Year	Area	OFL	ABC	TAC	Catch
2004	BSAI	1,280			
	EBS		960	960	317
	AI		634	634	337
2005	BSAI	1,870			
	EBS		810	460	178
	AI		590	590	286
2006*	BSAI	1,870			
	EBS		810	460	153
	AI		590	590	417
2007	BSAI	1,330			
	EBS		414	n/a	n/a
	AI		585	n/a	n/a

*catch through 11/04/06

Historical catches of other rockfish are shown in table 21 below. Peak catch in the EBS occurred in 1978 with a catch of 941 mt while peak catch in the AI was in 1982 with a harvest of 2,114 (Reuter and Spencer, 2006).

Table 22 Summary of catches (mt) of other rockfish in the eastern Bering Sea and Aleutian Islands regions. (from Reuter and Spencer, 2006) data from NMFS/AK regional website.

Year	<u>Eastern Bering Sea</u>						<u>Aleutian Islands</u>					
	<u>Domestic</u>						<u>Domestic</u>					
	<u>For.</u>	<u>JV</u>	<u>DAP</u>	<u>Total</u>	<u>ABC</u>	<u>OFL</u>	<u>For.</u>	<u>JV</u>	<u>DAP</u>	<u>Total</u>	<u>ABC</u>	<u>OFL</u>
1977*	112	--	--	112			700	--	--	700		
1978*	941	--	--	941			212	--	--	212		
1979*	759	--	--	759			1,039	--	--	1,039		
1980	456	3	--	459			420	--	--	420		
1981	331	--	25	356			328	--	--	328		
1982	262	11	3	276			2,114	--	--	2,114		
1983	212	8	--	220			1,041	4	--	1,045		
1984	121	8	47	176			42	14	--	56		
1985	33	3	56	92			2	14	83	99		
1986	4	12	86	102			Tr	15	154	169		
1987	3	4	467	474			0	6	141	147		
1988	0	8	333	341			0	68	210	278		
1989	0	4	188	192			0	0	481	481		
1990	0	0	418	418			0	0	858	858		
1991	0	0	422	422			0	0	343	343		
1992	0	0	600	600			0	0	664	664		
1993	0	0	192	192			0	0	496	496		
1994	0	0	133	133			0	0	292	292		
1995	0	0	288	288			0	0	219	219		
1996	0	0	170	170			0	0	282	282		
1997	0	0	163	163			0	0	305	305		
1998	0	0	188	188			0	0	364	364		
1999	0	0	135	135			0	0	631	631		
2000	0	0	232	232	369	492	0	0	563	563	685	913
2001	0	0	295	295	361	482	0	0	592	592	676	901
2002	0	0	398	398	361	482	0	0	518	518	676	901
2003†	0	0	293	293	960	1,280	0	0	366	366	634	846
2004†	0	0	289	289	960	1,280	0	0	314	314	634	846
2005†	0	0	157	157	809	1,865	0	0	275	275	590	1,865
2006§	0	0	139	139	809	1,865			389	389	590	1,865

These biomass estimates were revised (2001) to show the catch of those species currently in the other rockfish category.

† Catch estimates updated 2006

§ Estimated removals through October 16th, 2006.

Historically the majority of the catch in the fishery (both EBS and AI) has been of dusky rockfish and shortspine thornyhead (Table 22) which make up the majority of the biomass in the complex as well.

Table 23 The common and scientific names of rockfish in the “other rockfish” reporting category identified, 1990- 2001, by AFSC research surveys (at least one observation) and U.S. fishery observers (greater than 1% of hauls) in the eastern Bering Sea and Aleutian Islands. *Source: Reuter and Spencer, 2006*

Common name	Scientific name	EBS		AI	
		Survey	Fishery	Survey	Fishery
Red banded rockfish	<i>Sebastes babcocki</i>	~	~	1%	<1%
Dark rockfish	<i>Sebastes ciliatus</i>	~	1%	4%	3%
Dusky rockfish	<i>Sebastes variabilis</i>	18%	39%	22%	45%
Redstripe rockfish	<i>Sebastes proriger</i>	~	1%	~	1%
Yelloweye rockfish	<i>Sebastes ruberrimus</i>	~	1%	<1%	1%
Harlequin rockfish	<i>Sebastes variegatus</i>	~	1%	9%	5%
Sharpchin rockfish	<i>Sebastes zacentrus</i>	~	<1%	<1%	<1%
Shortspine thornyhead	<i>Sebastolobus alascanus</i>	62%	43%	61%	34%

Recent catches in both the AI and EBS show a similar trend (Table 22). There is no target fishery for the other rockfish complex. Target fisheries which catch these two species are primarily the Atka mackerel trawl fishery and Pacific cod longline fishery (for dusky rockfish catch) and the longline fisheries (sablefish, turbot, halibut) as well as rockfish trawl fishery (for shortspine thornyhead catch) (Reuter and Spencer, 2006). No specific information is currently available on the catch by fishery of the dark rockfish component of the catch in the AI or EBS.

Table 24 Total fishery catch (mt) of top species in other rockfish group in the Aleutian Islands and eastern Bering Sea from 2003-2006. *Source: Reuter and Spencer, 2006. data from Catch Accounting System, NMFS AK Regional Office.*

Aleutian Islands

	2006*	541	542	543	Total
Dusky	101	48	9		158
Shortspine	35	96	15		146
Rockfish unid.	7	54	>1		61
Harlequin	4	9	10		23
Total	147	207	34		388
	2005	541	542	543	Total
Dusky	66	53	14		133
Shortspine	40	46	27		113
Rockfish unid.	1	4	9		14
Harlequin	1	8	5		14
Total	108	111	55		274
	2004	541	542	543	Total
Dusky	33	81	18		132
Shortspine	42	36	18		96
Harlequin	1	17	18		36
Rockfish unid.	>1	26	21		47

Total	76	160	75	311
2003	541	542	543	Total
Dusky	62	73	17	152
Shortspine	67	69	41	177
Harlequin	1	22	11	34
Rockfish unid.	1	1	1	3
Total	130	165	70	366

*Total catch as of October 16, 2006

Eastern Bering Sea

2006*	EBS
Shortspine thornyhead	92
Dusky	40
Rockfish unid.	6
Total	139

2005	EBS
Shortspine thornyhead	119
Dusky	36
Rockfish unid.	1.5
Total	157

2004	EBS
Shortspine thornyhead	242
Dusky	32
Rockfish unid.	15
Total	289

2003	EBS
Shortspine thornyhead	256
Dusky	23
Rockfish unid.	13
Total	293

*Total catch as of October 16, 2006

Dark rockfish are included in the MRA for aggregate rockfish in the BSAI. MRAs for aggregate rockfish range from 5-15% by fishery except for arrowtooth flounder which remains at 0 (Appendix 1).

3.4 Other Groundfish Stocks

Groundfish stocks caught in conjunction with fisheries for pelagic shelf rockfish in the GOA include Pacific ocean perch, northern rockfish and species in the "other slope" rockfish complex. In the BSAI there are no targeted fisheries for other rockfish, but these fish are commonly caught in the Atka mackerel fishery (AI) and Pacific cod longline and trawl fisheries in the BSAI. Descriptions of these species and fisheries are contained in the annual Stock Assessment and Fishery Evaluation reports for the Gulf of Alaska (NPFMC 2005).

Dark rockfish are often caught in conjunction with black rockfish. Dark rockfish and black rockfish often co-occur in nearshore kelp beds of the Gulf of Alaska, and are superficially similar in appearance, especially in body color, which can lead to misidentification. Black rockfish are a nearshore, shallow water species that are commercially targeted using jig gear. Black and blue rockfish were both removed from the Federal FMP in 1998 under amendment 46 and turned over to the State of Alaska for management due to concerns of overfishing these species under the relatively high TAC for the pelagic shelf species complex (NPFMC 1998).

Black rockfish is now solely managed by the State of Alaska following removal from the GOA groundfish FMP of black and blue rockfish under amendment 46 to the FMP (NPFMC 1998). Commercial fisheries targeting black rockfish use jig gear.

3.4.1 GOA black rockfish fishery

In the GOA, the commercial fishery for black rockfish opens in all Westward districts on January 1st and remains open until December 31, or until GHGs are attained (Mattes and Failer-Rounds 2005). Harvests are monitored through fish ticket records, processor reports and dockside sampling of commercial catches. Some black rockfish is also landed as bycatch in other fisheries (Ruccio et al. 2004). Trip limits in the Kodiak District for black rockfish are 5,000 pounds per five day harvest and landing. Vessel operators must register specifically for the black rockfish fishery in this district. No trip limits are imposed in the Chignik or South Alaska Districts of the Westward Region.

Canneries processing black rockfish in Kodiak in 2003 noted that increased sorting efforts for dusky and dark rockfish led to estimates that many deliveries that were close to 5,000 pounds total for all rockfish species often contained ¼ to ½ “dusky” rockfish (combined light and dark dusky rockfish species) once sorted (Ruccio et al. 2004). Total harvest in 2003 as reported on fish tickets for Kodiak, Chignik and South Alaska Peninsula areas for black rockfish was 141,265 pounds and for combined dusky rockfish species 17,967 pounds. The majority of the dusky rockfish harvest (17,910 of the total 17,967 pounds) was taken in the Kodiak District.

Information from ADF&G has indicated that as much as 25% of the fish reported as black rockfish caught in the Kenai Peninsula jig fishery may have actually been dark rockfish (Lunsford et al 2005).

Catch and effort data for the Kodiak District from 1990-2004 are shown in Table 25.

Table 25 Catch and effort, excluding discards, for the Kodiak Area black rockfish fishery 1998-2004 (from Sagalkin and Spalinger 2005)

Year	Vessels	Number of Landings	Directed GHG	Total Harvest (lbs)	Price per pound
1998	76	355	190,000	195,623	0.32
1999	84	316	185,000	131,986	0.40
2000	92	282	185,000	255,044	0.41
2001	55	194	185,000	220,825	0.40
2002	41	143	185,000	204,547	0.43
2003	49	106	185,000	85,362	0.36
2004	52	140	185,000	123,231	0.36

A total of 76 vessels harvested 231,555 pounds (105 mt) of black rockfish from the combined Kodiak, Chignik and Eastern District of the South Alaska Peninsula Area in the 2004 fishery (Sagalkin and Spalinger 2005). Of those participating, 31 vessels harvested black rockfish in the directed commercial

fishery with jig gear while the remainder landed it as bycatch in other fisheries (Sagalkin and Spalinger 2005). The majority of the harvest was from the Kodiak District.

Dockside sampling efforts have increased in recent years and samplers have collected a range of data in addition to fish ticket records, fishing locations and effort. Recently data has been collected during the black rockfish jig fishery on fish length, sex, reproductive maturity, and otoliths for aging (Sagalkin and Spalinger 2005). Species composition data from dockside sampling indicates that the percentage of black rockfish identified as darks is higher in recent years (Figure 3-13 and Figure 3-14).




Figure 3-13 Percent species composition landed in the 2002 Black rockfish jig fishery (Source ADF&G)




Figure 3-14 Percent species composition landed in the Black rockfish jig fishery 2003, 2005 (Source ADF&G)

Figures 5 and 6 show the percent species composition from the directed black rockfish jig fishery from dockside sampling in the Westward Region. In all areas and years the catch is predominantly black rockfish, however in 2005 a substantial proportion of the catch was dark rockfish (Figure 3-14). Generally processors offer less money for dark rockfish than for black rockfish, thus there is limited incentive for the fishermen to separate the two species (N. Sagalkin pers. comm.).

Preliminary data from the Cook Inlet management region also shows the proportion of dark rockfish in the landed black rockfish catch (Table 26). The relative proportion of dark rockfish in the catch has ranged from 0.9 to 5.6%. The lower rates of 0.9 in 2001, as compared to the following three years, may be due to the higher relative percentage of unidentified dusky rockfish in that year that were likely dark rockfish (Table 26).

Table 26 Species composition of pelagic shelf rockfish sampled in the Cook Inlet Area jig fishery and surveys 2001-2004.

Species	2001	2002	2003	2004	Ave (01-04)
Black rockfish	94.4	94.7	93.5	96.4	94.5
Unspec. Dusky rockfish	4.3	0.5	0.2	0.0	1.9
Dark rockfish	0.9	4.2	5.6	3.3	3.0
Dusky rockfish	0.4	0.7	0.8	0.3	0.5

Source: W. Dunn, ADF&G preliminary data

Dockside sampling data in the 2004 fishery for the Cook Inlet Area indicated that from a total of 672 rockfish sampled in the ports of Homer and Seward, species composition were 79% black rockfish, 7% dusky rockfish, 1% quillback rockfish and 13% yelloweye rockfish (Trowbridge and Bechtol 2004). Dusky rockfish were not separated into dusky and dark by species. Of the samples collected 87% came from the directed jig fishery.

A research survey in 2004 in the Shumagins area using a chartered jig vessel caught approximately 900 black rockfish and 434 dark rockfish, which could show an indication of the species composition in that region (D. Urban pers. comm.). The Shumagins are also the region of the high biomass estimates from tows in the trawl surveys in 1999 and 2005 (Figure 3-5).

Thus while data are still limited there are indications that a relatively high proportion of dark rockfish are caught in the commercial fisheries for black rockfish. Dusky rockfish are not caught in high amounts in the black rockfish fishery (Figure 3-13, Figure 6 and Table 12).

3.4.2 BSAI black rockfish fishery

State waters of the Aleutian Islands District and the Western District of the South Alaska Peninsula Registration Area are managed jointly for black rockfish. This area consists of all waters south of a line extending west from Cape Sarichef (54 ° 36' N. lat) and west of a line extending south of Scotch Cap Light (164 ° 44' W. long.). For management purposes this is referred to as the Aleutian Islands black rockfish fishery. In the AI, the commercial fishery for black rockfish opens in all on January 1st and remains open until December 31, or until GHLS are attained (Mattes and Failer-Rounds 2005). Harvests are monitored through fish ticket records, processor reports and dockside sampling of commercial catches.

The GHLS for black rockfish was 100,000 pounds from 1994-1998 and 90,000 pounds from 1999-2006. Harvest has been far below the GHLS in recent years.

Landings and vessel participation are listed in Table 26. Most years landing information cannot be shown due to confidentiality restrictions.

Table 27 Black rockfish landings (in pounds) in the State Aleutian Islands fishery 1997-2006

Aleutian Islands Black Rockfish			
Year	Round Pounds	Unique Vessel Count	No. of Landings
1997	102,588	5	20
1998	confidential	confidential	confidential
1999	21,522	11	44
2000	confidential	confidential	confidential
2001	confidential	confidential	confidential
2002	confidential	confidential	confidential
2003	confidential	confidential	confidential
2004	2,801	15	34
2005	6,090	9	21
2006	confidential	confidential	confidential
	confidential	confidential	confidential

Dockside sampling data are not available for the black rockfish fishery in the Aleutian Islands thus the possible percentage of landings of dark rockfish in the black rockfish fishery are unknown.

3.5 Threatened and Endangered Species

The Endangered Species Act of 1973 as amended [16 U.S.C. 1531 et seq; ESA], provides for the conservation of endangered and threatened species of fish, wildlife, and plants. The program is administered jointly by the NMFS for most marine mammal species, marine and anadromous fish species, and marine plants species and by the USFWS for bird species, and terrestrial and freshwater wildlife and plant species.

The designation of an ESA listed species is based on the biological health of that species. The status determination is either threatened or endangered. Threatened species are those likely to become endangered in the foreseeable future [16 U.S.C. § 1532(20)]. Endangered species are those in danger of becoming extinct throughout all or a significant portion of their range [16 U.S.C. § 1532(20)]. Species can be listed as endangered without first being listed as threatened. The Secretary of Commerce, acting through NMFS, is authorized to list marine fish, plants, and mammals (except for walrus and sea otter) and anadromous fish species. The Secretary of the Interior, acting through the USFWS, is authorized to list walrus and sea otter, seabirds, terrestrial plants and wildlife, and freshwater fish and plant species.

In addition to listing species under the ESA, the critical habitat of a newly listed species must be designated concurrent with its listing to the "maximum extent prudent and determinable" [16 U.S.C. § 1533(b)(1)(A)]. The ESA defines critical habitat as those specific areas that are essential to the conservation of a listed species and that may be in need of special consideration. Federal agencies are prohibited from undertaking actions that destroy or adversely modify designated critical habitat. Some species, primarily the cetaceans, which were listed in 1969 under the Endangered Species Conservation Act and carried forward as endangered under the ESA, have not received critical habitat designations.

Table 28 Species listed as endangered and threatened under the ESA that may be present in the Federal waters off Alaska include:

Common Name	Scientific name	ESA status
Northern Right Whale	<i>Balaena glacialis</i>	Endangered
Bowhead Whale ¹	<i>Balaena mysticetus</i>	Endangered
Sei Whale	<i>Balaenoptera borealis</i>	Endangered
Blue Whale	<i>Balaenoptera musculus</i>	Endangered
Fin Whale	<i>Balaenoptera physalus</i>	Endangered
Humpback Whale	<i>Megaptera novaeangliae</i>	Endangered
Sperm Whale	<i>Physeter macrocephalus</i>	Endangered
Snake River Sockeye Salmon	<i>Onchorynchus nerka</i>	Endangered
Short-tailed Albatross	<i>Phoebastria albatrus</i>	Endangered
Steller Sea Lion	<i>Eumetopias jubatus</i>	Endangered and Threatened ²
Snake River Fall Chinook Salmon	<i>Onchorynchus tshawytscha</i>	Threatened
Snake River Spring/ Summer Chinook Salmon	<i>Onchorynchus tshawytscha</i>	Threatened
Puget Sound Chinook Salmon	<i>Onchorynchus tshawytscha</i>	Threatened
Lower Columbia River Chinook Salmon	<i>Onchorynchus tshawytscha</i>	Threatened
Upper Willamette River Chinook Salmon	<i>Onchorynchus tshawytscha</i>	Threatened
Upper Columbia River Spring Chinook Salmon	<i>Onchorynchus tshawytscha</i>	Endangered
Upper Columbia River Steelhead	<i>Onchorynchus mykiss</i>	Endangered
Snake River Basin Steelhead	<i>Onchorynchus mykiss</i>	Threatened
Lower Columbia River Steelhead	<i>Onchorynchus mykiss</i>	Threatened
Upper Willamette River Steelhead	<i>Onchorynchus mykiss</i>	Threatened
Middle Columbia River Steelhead	<i>Onchorynchus mykiss</i>	Threatened

Spectacled Eider	<i>Somateria fishcheri</i>	Threatened
Steller Eider	<i>Polysticta stelleri</i>	Threatened

¹ The bowhead whale is present in the Bering Sea area only.

² Steller sea lion are listed as endangered west of Cape Suckling and threatened east of Cape Suckling.

Of the species listed under the ESA and present in the action area, some may be negatively affected by commercial groundfish fishing. Section 7 consultations with respect to the actions of the Federal groundfish fisheries have been done for all the species listed above, either individually or in groups. Additional information on endangered and threatened species appears in the Alaska Groundfish Fisheries Programmatic Supplemental Environmental Impact Statement (NMFS 2004).

3.6 Marine Mammals

Marine mammals not listed under the ESA that may be present in the GOA and BSAI include cetaceans [minke whale (*Balaenoptera acutorostrata*), killer whale (*Orcinus orca*), Dall's porpoise (*Phocoenoides dalli*), harbor porpoise (*Phocoena phocoena*), Pacific white-sided dolphin (*Lagenorhynchus obliquidens*) and the beaked whales (e.g., *Berardius bairdii* and *Mesoplodon spp.*)] and pinnipeds [northern fur seals (*Callorhinus ursinus*) and Pacific harbor seals (*Phoca vitulina*)] and the sea otter (*Enhydra lutris*).

Direct and indirect interactions between marine mammals and groundfish harvest occur due to overlap in the size and species of groundfish harvested in the fisheries that are also important marine mammal prey and due to temporal and spatial overlap in marine mammal foraging and commercial fishing activities. A detailed analysis of the effects of commercial fisheries on marine mammals appears in the Alaska Groundfish Fisheries Programmatic Supplemental Environmental Impact Statement (NMFS 2004).

3.7 Seabirds


Many seabirds occur in Alaskan waters indicating a potential for interaction with commercial fisheries. The most numerous seabirds in Alaska are northern fulmars, storm petrels, kittiwakes, murre, auklets, and puffins. These groups, and others, represent 38 species of seabirds that breed in Alaska. Eight species of Alaska seabirds breed only in Alaska and in Siberia. Populations of five other species are concentrated in Alaska but range throughout the North Pacific region. Marine waters off Alaska provide critical feeding grounds for these species as well as others that do not breed in Alaska but migrate to Alaska during summer, and for other species that breed in Canada or Eurasia and overwinter in Alaska. A detailed analysis of the effects of commercial fisheries on seabirds appears in the Alaska Groundfish Fisheries Programmatic Supplemental Environmental Impact Statement (NMFS 2004).

3.8 Habitat and Essential Fish Habitat

Section 303(a)(7) of the Magnuson-Stevens Act requires all FMPs to describe and identify Essential Fish Habitat (EFH), defined as “those waters and substrate necessary to fish for spawning, breeding, feeding or growth to maturity.” In addition, FMPs must minimize to the extent practicable adverse effects of fishing on EFH and identify other actions to conserve and enhance EFH. To this end, the Environmental Impact Statement for Essential Fish Habitat Identification and Conservation in Alaska (NMFS, 2004) provides a detailed analysis of the interactions between fisheries and EFH. Most of the controversy surrounding EFH concerns the effects of fishing activities on sea floor habitats. The analysis concludes that there are long term effects of fishing on benthic habitat features off Alaska and acknowledges that considerable scientific uncertainty remains regarding the consequences of those effects on the sustained productivity of


managed species. Based on the best available scientific information, the EIS concludes that the effects on EFH are minimal because the analysis finds no indication that continued fishing activities at the current rate and intensity would alter the capacity of EFH to support health populations of managed species over the long term. The analysis concludes that no Council-managed fishing activities have more than a minimal adverse effect on EFH, which is the regulatory standard requiring action to minimize adverse effects under the MSA. Notwithstanding these findings, the Council elected to adopt a variety of new measures to conserve EFH, which are scheduled to be implemented by August 13, 2006.

Figure 3-15 shows the concentration of observed rockfish pelagic trawl hauls from 1990 to 2002. The figure suggests that slope rockfish pelagic trawl fisheries occur at relatively low effort levels (fewer than 33 observed hauls/25 square kilometers from 1990 to 2002) in all locations in the Gulf of Alaska. The areas of greatest concentration are on the slope south of the Kenai Peninsula, with fewer areas of concentration south of Kodiak Island. Figure 3-16 shows the concentration of observed rockfish non-pelagic (bottom) trawl hauls from 1990 to 2002. The figure suggests that bottom trawl fishery for slope rockfish has taken place at relatively low effort levels all along slope areas. As with the pelagic trawl effort, concentrations of bottom trawl effort (more than 71 observed hauls/25 square kilometers from 1990 to 2002) in the Central Gulf have occurred south of Kodiak Island and south of the Kenai Peninsula. The Pacific Ocean perch fishery occurs over sand, gravel, and mud at depths of 90 to 200 fathoms. The northern rockfish and pelagic shelf rockfish fisheries occur over rock, gravel, and hard sand at depths of 40 to 80 fathoms. The analysis of the EIS provides detailed descriptions of EFH and the effects of fishing on EFH (NMFS, 2004).



Source: EFH EIS (NMFS 2004).

Figure 3-15 Observed slope rockfish pelagic trawl effort (hauls/25 square kilometers), 1990 to 2002.



Source: EFH EIS (NMFS 2004).

Figure 3-16 Observed slope rockfish non-pelagic (bottom) trawl effort (hauls/25 square kilometers), 1990 to 2002.

3.9 Ecosystem Considerations

Ecosystem considerations from the pelagic shelf rockfish fishery are summarized annually in the Gulf of Alaska Stock Assessment and Fishery Evaluation (SAFE) report (NPFMC 2005). Ecosystem considerations from the other rockfish fishery are summarized annually in the Bering Sea Aleutian Islands SAFE report (NPFMC 2006). These considerations are summarized according to the ecosystem effects on the pelagic shelf rockfish stock as well as the potential fishery effects on the ecosystem. Generally, determination of ecosystem considerations is limited by lack of biological and habitat information for rockfish.

The following tables summarize the available information on indicators of ecosystem effects for these two fisheries (Table 13 GOA PSR fishery , Table 14 BSAI other rockfish fishery).

Table 29 Analysis of ecosystem considerations for pelagic shelf rockfish and the dusky rockfish fishery.

Ecosystem effects on GOA pelagic shelf rockfish			
Indicator	Observation	Interpretation	Evaluation
<i>Prey availability or abundance trends</i>			
Phytoplankton and Zooplankton	Important for larval and post-larval survival but no information known	May help determine year class strength, no time series	Possible concern if some information available
<i>Predator population trends</i>			
Marine mammals	Not commonly eaten by marine mammals	No effect	No concern
Birds	Stable, some increasing some decreasing	Affects young-of-year mortality	Probably no concern
Fish (Halibut, arrowtooth, lingcod)	Arrowtooth have increased, others stable	More predation on juvenile rockfish	Possible concern
<i>Changes in habitat quality</i>			
Temperature regime	Higher recruitment after 1977 regime shift	Contributed to rapid stock recovery	No concern
Winter-spring environmental conditions	Affects pre-recruit survival	Different phytoplankton bloom timing	Causes natural variability, rockfish have varying larval release to compensate
Production	Relaxed downwelling in summer brings in nutrients to Gulf shelf	Some years are highly variable like El Nino 1998	Probably no concern, contributes to high variability of rockfish recruitment
GOA pelagic rockfish fishery effects on ecosystem			
Indicator	Observation	Interpretation	Evaluation

Fishery contribution to bycatch

Prohibited species	Stable, heavily monitored	Minor contribution to mortality	No concern
Forage (including herring, Atka mackerel, cod, and pollock)	Stable, heavily monitored (P. cod most common)	Bycatch levels small relative to forage biomass	No concern
HAPC biota	Medium bycatch levels of sponge and corals	Bycatch levels small relative to total HAPC biota, but can be large in specific areas	Probably no concern
Marine mammals and birds	Very minor take of marine mammals, trawlers overall cause some bird mortality	Rockfish fishery is short compared to other fisheries	No concern
Sensitive non-target species	Likely minor impact on non-target rockfish	Data limited, likely to be harvested in proportion to their abundance	Probably no concern
Fishery concentration in space and time	Duration is short and in patchy areas	Not a major prey species for marine mammals	No concern, fishery is being extended for several months starting 2006
Fishery effects on amount of large size target fish	Depends on highly variable year-class strength	Natural fluctuation	Probably no concern
Fishery contribution to discards and offal production	Decreasing	Improving, but data limited	Possible concern with non-target rockfish
Fishery effects on age-at-maturity and fecundity	Black rockfish show older fish have more viable larvae	Inshore rockfish results may not apply to longer-lived slope rockfish	Definite concern, studies being initiated in 2005

Table 14 Analysis of ecosystem considerations for other rockfish

Ecosystem effects on <i>Other Rockfish</i>			
Indicator	Observation	Interpretation	Evaluation
<i>Prey availability or abundance trends</i>			
Zooplankton	Stomach contents, ichthyoplankton surveys, changes mean wt-at-age	Data non-existent	Unknown
<i>Predator population trends</i>			
Marine mammals	Fur seals declining, Steller sea lions increasing slightly	No affect	Probably no concern
Birds	Stable, some increasing some decreasing	No affect	Probably no concern
Fish (Pollock, Pacific cod, halibut)	Stable to increasing	Affects not known	Probably no concern
<i>b. Changes in habitat quality</i>			
Temperature regime	None	Affects not known	Unknown
Winter-spring environmental conditions	None	Probably a number of factors	Unknown
Production	Fairly stable nutrient flow from upwelled BS Basin	Inter-annual variability low	No concern

4.0 ENVIRONMENTAL IMPACTS

4.1 Action 1 GOA FMP

4.1.1 Alternative 1: Status quo

4.1.2 Impacts on Pelagic Shelf Rockfish Stocks

Under alternative 1, Status Quo, there would be no change to the management of the pelagic shelf rockfish assemblage. Complex-level ABCs, OFLs and TACs would continue to be specified. As the TAC for the complex as a whole is largely based upon the biomass of dusky rockfish, the dark rockfish stock would continue to be at risk for potential overfishing under this relatively high complex-level TAC. One change that is anticipated under the status quo alternative is that catch accounting would begin to identify dark rockfish in the catch records due to the segregation of dark and light dusky by species. Catch information for dark rockfish will improve. However no management measures to restrict the harvest of dark rockfish will be taken.

4.1.3 Impacts on Other Groundfish Stocks

Under alternative 1, Status Quo, there would be no change to management of the pelagic shelf rockfish assemblage, thus there is no anticipated change in the impact of this fishery on other groundfish stocks. Bycatch in the PSR fishery includes northern rockfish and species in the “other slope” rockfish complex (see Section 3.2 for more information). The pelagic shelf rockfish fishery will continue to concentrate on dusky rockfish and relative bycatch of species is unlikely to change.

The impact on the State-managed black rockfish fishery is unlikely to change under current management of the pelagic shelf rockfish complex. Dark rockfish will likely continue to be caught in conjunction with

the black rockfish fishery. Under the current management system there is limited incentive to report dark rockfish landings as separate from black rockfish landings. With the separation of dark and dusky rockfish by species, State reporting codes will change (as with Federal) and improved information on dark rockfish information is likely.

4.1.4 Impacts on Threatened or Endangered Species

This alternative is not expected to have negative impacts on endangered or threatened species beyond those identified in previous consultations under section 7 of the Endangered Species Act. No spatial or temporal dispersion of pelagic shelf rockfish catch is anticipated.

4.1.5 Impacts on Marine Mammals

Direct and indirect interactions between marine mammals and harvests from the pelagic shelf rockfish fisheries are not expected to differ under this alternative. Total catch is expected to be the same and the distribution of catch is not expected to differ in a way that will affect interactions.

4.1.6 Impacts on Seabirds

Direct and indirect interactions between seabirds and harvests from the pelagic shelf rockfish fisheries are not expected to differ under this alternative. Total catch is expected to be the same and the distribution of catch is not expected to differ in a way that will affect interactions.

4.1.7 Impacts on Habitat and EFH

The Status Quo alternative is not expected to have any additional impacts on habitat or EFH. Effort levels for rockfish fisheries in general (of which pelagic shelf rockfish fishery is a small portion) are considered low and occur in areas of less sensitive habitat (rock, gravel, mud, and sand). The current fishing has minimal effects on benthic habitat and essential fish habitat (EFH EIS). These effects are likely to continue, if current management is maintained.

4.1.8 Impacts on the Ecosystem

Effects of fishing on the Gulf of Alaska marine ecosystem are analyzed in detail in the Alaska Groundfish Fisheries Programmatic SEIS. Additional impacts on the ecosystem from the pelagic shelf rockfish fishery are summarized annually in the SAFE report. The status quo alternative is not anticipated to have any negative impact on the Gulf of Alaska ecosystem.

4.1.9 Socioeconomic Impacts

Socioeconomic impacts of maintaining the current pelagic shelf rockfish assemblage are anticipated to be minimal. Dark rockfish make up a small percentage of overall catch in the complex. Retaining them in the pelagic shelf rockfish complex has limited economic impact. Additional information on participation in the PSR fishery, ex-vessel values in the PSR fishery and economic impacts can be found in the Regulatory Impact Review in Chapter 5 of this document.

4.2 Alternative 2

4.2.1 Impacts on Pelagic Shelf Rockfish Stocks

Alternative 2, transferring dark rockfish to State management by removing it from the Federal FMP, is anticipated to result in better management of the dark rockfish stock. Currently dark rockfish are managed under a relatively high complex-level TAC which is set primarily for dusky rockfish. If dark rockfish are removed from the pelagic shelf rockfish assemblage, the State will manage them as a single stock in State and Federal waters. The majority of the dark rockfish stock are presumed to be located in near-shore, shallow waters. The biennial trawl survey conducted by NMFS does not adequately assess this habitat and thus does not adequately assess the biomass of dark rockfish stocks.

Dark rockfish are caught infrequently in the Federal PSR fishery but more frequently in the State jig fishery. Under State management, dark rockfish would be assessed and managed as a single stock and the potential would exist to manage on smaller regions than the Federal management of the complex. There would be a decrease in the overall annual ABCs (and TACs) for the pelagic shelf rockfish complex as a result of no longer including the fractional amount of biomass contributed by the dark rockfish stock.

In recent years (with the exception of 2005) this decrease in the overall ABC (and TAC) has been less than 2% (Table 30). As discussed in Sections 3.1 and 3.2, the ABC and TAC for the complex is primarily based on the much larger biomass of dusky rockfish thus the contribution from dark rockfish is very low in most years. Widow and yellowtail rockfish would continue to be managed within the pelagic shelf rockfish complex and the relative contribution to the ABC from these stocks will continue to be incorporated into the PSR ABC.

Table 30 ABC for the pelagic shelf rockfish complex 2002-2006 and the relative contribution from the dark rockfish stock to the overall complex ABC.

Year	PSR ABC	Dark rockfish ABC (mt)	% contribution to ABC
2002	5,490	90	1.64
2003	5,490	90	1.64
2004	4,470	88	1.99
2005	4,553	88	1.93
2006	5,436	436	8.02

As discussed in section 3.1, the trawl survey biomass estimate for dark rockfish in 2005 was much higher than previous years (12% of the 2005 biomass estimate was made up of dark rockfish). Again, this was due to one abnormally large tow in the survey. The ABC is based upon a three survey average due to fluctuations in biomass from one survey to the next (Lunsford et al. 2005). Thus, even with the three survey average taken into consideration, the percent contribution to the ABC in 2006 from dark rockfish is 8%.

In all fisheries (including Federal fisheries), State managers would set an MRA (or separate bycatch limit) to limit incidental catch. Although uncertain, these MRAs would likely allow minor amounts of dark rockfish to continue to be retained in the pelagic shelf rockfish fishery. Since historic catches are approximately 1 percent or less of pelagic shelf rockfish catch, it is unlikely that the MRA would compel substantial discarding or reduce catch. The MRA, however, would prevent targeting of dark rockfish, which could occur under current rules. In other directed fisheries, discards of dark rockfish required by the MRA are likely to be minor, as catch of the species is relatively small relative to target catch.

Under this alternative the State would assume all management responsibilities for dark rockfish. This would entail assessment of the stock, management and all recordkeeping and recording requirements. Both federal and State recordkeeping requirements would be adjusted to account for dark rockfish as a species separate from dusky rockfish. Catch information for dark rockfish will be improved by these changes in catch accounting. New reporting requirements will be necessary whether or not the Council chooses to move dark rockfish for State management. The State of Alaska reporting requirements and

catch processing coding changes will also be necessary. Creation of a State Fishery Management Plan for dark rockfish will also presumably be necessary as well as the reporting requirements (logbook requirements and other dockside sampling as per black rockfish) that are necessary for directed State fisheries.

A potential exists for exploiting the State management of this stock in Federal waters under this alternative. Hypothetically, a vessel could refuse to comply with State regulations for the State dark rockfish fishery (e.g., a permit and compliance with directed fishing according to State law) and then proceed to fish the species in Federal waters. A similar situation occurred in the scallop fishery in 1995, when a Federal Scallop FMP did not exist (for more information see the 2006 Scallop SAFE report, NPFMC 2006). The fishery was eventually closed in State and Federal waters by emergency order and re-opened when a Federal FMP officially delegating authority to the State was approved. However, given the limited interest in the dark rockfish fishery, coupled with the predominance of the biomass of the nearshore species in State waters, it appears highly unlikely that such a situation would develop. Nevertheless, if a situation as described were to develop, emergency State and Federal measures would be immediately taken to protect the dark rockfish stock and ameliorate the situation.

4.2.2 Impacts on other groundfish stocks

Transferal to State management under alternative 2 is expected to have no impact on other Federally managed groundfish stocks. As discussed in Chapter 3 dark rockfish make up a very small percentage of the overall biomass and catch in the pelagic shelf rockfish complex. Dusky rockfish make up the majority of all catch (and the biomass of the complex). Impacts to the bycatch of species such as northern rockfish are expected to be the same under alternative 2 as under the current status quo alternative.

State management of dark rockfish under this alternative would enhance reporting of dark rockfish in both the directed dark rockfish fishery as well as the black rockfish fishery. This would enhance data collection on dark rockfish and black rockfish stocks and improve catch accounting for both species.

4.2.3 Impacts on threatened or endangered species

This alternative is not expected to have negative impacts on endangered or threatened species beyond those identified in previous consultations under section 7 of the Endangered Species Act. No spatial or temporal dispersion of pelagic shelf rockfish catch is anticipated.

4.2.4 Impacts marine mammals

Direct and indirect interactions between marine mammals and harvests from the pelagic shelf rockfish fisheries are not expected to differ under this alternative. Total catch is expected to be the same or slightly decreased and the distribution of catch is not expected to differ in a way that will affect interactions.

4.2.5 Impacts on seabirds

Direct and indirect interactions between seabirds and harvests from the pelagic shelf rockfish fisheries are not expected to differ under this alternative. Total catch is expected to be the same or slightly decreased and the distribution of catch is not expected to differ in a way that will affect interactions.

4.2.6 Impacts on habitat and EFH

This alternative is not expected to have any additional impacts on habitat or EFH. Effort levels for rockfish fisheries in general (of which pelagic shelf rockfish fishery is a small portion) are considered low and occur in areas of less sensitive habitat (rock, gravel, mud, and sand). The current fishing has minimal effects on benthic habitat and essential fish habitat (EFH EIS). These effects are likely to continue, if current management is maintained.

4.2.7 Impacts on the ecosystem

Effects of fishing on the Gulf of Alaska marine ecosystem are analyzed in detail in the Alaska Groundfish Fisheries Programmatic SEIS. Additional impacts on the ecosystem from the pelagic shelf rockfish fishery are summarized annually in the SAFE report. This alternative is not anticipated to have any negative impact on the Gulf of Alaska ecosystem.

4.2.8 Socio-economic impacts

Removing dark rockfish from the Federal FMP and developing State management would convey additional protection for the species from overfishing and would allow for more conservative and potentially area (and species) specific management. Removal of dark rockfish from the pelagic shelf rockfish complex could result in decreases in the pelagic shelf rockfish TAC. As discussed in section 4.2.1, the contribution to the TAC from the dark rockfish portion of the PSR assemblage is variable from one survey year to the next. It has ranged from 2-8% of the total complex ABC from 2000-2006.

Additional information on participation in the PSR fishery, ex-vessel values in the PSR fishery and economic impacts can be found in the Regulatory Impact Review in Chapter 5 of this document.

4.3 Action 2 BSAI FMP

4.3.1 Alternative 1: Status quo

4.3.2 Impacts on Other Rockfish Stocks

Under alternative 1, Status Quo, there would be no change to the management of the other rockfish assemblage. Complex-level ABCs, OFLs and TACs would continue to be specified. The TAC for the complex as a whole is largely based upon the biomass of shortspine thornyhead and dusky rockfish, with limited contribution from the dark rockfish stock. One change that is anticipated under the status quo alternative is that catch accounting would begin to identify dark rockfish in the catch records due to the segregation of dark and light dusky by species. Catch information for dark rockfish will improve. However no management measures to restrict the harvest of dark rockfish will be taken.

4.3.3 Impacts on Other Groundfish Stocks

Under alternative 1, Status Quo, there would be no change to management of the other rockfish assemblage, thus there is no anticipated change in the impact of this fishery on other groundfish stocks.

The impact on the State-managed black rockfish fishery is unlikely to change under current management of the other rockfish complex. Dark rockfish will likely continue to be caught in conjunction with the black rockfish fishery. With the separation of dark and dusky rockfish by species, State reporting codes will change (as with Federal) and improved information on dark rockfish information is likely.

4.3.4 Impacts on Threatened or Endangered Species

This alternative is not expected to have negative impacts on endangered or threatened species beyond those identified in previous consultations under section 7 of the Endangered Species Act. No spatial or temporal dispersion of pelagic shelf rockfish catch is anticipated.

4.3.5 Impacts on Marine Mammals

Direct and indirect interactions between marine mammals and harvests from the incidental catch of other rockfish in directed fisheries are not expected to differ under this alternative. Total catch is expected to be the same and the distribution of catch is not expected to differ in a way that will affect interactions.

4.3.6 Impacts on Seabirds

Direct and indirect interactions between seabirds and harvests from the incidental catch of other rockfish in directed fisheries are not expected to differ under this alternative. Total catch is expected to be the same and the distribution of catch is not expected to differ in a way that will affect interactions.

4.3.7 Impacts on Habitat and EFH

The Status Quo alternative is not expected to have any additional impacts on habitat or EFH. The current fishing has minimal effects on benthic habitat and essential fish habitat (EFH EIS). These effects are likely to continue, if current management is maintained.

4.3.8 Impacts on the Ecosystem

Effects of fishing on the Bering Sea Aleutian Islands marine ecosystem are analyzed in detail in the Alaska Groundfish Fisheries Programmatic SEIS. The status quo alternative is not anticipated to have any negative impact on the Bering Sea and Aleutian Islands ecosystem.

4.3.9 Socioeconomic Impacts

Socioeconomic impacts of maintaining the current other rockfish assemblage are anticipated to be minimal. Dark rockfish make up a small percentage of overall catch in the complex and are only incidentally caught in other directed fisheries. Retaining them in the other rockfish complex has limited economic impact. Additional information on participation in the rockfish fisheries, ex-vessel values in the rockfish fisheries in the BSAI and economic impacts can be found in the Regulatory Impact Review in Chapter 5 of this document.

4.4 Alternative 2

4.4.1 Impacts on Other Rockfish Stocks

Alternative 2, transferring dark rockfish to State management by removing it from the Federal FMP, is anticipated to result in better management of the dark rockfish stock by managing it in conjunction with black rockfish. Black rockfish are a target fishery in the State in the Aleutian Islands region. If dark rockfish are removed from the other rockfish assemblage, the State will manage them as a single stock in State and Federal waters. The majority of the dark rockfish stock are presumed to be located in near-shore, shallow waters. The trawl surveys conducted by NMFS does not adequately assess this habitat and thus does not adequately assess the biomass of dark rockfish stocks.

Dark rockfish are caught infrequently as incidental catch in other target fisheries in the BSAI. Under State management, dark rockfish would be assessed and managed as a single stock and the potential

would exist to manage on smaller regions than the Federal management of the complex. There would be a minimal decrease in the overall annual ABCs (and TACs) for the other rockfish complex as a result of no longer including the fractional amount of biomass contributed by the dark rockfish stock.

In all fisheries (including Federal fisheries), State managers would set an MRA (or separate bycatch limit) to limit incidental catch. Although uncertain, these MRAs would likely allow minor amounts of dark rockfish to continue to be retained BSAI fisheries. Since historic catches are approximately 1 percent or less of other rockfish catch in the EBS and 3% or less in the AI, it is unlikely that the MRA would compel substantial discarding or reduce catch. The MRA, however, would prevent targeting of dark rockfish, which could occur under current rules. In other directed fisheries, discards of dark rockfish required by the MRA are likely to be minor, as catch of the species is relatively small relative to target catch.

Under this alternative the State would assume all management responsibilities for dark rockfish. This would entail assessment of the stock, management and all recordkeeping and recording requirements. Both federal and State recordkeeping requirements would be adjusted to account for dark rockfish as a species separate from dusky rockfish. Catch information for dark rockfish will be improved by these changes in catch accounting. New reporting requirements will be necessary whether or not the Council chooses to move dark rockfish for State management. The State of Alaska reporting requirements and catch processing coding changes will also be necessary. Creation of a State Fishery Management Plan for dark rockfish will also presumably be necessary as well as the reporting requirements (logbook requirements and other dockside sampling as per black rockfish) that are necessary for directed State fisheries.

A potential exists for exploiting the State management of this stock in Federal waters under this alternative. Hypothetically, a vessel could refuse to comply with State regulations for the State dark rockfish fishery (e.g., a permit and compliance with directed fishing according to State law) and then proceed to fish the species in Federal waters. A similar situation occurred in the scallop fishery in 1995, when a Federal Scallop FMP did not exist (for more information see the 2006 Scallop SAFE report, NPFMC 2006). The fishery was eventually closed in State and Federal waters by emergency order and re-opened when a Federal FMP officially delegating authority to the State was approved. However, given the limited interest in the dark rockfish fishery, coupled with the predominance of the biomass of the nearshore species in State waters, it appears highly unlikely that such a situation would develop. Nevertheless, if a situation as described were to develop, emergency State and Federal measures would be immediately taken to protect the dark rockfish stock and ameliorate the situation.

4.4.2 Impacts on other groundfish stocks

Transferal to State management under alternative 2 is expected to have no impact on other Federally managed groundfish stocks. As discussed in Chapter 3 dark rockfish make up a very small percentage of the overall biomass and catch in the other rockfish complex. Shortspine thornyhead and dusky rockfish make up the majority of all catch (and the biomass of the complex).

State management of dark rockfish under this alternative would enhance reporting of dark rockfish in both the directed dark rockfish fishery as well as the black rockfish fishery. This would enhance data collection on dark rockfish and black rockfish stocks and improve catch accounting for both species.

4.4.3 Impacts on threatened or endangered species

This alternative is not expected to have negative impacts on endangered or threatened species beyond those identified in previous consultations under section 7 of the Endangered Species Act. No spatial or temporal dispersion of catch is anticipated.

4.4.4 Impacts marine mammals

Direct and indirect interactions between marine mammals and harvests from fisheries are not expected to differ under this alternative. Total catch is expected to be the same or slightly decreased and the distribution of catch is not expected to differ in a way that will affect interactions.

4.4.5 Impacts on seabirds

Direct and indirect interactions between seabirds and harvests from fisheries are not expected to differ under this alternative. Total catch is expected to be the same or slightly decreased and the distribution of catch is not expected to differ in a way that will affect interactions.

4.4.6 Impacts on habitat and EFH

This alternative is not expected to have any additional impacts on habitat or EFH. Effort levels for rockfish fisheries in general are considered low and occur in areas of less sensitive habitat (rock, gravel, mud, and sand). The current fishing has minimal effects on benthic habitat and essential fish habitat (EFH EIS). These effects are likely to continue, if current management is maintained.

4.4.7 Impacts on the ecosystem

Effects of fishing on the Bering Sea Aleutian Islands marine ecosystem are analyzed in detail in the Alaska Groundfish Fisheries Programmatic SEIS. Additional impacts on the ecosystem from the pelagic shelf rockfish fishery are summarized annually in the SAFE report. This alternative is not anticipated to have any negative impact on the Bering Sea Aleutian Islands ecosystem.

4.4.8 Socio-economic impacts

Removing dark rockfish from the Federal FMP and developing State management would convey additional protection for the species from overfishing and would allow for more conservative and potentially area (and species) specific management. Removal of dark rockfish from the other rockfish complex could result in minimal decreases in the other rockfish ABC and TAC.

Additional information on participation in the rockfish fisheries, ex-vessel values in the rockfish fisheries and economic impacts can be found in the Regulatory Impact Review in Chapter 5 of this document.

4.4.9 Cumulative Impacts

This section describes the cumulative effects of the various alternatives. Cumulative effects of an alternative are the impacts on the environment resulting from the incremental effect of the alternative when added to other past, present or reasonably foreseeable future actions.

Direct and indirect effects of this action have been discussed in previous sections of this analysis. Additional actions considered here are ones which are reasonably foreseeable and may in conjunction with the proposed action have an additional impact.

One action of this nature is the Central GOA pilot rockfish program, a five-year management program approved by the Council under Amendment 68 to the GOA groundfish FMP. This program will allocate rockfish species in the Central GOA management area in order to convey short-term economic stability to the region while comprehensive GOA groundfish rationalization initiatives are undertaken by the Council

and NMFS. The pelagic shelf rockfish assemblage is among the species to be allocated under this program. A direct allocation of PSR will be specified. If dark rockfish are removed from that assemblage, it will likely have either a separate MRA or be included under the aggregate rockfish MRA. In either case the incremental effect of implementing this program with dark rockfish excluded from the PSR allocation is expected to be minimal. The cumulative greatest effect will be realized by harvesters in the non-trawl sector who will benefit from a separate federal allocation of rockfish under the program (which will include primarily PSR and northern rockfish), while still having access to dark rockfish under State management. Since trawl vessels have little catch of dark rockfish, the cumulative effect of pilot program and the action to separate dark rockfish from the PSR assemblage will be minimal. The pilot program is anticipated to be implemented in 2008.

As with implementation of the pilot rockfish program, any incremental effect of implementing long-term comprehensive rationalization of the GOA groundfish fishery with dark rockfish removed from the PSR assemblage is likewise expected to be minimal. The specific effects of that possible action on any sector are not predictable, given the current hiatus in the development of that program.

5.0 REGULATORY IMPACT REVIEW

5.1 Introduction

This Regulatory Impact Review (RIR) examines the costs and benefits of a proposed amendment to remove dark rockfish from the Gulf of Alaska groundfish FMP.

5.2 What is a Regulatory Impact Review?

The preparation of an RIR is required under Presidential Executive Order (E.O.) 12866 (58 *FR* 51735: October 4, 1993). The requirements for all regulatory actions specified in E.O. 12866 are summarized in the following Statement from the E.O.:

In deciding whether and how to regulate, agencies should assess all costs and benefits of available regulatory alternatives, including the alternative of not regulating. Costs and Benefits shall be understood to include both quantifiable measures (to the fullest extent that these can be usefully estimated) and qualitative measures of costs and benefits that are difficult to quantify, but nonetheless essential to consider. Further, in choosing among alternative regulatory approaches agencies should select those approaches that maximize net benefits (including potential economic, environmental, public health and safety, and other advantages; distributive impacts; and equity), unless a statute requires another regulatory approach.

E.O. 12866 requires that the Office of Management and Budget (OMB) review proposed regulatory programs that are considered to be “significant.” A “significant regulatory action” is one that is likely to:

Have an annual effect on the economy of \$100 million or more or adversely affect in a material way the economy, a sector of the economy, productivity, competition, jobs, local or tribal governments or communities;

Create a serious inconsistency or otherwise interfere with an action taken or planned by another agency; Materially alter the budgetary impact of entitlements, grants, user fees, or loan programs or the rights and obligations of recipients thereof; or

Raise novel legal or policy issues arising out of legal mandates, the President’s priorities, or the principles set forth in this Executive Order.

5.3 Statutory Authority

Under the Magnuson-Stevens Act, the United States has exclusive fishery management authority over all marine fishery resources found within the exclusive economic zone (EEZ). The management of these marine resources is vested in the Secretary of Commerce (Secretary) and in the Regional Fishery Management Councils. The groundfish fisheries in the EEZ off Alaska are managed under the Fishery Management Plan (FMP) for Groundfish of the GOA.

5.4 Purpose and Need for Action

Dark rockfish are part of the pelagic shelf rockfish (PSR) assemblage in the Gulf of Alaska Groundfish Fishery Management Plan (FMP). Members of this assemblage include the following four species: dusky rockfish (*Sebastes variabilis*), dark rockfish (*S. ciliatus*), yellowtail rockfish (*S. flavidus*), and widow rockfish (*S. entomelas*). In the Bering Sea Aleutian Islands FMP dark rockfish are contained within the “other rockfish” complex which contains the following eight species: red banded rockfish (*Sebastes babcocki*), dark rockfish, dusky rockfish, redstripe rockfish (*S. proriger*), yelloweye rockfish (*S.*

ruberrimus), harlequin rockfish (*S. variegates*), sharpchin rockfish (*S. zacentrus*), shortspine thornyhead (*Sebatolobus alascanus*).

The forms of dusky rockfish commonly recognized as “light dusky rockfish” and “dark dusky rockfish” are now officially recognized as two species (Orr and Blackburn 2004). *S. ciliatus* applies to the dark shallow-water species with a common name dark rockfish, and *S. variabilis* applies to variably colored deeper-water species with a common name dusky rockfish.

Dark rockfish are found predominantly in nearshore, shallow waters. Assessment authors have suggested for years that dark rockfish be turned over the State of Alaska for management in the GOA as data in the stock assessment for PSR are predominantly from dusky rockfish (the offshore variety) not dark rockfish (the nearshore, shallow water variety). Most of the available information is from the offshore trawl surveys and offshore commercial fishery and dusky rockfish makes up the majority of the exploitable biomass and catch from the assemblage. A similar concern has been raised by the BSAI plan team for dark rockfish in the overall other rockfish assemblage.

5.5 Alternatives Considered

Two actions are analyzed in this document with two alternatives for each action: Action 1 refers to the GOA groundfish FMP. Under this action there are two alternatives: alternative 1, to continue managing dark rockfish within the larger pelagic shelf rockfish complex; and alternative 2, to remove dark rockfish from the GOA FMP and turn over to the State of Alaska for management. Action 2 refers to the BSAI groundfish FMP. Under this action there are two alternatives: alternative 1, to continue managing dark rockfish within the other rockfish complex; and alternative 2, to remove dark rockfish from the BSAI FMP and turn over to the State of Alaska for management.

5.6 Action 1: GOA groundfish FMP

5.6.1 Alternative 1: Status quo

Under this alternative, dark rockfish would continue to be managed within the pelagic shelf rockfish assemblage. The Council and the National Marine Fisheries Service would retain management authority for dark rockfish within the PSR complex in the EEZ. Overfishing limits (OFLs), acceptable biological catch (ABC) limits and total allowable catch (TAC) limits are established for the complex as a whole and managed accordingly. In season, catch is managed through monitoring directed fishing, with the fishery closed when directed fishing is estimated to leave only the portion of the TAC necessary to support incidental catch in other directed fisheries. Once the directed fishery is closed, incidental catch is managed under the aggregate rockfish MRA, which limits catch of all rockfish of the genera *Sebastes* and *Sebatolobus* (which includes Pacific ocean perch, northern rockfish, pelagic shelf rockfish, demersal shelf rockfish, and “other rockfish”) to 15 percent of directed fishing harvests.

5.6.2 Alternative 2: Remove dark rockfish from the Gulf of Alaska FMP

Under this alternative, management authority for dark rockfish is redefined by withdrawing dark rockfish from the Federal GOA groundfish FMP. Under the Magnuson-Stevens Act, State management authority may be extended into Federal waters off Alaska in the absence of Federal management of the species in question. Under this alternative, the State of Alaska could assume management authority for dark rockfish. Management plans for this species would be prepared by ADF&G staff for the Gulf of Alaska state management regions and reviewed by the Board of Fisheries.

OFLs, ABCs and TACs would continue to be specified for the PSR complex, but this complex would no longer include dark rockfish. The State would take on the responsibility for assessment and management of the dark rockfish stock.

In managing dark rockfish, the State of Alaska would develop a fishery management plan for the species under which gear type, season and guideline harvest level (GHL) for the species would be specified. The State may impose on State-registered vessels fishing in Federal fisheries only such additional State measures such as bycatch retention limits for dark rockfish, as are consistent with the applicable Federal fishing regulations for the fishery in which the vessel is operating. It is not the intention of the Council or NMFS to give the State authority to indirectly regulate other Federal fisheries through State implementation of gear restrictions, area closures or other bycatch control measures. Most likely, State management of dark rockfish would include regulation of any directed fishing for dark rockfish. Dark rockfish catch in Federal fisheries would be limited by the current MRA for aggregate rockfish or a separate bycatch limit as established by the State.

While specific management plans have not yet been formulated by the State, it is likely that measures used currently (e.g., in management of black rockfish) would be among those considered for dark rockfish management by the State (D. Carlile, pers. comm.).

These candidate measures would include, but not necessarily be limited to the following:

- Guideline harvest limits (GHLs, or quotas)
- Gear-, area- and directed-fishery-specific bycatch limits, wherein catch in excess of bycatch limits would be reported as bycatch overage on an ADF&G fish ticket, the excess bycatch would be required to be landed, with all proceeds from the sale of excess dark rockfish bycatch surrendered to the State.
- Full retention of all rockfish caught, with proceeds of the sale of any bycatch overage paid to the State of Alaska.
- Directed fisheries for dark rockfish in some areas of the State; in others perhaps bycatch only.
- No-take zones, wherein dark rockfish might not be allowed to be taken in a directed fishery and proceeds from any bycatch would be surrendered to the State.
- Gear restrictions (e.g. jig only) for directed fisheries.
- Trip limits.
- Reporting requirements such as submission of ADF&G fish tickets and/or logbooks.
- Vessel registrations for specific directed dark rockfish fishery areas.

5.7 Action 2: BSAI groundfish FMP

5.7.1 Alternative 1: Status Quo

Under this alternative, dark rockfish would continue to be managed within the other rockfish assemblage in the BSAI. The Council and the National Marine Fisheries Service would retain management authority for dark rockfish within the other rockfish complex in the EEZ. Overfishing limits (OFLs), acceptable biological catch (ABC) limits and total allowable catch (TAC) limits are established for the complex as a whole and managed accordingly. In season, catch is managed through monitoring directed fishing, with the fishery closed when directed fishing is estimated to leave only the portion of the TAC necessary to support incidental catch in other directed fisheries. Once the directed fishery is closed, incidental catch is managed under the aggregate rockfish MRA, which limits catch of all rockfish of the genera *Sebastes* and *Sebastolobus* (which includes Pacific ocean perch, northern rockfish, pelagic shelf rockfish, demersal shelf rockfish, and “other rockfish”) to 15 percent of directed fishing harvests.

5.7.2 Alternative 2: Remove dark rockfish from the BSAI FMP

Under this alternative, management authority for dark rockfish is redefined by withdrawing dark rockfish from the Federal GOA groundfish FMP. Under the Magnuson-Stevens Act, State management authority may be extended into Federal waters off Alaska in the absence of Federal management of the species in question. Under this alternative, the State of Alaska could assume management authority for dark rockfish. Management plans for this species would be prepared by ADF&G staff for the Gulf of Alaska state management regions and reviewed by the Board of Fisheries.

OFLs, ABCs and TACs would continue to be specified for the other rockfish complex, but this complex would no longer include dark rockfish. The State would take on the responsibility for assessment and management of the dark rockfish stock.

In managing dark rockfish, the State of Alaska would develop a fishery management plan for the species under which gear type, season and guideline harvest level (GHL) for the species would be specified. Candidate measures to be included in any State management plan would be similar to those listed for the GOA FMP (see section 2.1.2).

5.8 Background

The 2005 Economic SAFE report gives summary information on the ex-vessel value of the rockfish fishery as a whole (Hiatt et al. 2005). Information from this document for the GOA aggregate rockfish fishery is summarized below in Table 31 and Table 32. Note this includes all rockfish catches, of which pelagic shelf rockfish is only a small fraction. Dark rockfish, in turn, are a small portion of the pelagic shelf rockfish catch indicates that catcher vessels catch a significantly higher proportion of the catch in this fishery than catcher processors.

Table 31 Ex-vessel value of rockfish catch in the GOA by vessel category and year (\$ millions) from Hiatt et al. 2006

Gear	Year	Catcher vessel	Catcher processor	Total
Trawl	2000	2.7	2.7	5.4
	2001	1.4	2.0	3.5
	2002	2.4	3.0	5.4
	2003	3.2	2.8	6.0
	2004	3.0	3.5	6.5
	2005	3.8	5.3	9.2
Hook and Line	2000	2.2	.2	2.4
	2001	1.9	.2	2.1
	2002	2.0	.2	2.1
	2003	1.6	.2	1.8
	2004	1.7	.2	2.0
	2005	1.5	.2	1.7

Table 32 Ex-vessel value of rockfish catch in the BSAI by vessel category and year (\$ millions)

Gear	Year	Catcher vessel	Catcher processor	Total
Trawl	2000	.0	2.7	2.7
	2001	.0	2.4	2.4
	2002	.1	2.9	2.9
	2003	.0	3.6	3.6
	2004	.1	3.6	3.7
	2005	.2	4.9	5.1
Hook and Line	2000	.1	.3	.4
	2001	.2	.2	.4

	2002	.2	.2	.3
	2003	.1	.2	.3
	2004	.1	.2	.3
	2005	.1	.2	.3

Source: Hiatt et al. 2006

Data for the dusky rockfish landings by all gear types from 2003-2005 (includes both dusky rockfish and dark rockfish) indicates that catcher vessels catch a significant higher proportion of the catch in this fishery than catcher processors (Table 33). Unfortunately, the same level of data for the BSAI is not available at this time. In order to provide some indication of the fishery, aggregated rockfish data was included (Table 33 and Table 34). The data indicates that trawl gear and catcher processors are the primary participants in the BSAI rockfish fishery. Other rockfish are not a target fishery and are instead caught incidentally in other directed fisheries, notable in the longline fisheries for Pacific cod (where dusky rockfish is retained), Atka mackerel trawl fishery (retaining dusky rockfish), longline fisheries for sablefish, turbot and halibut (retaining thornyheads) and the rockfish trawl fishery (retaining thornyhead rockfish).

Table 33 Number of vessels and retained catch of pelagic shelf rockfish by vessel category in the GOA

Year	Vessel category	Number of Vessels	Retain Catch (mt)
2003	Catcher processor	17	926
	Catcher Vessel	148	1,466
2004	Catcher processor	19	985
	Catcher Vessel	134	1,381
2005	Catcher processor	18	777
	Catcher Vessel	89	1,104

Source: NPFMC, 2005

Table 34 Number of vessels by gear that caught rock fish by vessel category in the BSAI

Year	Vessel category	Trawl	Hook and Line
2003	Catcher processor	11	2
	Catcher Vessel	1	4
2004	Catcher processor	10	2
	Catcher Vessel	1	1
2005	Catcher processor	6	3
	Catcher Vessel	0	1

Source: Hiatt et al. 2006

Table 35 Total catch of rockfish by gear by vessel category in the BSAI (1,000 metric tons, round weight)

Year	Vessel category	Trawl	Hook and Line
2003	Catcher processor	20	0
	Catcher Vessel	0	0
2004	Catcher processor	17	0
	Catcher Vessel	0	0
2005	Catcher processor	14	0
	Catcher Vessel	1	0

Source: Hiatt et al. 2006

Data are not available at this time for ex-vessel value in price per pound specifically for the dusky rockfish fishery. However, data analyzed for the Central GOA pilot rockfish project (Amendment 68 to the GOA groundfish FMP) show ex-vessel price per pound for catcher vessels in the Central GOA pelagic shelf rockfish fishery (Table 36). The table shows that trawl ex-vessel prices ranged from 5 cents per pound to 7 cents per pound, while non-trawl ex-vessel prices ranged from 17 cents per pound to 26 cents per pound.

Table 36 Number of catcher vessels, landings, ex-vessel revenues and average ex vessel prices in the Central GOA pelagic shelf rockfish fishery (1998-2002)

Year	Gear	Number of vessels	Landings (mt)	Ex-vessel gross revenues (\$)	Average ex-vessel price (\$/lb)
1998	Non-Trawl	2	*	*	*
	Trawl	29	615.8	81,450	0.60
1999	Non-Trawl	2	*	*	*
	Trawl	32	1,293.2	199,577	0.070
2000	Non-Trawl	2	*	*	*
	Trawl	31	2,240.9	301,359	0.061
2001	Non-Trawl	6	4.0	2,374	0.272
	Trawl	33	1,232.6	138,534	0.051
2002	Non-Trawl	8	2.1	1,224	0.261
	Trawl	33	1,265.6	147,873	0.053

Source: NPFMC, 2005

A further look at participation by year, gear and management area gives some indication of the relative participation for each gear type in the overall pelagic shelf rockfish fishery (Table 37).

Table 37 Number of vessels operating by NMFS management area and gear types for pelagic rockfish (primarily dusky, dark, yellowtail, and widow rockfish)

Year/Sum of # of vessels	Gear	Area							GOA Total
		610	620	630	640	649	650	659	
1998	Jig		2	11	2	1	11	1	28
	Fixed	15	18	70	20	12	33	30	198
	Trawl	26	37	53	16				132
<i>1998 Total</i>		<i>41</i>	<i>57</i>	<i>134</i>	<i>38</i>	<i>13</i>	<i>44</i>	<i>31</i>	<i>358</i>
1999	Jig			10		2	13	1	26
	Fixed	27	27	60	19	16	33	38	220
	Trawl	22	39	52	20				133
<i>1999 Total</i>		<i>49</i>	<i>66</i>	<i>122</i>	<i>39</i>	<i>18</i>	<i>46</i>	<i>39</i>	<i>379</i>
2000	Jig		6	12		2	13	5	38
	Fixed	25	30	79	24	13	39	39	249
	Trawl	27	26	39	9	2			103
<i>2000 Total</i>		<i>52</i>	<i>62</i>	<i>130</i>	<i>33</i>	<i>17</i>	<i>52</i>	<i>44</i>	<i>390</i>
2001	Jig		4	13			12	5	34
	Fixed	29	21	55	11	6	36	26	184
	Trawl	27	27	38	11				103
<i>2001 Total</i>		<i>56</i>	<i>52</i>	<i>106</i>	<i>22</i>	<i>6</i>	<i>48</i>	<i>31</i>	<i>321</i>
2002	Jig	2	6	18	3		11	9	49
	Fixed	22	14	37	7		28	21	129
	Trawl	20	19	33	4				76
<i>2002 Total</i>		<i>44</i>	<i>39</i>	<i>88</i>	<i>14</i>		<i>39</i>	<i>30</i>	<i>254</i>
2003	Jig			10	0		57	3	70
	Fixed	0	0	3			35	8	46
	Trawl	9	3	37	0				49
<i>2003 Total</i>		<i>9</i>	<i>3</i>	<i>50</i>	<i>0</i>		<i>92</i>	<i>11</i>	<i>165</i>
2004	Jig		0	22			43	4	69
	Fixed			3	0		25	5	33
	Trawl	10	5	36	0				51
<i>2004 Total</i>		<i>10</i>	<i>5</i>	<i>61</i>	<i>0</i>		<i>68</i>	<i>9</i>	<i>153</i>
2005	Jig	0	0	16			27	0	43
	Fixed			0			10	7	17
	Trawl	8	6	33	0				47
<i>2005 Total</i>		<i>8</i>	<i>6</i>	<i>49</i>	<i>0</i>		<i>37</i>	<i>7</i>	<i>107</i>

Source: NMFS Catch Accounting (preliminary data)

The State fishery for black rockfish catches a higher proportion of dark rockfish than the Federal fishery for pelagic shelf rockfish (see section 3.4 for additional information on the black rockfish fishery and relative species composition of dark rockfish). The majority of the black rockfish harvest occurs in the Kodiak District. Catch and effort data for this fishery indicate that price per pound for landed black rockfish has varied from 32 cents per pound to 40 cents per pound since 1998 (Table 38).

Table 38 Catch and effort, excluding discards, for the Kodiak Area black rockfish fishery 1998-2004

Year	Vessels	Number of Landings	Directed GHL	Total Harvest (lbs)	Price per pound
1998	76	355	190,000	195,623	0.32
1999	84	316	185,000	131,986	0.40
2000	92	282	185,000	255,044	0.41
2001	55	194	185,000	220,825	0.40
2002	41	143	185,000	204,547	0.43
2003	49	106	185,000	85,362	0.36
2004	52	140	185,000	123,231	0.36

Source: Sagalkin and Spalinger 2005

No price information is available for dark rockfish, but anecdotal reports indicate that price per pound is lower than for black rockfish. Currently, fishermen have limited incentive to report catches of dark rockfish separately from black rockfish, given the lower price potential for dark rockfish.

5.9 Analysis of the Alternatives

Under the current management, dark rockfish is managed by federal managers as a part of the pelagic shelf rockfish complex. Continued management as part of the complex is likely to maintain current impacts. As described in Section 3.3, the majority of the pelagic shelf rockfish catch is dusky rockfish. Percent contribution by dark rockfish to total catch ranges from 0.4 to 1.1 percent of the total catch between 2000 and 2005, inclusive (Table 4). Catch of dark rockfish is likely to remain a relatively small portion of the pelagic shelf rockfish catch, with little affect on fishermen.

Removing dark rockfish from the Federal FMP and developing State management would convey additional protection for the species from overfishing and would allow for more conservative and potentially area (and species) specific management. Removal of dark rockfish from the pelagic shelf rockfish complex in the GOA and the other rockfish complex in the BSAI will likely result in decreases in the associated TACs. As discussed in section 4.2.1, the contribution to the TAC from the dark rockfish portion of the PSR assemblage is variable from one survey year to the next. From 2000-2005, the dark rockfish stock contributed less than 2% of the total ABC (and TAC). However, due to a large survey biomass estimate in 2005, the resulting relative contribution of dark rockfish to the PSR ABC (and TAC) rose to 8% in 2006.

Landing trends indicate a decline in trawl catch over the period considered 1998-2005 (Table 18). Removal of dark rockfish from the PSR complex in the GOA is unlikely to result in catch exceeding the revised MRA. Removal of dark rockfish from the other rockfish complex in the BSAI is also unlikely to result in catch exceeding a revised MRA for this species.

In all fisheries (including Federal fisheries), State managers would likely set an MRA (or separate bycatch limit) to limit incidental catch. Although uncertain, these MRAs would likely allow minor amounts of dark rockfish to continue to be retained in the pelagic shelf rockfish fishery. Since historic catches are approximately 1 percent or less of pelagic shelf rockfish catch, it is unlikely that the MRA would compel substantial discarding or reduce catch. The MRA, however, would prevent targeting of dark rockfish, which could occur under current rules. In other directed fisheries, discards of dark rockfish required by the MRA are likely to be negligible, as catch of the species is relatively small relative to target catch.

Depending on management choices of the State, it is possible that a directed fishery for dark rockfish could develop in State waters. Most likely such a fishery would be prosecuted with fixed gear, which

could increase the value of catch, as rockfish harvested with fixed gear have typically brought higher ex vessel prices than trawl caught rockfish. Whether such a fishery would grow to the current level of trawl catch is uncertain. Since the current directed fishery for pelagic shelf rockfish opens on January 1st and closes with the closing of the trawl fishery, which starts in the first week of July, opportunity exists for targeting dark rockfish with fixed gear under current management. In any case, due to the relatively low abundance of dark rockfish, such a fishery is likely to be relatively small. Anecdotal reports from fishermen indicate that dark rockfish can be targeted, either together with black rockfish or separately. In either case, the development of a directed fixed gear fishery for the dark rockfish could increase revenues generated from harvest of the species, but depends heavily on the growth of the fishery.

6.0 INITIAL REGULATORY FLEXIBILITY ANALYSIS

6.1 Introduction

This Initial Regulatory Flexibility Analysis (IRFA) evaluates the impacts, on small entities, of a proposed amendment to remove dark rockfish from the Gulf of Alaska groundfish FMP.

This IRFA addresses the statutory requirements of the Regulatory Flexibility Act (RFA) of 1980, as amended by the Small Business Regulatory Enforcement Fairness Act (SBREFA) of 1996 (5 U.S.C. 601-612).

6.2 The Purpose of an IRFA

The Regulatory Flexibility Act (RFA), first enacted in 1980, was designed to place the burden on the government to review all regulations to ensure that, while accomplishing their intended purposes, they do not unduly inhibit the ability of small entities to compete. The RFA recognizes that the size of a business, unit of government, or nonprofit organization frequently has a bearing on its ability to comply with a Federal regulation. Major goals of the RFA are: (1) to increase agency awareness and understanding of the impact of their regulations on small business, (2) to require that agencies communicate and explain their findings to the public, and (3) to encourage agencies to use flexibility and to provide regulatory relief to small entities. The RFA emphasizes predicting impacts on small entities as a group distinct from other entities and on the consideration of alternatives that may minimize the impacts while still achieving the stated objective of the action.

On March 29, 1996, President Clinton signed the Small Business Regulatory Enforcement Fairness Act. Among other things, the new law amended the RFA to allow judicial review of an agency's compliance with the RFA. The 1996 amendments also updated the requirements for a final regulatory flexibility analysis, including a description of the steps an agency must take to minimize the significant economic impact on small entities. Finally, the 1996 amendments expanded the authority of the Chief Counsel for Advocacy of the Small Business Administration (SBA) to file *amicus* briefs in court proceedings involving an agency's violation of the RFA.

In determining the scope, or 'universe', of the entities to be considered in an IRFA, NMFS generally includes only those entities that can reasonably be expected to be directly regulated by the proposed action. If the effects of the rule fall primarily on a distinct segment, or portion thereof, of the industry (e.g., user group, gear type, geographic area), that segment would be considered the universe for the purpose of this analysis. NMFS interprets the intent of the RFA to address negative economic impacts, not beneficial impacts, and thus such a focus exists in analyses that are designed to address RFA compliance.

Data on cost structure, affiliation, and operational procedures and strategies in the fishing sectors subject to the proposed regulatory action are insufficient, at present, to permit preparation of a "factual basis" upon which to certify that the preferred alternative does not have the potential to result in "significant adverse impacts on a substantial number of small entities" (as those terms are defined under RFA).

Because, based on all available information, it is not possible to 'certify' this outcome, should the proposed action be adopted, a formal IRFA has been prepared and is included in this package for initial review.

6.3 What is required in an IRFA?

Under 5 U.S.C., Section 603(b) of the RFA, each IRFA is required to contain:

- A description of the reasons why action by the agency is being considered;
- A succinct statement of the objectives of, and the legal basis for, the proposed rule;
- A description of and, where feasible, an estimate of the number of small entities to which the proposed rule will apply (including a profile of the industry divided into industry segments, if appropriate);
- A description of the projected reporting, record keeping and other compliance requirements of the proposed rule, including an estimate of the classes of small entities that will be subject to the requirement and the type of professional skills necessary for preparation of the report or record;
- An identification, to the extent practicable, of all relevant Federal rules that may duplicate, overlap or conflict with the proposed rule;
- A description of any significant alternatives to the proposed rule that accomplish the stated objectives of the proposed action, consistent with applicable statutes, and that would minimize any significant economic impact of the proposed rule on small entities. Consistent with the stated objectives of applicable statutes, the analysis shall discuss significant alternatives, such as:
 1. The establishment of differing compliance or reporting requirements or timetables that take into account the resources available to small entities;
 2. The clarification, consolidation, or simplification of compliance and reporting requirements under the rule for such small entities;
 3. The use of performance rather than design standards;
 4. An exemption from coverage of the rule, or any part thereof, for such small entities.

6.4 What is a small entity?

The RFA recognizes and defines three kinds of small entities: (1) small businesses, (2) small non-profit organizations, and (3) small government jurisdictions.

Small business. Section 601(3) of the RFA defines a ‘small business’ as having the same meaning as ‘small business concern’, which is defined under Section 3 of the Small Business Act. ‘Small business’ or ‘small business concern’ includes any firm that is independently owned and operated and not dominant in its field of operation. The SBA has further defined a “small business concern” as one “organized for profit, with a place of business located in the United States, and which operates primarily within the United States or which makes a significant contribution to the U.S. economy through payment of taxes or use of American products, materials or labor... A small business concern may be in the legal form of an individual proprietorship, partnership, limited liability company, corporation, joint venture, association, trust or cooperative, except that where the firm is a joint venture there can be no more than 49 percent participation by foreign business entities in the joint venture.”

The SBA has established size criteria for all major industry sectors in the United States, including fish harvesting and fish processing businesses. A business involved in fish harvesting is a small business if it is independently owned and operated and not dominant in its field of operation (including its affiliates) and if it has combined annual receipts not in excess of \$3.5 million for all its affiliated operations worldwide. A seafood processor is a small business if it is independently owned and operated, not dominant in its field of operation, and employs 500 or fewer persons on a full-time, part-time, temporary, or other basis, at all its affiliated operations worldwide. A business involved in both the harvesting and

processing of seafood products is a small business if it meets the \$3.5 million criterion for fish harvesting operations. Finally, a wholesale business servicing the fishing industry is a small business if it employs 100 or fewer persons on a full-time, part-time, temporary, or other basis, at all its affiliated operations worldwide.

The SBA has established “principles of affiliation” to determine whether a business concern is “independently owned and operated.” In general, business concerns are affiliates of each other when one concern controls or has the power to control the other, or a third party controls or has the power to control both. The SBA considers factors such as ownership, management, previous relationships with or ties to another concern, and contractual relationships, in determining whether affiliation exists. Individuals or firms that have identical or substantially identical business or economic interests, such as family members, persons with common investments, or firms that are economically dependent through contractual or other relationships, are treated as one party with such interests aggregated when measuring the size of the concern in question. The SBA counts the receipts or employees of the concern whose size is at issue and those of all its domestic and foreign affiliates, regardless of whether the affiliates are organized for profit, in determining the concern’s size. However, business concerns owned and controlled by Indian Tribes, Alaska Regional or Village Corporations organized pursuant to the Alaska Native Claims Settlement Act (43 U.S.C. 1601), Native Hawaiian Organizations, or Community Development Corporations authorized by 42 U.S.C. 9805 are not considered affiliates of such entities, or with other concerns owned by these entities solely because of their common ownership.

Affiliation may be based on stock ownership when, (1) a person is an affiliate of a concern if the person owns or controls, or has the power to control 50 percent or more of its voting stock, or a block of stock which affords control because it is large compared to other outstanding blocks of stock, or (2) if two or more persons each owns, controls or has the power to control less than 50 percent of the voting stock of a concern, with minority holdings that are equal or approximately equal in size, but the aggregate of these minority holdings is large as compared with any other stock holding, each such person is presumed to be an affiliate of the concern.

Affiliation may be based on common management or joint venture arrangements. Affiliation arises where one or more officers, directors, or general partners, controls the board of directors and/or the management of another concern. Parties to a joint venture also may be affiliates. A contractor and subcontractor are treated as joint venturers if the ostensible subcontractor will perform primary and vital requirements of a contract or if the prime contractor is unusually reliant upon the ostensible subcontractor. All requirements of the contract are considered in reviewing such relationship, including contract management, technical responsibilities, and the percentage of subcontracted work.

Small organizations. The RFA defines “small organizations” as any not-for-profit enterprise that is independently owned and operated and is not dominant in its field.

Small governmental jurisdictions. The RFA defines small governmental jurisdictions as governments of cities, counties, towns, townships, villages, school districts, or special districts with populations of fewer than 50,000.

6.5 Reason for considering the action

As discussed in the EA and RIR sections of this analysis, the Council is considering this action in order to provide better protection for the small inshore stock of dark rockfish. This stock is currently contained in the pelagic shelf rockfish assemblage which is dominated by the biomass of the offshore dusky rockfish. Detailed descriptions of each alternative analyzed in this EA/RIR/IRFA can be found in Section 2.0.

6.6 Objectives of, and legal basis for, the proposed action

Under the Magnuson-Stevens Act, the United States has exclusive fishery management authority over all marine fishery resources found within the exclusive economic zone (EEZ). The management of these marine resources is vested in the Secretary of Commerce (Secretary) and in the Regional Fishery Management Councils. The groundfish fisheries in the EEZ off Alaska are managed under the Fishery Management Plan (FMP) for Groundfish of the GOA.

Statutory authority for measures designed to prevent overfishing is specifically addressed in Sec. 600.350 of the Magnuson-Stevens Act. That section establishes National Standard 1—Prevent Overfishing.

The objective of the proposed action is to prevent overfishing of the dark rockfish stock.

6.7 Number and description of small entities regulated by the proposed action

6.8 Impacts on Regulated Small Entities

Transfer of management of dark rockfish to the State is likely to result in some changes in regulation of catch. The State could develop a directed fishery for dark rockfish, most likely for fixed gear vessels. Since fixed gear vessels tend to be small, it is possible that the development of such a directed fishery would have a positive impact on small entities, by increasing fishing opportunities. The increase over current opportunities is likely to be relatively minor, as fixed gear vessels currently have little catch despite an extended season.¹

Small entities that own trawl vessels are unlikely to realize any noticeable adverse effects from this action. Although trawl vessels would no longer be permitted to directed fish for dark rockfish (as a part of the pelagic shelf rockfish complex) in federal waters, since dark rockfish make up a very small part of the pelagic shelf rockfish catch, it is unlikely that any vessels would be required to discard dark rockfish catch in that fishery because of the MRA. A decline in the pelagic shelf rockfish TAC could occur, but that decline is likely to be small since the dark rockfish stock is a very small part of the combined stock of pelagic shelf rockfish. Since some of the decline in TAC is likely to be offset by incidental catch under the MRA, it is unlikely that catches will be noticeably affected.

6.9 Recordkeeping and Reporting Requirements

No additional recordkeeping will be required by the change in management proposed by this action.

6.10 Federal rules that may duplicate, overlap, or conflict with proposed action

This analysis did not reveal any federal rules that duplicate, overlap or conflict with the proposed action.

7.0 CONSISTENCY WITH APPLICABLE LAW AND POLICY

7.1 Magnuson-Stevens Act

7.1.1 National Standards

¹ Current regulations allow fixed gear vessels to begin fishing for pelagic shelf rockfish (including dark rockfish) in January, with the fixed gear fishery typically closing in mid-July with the harvest of the TAC by trawl vessels, which begin fishing in early July.

The Council's over-arching mandate to guide it in the prevention of overfishing is National Standard 1. This national standard states that:

Conservation and management measures shall prevent overfishing while achieving, on a continuing basis, the optimum yield from each fishery.

This action is specifically being considered in order to limit the current potential for overfishing of the dark rockfish stock. Under the current pelagic shelf rockfish assemblage, the dark rockfish stock is vulnerable to overfishing given the relatively high complex-level TAC that could be taken on any member of the assemblage. Dark rockfish as discussed in the analysis makes up a small fraction of the biomass in the assemblage and is generally found in shallow, in-shore waters. Transferal of management to the State of Alaska is anticipated to be better responsive to protection of this stock.

7.1.2 Section 303(a)(9) – Fisheries Impact Statement

Section 303(a)(9) of the Magnuson-Stevens Act requires that any plan or amendment include a fishery impact statement which shall assess and describe the likely effects, if any, of the conservation and management measures on a) participants in the fisheries and fishing communities affected by the plan or amendment; and b) participants in the fisheries conducted in adjacent areas under the authority of another Council, after consultation with such Council and representatives of those participants taking into account potential impacts on the participants in the fisheries, as well as participants in adjacent fisheries.

The alternative actions considered in this analysis are described in Chapter 2 of this document. The impacts of these actions on participants in the fisheries and fishing communities are evaluated in the Regulatory Impact Review, Chapter 5.

7.2 Marine Mammal Protection Act

The alternatives analyzed in this action are not likely to result in any significant impacts to marine mammals.

7.3 Coastal Zone Management Act

This action is consistent with the Coastal Zone Management Act.

8.0 CONSULTATION AND PREPARERS

8.1 List of Persons and Agencies Consulted

NOAA Fisheries (RO): Mary Furuness
Tom Pearson
Andy Smoker
Lew Queirolo

NOAA Fisheries (AFSC): Dave Clausen
Dana Hanselman
James Orr

Alaska Department of Fish and Game:
David Carlile
Gail Smith

8.2 List of Preparers

North Pacific Fishery
Management Council: Diana Stram
Mark Fina
Jon McCracken
Peggy Kircher

NOAA Fisheries (AFSC): Jim Ianelli
Chris Lunsford
Rebecca Reuter
Kalei Shotwell

Alaska Department of Fish and Game:
Nick Sagalkin
Dan Urban
Carrie Worton
Krista Milani

9.0 REFERENCES

Ackley, D. R., and J. Heifetz. 2001. Fishing practices under maximum retainable bycatch rates in Alaska=s groundfish fisheries. *Alaska Fish. Res. Bull.* 8(1): 22-44.

Chilton, L. *In Review*. Growth and natural mortality of dark rockfish (*Sebastes ciliatus*) in the western Gulf of Alaska. 23rd Lowell Wakefield Fisheries Symposium on Biology, Assessment and Management of North Pacific Rockfishes.

Clausen, D. M., and J. Heifetz. 1991. Pelagic shelf rockfish. In Stock assessment and fishery evaluation report for the 1992 Gulf of Alaska groundfish fishery, p. 7-1 - 7-12. North Pacific Fishery Management Council, 605 W. 4th. Avenue, Suite 306, Anchorage, AK 99501-2252.

- Clausen, D. M., and J. Heifetz. 1996. Pelagic shelf rockfish. *In* Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska, p. 271-288. North Pacific Fishery Management Council, 605 W. 4th. Avenue, Suite 306, Anchorage, AK 99501-2252.
- Clausen, D. M., C. R. Lunsford, and J. T. Fujioka. 2002. Pelagic shelf rockfish. *In* Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska, p. 383-418. North Pacific Fishery Management Council, 605 W. 4th. Avenue, Suite 306, Anchorage, AK 99501-2252.
- Cortés, E. 1997. A critical review of methods of studying fish feeding based on analysis of stomach contents: application to elasmobranch fishes. *Can. J. Fish. Aquat. Sci.* 54:726-738.
- Hoening, J. 1983. Empirical use of longevity data to estimate mortality rates. *Fish. Bull.* 82:898-903.
- Leaman, B.M. and D.A. Nagtegaal. 1987. Age validation and revised natural mortality rate for yellowtail rockfish. *Trans. Am. Fish. Soc.* 116:171-175.
- Lunsford, C.R., S.K. Shotwell, D.H. Hanselman, D.M. Clausen, and D.L. Courtney. 2005. Gulf of Alaska Pelagic Shelf Rockfish. *In*, Stock Assessment and Fishery Evaluation Report for the groundfish fisheries of the Gulf of Alaska. P 727 – 780. Compiled by the GOA Groundfish Plan Team. North Pacific Fishery Management Council, 605 West 4th Ave., Anchorage, AK, 99501.
- Malecha, P.W., and J. Heifetz. 2004. Growth and mortality of rockfish (Scorpaenidae) from Alaska waters. In Review, 39 p. Available from the Auke Bay Laboratory, NMFS, NOAA, 11305 Glacier Hwy, Juneau, AK 99801.
- National Marine Fisheries Service. 2004. Alaska groundfish fisheries final programmatic supplemental environmental impact statement. Available from Alaska Region, National Marine Fisheries Service, Box 21668, Juneau, AK 99802.
- National Marine Fisheries Service. 2005. Final Environmental Impact Statement for Essential Fish Habitat Identification and Conservation in Alaska. Available from Alaska Region, National Marine Fisheries Service, Box 21668, Juneau, AK 99802.
- North Pacific Fishery Management Council. 1998. EA/RIR/IRFA for Amendment 46 to the Fishery Management Plan for the Groundfish Fishery of the Gulf of Alaska for Revise Management Authority of Pelagic Shelf Rockfish. North Pacific Fishery Management Council, 604 West 4th Ave., Anchorage, AK, 99501.
- North Pacific Fishery Management Council. 2005. EA/RIR/IRFA for Amendment 68 to the Fishery Management Plan for the Groundfish Fishery of the Gulf of Alaska for Central Gulf of Alaska Rockfish Demonstration Program. North Pacific Fishery Management Council, 604 West 4th Ave., Anchorage, AK, 99501
- Orr, J. W., and J. E. Blackburn. 2004. The dusky rockfishes (Teleostei: Scorpaeniformes) of the North Pacific Ocean: resurrection of *Sebastes variabilis* (Pallas, 1814) and a redescription of *Sebastes ciliatus* (Tilesius, 1813). *Fish. Bull.*, U.S. 102:328-348. Online. (.pdf, 569KB).
- Reuter, R. F. 1999. Describing dusky rockfish (*Sebastes ciliatus*) habitat in the Gulf of Alaska using historical data. M.S. Thesis, California State University, Hayward CA. 83 p.

Seeb, L. W. 1986. Biochemical systematics and evolution of the Scorpaenid genus *Sebastes*. Ph.D. Thesis, Univ. Washington, Seattle, WA. 177 p.

Yang, M-S. 1993. Food habits of the commercially important groundfishes in the Gulf of Alaska in 1990. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-22. 150 p.

APPENDIX 1. MRA TABLES

Table 10 to Part 679--Gulf of Alaska Retainable Percentages

BASIS SPECIES		INCIDENTAL CATCH SPECIES (for DSR caught on catcher vessels in the SEO, see § 679.20 (j) ⁶)														
Code	Species	Pollock	Pacific cod	DW flat ⁽²⁾	Rex sole	Flathead sole	SW Flat ⁽³⁾	Arrowtooth	Sablefish	Aggregated rockfish ⁽⁸⁾	SR/RE ERA ⁽¹⁾	DSR SEO (C/Ps only) ⁽⁶⁾	Atka mackerel	Aggregated forage fish ⁽¹⁰⁾	Skates ⁽¹¹⁾	Other species ⁽⁷⁾
110	Pacific cod	20	na ⁹	20	20	20	20	35	1	5	⁽¹⁾	10	20	2	20	20
121	Arrowtooth	5	5	0	0	0	0	na ⁹	0	0	0	0	0	2	0	20
122	Flathead sole	20	20	20	20	na ⁹	20	35	7	15	7	1	20	2	20	20
125	Rex sole	20	20	20	na ⁹	20	20	35	7	15	7	1	20	2	20	20
136	Northern rockfish	20	20	20	20	20	20	35	7	15	7	1	20	2	20	20
141	Pacific ocean perch	20	20	20	20	20	20	35	7	15	7	1	20	2	20	20
143	Thornyhead	20	20	20	20	20	20	35	7	15	7	1	20	2	20	20
152/ 151	Shortraker/ rougheye ⁽¹⁾	20	20	20	20	20	20	35	7	15	na ⁹	1	20	2	20	20
193	Atka mackerel	20	20	20	20	20	20	35	1	5	⁽¹⁾	10	na ⁹	2	20	20
270	Pollock	na ⁹	20	20	20	20	20	35	1	5	⁽¹⁾	10	20	2	20	20
710	Sablefish	20	20	20	20	20	20	35	na ⁹	15	7	1	20	2	20	20
Flatfish, deep water ⁽²⁾		20	20	na ⁹	20	20	20	35	7	15	7	1	20	2	20	20
Flatfish, shallow water ⁽³⁾		20	20	20	20	20	na ⁹	35	1	5	⁽¹⁾	10	20	2	20	20
Rockfish, other ⁽⁴⁾		20	20	20	20	20	20	35	7	15	7	1	20	2	20	20
Rockfish, pelagic ⁽⁵⁾		20	20	20	20	20	20	35	7	15	7	1	20	2	20	20
Rockfish, DSR-SEO ⁽⁶⁾		20	20	20	20	20	20	35	7	15	7	na ⁹	20	2	20	20
Skates ⁽¹¹⁾		20	20	20	20	20	20	35	1	5	⁽¹⁾	10	20	2	na ⁹	20
Other species ⁽⁷⁾		20	20	20	20	20	20	35	1	5	⁽¹⁾	10	20	2	20	na ⁹
Aggregated amount of non-groundfish species		20	20	20	20	20	20	35	1	5	⁽¹⁾	10	20	2	20	20

Notes to Table 10 to Part 679					
1	Shortraker/rougheye rockfish				
		SR/RE	shortraker/rougheye rockfish (171)		
			shortraker rockfish (152)		
			rougheye rockfish (151)		
	SR/RE ERA	shortraker/rougheye rockfish in the Eastern Regulatory Area.			
Where numerical percentage is not indicated, the retainable percentage of SR/RE is included under Aggregated Rockfish					
2	Deep-water flatfish	Dover sole, Greenland turbot, and deep-sea sole			
3	Shallow water flatfish	Flatfish not including deep water flatfish, flathead sole, rex sole, or arrowtooth flounder			
4	Other rockfish	Western Regulatory Area	means slope rockfish and demersal shelf rockfish		
		Central Regulatory Area			
		West Yakutat District			
		Southeast Outside District			means slope rockfish
	Slope rockfish				
		<i>S. aurora</i> (aurora)	<i>S. variegatus</i> (harlequin)	<i>S. brevispinis</i> (silvergrey)	
		<i>S. melanostomus</i> (blackgill)	<i>S. wilsoni</i> (pygmy)	<i>S. diploproa</i> (splitnose)	
		<i>S. paucispinis</i> (bocaccio)	<i>S. babcocki</i> (redbanded)	<i>S. saxicola</i> (stripetail)	
		<i>S. goodei</i> (chilipepper)	<i>S. proriger</i> (redstripe)	<i>S. miniatus</i> (vermillion)	
		<i>S. crameri</i> (darkblotch)	<i>S. zacentrus</i> (sharpchin)	<i>S. reedi</i> (yellowmouth)	
	<i>S. elongatus</i> (greenstriped)	<i>S. jordani</i> (shortbelly)			
In the Eastern GOA only, Slope rockfish also includes <i>S. polyspinous</i> . (Northern)					
5	Pelagic shelf rockfish	<i>S. ciliatus</i> (dusky)	<i>S. entomelas</i> (widow)	<i>S. flavidus</i> (yellowtail)	
6	Demersal shelf rockfish (DSR)	<i>S. pinniger</i> (canary)	<i>S. maliger</i> (quillback)	<i>S. ruberrimus</i> (yelloweye)	
		<i>S. nebulosus</i> (china)	<i>S. helvomaculatus</i> (rosethorn)		
		<i>S. caurinus</i> (copper)	<i>S. nigrocinctus</i> (tiger)		
		DSR-SEO = Demersal shelf rockfish in the Southeast Outside District The operator of a catcher vessel that is required to have a Federal fisheries permit, or that harvests IFQ halibut with hook and line or jig gear, must retain and land all DSR that is caught while fishing for groundfish or IFQ halibut in the SEO. Limits on sale and requirements for disposal of DSR are set out at § 679.20 (j).			
7	Other species	sculpins	octopus	sharks	Squid
8	Aggregated rockfish	Means rockfish of the genera <i>Sebastes</i> and <i>Sebastobus</i> defined at § 679.2 except in:			
		Southeast Outside District (SEO)	where DSR is a separate category for those species marked with a numerical percentage		
		Eastern Regulatory Area (ERA)	where SR/RE is a separate category for those species marked with a numerical percentage		

Table 10 to part 679
Updated April 12, 2006

Notes to Table 10 to Part 679		
9	N/A	not applicable
10	Aggregated forage fish (all species of the following families)	
	Bristlemouths, lightfishes, and anglemouths (family <i>Gonostomatidae</i>)	209
	Capelin smelt (family <i>Osmeridae</i>)	516
	Deep-sea smelts (family <i>Bathylagidae</i>)	773
	Eulachon smelt (family <i>Osmeridae</i>)	511
	Gunnels (family <i>Pholidae</i>)	207
	Krill (order <i>Euphausiacea</i>)	800
	Laternfishes (family <i>Myctophidae</i>)	772
	Pacific herring (family <i>Clupeidae</i>)	235
	Pacific Sand fish (family <i>Trichodontidae</i>)	206
	Pacific Sand lance (family <i>Ammodytidae</i>)	774
	Pricklebacks, war-bonnets, eelblennys, cockscombs and Shannys (family <i>Stichaeidae</i>)	208
	Surf smelt (family <i>Osmeridae</i>)	515
11	Skates Species and Groups	
	Big Skates	702
	Longnose Skates	701
	Other Skates	700

Table 11 to Part 679–BSAI Retainable Percentages (*Updated 10/18/02*)

BASIS SPECIES		INCIDENTAL CATCH SPECIES ⁵															
		Pollock	Pacific cod	Atka mackerel	Alaska plaice	Arrow-tooth	Yellow fin sole	Other flatfish ²	Rock sole	Flathead sole	Green-land turbot	Sable-fish ¹	Short-raker/rougheye	Aggregated rockfish ⁶	Squid	Aggregated forage fish ⁷	Other species ⁴
110	Pacific cod	20	na ⁵	20	20	35	20	20	20	20	1	1	2	5	20	2	20
121	Arrow-tooth	0	0	0	0	na ⁵	0	0	0	0	0	0	0	0	0	2	0
122	Flathead sole	20	20	20	35	35	35	35	35	na5	35	15	7	15	20	2	20
123	Rock sole	20	20	20	35	35	35	35	na ⁵	35	1	1	2	15	20	2	20
127	Yellowfin sole	20	20	20	35	35	na ⁵	35	35	35	1	1	2	5	20	2	20
133	Alaska Plaice	20	20	20	na ⁵	35	35	35	35	35	1	1	2	5	20	2	20
134	Greenland turbot	20	20	20	20	35	20	20	20	20	na ⁵	15	7	15	20	2	20
136	Northern	20	20	20	20	35	20	20	20	20	35	15	7	15	20	2	20
141	Pacific Ocean perch	20	20	20	20	35	20	20	20	20	35	15	7	15	20	2	20
152/ 151	Shortraker/ Rougheye	20	20	20	20	35	20	20	20	20	35	15	na ⁵	5	20	2	20
193	Atka mackerel	20	20	na ⁵	20	35	20	20	20	20	1	1	2	5	20	2	20
270	Pollock	na ⁵	20	20	20	35	20	20	20	20	1	1	2	5	20	2	20
710	Sablefish1	20	20	20	20	35	20	20	20	20	35	na ⁵	7	15	20	2	20
875	Squid	20	20	20	20	35	20	20	20	20	1	1	2	5	na ⁵	2	20
Other flatfish ²		20	20	20	35	35	35	na ⁵	35	35	1	1	2	5	20	2	20
Other rockfish ³		20	20	20	20	35	20	20	20	20	35	15	7	15	20	2	20
Other species ⁴		20	20	20	20	35	20	20	20	20	1	1	2	5	20	2	na ⁵
Aggregated amount non-groundfish species		20	20	20	20	35	20	20	20	20	1	1	2	5	20	2	20

NOTES to Table 11	
1	Sablefish: for fixed gear restrictions, see 50 CFR 679.7(f)(3)(ii) and 679.7(f)(11).
2	Other flatfish includes all flatfish species, except for Pacific halibut (a prohibited species), flathead sole, Greenland turbot, rock sole, yellowfin sole, Alaska plaice, and arrowtooth flounder.
3	Other rockfish includes all <i>Sebastes</i> and <i>Sebastolobus</i> species except for Pacific ocean perch; and northern, shortraker, and roughey rockfish. The CDQ reserves for shortraker, roughey, and northern rockfish will continue to be managed as the “other red rockfish” complex for the BS.
4	Other species includes sculpins, sharks, skates and octopus. Forage fish, as defined at Table 2 to this part are not included in the “other species” category.
5	na = not applicable
6	Aggregated rockfish includes all of the genera <i>Sebastes</i> and <i>Sebastolobus</i> , except shortraker and roughey rockfish.
7	Forage fish are defined at Table 2 to this part.