

TABLE 8

**The 30 most frequently reported *Salmonella* serotypes from Human sources
Percent Change in reported isolates**

Rank			Serotype	Reported Isolates			Percent Change		
1995	2000	2005		1995	2000	2005	1995-2000	2000-2005	1995-2005
2	1	1	Typhimurium *	9702	7428	6982	-23	-6	-28
1	2	2	Enteritidis	10201	6487	6730	-36	4	-34
3	3	3	Newport	2566	3074	3295	20	7	28
4	4	4	Heidelberg	2095	1772	1903	-15	7	-9
6	5	5	Javiana	758	1204	1324	59	10	75
64	21	6	I 4,[5],12:i:-	45	233	822	418	253	1727
8	6	7	Montevideo	685	841	809	23	-4	18
7	7	8	Muenchen	754	642	733	-15	14	-3
16	11	9	Saintpaul	467	548	683	17	25	46
12	12	10	Braenderup	588	531	603	-10	14	3
11	10	11	Oranienburg	595	563	590	-5	5	-1
23	19	12	Mississippi	199	286	565	44	98	184
14	8	13	Infantis	521	613	505	18	-18	-3
20	13	14	Paratyphi B var. L(+) tartrate+	268	468	460	75	-2	72
10	9	15	Thompson	625	609	428	-3	-30	-32
9	14	16	Agona	683	406	367	-41	-10	-46
17	15	17	Typhi	442	399	348	-10	-13	-21
27	27	18	Hartford	164	150	239	-9	59	46
15	20	19	Stanley	481	239	224	-50	-6	-53
18	18	20	Berta	367	312	209	-15	-33	-43
5	16	21	Hadar	814	354	205	-57	-42	-75
33	23	22	Bareilly	109	182	201	67	10	84
25	24	23	Anatum	174	177	197	2	11	13
13	17	24	Poona	531	337	196	-37	-42	-63
29	26	25	Mbandaka	154	157	190	2	21	23
26	25	26	Panama	173	158	148	-9	-6	-14
31	31	27	Litchfield	115	119	141	3	18	23
30	29	28	Sandiego	117	142	138	21	-3	18
28	33	29	Schwarzengrund	162	113	138	-30	22	-15
19	40	30	Brandenburg	284	84	134	-70	60	-53

NOTE:

* Typhimurium includes var. 5- (Formerly var. Copenhagen)