

Unspecified and other pneumoconioses: Death rates (per million population) by race and sex, U.S. residents age 15 and over, 1996–2005

| Year | Overall | White | | Black | | Other | |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | | Male | Female | Male | Female | Male | Female |
| Crude Death Rate | | | | | | | |
| 1996 | 1.68 | 3.72 | 0.12 | 1.58 | - | 1.16 | - |
| 1997 | 1.37 | 3.05 | 0.11 | 1.38 | - | 0.23 | - |
| 1998 | 1.55 | 3.51 | 0.10 | 1.19 | 0.07 | 0.22 | 0.39 |
| 1999 | 1.48 | 3.38 | 0.08 | 1.34 | - | 0.42 | - |
| 2000 | 1.34 | 3.02 | 0.06 | 1.45 | 0.07 | 0.18 | - |
| 2001 | 1.19 | 2.66 | 0.09 | 1.19 | 0.14 | - | - |
| 2002 | 1.13 | 2.43 | 0.15 | 1.47 | - | 0.17 | - |
| 2003 | 1.01 | 2.37 | 0.06 | 0.15 | 0.13 | 0.49 | - |
| 2004 | 0.91 | 2.10 | 0.04 | 0.82 | - | - | - |
| 2005 | 0.92 | 1.99 | 0.12 | 1.10 | - | - | - |
| 1996–2005 | 1.23 | 2.77 | 0.09 | 1.14 | 0.04 | 0.24 | 0.03 |
| Age-Adjusted Death Rate | | | | | | | |
| 1996 | 1.73 | 4.70 | 0.11 | 3.09 | - | 2.13 | - |
| 1997 | 1.40 | 3.82 | 0.09 | 2.88 | - | 0.37 | - |
| 1998 | 1.57 | 4.22 | 0.09 | 2.31 | 0.09 | 0.62 | 0.50 |
| 1999 | 1.49 | 4.06 | 0.06 | 2.33 | - | 0.72 | - |
| 2000 | 1.36 | 3.83 | 0.05 | 3.13 | 0.07 | 0.43 | - |
| 2001 | 1.20 | 3.29 | 0.08 | 2.41 | 0.15 | - | - |
| 2002 | 1.14 | 2.93 | 0.12 | 2.95 | - | 0.38 | - |
| 2003 | 1.01 | 2.82 | 0.05 | 0.28 | 0.14 | 0.71 | - |
| 2004 | 0.91 | 2.46 | 0.04 | 1.74 | - | - | - |
| 2005 | 0.89 | 2.25 | 0.10 | 2.23 | - | - | - |
| 1996–2005 | 1.25 | 3.36 | 0.08 | 2.33 | 0.04 | 0.46 | 0.04 |

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.