

Unspecified and other pneumoconioses: Years of potential life lost to age 65 and to life expectancy by race and sex, U.S. residents age 15 and over, 1996–2005

Year	White		Black		Other		Total
	Male	Female	Male	Female	Male	Female	
Years of Potential Life Lost to Age 65							
1996	245	40	5	-	15	-	305
1997	135	15	10	-	-	-	160
1998	220	10	30	5	-	5	270
1999	155	-	45	-	5	-	205
2000	110	5	50	25	-	-	190
2001	140	15	15	20	-	-	190
2002	100	5	25	-	-	-	130
2003	130	-	-	50	20	-	200
2004	135	-	5	-	-	-	140
2005	220	30	15	-	-	-	265
TOTAL	1,590	120	200	100	40	5	2,055
Years of Potential Life Lost to Life Expectancy							
1996	2,973	172	154	-	78	-	3,377
1997	2,327	117	137	-	14	-	2,595
1998	2,829	140	144	21	6	36	3,176
1999	2,683	59	189	-	30	-	2,961
2000	2,344	77	186	38	9	-	2,654
2001	2,165	118	145	51	-	-	2,479
2002	2,047	154	189	-	9	-	2,399
2003	2,081	69	20	77	62	-	2,309
2004	1,977	52	100	-	-	-	2,129
2005	2,085	162	145	-	-	-	2,392
TOTAL	23,511	1,120	1,409	187	208	36	26,471

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data.