

All pneumoconioses: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1996–2005

State	No. of Deaths Rank		Death Rate				YPLL to Life Expectancy			
			Crude	Rank	Age-Adjusted	Rank	Total	Rank	YPLL/death	Rank
Alabama	673	12	18.98	7	18.72	8	7,812	12	11.6	17
Alaska	29	49	6.08	34	14.53	12	301	49	10.4	43
Arizona	220	28	5.37	39	5.32	40	2,334	28	10.6	38
Arkansas	163	32	7.65	28	7.02	30	2,037	31	12.5	7
California	1,268	5	4.75	42	5.49	38	13,959	6	11.0	29
Colorado	299	23	8.58	20	11.26	18	3,219	24	10.8	35
Connecticut	206	29	7.59	29	6.80	31	2,306	29	11.2	24
Delaware	148	33	23.22	5	23.52	5	1,621	33	11.0	29
District of Columbia	10	51	2.09	51	2.30	51	137	51	13.7	2
Florida	1,088	8	8.23	25	6.24	34	11,923	8	11.0	29
Georgia	245	25	3.75	48	4.89	42	2,958	25	12.1	10
Hawaii	30	48	3.06	50	2.94	49	392	47	13.1	5
Idaho	86	39	8.47	23	9.14	24	964	40	11.2	24
Illinois	589	14	6.01	35	6.17	35	6,567	15	11.1	26
Indiana	302	22	6.29	33	6.37	33	3,401	22	11.3	22
Iowa	121	36	5.15	40	4.25	45	1,279	36	10.6	38
Kansas	81	40	3.81	47	3.62	47	1,035	39	12.8	6
Kentucky	1,229	6	37.50	3	39.02	3	14,492	5	11.8	12
Louisiana	351	20	10.08	17	10.83	19	4,723	19	13.5	3
Maine	144	34	13.67	12	12.37	15	1,511	35	10.5	41
Maryland	538	15	12.68	14	14.08	13	6,111	16	11.4	19
Massachusetts	416	19	8.04	27	7.51	27	4,338	20	10.4	43
Michigan	465	17	5.91	36	6.06	36	5,340	18	11.5	18
Minnesota	228	27	5.78	37	5.90	37	2,453	27	10.8	35
Mississippi	477	16	21.42	6	22.13	6	8,478	10	17.8	1
Missouri	193	31	4.31	44	4.09	46	2,286	30	11.8	12
Montana	114	38	15.65	9	14.79	11	1,268	37	11.1	26
Nebraska	54	45	3.98	46	3.60	48	505	45	9.4	49
Nevada	68	44	4.17	45	4.93	41	898	41	13.2	4
New Hampshire	71	42	7.06	31	7.55	26	786	42	11.1	26
New Jersey	950	9	14.10	11	13.44	14	9,789	9	10.3	45
New Mexico	121	36	8.53	22	9.32	23	1,189	38	9.8	48
New York	749	10	4.92	41	4.83	43	8,467	11	11.3	22
North Carolina	606	13	9.33	19	9.97	22	7,112	13	11.7	16
North Dakota	38	46	7.38	30	6.38	32	400	46	10.5	41
Ohio	1,155	7	12.83	13	12.24	16	12,754	7	11.0	29
Oklahoma	123	35	4.48	43	4.33	44	1,528	34	12.4	8
Oregon	336	21	12.14	15	11.83	17	3,597	21	10.7	37
Pennsylvania	5,668	1	57.09	2	46.10	2	53,203	1	9.4	49
Rhode Island	71	42	8.28	24	7.24	28	753	43	10.6	38
South Carolina	278	24	8.58	20	9.09	25	3,297	23	11.9	11
South Dakota	20	50	3.35	49	2.88	50	237	50	11.8	12
Tennessee	458	18	9.99	18	10.40	20	5,399	17	11.8	12
Texas	1,339	4	8.20	26	10.15	21	16,531	4	12.3	9
Utah	195	30	11.60	16	16.16	9	1,973	32	10.1	46
Vermont	35	47	7.02	32	7.11	29	386	48	11.0	29
Virginia	1,705	3	29.77	4	34.00	4	19,463	3	11.4	19
Washington	693	11	14.61	10	16.02	10	7,008	14	10.1	46
West Virginia	2,747	2	185.78	1	161.36	1	30,266	2	11.0	29
Wisconsin	244	26	5.68	38	5.48	39	2,781	26	11.4	19
Wyoming	73	41	18.51	8	20.27	7	664	44	9.1	51

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.