

All pneumoconioses: Number of deaths by sex, race, and age, and median age at death, U.S. residents age 15 and over, 1996–2005

Year	No. of Deaths	Underlying Cause (%)	Sex		Race			Age Group (yrs)								Median Age (yrs)
			Male	Female	White	Black	Other	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	
1996	3,115	35.8	3,031	84	2,937	164	14	-	1	11	51	210	883	1,364	595	78
1997	2,928	37.1	2,851	77	2,780	139	9	-	2	12	48	191	751	1,313	611	78
1998	2,791	39.4	2,720	71	2,651	127	13	-	3	11	41	189	733	1,226	588	78
1999	2,745	39.5	2,683	62	2,592	136	17	-	1	9	38	177	683	1,210	627	78
2000	2,864	39.9	2,783	81	2,704	144	16	2	4	14	42	170	651	1,236	745	79
2001	2,747	40.4	2,674	73	2,585	148	14	2	-	11	47	194	632	1,195	666	78
2002	2,720	40.3	2,633	87	2,572	137	11	1	3	10	41	186	599	1,214	666	79
2003	2,639	41.8	2,546	93	2,488	134	17	2	4	15	46	184	544	1,180	664	79
2004	2,531	40.1	2,454	77	2,372	149	10	-	4	14	49	175	501	1,110	678	80
2005	2,430	40.5	2,337	93	2,294	122	14	3	3	14	57	175	469	1,051	658	80
TOTAL	27,510	39.4	26,712	798	25,975	1,400	135	10	25	121	460	1,851	6,446	12,099	6,498	79

- indicates no deaths listed.
 NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.
 SOURCE: National Center for Health Statistics multiple cause-of-death data.