

Unspecified and other pneumoconioses: Number of deaths by sex, race, and age, and median age at death, U.S. residents age 15 and over, 1996–2005

Year	No. of Deaths	Underlying Cause (%)	Sex		Race			Age Group (yrs)								Median Age (yrs)
			Male	Female	White	Black	Other	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	
1996	349	34.1	338	11	326	18	5	-	-	3	6	28	81	147	84	78
1997	287	32.1	277	10	270	16	1	-	-	1	3	18	57	130	78	80
1998	329	37.4	317	12	311	15	3	-	1	1	6	24	70	151	76	79
1999	318	36.2	311	7	300	16	2	-	-	2	4	19	70	138	85	79
2000	297	39.4	290	7	277	19	1	-	-	4	2	12	58	114	107	82
2001	267	38.2	256	11	250	17	-	-	-	1	9	6	58	101	92	81
2002	257	44.0	243	14	237	19	1	-	-	-	3	17	52	106	79	80
2003	233	40.3	225	8	226	4	3	-	-	2	4	18	38	100	71	80
2004	212	46.2	208	4	201	11	-	-	-	1	4	11	51	82	63	79
2005	216	45.8	204	12	201	15	-	-	-	1	7	27	45	65	71	79
TOTAL	2,765	38.8	2,669	96	2,599	150	16	-	1	16	48	180	580	1,134	806	80

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data.