

Hypersensitivity pneumonitis: Death rates (per million population) by race and sex, U.S. residents age 15 and over, 1996–2005

Year	Overall	White		Black		Other	
		Male	Female	Male	Female	Male	Female
Crude Death Rate							
1996	0.25	0.40	0.17	0.09	0.08	-	-
1997	0.18	0.29	0.14	-	-	-	-
1998	0.18	0.35	0.08	0.08	-	-	-
1999	0.27	0.40	0.23	0.08	-	-	-
2000	0.22	0.29	0.20	0.16	0.14	-	-
2001	0.30	0.51	0.20	0.08	0.07	-	-
2002	0.29	0.33	0.35	0.08	-	-	-
2003	0.32	0.41	0.30	-	0.13	0.32	0.30
2004	0.28	0.41	0.26	-	0.07	-	0.15
2005	0.28	0.42	0.24	0.07	-	0.15	0.14
1996–2005	0.25	0.38	0.22	0.06	0.05	0.05	0.06
Age-Adjusted Death Rate							
1996	0.25	0.46	0.14	0.07	0.09	-	-
1997	0.19	0.35	0.12	-	-	-	-
1998	0.18	0.40	0.06	0.10	-	-	-
1999	0.27	0.46	0.20	0.10	-	-	-
2000	0.22	0.34	0.18	0.20	0.17	-	-
2001	0.30	0.60	0.18	0.08	0.09	-	-
2002	0.29	0.36	0.32	0.12	-	-	-
2003	0.32	0.47	0.26	-	0.15	0.71	0.41
2004	0.28	0.46	0.23	-	0.08	-	0.20
2005	0.28	0.47	0.21	0.06	-	0.16	0.19
1996–2005	0.26	0.43	0.19	0.07	0.06	0.10	0.09

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.