

Hypersensitivity pneumonitis: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1996–2005

State	No. of Deaths		Death Rate				YPLL to Life Expectancy			
	Deaths	Rank	Crude	Rank	Age-Adjusted	Rank	Total	Rank	YPLL/death	Rank
Alabama	4	35	0.11	42	0.11	42	60	37	15.1	22
Alaska	-	-	-	-	-	-	-	-	-	-
Arizona	7	21	0.17	25	0.16	30	152	17	21.7	2
Arkansas	1	43	0.05	46	0.04	47	15	42	14.6	29
California	60	2	0.22	20	0.25	18	1,157	1	19.3	11
Colorado	8	17	0.23	19	0.28	16	127	23	15.9	18
Connecticut	7	21	0.26	17	0.25	18	143	19	20.4	6
Delaware	-	-	-	-	-	-	-	-	-	-
District of Columbia	1	43	0.21	21	0.23	22	9	47	8.6	46
Florida	21	9	0.16	30	0.13	37	321	9	15.3	21
Georgia	2	40	0.03	48	0.04	47	15	42	7.7	48
Hawaii	-	-	-	-	-	-	-	-	-	-
Idaho	9	15	0.89	5	0.95	4	129	22	14.3	32
Illinois	17	12	0.17	25	0.18	25	211	12	12.4	40
Indiana	8	17	0.17	25	0.17	27	121	24	15.1	22
Iowa	23	8	0.98	4	0.84	5	271	10	11.8	42
Kansas	10	14	0.47	11	0.47	10	191	14	19.1	12
Kentucky	5	30	0.15	32	0.15	34	75	29	14.9	25
Louisiana	6	28	0.17	25	0.18	25	121	24	20.1	8
Maine	5	30	0.47	11	0.44	11	61	35	12.2	41
Maryland	7	21	0.16	30	0.17	27	139	20	19.8	10
Massachusetts	7	21	0.14	36	0.13	37	72	32	10.2	43
Michigan	27	7	0.34	14	0.35	14	379	7	14.1	34
Minnesota	31	5	0.79	7	0.79	6	405	6	13.1	37
Mississippi	3	37	0.13	39	0.14	35	61	35	20.2	7
Missouri	9	15	0.20	24	0.19	24	139	20	15.4	20
Montana	1	43	0.14	36	0.13	37	8	48	8.4	47
Nebraska	6	28	0.44	13	0.42	13	82	28	13.7	36
Nevada	2	40	0.12	40	0.16	30	56	38	28.1	1
New Hampshire	1	43	0.10	43	0.11	42	14	45	14.3	32
New Jersey	7	21	0.10	43	0.10	44	150	18	21.4	4
New Mexico	1	43	0.07	45	0.07	45	14	45	14.4	31
New York	37	3	0.24	18	0.24	21	538	4	14.5	30
North Carolina	19	10	0.29	15	0.31	15	249	11	13.1	37
North Dakota	3	37	0.58	8	0.44	11	27	41	9.0	45
Ohio	19	10	0.21	21	0.20	23	356	8	18.7	13
Oklahoma	4	35	0.15	32	0.14	35	68	34	17.1	16
Oregon	14	13	0.51	9	0.49	9	207	13	14.8	28
Pennsylvania	29	6	0.29	15	0.26	17	505	5	17.4	15
Rhode Island	1	43	0.12	40	0.12	40	15	42	14.9	25
South Carolina	5	30	0.15	32	0.16	30	91	26	18.1	14
South Dakota	5	30	0.84	6	0.75	7	75	29	14.9	25
Tennessee	8	17	0.17	25	0.17	27	166	15	20.8	5
Texas	34	4	0.21	21	0.25	18	550	3	16.2	17
Utah	8	17	0.48	10	0.62	8	159	16	19.9	9
Vermont	7	21	1.40	2	1.42	2	70	33	10.0	44
Virginia	3	37	0.05	46	0.06	46	47	39	15.7	19
Washington	7	21	0.15	32	0.16	30	91	26	12.9	39
West Virginia	2	40	0.14	36	0.12	40	43	40	21.6	3
Wisconsin	64	1	1.49	1	1.44	1	891	2	13.9	35
Wyoming	5	30	1.27	3	1.32	3	75	29	15.0	24

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.