

Women's Health USA 2008



Please note that *Women's Health USA 2008* is not copyrighted. Readers are free to duplicate and use all or part of the information contained in this publication.

Suggested Citation:

U.S. Department of Health and Human Services,
Health Resources and Services Administration. *Women's Health USA 2008*.
Rockville, Maryland: U.S. Department of Health and Human Services, 2008.

This publication is available online at www.hrsa.gov/womenshealth
Single copies of this publication are also available at no charge from the
HRSA Information Center
P.O. Box 2910
Merrifield, VA 22116
1-888-ASK-HRSA or ask@hrsa.gov

The data book is available in limited quantities in CD format.

PREFACE AND READER'S GUIDE	4	Chronic Fatigue Syndrome	31	<i>Special Populations</i>	
INTRODUCTION	6	Cancer	32	Older Women	58
POPULATION CHARACTERISTICS	9	Diabetes	34	Rural and Urban Women	59
U.S. Population	10	Overweight and Obesity	35	HEALTH SERVICES UTILIZATION	60
U.S. Female Population	11	Heart Disease and Stroke	36	Usual Source of Care	61
Household Composition	12	Hypertension	37	Health Insurance	62
Women and Poverty	13	Oral Health and Dental Care	38	Medicare and Medicaid	63
Food Security	14	Eye Health	39	Preventive Care	64
Women and Federal Nutrition Programs	15	Osteoporosis	40	Vaccination	65
Women in Health Profession Schools	16	Digestive Disorders	41	Hospitalizations	66
Educational Degrees and Instructional Staff	17	Endocrine and Metabolic Disorders	42	Health Care Expenditures	67
Women in the Labor Force	18	Genetics and Women's Health	43	Medication Use	68
HEALTH STATUS	19	HIV/AIDS	44	Mental Health Care Utilization	69
<i>Health Behaviors</i>		Sexually Transmitted Infections	45	HIV Testing	70
Physical Activity	20	Injury	46	Organ Transplantation	71
Nutrition	21	Occupational Injury	47	Quality of Women's Health Care	72
Alcohol Use	22	Attention Deficit Hyperactivity Disorder	48	Satisfaction with Health Care	73
Cigarette Smoking	23	Mental Illness and Suicide	49	HRSA Programs Related to Women's Health	74
Illicit Drug Use	24	Intimate Partner Violence	50	INDICATORS IN PREVIOUS EDITIONS	75
<i>Health Indicators</i>		Urologic Disorders	51	ENDNOTES	76
Self-Reported Health Status	25	Gynecological and Reproductive Disorders	52	DATA SOURCES	78
Life Expectancy	26	<i>Maternal Health</i>		CONTRIBUTORS	80
Leading Causes of Death	27	Live Births	53		
Activity Limitations and Disabilities	28	Breastfeeding	54		
Arthritis	29	Smoking During Pregnancy	55		
Asthma	30	Maternal Morbidity and Risk Factors in Pregnancy	56		
		Maternal Mortality	57		

PREFACE AND READER'S GUIDE

The U.S. Department of Health and Human Services, Health Resources and Services Administration (HRSA) supports healthy women building healthy communities. HRSA is charged with ensuring access to quality health care through a network of community-based health centers, maternal and child health programs, and community HIV/AIDS programs throughout the States and Territories. In addition, HRSA's mission includes supporting individuals pursuing careers in medicine, nursing, and many other health disciplines. HRSA fulfills these responsibilities by collecting and analyzing timely, topical information that identifies health priorities and trends that can be addressed through program interventions and capacity building.

HRSA is pleased to present *Women's Health USA 2008*, the seventh edition of the Women's Health USA data book. To reflect the ever-changing, increasingly diverse population and its characteristics, *Women's Health USA* selectively highlights emerging issues and trends in women's health. Data and information on occupational injury, maternal mortality, digestive disorders, oral health, eye health, and urologic disorders are only a few of the new topics included in this edition. Every effort has been made to highlight racial and ethnic, sex/gender, and socioeconomic



disparities where possible. Where race and ethnicity data are reported, every effort was made to ensure that groups are mutually exclusive; when groups of Blacks and Whites exclude Hispanics they are described as non-Hispanic, and Asian/Pacific Islanders and American Indian/Alaska Natives are also generally non-Hispanic. In some instances, it was not possible to provide data for all races due to the design of the original data source or the size of the sample population; therefore, data with a relative standard error of 30 percent or greater were considered unreliable and were not reported.

The data book was developed by HRSA to provide readers with an easy-to-use collection of current and historical data on some of the most pressing health challenges facing women, their families, and their communities. *Women's Health USA 2008* is intended to be a concise reference for policymakers and program managers at the Federal, State, and local levels to identify and clarify issues affecting the health of women. In these pages, readers will find a profile of women's health from a variety of data sources. The data book brings together the latest available information from various agencies within the Federal government, including the U.S. Department of Health and Human Services, U.S. Department of Agriculture, U.S. Department of Labor, and U.S. Department of Justice. Non-Federal data

sources were used when no Federal source was available. Every attempt has been made to use data collected in the past 5 years. It is important to note that the incidence data included are generally not age-adjusted to the 2000 population standard of the United States. This affects the comparability of data from year to year, and the interpretation of differences across various groups, especially those of different races and ethnicities. Without age adjustment, it is difficult to know how much of the difference in incidence rates between groups can be attributed to differences in the groups' age distributions.

Women's Health USA 2008 is available online through either the HRSA Office of Women's Health Web site at www.hrsa.gov/womenshealth or the Office of Data and Program Development's Web site at www.mchb.hrsa.gov/data. In an effort to produce a timely document, some of the topics covered in *Women's Health USA 2007* were not included in this year's edition because new data were not available. For coverage of these issues, please refer to *Women's Health USA 2007*, also available online. The National Women's Health Information Center, located online at www.womenshealth.gov, has updated and detailed women's and minority health data and maps. These data are available through Quick Health Data Online at www.4woman.gov/quickhealthdata. Data are available at the State

and county levels, by age, race and ethnicity, and sex/gender.

Women's Health USA 2008 is not copyrighted. Readers are free to duplicate and use any of the information contained in this publication. Please provide any feedback on this publication to the HRSA Information Center which offers single copies of the data book in print or on CD at no charge:

HRSA Information Center
P.O. Box 2910
Merrifield, VA 22116
Phone: 703-442-9051
Toll-free: 1-888-ASK-HRSA
TTY: 1-877-4TY-HRSA
Fax: 703-821-2098
ask@hrsa.gov
www.ask.hrsa.gov

INTRODUCTION

In 2006, women represented 50.8 percent of the 299 million people residing in the United States. In most age groups, women accounted for approximately half of the population, with the exception of people aged 65 years and older; within this age group, women represented 58 percent of the population. The growing diversity of the U.S. population is reflected in the racial and ethnic distribution of women across age groups. Black and Hispanic women accounted for 8.9 and 6.3 percent of the female population aged 65 years and older, respectively, but they represented 14.7 and 21.0 percent of females under 15 years of age. Non-Hispanic Whites accounted for 80.6 percent of women aged 65 years and older, but only 56.4 percent of those under 15 years of age.

America's growing diversity underscores the importance of examining and addressing racial and ethnic disparities in health status and the use of health care services. In 2006, 62.3 percent of non-Hispanic White women reported themselves to be in excellent or very good health, compared to only 53.4 percent of Hispanic women and 50.2 percent of non-Hispanic Black women.

Minority women are disproportionately affected by a number of diseases and health conditions, including HIV/AIDS, sexually transmitted infections, diabetes, and asthma. For



instance, in 2006, non-Hispanic Black and Hispanic women accounted for more than three-fourths of women living with HIV/AIDS (63.9 and 15.2 percent, respectively). One-third of non-Hispanic White women had ever been tested for HIV, compared to 53.7 percent of non-Hispanic Black women and 46.1 percent of Hispanic women.

Diabetes is a chronic condition and a leading cause of death and disability in the United States, and is especially prevalent among minority populations. Among non-Hispanic Black and Hispanic women, diabetes occurred at a rate of 117.6 and 111.8 per 1,000 women, respectively, compared to 69.4 per 1,000 non-Hispanic White women. Hypertension, or high blood pressure, was also more prevalent among non-Hispanic Black women than women of other races. In 2005–2006, this condition occurred at a rate of 179.2 per 1,000 non-Hispanic Black women, compared to 157.0 per 1,000 non-Hispanic White women and 113.1 per 1,000 Hispanic women.

Some conditions, such as arthritis and heart disease, disproportionately affect non-Hispanic White women. For instance, in 2006, more than 27 percent of non-Hispanic White women had arthritis, compared to 23.5 percent of non-Hispanic Black women and 14.3 percent of Hispanic women.

In addition to race and ethnicity, income and education are important factors that contribute to women's health and access to health care. Regardless of family structure, women are more likely than men to live in poverty. Poverty rates were highest among women who were heads of their households (25.1 percent). Poverty rates were also highest among American Indian/Alaska Native women (27.6 percent), followed by non-Hispanic Black and Hispanic women (23.4 and 20.2 percent, respectively). Non-Hispanic Black and Hispanic women were also more likely to be heads of households than their non-Hispanic White and Asian counterparts.

Some conditions and health risks are more closely linked to family income than to race and ethnicity, such as asthma. Rates of asthma decline as income increases and women with higher incomes are more likely to effectively manage their asthma. Among women with asthma whose incomes were below 100 percent of poverty, nearly 36 percent had an asthma-related emergency room visit in the past year, compared to 24.8 percent of women with family incomes of 300 percent or more of poverty.

Mental health is another important aspect of women's overall health. A range of mental health problems, including depression, anxiety, phobias, and post-traumatic stress disorder, disproportionately affect women. Unlike many other health

concerns, younger women are more likely than older women to suffer from serious psychological stress and major depressive episodes.

Physical disabilities are more prevalent among women as well. Disability can be defined as impairment of the ability to perform common activities like walking up stairs, sitting or standing for 2 hours or more, grasping small objects, or carrying items like groceries. Therefore, the terms "activity limitations" and "disabilities" are used interchangeably throughout this book. Overall, 15.0 percent of women and 12.6 percent of men reported having activity limitations in 2006.

Men, however, bear a disproportionate burden of some health conditions, such as HIV/AIDS, hypertension and heart disease. In 2006, for instance, adolescent and adult males accounted for 72.9 percent of new AIDS cases, though a smaller proportion of men had ever been tested for HIV than women (33.7 versus 37.8 percent, respectively).

Certain health risks, such as cigarette use and injury, occur more commonly among men than women. In 2006, 27.8 percent of males smoked cigarettes, compared to 22.4 percent of females. Among men, 30.2 percent of emergency department visits were injury related, while only 21.3 percent of women's visits were due to injury. In addition, men were more likely than women to lack health insurance.

Many diseases and health conditions, such as those mentioned above, can be avoided or minimized through good nutrition, regular physical activity, and preventive health care. In 2005, 19.7 percent of women's visits to physicians were for preventive care, including prenatal care, preventive screenings, and immunizations. Overall, 64.6 percent of older women reported receiving a flu shot in 2006; however, this percentage ranged from 46.6 percent among Hispanic women to 67.3 percent of non-Hispanic White women.

In addition to preventive health care, preventive dental care is also important to prevent dental caries and gum disease. In 2003–2004, 74.6 percent of women with incomes of 300 percent or more of poverty saw a dentist in the past year, compared to 51.4 percent of women with incomes below 100 percent of poverty, and 44.9 percent of women with incomes of 200–299 percent of poverty.

There are many ways women (and men) can promote health and help prevent disease and disability. Regular physical activity is one of these. In 2006, 10.3 percent of women participated in adequate physical activity—30 minutes of moderate-intensity physical activity on most days of the week or 20 minutes of vigorous-intensity activity on 3 or more days per week. Non-Hispanic White women and women with higher

incomes were most likely to meet the recommended levels of physical activity.

Healthy eating habits can also be a major contributor to long-term health and prevention of chronic disease. In 2003–2004, however, more than half of all adult women had diets that included more than the recommended amount of saturated fat and sodium and less than the recommended amount of folate and calcium. Overall, 63.5 percent of women exceeded the maximum daily intake of saturated fat, and 70 percent exceeded the maximum amount of sodium.

While some behaviors have a positive effect on health, a number of others, such as smoking, illicit drug use, and excessive alcohol use can have a negative effect. In 2006, 22.4 percent of women smoked. However, nearly 46 percent of female smokers tried to quit at some point in the past year. During the same year, 44.9 percent of women reported any alcohol use in the past month, but relatively few women (15.6 percent) reported binge drinking (five or more drinks on the same occasion) and even fewer (3.5 percent) reported heavy alcohol use (binge drinking on 5 days or more in the past month).

Cigarette, alcohol, and illicit drug use is particularly harmful during pregnancy. The use of tobacco during pregnancy has declined steadily since 1989. Based on data from 36 states, 10.5

percent of pregnant women reported smoking during pregnancy in 2005. This rate was highest among American Indian/Alaska Native women (18.1 percent) and lowest among Asian/Pacific Islander women (2.1 percent).

Women's Health USA 2008 can be an important tool for emphasizing the importance of preventive care, counseling, and education, and for illustrating disparities in the health status of women from all age groups and racial and ethnic backgrounds. Health problems can only be remedied if they are recognized. This data book provides information on a range of indicators that can help us track the health behaviors, risk factors, and health care utilization practices of women throughout the United States.

