



SAFER • HEALTHIER • PEOPLE™

Vital and Health Statistics

Series 13, Number 165

December 2007



National Hospital Discharge Survey: 2005 Annual Summary With Detailed Diagnosis and Procedure Data

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Suggested citation

DeFrances CJ, Cullen KA, Kozak LJ. National Hospital Discharge Survey: 2005 annual summary with detailed diagnosis and procedure data. National Center for Health Statistics. Vital Health Stat 13(165). 2007.

Library of Congress Catalog Card Number 97-10617

For sale by the U.S. Government Printing Office
Superintendent of Documents
Mail Stop: SSOP
Washington, DC 20402-9328
Printed on acid-free paper.

Vital and Health Statistics

Series 13, Number 165

National Hospital Discharge Survey: 2005 Annual Summary With Detailed Diagnosis and Procedure Data

Data From the National Health Care
Survey

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Hyattsville, Maryland
December 2007
DHHS Publication No. (PHS) 2008-1736

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*

Jennifer H. Madans, Ph.D., *Acting Co-Deputy Director*

Michael H. Sadagursky, *Acting Co-Deputy Director*

Jennifer H. Madans, Ph.D., *Associate Director for Science*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Planning, Budget, and Legislation*

Michael H. Sadagursky, *Associate Director for Management and Operations*

Lawrence H. Cox, Ph.D., *Associate Director for Research and Methodology*

Linda B. Torian, *Acting Director for Information Technology*

Linda B. Torian, *Acting Director for Information Services*

Linda T. Bilheimer, Ph.D., *Associate Director for Analysis and Epidemiology*

Charles J. Rothwell, M.S., *Director for Vital Statistics*

Jane E. Sisk, Ph.D., *Director for Health Care Statistics*

Jane F. Gentleman, Ph.D., *Director for Health Interview Statistics*

Clifford L. Johnson, M.S.P.H., *Director for Health and Nutrition Examination Surveys*

Division of Health Care Statistics

Jane E. Sisk, Ph.D., *Director*

Robin E. Remsburg, Ph.D., A.P.R.N., B.C., *Deputy Director*

Lola Jean Kozak, Ph.D., *Acting Chief, Hospital Care Statistics Branch*

Charles Adams, *Chief, Technical Services Branch*

Contents

Abstract	1
Highlights.....	1
Patient and Hospital Characteristics	1
Diagnoses.....	1
Procedures	2
Deliveries and Newborn Infants	2
Introduction	3
Methods	4
Data Source	4
Estimation	5
Use of Tables.....	5
References	6
Appendix	193
Technical Notes	193

Text Figures

1. Percent distribution of hospital discharges, days of care, and the civilian population, by age: United States, 2005	2
2. Percentage of inpatients with a routine discharge or discharged home, by age: United States, 2005	2
3. Rate of hospitalization for malignant neoplasms by age: United States, 1990–2005	3
4. Rate of discharges with a first-listed diagnoses of asthma, by age: United States, 2005.....	3
5. Distribution of discharges by number of procedures: United States, 2005	4
6. Rate of episiotomy for females with deliveries: United States, 1980–2005	4

Detailed Tables

Tables on Patient and Hospital Characteristics

1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2005	8
2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2005	9
3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2005	10
4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2005	12
5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, according to sex, age, and geographic region: United States, 2005	15
6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, according to sex and age: United States, 2005	17
7. Number and percent distribution of discharges by type and source of admission to short-stay hospitals, according to sex and age: United States, 2005	18
8. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by disposition of patient, according to sex, age, and geographic region: United States, 2005	19

Diagnosis Tables

9.	Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2005	21
10.	Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2005	22
11.	Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2005	24
12.	Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2005	25
13.	Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2005..	26
14.	Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2005	27
15.	Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2005	28
16.	Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States..	29
17.	Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2005 ..	30
18.	Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2005 ..	31
19.	Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2005.....	32
20.	Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2005....	33
21.	Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2005 ...	34
22.	Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2005.....	35
23.	Number and rate of discharges from short-stay hospitals with a human immunodeficiency virus diagnosis, by selected characteristics: United States, selected years 1990–2005.....	36
24.	Annual average number and rate of discharges from short-stay hospitals with a first-listed injury diagnosis, by Barell injury diagnosis matrix categories: United States, 2003–2005.....	37
25.	Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnosis: United States, 2005.....	38

Procedure Tables

26.	Number of discharges from short-stay hospitals with and without procedures and percentage with procedures, by selected characteristics: United States, 2005	38
27.	Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals by selected procedure categories: United States, 2005.....	39
28.	Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2005	40
29.	Number and rate of discharges from short-stay hospitals with a coronary artery bypass graft and number and rate of procedures, by sex, age, and geographic region: United States, 2005.....	41
30.	Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2005	42
31.	Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2005.....	43
32.	Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2005.....	44
33.	Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2005.....	45
34.	Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2005.....	46
35.	Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2005.....	47

Tables on Deliveries and Newborn Infants

36.	Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States	48
37.	Number and percent distribution of females with deliveries discharged from short-stay hospitals, by length of stay and type of delivery: United States, selected years 1980–2005	49
38.	Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–2005.....	50
39.	Number and rate of females with deliveries discharged from short-stay hospitals, by type of delivery: United States, selected years 1980–2005.....	51

40.	Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery, age, and geographic region: United States, 2005.....	51
41.	Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals, by sex, geographic region, and health status: United States, 2005	52
42.	Number and percent distribution of newborn infants discharged from short-stay hospitals, by length of stay, and average length of stay, by health status: United States, selected years 1980–2005	52
43.	Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2005	53
44.	Number and percentage of male newborn infants circumcised during hospitalization, by geographic region: United States, selected years 1980–2005.....	53

Diagnosis, Days of Care, and Procedure Tables by ICD-9-CM Code

45.	Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005.....	54
46.	Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005	82
47.	Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005	110
48.	Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005.....	173

Appendix Tables

I.	Civilian population by age, geographic region, and sex: United States, 2005	193
II.	Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics, by selected characteristics, 2005	196
III.	Code numbers for Barell injury diagnosis matrix categories in Table 24	198
IV.	Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey, by ICD-9-CM category, 2005.....	199
V.	Changes in ICD-9-CM diagnosis codes, 1986–2004.....	201
VI.	Changes in ICD-9-CM in procedure codes, 1986–2004	207

Appendix Figure

I.	Medical abstract form for the National Hospital Discharge Survey, 2005	194
----	--	-----

Abstract

Objectives

This report presents 2005 national estimates and selected trend data on the use of nonfederal short-stay hospitals in the United States. Estimates are provided by selected patient and hospital characteristics, diagnoses, and surgical and nonsurgical procedures performed. Estimates of diagnoses and procedures are presented according to *International Classification of Diseases, Ninth Revision, Clinical Modification* codes.

Methods

The estimates are based on data collected through the National Hospital Discharge Survey. The survey has been conducted annually since 1965. In 2005, data were collected for approximately 375,000 discharges. Of the 473 eligible nonfederal short-stay hospitals in the sample, 444 (94 percent) responded to the survey.

Results

An estimated 34.7 million discharges from nonfederal short-stay hospitals occurred in 2005. Discharges used 165.9 million days of care and had an average length of stay of 4.8 days. Persons 65 years and over accounted for 38 percent of the hospital discharges and 44 percent of the days of care. The proportion of discharges whose status was described as routine discharge or discharged to the patient's home declined with age, from 91 percent for inpatients under 45 years of age to 41 percent for those 85 years and over. Hospitalization for malignant neoplasms decreased from 1990–2005. The hospitalization rate for asthma was the highest for children under 15 years of age and those 65 years of age and over. The rate was lowest for those 15–44 years of age. Thirty-eight percent of hospital discharges had no procedures performed, whereas 12 percent had four or more procedures performed. An episiotomy was performed during a majority of vaginal deliveries in 1980 (64 percent), but by 2005, it was performed during less than one of every five vaginal deliveries (19 percent).

Keywords: hospitalization • inpatient • diagnoses • procedures

National Hospital Discharge Survey: 2005 Annual Summary With Detailed Diagnosis and Procedure Data

by Carol J. DeFrances, Ph.D.; Karen A. Cullen, Ph.D., M.P.H.; and Lola Jean Kozak, Ph.D., Division of Health Care Statistics

Highlights

Patient and Hospital Characteristics

- There were an estimated 34.7 million discharges from nonfederal short-stay hospitals in 2005. Discharges used an estimated 165.9 million days of care and were hospitalized for an average of 4.8 days.
- In 2005, children under 15 years of age comprised 21 percent of the U.S. population but accounted for only 7 percent of hospital discharges and days of care. In contrast, persons 65 years of age and over comprised 12 percent of the U.S. population but accounted for 38 percent of hospital discharges and 44 percent of days of care ([Figure 1](#)).
- The average length of a hospital stay ranged from 4.2 days in the Midwest to 5.4 days in the Northeast.
- Discharges 15–44 years of age made up 56 percent of Medicaid discharges and 56 percent of discharges in the self-pay category.

but they were only 31 percent of total discharges.

- Nonprofit hospitals treated 76 percent of discharges from short-stay hospitals in the country as a whole, but the proportion varied from 58 percent in the South region to 95 percent in the Northeast region.
- The proportion of discharges whose status was described as a routine discharge or discharged to the patient's home declined with age, from 91 percent for inpatients under 45 years of age to 41 percent for those 85 years of age and over ([Figure 2](#)).

Diagnoses

- In 2005, diseases of the circulatory system accounted for 6.2 million hospitalizations, of which 4.2 million were for heart disease. Childbirth was the first-listed diagnosis for 4.0 million hospitalizations, diseases of the respiratory system for 3.7 million, and diseases of the digestive system for 3.4 million.
- During the period from 1990–2005, the rate of hospitalization for malignant neoplasms for inpatients

This report was prepared in the Division of Health Care Statistics. Rong Cai, in the Technical Services Branch, did the computer programming for the report and produced the estimated parameters for approximate relative standard error equations. Alexander Schwartzman, contractor with Lockheed Martin Information Technology, converted the ASCII tables to Excel. This report was edited by Demarius V. Miller and Megan M. Cox, CDC/CCHIS/NCHM/Division of Creative Services, Writer-Editor Services Branch, and typeset by Annette F. Holman, CDC/CCHIS/NCHM/Division of Creative Services. Graphics were produced by Jarmila G. Ogburn, CDC/CCHIS/NCHM/Division of Creative Services.




Figure 1. Percent distribution of hospital discharges, days of care, and the civilian population, by age: United States, 2005




Figure 2. Percentage of inpatients with a routine discharge or discharged home, by age: United States, 2005

65 years of age and over decreased 37 percent, but the rate remained significantly higher than the rates for other age groups. The rates for inpatients under 45 years of age and those aged 45–64 years also decreased during the same period (Figure 3).

- The hospitalization rate for asthma was highest for children under 15 years of age and for those aged 65

years and over. The rate was the lowest for those 15–44 years of age (Figure 4).

- The average annual number of discharges with a first-listed diagnosis of injury was 1,934,000 in 2003–2005. Using the Barell injury diagnosis matrix, fracture was the most common type of injury reported (1,033,000), and the most common site of fracture was lower

extremity, including the hip (590,000).

Procedures

- In 2005, 26.6 million surgical procedures and 18.4 million nonsurgical procedures were performed on hospitalized patients.
- Thirty-eight percent of hospital discharges had no procedures performed, whereas 12 percent had four or more procedures performed (Figure 5).
- Females had 6.9 million obstetrical procedures, including 1.3 million cesarean sections and 1.3 million repairs of current obstetric laceration.
- Almost 7 million inpatient operations were performed on the cardiovascular system in 2005, including 1.2 million cardiac catheterizations, and 5.6 million operations were performed on the digestive system, including 1.1 million endoscopies of the small intestine.

Deliveries and Newborn Infants

- Approximately 4.0 million women were hospitalized for delivery in 2005, and their average stay was 2.6 days.
- Cesarean sections per 100 deliveries increased from 20.8 in 1995 to 31.3 in 2005, and the rate of primary cesareans per 100 deliveries without a previous cesarean increased from 15.5 to 21.7 during this period.
- An episiotomy was performed during a majority of vaginal deliveries in 1980 (64 percent), but by 2005, it was performed during less than one of every five vaginal deliveries (19 percent) (Figure 6).
- In 2005, 45 percent of newborn infants had at least one illness or risk-related diagnosis. Common diagnoses were perinatal jaundice (798,000), respiratory conditions (437,000), disorders related to prematurity (315,000), and congenital anomalies (311,000).




Figure 3. Rate of hospitalization for malignant neoplasms by age: United States, 1990–2005




Figure 4. Rate of discharges with a first-listed diagnosis of asthma, by age: United States, 2005

- The proportion of male newborn infants circumcised during the birth hospitalization was 57 percent nationally, but ranged from 32 percent in the West to 79 percent in the Midwest.



Introduction

This report presents data from the 2005 National Hospital Discharge Survey (NHDS). The survey has been conducted continuously by the

Centers for Disease Control and Prevention's (CDC) National Center for Health Statistics (NCHS) since 1965. National estimates of hospital use derived from NHDS are published for each calendar year by NCHS. This report provides a summary of the 2005 data, including estimates of diagnoses and procedures by *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM) codes (1).

NHDS is the principal source for national data on the characteristics of patients discharged from nonfederal short-stay hospitals. Data from NHDS are used for evaluating the health status of the population, planning programs to improve health status, studying trends in morbidity, and carrying out research activities in the health field (2–6). The Department of Health and Human Services uses NHDS data in the development and monitoring of goals for the Healthy People 2010 health objectives. Other CDC centers, including the National Center for Injury Prevention and Control and the National Center for Chronic Disease Prevention and Health Promotion, use NHDS data to track a variety of conditions, including injuries and chronic diseases. The National Institutes of Health make extensive use of NHDS data on heart disease and cancer. The Veterans Administration and Department of Defense compare NHDS data with statistics on the inpatient care provided in their hospitals (7–8).

NHDS produces estimates of utilization by inpatients in nonfederal short-stay hospitals. Data on ambulatory medical care services are collected in other NCHS surveys. Data on ambulatory surgery performed in hospitals and freestanding surgery centers were collected in the National Survey of Ambulatory Surgery from 1994–1996 and in 2006 (9). The 2006 data will be available in early 2008. The National Ambulatory Medical Care Survey obtains information on visits to physicians' offices (10). The National Hospital Ambulatory Medical Care Survey collects data on visits to hospital outpatient and emergency departments (11,12). Information about long-term care is collected in the National Nursing

**Figure 5. Distribution of discharges by number of procedures: United States, 2005****Figure 6. Rate of episiotomy for females with deliveries: United States, 1980–2005**

Home Survey (13) and the National Home and Hospice Care Survey (14). NCHS will conduct the National Survey of Residential Care Facilities for the first time in 2008.

Methods

Data Source

NHDS data are collected from a sample of inpatient records acquired from a national probability sample of hospitals. Because persons with multiple discharges during the year can be sampled more than once, NHDS produces estimates for discharges, not persons. General hospitals, children's general hospitals, or hospitals with an average length of stay of fewer than 30 days for all patients are eligible for inclusion in the survey. Federal, military, and Department of Veterans Affairs hospitals, as well as hospital units of institutions (such as prison hospitals) and hospitals with fewer than six beds staffed for patient use, are excluded.

From 1988 until 2003, the hospital sampling frame for NHDS was constructed from the SMG Hospital Market Database (15). Beginning in 2003, the sampling frame was constructed from products of VeriSpan, L.L.C., specifically their "Healthcare Market Index," updated May 15, 2003, and their "Hospital Market Profiling Solution," second quarter, 2003 (16). These two products were formerly known as the SMG Hospital Market Database. The sampling frame and sample are updated every three years to include hospitals that opened or changed their eligibility status since the previous update (17).

Since 1988, NHDS has used a modified three-stage sample design. Units selected at the first stage consist of either hospitals or geographic areas, such as counties, groups of counties, or metropolitan statistical areas in the 50 states and the District of Columbia. Within sampled geographic areas, additional hospitals are selected. Finally, at the last stage, discharges are selected

within sampled hospitals using systematic random sampling.

For 2005, the sample consisted of 501 hospitals, of which 28 were out of scope because they had gone out of business or were otherwise ineligible to be in the NHDS universe. Of the 473 in-scope (eligible) hospitals, 444 (94 percent) responded to the survey. Data were collected for approximately 375,000 discharges from the responding hospitals.

The survey uses two data collection procedures. One is a manual system in which sample selection and transcription of information from medical records are performed by hospital staff or by staff of the U.S. Census Bureau on behalf of NCHS. For the other data collection procedure, NCHS purchases electronic files containing medical record data from commercial organizations, state data systems, hospitals, or hospital associations. Discharges from these files are systematically sampled.

Approximately 44 percent of respondent hospitals provided data through this automated system for the 2005 survey.

The NHDS medical abstract form ([Figure I](#)) and the automated data contain items that relate to the personal characteristics of the patient. These include birth date (converted to age), sex, race, ethnicity, marital status, ZIP Code, and expected sources of payment. The form also includes administrative items, such as admission and discharge dates, admission type and source, and discharge status. The medical information includes final medical diagnoses, procedures performed, and dates of surgery. Medical data are coded according to ICD-9-CM (1). The date of birth and ZIP Code are confidential information and are not available to the public, though they may be used, with privacy safeguards, through the NCHS Research Data Center.

For the manual data collection in NHDS, an ongoing quality control program is undertaken for the medical coding and data entry from abstracts to electronic form. Approximately 10 percent of abstracts are independently recoded, with discrepancies resolved by a chief coder. The overall error rate for records manually coded for the 2005 survey year was 0.3 percent for medical

(ICD-9-CM) coding and data entry and 0.2 percent for demographic coding and data entry.

Estimation

Because they are sample data, the survey data must be inflated or weighted to produce national estimates. The estimation procedure produces essentially unbiased national estimates and has three basic components: inflation by reciprocals of the probabilities of sample selection, adjustment for nonresponse, and population weighting ratio adjustments.

These three components of the final weight are described in more detail in another report (17). Information about the standard errors of statistics for the 2005 NHDS is found in [Table II](#).

Use of Tables

Sociodemographic information about patients and utilization estimates by hospital characteristics is shown in [Tables 1–8](#). Diagnostic data are included in [Tables 9–25](#) and information about procedures is found in [Tables 26–35](#). [Tables 36–44](#) present data on women with deliveries and newborn infants. Estimates of diagnoses, days of care by diagnosis, and procedures are shown by ICD-9-CM code in [Tables 45–48](#). [Figures 1–6](#) illustrate highlights from the survey data. Definitions of terms used in these tables and figures are presented in the appendix.

Diagnostic and procedure data are presented by chapter of the ICD-9-CM with frequently occurring broad and specific subcategories within chapters also shown (see [Tables 11–22](#) and [30–35](#)). The diagnostic categories shown in [Table 9](#) are the same as the subcategories within chapters shown in [Tables 11–18](#). Categories shown in [Table 10](#) are the subcategories in [Tables 11–18](#), with estimates of 100,000 or more for patients 65 years of age and over.

The procedure categories shown in [Table 27](#) are the same as the subcategories in [Table 30–35](#), except that they are divided into surgical and nonsurgical components. The categories in [Table 28](#) are the subcategories in

[Tables 30–35](#) for patients 65 years of age and over divided into surgical and nonsurgical components with estimates of 100,000 or more.

In 2002, the ICD-9-CM Coordination and Maintenance Committee created Chapter 00, “Procedures Not Elsewhere Classified” to provide additional codes for procedures. Beginning with the 2003 NHDS, this chapter is included in the miscellaneous diagnostic and therapeutic procedures and new technologies category shown in procedure [Tables 30–35](#).

Diagnoses assigned ICD-9-CM codes E800–E999 (supplementary classification of external causes of injury and poisoning) are not included in this report. However, detailed information about injury diagnoses and E codes has been published (18).

Injury data are presented in this report using the Barell injury diagnosis matrix ([Table 24](#)). This matrix is a two-dimensional array of ICD-9-CM codes (excluding external cause codes) that groups injuries by body site and type of injury. The codes for the matrix are shown in [Table III](#). The matrix provides a standard format for reporting injury data that improves the usefulness and comparability of epidemiological and clinical analyses (19). Data categorized using the Barell matrix are included in this report to provide a national benchmark for states and localities.

[Table 29](#) compares the number of coronary artery bypass grafts with the number of inpatients who receive these grafts. More than one code is often assigned to a patient to indicate the number and type of bypasses performed during a single operation. This results in a much larger number of procedures than the number of inpatients who have the procedures.

Beginning with the 2003 NHDS data, estimates for the procedure of “Insertion of coronary artery stent(s)” are shown separately from estimates for “Balloon angioplasty of coronary artery or coronary atherectomy” ([Tables 27–28](#) and [30–35](#)). Previously, these were combined in the category “Removal of coronary artery obstruction and insertion of stent(s).”

Tables 8 and **25** show death rates per 100 hospital discharges. These death rates are different from mortality rates published regularly by NCHS (20), which are ratios of deaths to the population. The mortality rates include deaths that occur in hospitals and other settings and are for the underlying cause of death, which may differ from the first-listed diagnosis of a patient who dies in the hospital.

Data for newborn infants are included only in **Tables 41–44**. Because estimates of newborn infants are based on a sample and because they do not include out-of-hospital births, these estimates may not agree with data on births published in the *National Vital Statistics Reports* (21), which are based on birth certificate data.

Tables 45–47 provide information by three-digit, four-digit, and five-digit ICD–9–CM diagnostic codes, and **Table 48** shows two-, three-, and four-digit procedure codes. Only ICD–9–CM code numbers appear in **Tables 45–48**. Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD–9–CM (1). If the estimate of the total for the code is 5,000 or more, a specific diagnosis or procedure code appears in **Tables 45, 47, and 48**. The codes that appear in **Table 46** for days of care are identical to those in **Table 45** for first-listed diagnoses.

Estimates in all the tables are rounded to the nearest thousand, and totals may include data for categories that are not shown individually in the tables. For these reasons, figures within tables do not always add to the totals. Rates, percentages, and average lengths of stay are calculated from unrounded figures and may not precisely agree with measures calculated from rounded data.

Because of low reliability, estimates are not presented in **Tables 1–44** if they have relative standard errors of more than 30 percent or are based on fewer than 30 records; instead, an asterisk (*) appears in the tables. Estimates that are based on 30–59 records are presented but are preceded by an asterisk (*) to indicate that they also have low reliability.

Because of the large number of estimates in **Tables 45–48**, more generalized criteria for reliability were used. An asterisk (*) is shown in **Tables 45, 47, and 48** if the estimate of the diagnosis or procedure is less than 5,000, and an asterisk (*) is shown in **Table 46** for estimates of days of care derived from fewer than 5,000 estimated discharges. These estimates generally have a relative standard error of more than 30 percent or a sample size of less than 30. Estimates of diagnoses and procedures of less than 9,000 are preceded by an asterisk (*) because they are generally based on fewer than 60 records. Days of care estimates in **Table 46** that are derived from fewer than 9,000 estimated discharges are also preceded by an asterisk (*).

The population estimates used to compute rates are shown in **Table I**. These are postcensal estimates for July 1, 2005, that are based on the 2000 census.

A weighted least squares regression method (22) was used to test the significance of the trends shown in **Figures 3 and 6**. Differences mentioned between NHDS estimates were tested using the two-sided *t*-test. Terms that express differences such as higher, lower, largest, smallest, leading, increased or decreased were used only when the differences were statistically significant at the 0.05 level.

References

1. National Center for Health Statistics. International Classification of Diseases, Ninth Revision, Clinical Modification. 6th ed. Washington, DC: Public Health Service. Available from: <http://www.cdc.gov/nchs/icd9.htm#ICD-9-CM>. 2004.
2. Bewtra M, Su C, Lewis JD. Trends in hospitalization rates for inflammatory bowel disease in the United States [online]. Clin Gastroenterol Hepatol 5(5):597–601.e1. 2007.
3. Stein PD, Beemath A, Meyers FA, Olson RE. Pulmonary embolism and deep venous thrombosis in hospitalized adults with chronic obstructive pulmonary disease. J Cardiovasc Med 8(4):253–7. 2007.
4. Bundy DG. Hospitalizations with primary versus secondary discharge diagnoses of asthma: Implications for pediatric asthma surveillance. J Pediatr 150(4):446–9.e1. 2007.
5. Klevens RM, Edwards JR, Richards CL Jr, Horan TC, Gaynes RP, Pollock DA, et al. Estimating health care-associated infections and deaths in U.S. hospitals, 2002. Public Health Rep 122(2):160–6. 2007.
6. Stanley A, DeLia D, Cantor JC. Racial disparity and technology diffusion: The case of cardioverter defibrillator implants. J Natl Med Assoc 99(3):201–7. 2007.
7. Rosenthal GE, Kaboli PJ, Barnett MJ. Differences in length of stay in Veterans Health Administration and other United States hospitals: Is the gap closing? Med Care 41(8):882–94. 2003.
8. Jackson JL, Cheng EY, Jones DL, Meyer G. Comparison of discharge diagnoses and inpatient procedures between military and civilian health care systems. Mil Med 164(10):701–4. 1999.
9. Owings MF, Kozak LJ. Ambulatory and inpatient procedures in the United States, 1996. National Center for Health Statistics. Vital Health Stat 13(139). Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_139.pdf. 1998.
10. Cherry DK, Woodwell DA, Rechtsteiner EA. National Ambulatory Medical Care Survey: 2005 summary. Advance data from vital and health statistics; no 387. Hyattsville, MD: National Center for Health Statistics. Available from: <http://www.cdc.gov/nchs/data/ad/ad387.pdf>. 2007.
11. Middleton K, Hing E, Xu J. National Hospital Ambulatory Medical Care Survey: 2005 outpatient department summary. Advance data from vital and health statistics; no 389. Hyattsville, MD: National Center for Health Statistics. Available from: <http://www.cdc.gov/nchs/data/ad/ad389.pdf>. 2007.
12. Nawar EW, Niska RW, Xu J. National Hospital Ambulatory Medical Care Survey: 2005 emergency department summary. Advance data from vital and health statistics; no 386. Hyattsville, MD: National Center for Health Statistics. Available from: <http://www.cdc.gov/nchs/data/ad/ad386.pdf>. 2007.
13. Jones A. The National Nursing Home Survey: 1999 summary. National Center for Health Statistics. Vital

- Health Stat 13(152). Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_152.pdf. 2002.
14. Haupt BJ. Characteristics of hospice care discharges and their length of service: United States, 2000. National Center for Health Statistics. Vital Health Stat 13(154). Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_154.pdf. 2003.
15. SMG Marketing Group, Inc. Hospital Market Database. Chicago, IL: Healthcare Information Specialists. April 1987, April 1991, April 1994, April 1997, April 2000.
16. Verispan, L.L.C. Healthcare Market Index, updated May 15, 2003, and Hospital Market Profiling Solution, second quarter, 2003. Available from: <http://www.verispan.com>.
17. Dennison CF, Pokras R. Design and operation of the National Hospital Discharge Survey: 1988 redesign. National Center for Health Statistics. Vital Health Stat 1(39). Available from: http://www.cdc.gov/nchs/data/series/sr_01/sr01_039.pdf. 2000.
18. Heinen M, Hall MJ, Boudreault MA, Fingerhut LA. National trends in injury hospitalizations, 1979–2001. Hyattsville, MD: National Center for Health Statistics. Available from: <http://www.cdc.gov/nchs/data/injury/InjuryChartbook79-01.pdf>. 2005.
19. Barell V, Aharonson-Daniel L, Fingerhut LA, et al. An introduction to the Barell body region by nature of injury diagnosis matrix. Inj Prev 8:91–6. 2002.
20. Minino AM, Heron M, Murphy SL, Kochanek KD. Deaths: Final data from 2004. Health E-Stats. Available from: <http://www.cdc.gov/nchs/products/pubs/pubd/hestats/finaldeaths04/finaldeaths04.htm>. 2006.
21. Hamilton BE, Martin JA, Ventura SJ, et al. Births: Preliminary data for 2005. National vital statistics reports 55(11). Hyattsville, MD: National Center for Health Statistics. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr55/nvsr55_11.pdf. 2006.
22. Gillum BS, Graves EJ, Kozak LJ. Trends in hospital utilization: United States, 1988–92. National Center for Health Statistics. Vital Health Stat 13(124). Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_124.pdf. 1996.
23. Research Triangle Institute. SUDAAN language manual, release 9.0. Research Triangle Park, NC: Research Triangle Institute. 2004.
24. Hansen MH, Hurwitz WN, Madow WG. Sample survey methods and theory, vol 1. Wiley Classics Library ed. New York, NY: John Wiley & Sons. 1993.
25. Cochran WG. Sampling techniques. 3d ed. New York: John Wiley & Sons. 1977.

Table 1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Age	1970	1980	1985	1990	1995	2000	2005
Number of discharges in thousands							
All ages	29,127	37,832	35,056	30,788	30,722	31,706	34,667
Under 15 years	3,873	3,672	2,972	2,412	2,405	2,383	2,431
15–44 years	12,664	15,635	13,966	11,799	10,593	9,969	10,659
45–64 years	6,693	8,660	7,610	6,244	6,168	6,958	8,349
65 years and over	5,897	9,864	10,508	10,333	11,556	12,396	13,228
65–74 years	3,163	4,943	5,011	4,689	4,832	4,678	4,900
75–84 years	2,099	3,611	3,969	3,949	4,590	5,119	5,394
85 years and over	635	1,310	1,528	1,694	2,134	2,599	2,934
Rate of discharges per 1,000 population ¹							
All ages	144.3	167.7	148.4	122.3	115.7	112.8	117.4
Under 15 years	66.8	71.6	57.7	43.1	40.4	39.5	40.0
15–44 years	154.6	150.1	125.0	99.3	87.8	80.9	85.3
45–64 years	159.6	194.8	170.8	135.5	118.5	111.4	114.7
65 years and over	293.3	383.7	369.8	334.1	347.7	353.4	359.6
65–74 years	253.2	315.8	297.3	261.6	260.0	254.6	262.9
75–84 years	339.5	463.9	446.4	395.7	415.6	412.4	413.2
85 years and over	443.7	576.4	572.9	560.6	592.5	605.1	575.8
Number of days of care in thousands							
All ages	226,445	274,508	226,217	197,422	164,627	155,857	165,925
Under 15 years	18,189	16,191	13,554	11,655	10,715	10,734	11,371
15–44 years	71,605	81,951	67,397	54,062	40,825	36,593	39,786
45–64 years	62,419	71,008	53,541	42,153	34,207	34,443	41,619
65 years and over	74,232	105,358	91,726	89,552	78,880	74,086	73,149
65–74 years	37,873	49,253	41,090	37,422	31,310	26,438	26,068
75–84 years	27,633	40,355	36,024	35,926	31,974	31,525	30,228
85 years and over	8,726	15,750	14,612	16,204	15,597	16,123	16,853
Rate of days of care per 1,000 population ¹							
All ages	1,121.6	1,216.7	957.7	784.0	620.2	554.6	562.1
Under 15 years	313.9	315.7	263.0	208.4	179.9	178.0	187.3
15–44 years	874.0	786.8	603.3	454.9	338.4	297.1	318.5
45–64 years	1,488.8	1,596.9	1,201.6	914.4	657.5	551.5	571.7
65 years and over	3,691.9	4,098.4	3,228.0	2,895.6	2,373.7	2,111.9	1,988.3
65–74 years	3,031.6	3,147.0	2,437.3	2,087.8	1,684.7	1,439.0	1,398.5
75–84 years	4,469.2	5,183.7	4,052.2	3,599.5	2,894.7	2,539.8	2,315.5
85 years and over	6,102.1	6,932.1	5,478.7	5,361.7	4,330.6	3,753.8	3,307.1
Average length of stay in days							
All ages	7.8	7.3	6.5	6.4	5.4	4.9	4.8
Under 15 years	4.7	4.4	4.6	4.8	4.5	4.5	4.7
15–44 years	5.7	5.2	4.8	4.6	3.9	3.7	3.7
45–64 years	9.3	8.2	7.0	6.8	5.5	5.0	5.0
65 years and over	12.6	10.7	8.7	8.7	6.8	6.0	5.5
65–74 years	12.0	10.0	8.2	8.0	6.5	5.7	5.3
75–84 years	13.2	11.2	9.1	9.1	7.0	6.2	5.6
85 years and over	13.7	12.0	9.6	9.6	7.3	6.2	5.7

¹Rates were calculated using the U.S. Census Bureau estimates of the civilian population. Rates for 1990 and 1995 were based on population estimates adjusted for the net underenumeration in the 1990 census. Rates for 2000 and 2005 were calculated using 2000-based postcensal estimates. Rates for 2000 differ from the ones shown in earlier reports because they were calculated using revised population estimates.

Table 2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex and age	Discharges			Days of care			
	Number in thousands	Percent distribution	Rate per 1,000 population ¹	Number in thousands	Percent distribution	Rate per 1,000 population ¹	Average length of stay in days
Both sexes							
All ages	34,667	100.0	117.4	165,925	100.0	562.1	4.8
Under 15 years	2,431	7.0	40.0	11,371	6.9	187.3	4.7
Under 1 year	800	2.3	194.9	4,982	3.0	1,213.1	6.2
1–4 years	696	2.0	43.0	2,195	1.3	135.5	3.2
5–14 years	934	2.7	23.1	4,194	2.5	103.8	4.5
15–44 years	10,659	30.7	85.3	39,786	24.0	318.5	3.7
15–19 years	1,171	3.4	55.9	4,712	2.8	224.8	4.0
20–24 years	1,903	5.5	92.2	6,137	3.7	297.5	3.2
25–34 years	4,001	11.5	100.8	13,636	8.2	343.4	3.4
35–44 years	3,584	10.3	82.2	15,301	9.2	350.8	4.3
45–64 years	8,349	24.1	114.7	41,619	25.1	571.7	5.0
45–54 years	4,093	11.8	96.4	19,995	12.1	471.1	4.9
55–64 years	4,257	12.3	140.2	21,624	13.0	712.4	5.1
65 years and over	13,228	38.2	359.6	73,149	44.1	1,988.3	5.5
65–74 years	4,900	14.1	262.9	26,068	15.7	1,398.5	5.3
75–84 years	5,394	15.6	413.2	30,228	18.2	2,315.5	5.6
85 years and over	2,934	8.5	575.8	16,853	10.2	3,307.1	5.7
Male							
All ages	13,902	100.0	95.9	72,186	100.0	498.0	5.2
Under 15 years	1,346	9.7	43.3	6,443	8.9	207.5	4.8
Under 1 year	458	3.3	218.1	2,894	4.0	1,377.4	6.3
1–4 years	385	2.8	46.5	1,317	1.8	159.0	3.4
5–14 years	502	3.6	24.3	2,232	3.1	107.9	4.4
15–44 years	2,864	20.6	45.5	13,989	19.4	222.0	4.9
15–19 years	339	2.4	31.7	1,749	2.4	163.1	5.2
20–24 years	359	2.6	34.2	1,764	2.4	167.8	4.9
25–34 years	813	5.8	40.5	3,908	5.4	194.9	4.8
35–44 years	1,352	9.7	62.3	6,568	9.1	302.3	4.9
45–64 years	4,076	29.3	114.9	20,487	28.4	577.3	5.0
45–54 years	1,957	14.1	93.8	9,759	13.5	467.8	5.0
55–64 years	2,120	15.2	144.9	10,728	14.9	733.6	5.1
65 years and over	5,616	40.4	364.4	31,267	43.3	2,028.6	5.6
65–74 years	2,339	16.8	274.3	12,370	17.1	1,450.3	5.3
75–84 years	2,317	16.7	438.8	13,255	18.4	2,510.7	5.7
85 years and over	960	6.9	598.4	5,642	7.8	3,517.9	5.9
Female							
All ages	20,766	100.0	138.2	93,739	100.0	623.9	4.5
Under 15 years	1,085	5.2	36.6	4,928	5.3	166.2	4.5
Under 1 year	342	1.6	170.6	2,088	2.2	1,041.0	6.1
1–4 years	311	1.5	39.3	878	0.9	111.0	2.8
5–14 years	432	2.1	21.9	1,962	2.1	99.5	4.5
15–44 years	7,795	37.5	125.9	25,797	27.5	416.7	3.3
15–19 years	832	4.0	81.2	2,962	3.2	289.4	3.6
20–24 years	1,543	7.4	152.6	4,373	4.7	432.3	2.8
25–34 years	3,188	15.4	162.2	9,728	10.4	494.9	3.1
35–44 years	2,231	10.7	101.9	8,733	9.3	398.9	3.9
45–64 years	4,273	20.6	114.5	21,132	22.5	566.4	4.9
45–54 years	2,136	10.3	99.0	10,236	10.9	474.3	4.8
55–64 years	2,137	10.3	135.9	10,896	11.6	692.7	5.1
65 years and over	7,613	36.7	356.1	41,882	44.7	1,959.2	5.5
65–74 years	2,561	12.3	253.3	13,699	14.6	1,354.9	5.3
75–84 years	3,077	14.8	395.8	16,973	18.1	2,183.0	5.5
85 years and over	1,975	9.5	565.4	11,211	12.0	3,210.4	5.7

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population ¹	Number in thousands	Rate per 1,000 population ¹	
Both sexes					
All ages:					
United States	34,667	117.4	165,925	562.1	4.8
Northeast	7,192	131.8	38,730	709.6	5.4
Midwest	7,948	120.6	33,741	512.1	4.2
South	12,907	120.8	62,797	587.9	4.9
West	6,621	97.5	30,656	451.5	4.6
Under 15 years:					
United States	2,431	40.0	11,371	187.3	4.7
Northeast	407	38.9	2,205	210.6	5.4
Midwest	*	*	*	*	3.9
South	960	43.3	4,285	193.4	4.5
West	*	*	*	*	5.1
15–44 years:					
United States	10,659	85.3	39,786	318.5	3.7
Northeast	2,007	88.7	8,885	392.8	4.4
Midwest	2,244	81.0	7,379	266.4	3.3
South	4,286	94.9	16,465	364.4	3.8
West	2,122	72.1	7,056	239.9	3.3
45–64 years:					
United States	8,349	114.7	41,619	571.7	5.0
Northeast	1,762	125.5	9,509	677.2	5.4
Midwest	1,834	111.2	7,950	482.1	4.3
South	3,204	122.5	16,738	639.8	5.2
West	1,550	96.2	7,423	460.9	4.8
65 years and over:					
United States	13,228	359.6	73,149	1988.3	5.5
Northeast	3,016	404.7	18,131	2433.2	6.0
Midwest	3,403	403.0	16,573	1962.8	4.9
South	4,456	334.7	25,309	1900.8	5.7
West	2,353	310.4	13,136	1732.9	5.6
Male					
All ages:					
United States	13,902	95.9	72,186	498.0	5.2
Northeast	3,018	114.0	17,506	661.1	5.8
Midwest	3,160	97.6	14,136	436.6	4.5
South	5,054	96.7	26,758	511.9	5.3
West	2,670	78.9	13,786	407.5	5.2
Under 15 years:					
United States	1,346	43.3	6,443	207.5	4.8
Northeast	224	41.8	1,232	230.0	5.5
Midwest	*	*	*	*	3.8
South	529	46.7	2,423	213.8	4.6
West	*	*	*	*	5.4
15–44 years:					
United States	2,864	45.5	13,989	222.0	4.9
Northeast	676	59.8	3,812	337.0	5.6
Midwest	610	43.6	2,418	172.7	4.0
South	1,112	49.1	5,667	250.2	5.1
West	466	31.0	2,092	139.1	4.5
45–64 years:					
United States	4,076	114.9	20,487	577.3	5.0
Northeast	869	128.2	4,768	703.1	5.5
Midwest	877	108.3	3,743	462.3	4.3
South	1,557	122.8	8,217	648.3	5.3
West	773	97.5	3,758	473.8	4.9

See footnotes at end of table.

Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population ¹	Number in thousands	Rate per 1,000 population ¹	
65 years and over:					
United States	5,616	364.4	31,267	2028.6	5.6
Northeast	1,248	412.2	7,693	2540.6	6.2
Midwest	1,419	406.1	7,007	2004.4	4.9
South	1,857	330.7	10,450	1861.4	5.6
West	1,091	333.3	6,117	1867.8	5.6
Female					
All ages:					
United States	20,766	138.2	93,739	623.9	4.5
Northeast	4,174	148.5	21,225	755.2	5.1
Midwest	4,788	142.9	19,605	585.0	4.1
South	7,852	144.0	36,039	660.7	4.6
West	3,951	116.0	16,870	495.1	4.3
Under 15 years:					
United States	1,085	36.6	4,928	166.2	4.5
Northeast	183	35.8	973	190.2	5.3
Midwest	*	*	*	*	4.1
South	431	39.8	1,862	172.0	4.3
West	*	*	*	*	4.7
15–44 years:					
United States	7,795	125.9	25,797	416.7	3.3
Northeast	1,330	117.7	5,073	448.8	3.8
Midwest	1,634	119.3	4,962	362.2	3.0
South	3,175	140.9	10,798	479.3	3.4
West	1,655	115.2	4,965	345.4	3.0
45–64 years:					
United States	4,273	114.5	21,132	566.4	4.9
Northeast	893	123.0	4,740	653.0	5.3
Midwest	957	114.0	4,207	501.2	4.4
South	1,647	122.1	8,520	631.8	5.2
West	776	95.0	3,664	448.4	4.7
65 years and over:					
United States	7,613	356.1	41,882	1959.2	5.5
Northeast	1,768	399.6	10,438	2359.6	5.9
Midwest	1,984	400.9	9,567	1933.4	4.8
South	2,600	337.6	14,859	1929.5	5.7
West	1,262	293.1	7,019	1630.4	5.6

* Figure does not meet standards of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO or PPO ²	Blue Cross Blue Shield and other private insurance	Self-pay	Other payments and no charge
Number of discharges in thousands								
United States								
All	34,667	13,788	5,836	740	4,793	6,412	1,500	1,030
Male	13,902	5,955	1,882	423	1,695	2,521	729	424
Female	20,766	7,834	3,953	318	3,098	3,891	771	606
Under 15 years	2,431	*	1,107	61	399	625	52	*
15–44 years	10,659	613	3,270	310	2,164	2,729	842	491
45–64 years	8,349	1,672	1,214	290	1,692	2,444	529	318
65 years and over	13,228	11,494	245	80	538	615	77	80
Northeast								
All	7,192	2,944	990	91	1,061	1,470	258	97
Male	3,018	1,252	410	56	393	597	134	39
Female	4,174	1,693	580	35	668	873	125	58
Under 15 years	407	*	137	*	96	135	14	*
15–44 years	2,007	121	523	37	426	592	138	44
45–64 years	1,762	338	284	30	360	552	84	30
65 years and over	3,016	2,483	46	*	178	191	*	*
Midwest								
All	7,948	3,659	1,110	*	1,054	1,645	253	57
Male	3,160	1,547	357	68	370	634	128	31
Female	4,788	2,112	753	*	684	1,011	126	26
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	2,244	140	681	*	493	698	138	24
45–64 years	1,834	380	240	39	432	618	89	17
65 years and over	3,403	3,139	35	*	61	122	*	*
South								
All	12,907	4,917	2,346	343	1,466	2,411	758	*
Male	5,054	2,090	678	195	504	934	354	*
Female	7,852	2,827	1,667	148	962	1,476	405	*
Under 15 years	960	*	488	18	104	208	18	*
15–44 years	4,286	281	1,345	140	711	1,073	430	*
45–64 years	3,204	684	442	156	507	945	282	*
65 years and over	4,456	3,947	70	29	144	184	29	*
West								
All	6,621	2,268	1,390	189	1,212	886	230	368
Male	2,670	1,066	437	105	429	356	114	127
Female	3,951	1,202	954	85	783	530	116	240
Under 15 years	*	*	*	*	*	74	9	*
15–44 years	2,122	71	721	84	533	366	137	176
45–64 years	1,550	271	248	64	393	329	74	*
65 years and over	2,353	1,925	*	*19	154	118	*11	*
Number of days of care in thousands								
United States								
All	165,925	77,640	27,448	3,265	18,328	25,612	6,046	4,353
Male	72,186	33,815	11,505	1,962	7,235	10,960	3,072	2,006
Female	93,739	43,825	15,943	1,304	11,093	14,652	2,974	2,347
Under 15 years	11,371	*	5,877	*	1,619	2,636	168	*
15–44 years	39,786	3,858	12,331	1,210	6,791	9,388	2,989	1,893
45–64 years	41,619	9,971	7,665	1,359	7,143	10,458	2,438	1,472
65 years and over	73,149	63,764	1,575	449	2,775	3,130	450	*

See footnotes at end of table.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO or PPO ²	Blue Cross Blue Shield and other private insurance	Self-pay	Other payments and no charge
Northeast								
All	38,730	18,117	5,848	463	4,399	6,329	1,299	509
Male	17,506	7,890	2,998	304	1,757	2,730	725	192
Female	21,225	10,227	2,850	159	2,642	3,599	574	*
Under 15 years	2,205	*	953	*	*	*	53	*
15–44 years	8,885	863	2,603	195	1,440	2,148	606	*
45–64 years	9,509	2,161	1,928	145	1,619	2,488	499	147
65 years and over	18,131	15,085	*	*	941	1,026	*	*
Midwest								
All	33,741	17,965	4,497	*	3,778	5,723	889	230
Male	14,136	7,688	1,821	*	1,411	2,273	448	124
Female	19,605	10,277	2,676	*	2,367	3,450	441	106
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	7,379	627	2,339	*	1,550	2,109	401	*
45–64 years	7,950	1,956	1,244	*	1,737	2,368	331	*
65 years and over	16,573	15,376	167	*	282	516	*	*
South								
All	62,797	28,856	10,522	1,532	5,588	10,201	2,972	*
Male	26,758	12,319	3,918	916	2,220	4,451	1,429	*
Female	36,039	16,538	6,604	616	3,368	5,750	1,543	*
Under 15 years	4,285	*	2,252	57	394	960	52	*
15–44 years	16,465	1,950	4,924	557	2,223	3,964	1,530	*
45–64 years	16,738	4,397	2,931	772	2,183	4,267	1,236	638
65 years and over	25,309	22,484	415	145	788	1,009	153	*
West								
All	30,656	12,701	6,582	817	4,563	3,360	886	1,299
Male	13,786	5,919	2,768	476	1,848	1,505	471	556
Female	16,870	6,783	3,814	341	2,715	1,854	415	743
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	7,056	417	2,465	287	1,579	1,167	452	509
45–64 years	7,423	1,457	1,563	289	1,603	1,335	372	*
65 years and over	13,136	10,819	*	*	764	578	*37	*
United States								
All	4.8	5.6	4.7	4.4	3.8	4.0	4.0	4.2
Male	5.2	5.7	6.1	4.6	4.3	4.3	4.2	4.7
Female	4.5	5.6	4.0	4.1	3.6	3.8	3.9	3.9
Under 15 years	4.7	5.3	5.3	4.1	4.1	4.2	3.3	3.8
15–44 years	3.7	6.3	3.8	3.9	3.1	3.4	3.5	3.9
45–64 years	5.0	6.0	6.3	4.7	4.2	4.3	4.6	4.6
65 years and over	5.5	5.5	6.4	5.6	5.2	5.1	5.8	5.5
Northeast								
All	5.4	6.2	5.9	5.1	4.1	4.3	5.0	5.3
Male	5.8	6.3	7.3	5.5	4.5	4.6	5.4	4.9
Female	5.1	6.0	4.9	4.5	4.0	4.1	4.6	5.5
Under 15 years	5.4	*	7.0	4.4	4.1	4.9	3.8	*3.3
15–44 years	4.4	7.1	5.0	5.2	3.4	3.6	4.4	5.5
45–64 years	5.4	6.4	6.8	4.8	4.5	4.5	5.9	4.9
65 years and over	6.0	6.1	7.9	5.5	5.3	5.4	6.3	6.4

See footnotes at end of table.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO or PPO ²	Blue Cross Blue Shield and other private insurance	Self-pay	Other payments and no charge
Midwest								
All	4.2	4.9	4.1	3.9	3.6	3.5	3.5	4.0
Male	4.5	5.0	5.1	3.9	3.8	3.6	3.5	4.0
Female	4.1	4.9	3.6	3.8	3.5	3.4	3.5	4.1
Under 15 years	3.9	*	4.8	4.9	3.1	3.5	3.4	3.2
15–44 years	3.3	4.5	3.4	3.5	3.1	3.0	2.9	4.0
45–64 years	4.3	5.1	5.2	3.9	4.0	3.8	3.7	4.7
65 years and over	4.9	4.9	4.8	*3.9	4.6	4.2	*7.5	*3.6
South								
All	4.9	5.9	4.5	4.5	3.8	4.2	3.9	4.6
Male	5.3	5.9	5.8	4.7	4.4	4.8	4.0	5.0
Female	4.6	5.8	4.0	4.2	3.5	3.9	3.8	4.2
Under 15 years	4.5	4.7	4.6	3.1	3.8	4.6	2.9	4.2
15–44 years	3.8	6.9	3.7	4.0	3.1	3.7	3.6	4.3
45–64 years	5.2	6.4	6.6	4.9	4.3	4.5	4.4	4.9
65 years and over	5.7	5.7	5.9	5.1	5.5	5.5	5.3	6.2
West								
All	4.6	5.6	4.7	4.3	3.8	3.8	3.9	3.5
Male	5.2	5.6	6.3	4.6	4.3	4.2	4.1	4.4
Female	4.3	5.6	4.0	4.0	3.5	3.5	3.6	3.1
Under 15 years	5.1	*	5.9	4.2	4.7	3.8	2.9	2.7
15–44 years	3.3	5.9	3.4	3.4	3.0	3.2	3.3	2.9
45–64 years	4.8	5.4	6.3	4.5	4.1	4.1	5.1	4.3
65 years and over	5.6	5.6	6.7	*	4.9	4.9	*3.5	4.7

* Figure does not meet standards of reliability or precision.

¹Includes discharges for which no expected source of payment was provided.²HMO is health maintenance organization, and PPO is preferred provider organization.

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, according to sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All bed sizes	6–99 beds	100–199 beds	200–299 beds	300–499 beds	500 beds or more
Number of discharges in thousands						
All	34,667	8,170	8,271	6,687	7,420	4,118
Sex						
Male	13,902	3,176	3,235	2,664	3,029	1,798
Female	20,766	4,995	5,036	4,024	4,392	2,319
Age						
Under 15 years	2,431	377	*	*	581	238
15–44 years	10,659	2,310	2,643	1,949	2,421	1,337
45–64 years	8,349	1,781	1,949	1,536	1,893	1,191
65 years and over	13,228	3,703	3,091	2,557	2,526	1,352
Region						
Northeast	7,192	1,639	1,692	1,156	1,574	1,132
Midwest	7,948	3,040	1,306	1,189	*	850
South	12,907	2,780	3,213	2,746	2,236	1,931
West	6,621	*	2,060	1,597	2,049	*
Percent distribution of discharges						
All	100.0	23.6	23.9	19.3	21.4	11.9
Sex						
Male	100.0	22.8	23.3	19.2	21.8	12.9
Female	100.0	24.1	24.3	19.4	21.2	11.2
Age						
Under 15 years	100.0	15.5	24.2	26.6	23.9	9.8
15–44 years	100.0	21.7	24.8	18.3	22.7	12.5
45–64 years	100.0	21.3	23.3	18.4	22.7	14.3
65 years and over	100.0	28.0	23.4	19.3	19.1	10.2
Region						
Northeast	100.0	22.8	23.5	16.1	21.9	15.7
Midwest	100.0	38.3	16.4	15.0	*	10.7
South	100.0	21.5	24.9	21.3	17.3	15.0
West	100.0	*	31.1	24.1	30.9	*
Number of days of care in thousands						
All	165,925	35,195	38,773	32,026	37,039	22,891
Sex						
Male	72,186	14,568	16,383	13,936	16,495	10,803
Female	93,739	20,628	22,390	18,089	20,543	12,088
Age						
Under						
15 years	11,371	1,517	*	*	2,839	1,313
15–44 years	39,786	8,491	9,238	6,755	9,198	6,104
45–64 years	41,619	7,664	9,382	7,548	9,949	7,075
65 years and over	73,149	17,523	17,837	14,338	15,052	8,400
Region						
Northeast	38,730	8,590	9,077	6,105	8,616	6,341
Midwest	33,741	11,176	5,952	5,309	*	4,109
South	62,797	12,160	14,954	13,176	11,171	11,336
West	30,656	*	8,790	7,435	10,058	*

See footnotes at end of table.

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, according to sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All bed sizes	6–99 beds	100–199 beds	200–299 beds	300–499 beds	500 beds or more
Percent distribution of days of care						
All	100.0	21.2	23.4	19.3	22.3	13.8
Sex						
Male	100.0	20.2	22.7	19.3	22.9	15.0
Female	100.0	22.0	23.9	19.3	21.9	12.9
Age						
Under 15 years	100.0	13.3	*	*	25.0	11.5
15–44 years	100.0	21.3	23.2	17.0	23.1	15.3
45–64 years	100.0	18.4	22.5	18.1	23.9	17.0
65 years and over	100.0	24.0	24.4	19.6	20.6	11.5
Region						
Northeast	100.0	22.2	23.4	15.8	22.2	16.4
Midwest	100.0	33.1	17.6	15.7	*	12.2
South	100.0	19.4	23.8	21.0	17.8	18.1
West	100.0	*	28.7	24.3	32.8	*
Average length of stay in days						
All	4.8	4.3	4.7	4.8	5.0	5.6
Sex						
Male	5.2	4.6	5.1	5.2	5.4	6.0
Female	4.5	4.1	4.4	4.5	4.7	5.2
Age						
Under 15 years	4.7	4.0	3.9	5.2	4.9	5.5
15–44 years	3.7	3.7	3.5	3.5	3.8	4.6
45–64 years	5.0	4.3	4.8	4.9	5.3	5.9
65 years and over	5.5	4.7	5.8	5.6	6.0	6.2
Region						
Northeast	5.4	5.2	5.4	5.3	5.5	5.6
Midwest	4.2	3.7	4.6	4.5	4.6	4.8
South	4.9	4.4	4.7	4.8	5.0	5.9
West	4.6	4.6	4.3	4.7	4.9	5.4

* Figure does not meet standards of reliability or precision.

Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, according to sex and age: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex and age	All	Proprietary	Government	Nonprofit
Number of discharges in thousands				
All	34,667	4,190	4,132	26,346
Percent distribution of discharges				
All	100.0	12.1	11.9	76.0
Number of days of care in thousands				
All	165,925	21,190	20,184	124,551
Percent distribution of days of care				
All	100	12.8	12.2	75.1
Average length of stay in days				
All	4.8	5.1	4.9	4.7
Age				
Under 15 years	2,431	*	272	1,880
15–44 years	10,659	1,593	1,400	7,665
45–64 years	8,349	970	1,096	6,283
65 years and over	13,228	1,348	1,363	10,518
Sex				
Male	100.0	11.4	12.1	76.4
Female	100.0	12.5	11.8	75.7
Age				
Under 15 years	100.0	*	11.2	77.3
15–44 years	100.0	14.9	13.1	71.9
45–64 years	100.0	11.6	13.1	75.2
65 years and over	100.0	10.2	10.3	79.5
Sex				
Male	72,186	8,872	9,146	54,167
Female	93,739	12,318	11,038	70,384
Age				
Under 15 years	11,371	*	1,115	9,119
15–44 years	39,786	6,389	5,719	27,678
45–64 years	41,619	5,364	5,973	30,282
65 years and over	73,149	8,300	7,377	57,472
Sex				
Male	100.0	12.3	12.7	75.0
Female	100.0	13.1	11.8	75.1
Age				
Under 15 years	100.0	*	9.8	80.2
15–44 years	100.0	16.1	14.4	69.6
45–64 years	100.0	12.9	14.4	72.8
65 years and over	100.0	11.3	10.1	78.6

* Figure does not meet standards of reliability or precision.

Table 7. Number and percent distribution of discharges by type and source of admission to short-stay hospitals, according to sex and age: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Admission type and source	All ages	Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over
Number of discharges in thousands							
All types and sources	34,667	13,902	20,766	2,431	10,659	8,349	13,228
Type of admission							
Emergency	14,131	6,528	7,603	923	3,246	3,658	6,305
Urgent	8,755	3,124	5,631	794	3,159	1,732	3,069
Elective.	8,317	2,900	5,417	493	2,946	2,082	2,796
Item not available	3,464	1,349	2,114	222	1,307	878	1,057
Source of admission							
Physician referral ¹	13,332	4,443	8,889	806	5,177	3,039	4,311
Emergency department	14,340	6,533	7,807	945	3,138	3,716	6,541
Transfer from a hospital.	1,068	544	524	146	192	257	474
Transfer from other health facility ²	485	208	277	18	59	75	333
Other sources ³	662	237	425	*	365	131	102
Item not available	4,780	1,937	2,843	453	1,728	1,131	1,467
Percent distribution							
All types and sources	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Type of admission							
Emergency	40.8	47.0	36.6	38.0	30.5	43.8	47.7
Urgent	25.3	22.5	27.1	32.7	29.6	20.7	23.2
Elective.	24.0	20.9	26.1	20.3	27.6	24.9	21.1
Item not available	10.0	9.7	10.2	9.1	12.3	10.5	8.0
Source of admission							
Physician referral ¹	38.5	32.0	42.8	33.2	48.6	36.4	32.6
Emergency department	41.4	47.0	37.6	38.9	29.4	44.5	49.4
Transfer from a hospital.	3.1	3.9	2.5	6.0	1.8	3.1	3.6
Transfer from other health facility ²	1.4	1.5	1.3	0.7	0.6	0.9	2.5
Other sources ³	1.9	1.7	2.0	*	3.4	1.6	0.8
Item not available	13.8	13.9	13.7	18.6	16.2	13.5	11.1

^{*} Figure does not meet standards of reliability or precision.¹Referral from personal physician, clinic physician, or health maintenance organization physician.²Transfer from skilled nursing facility or other health facility.³Includes admissions directed by a court of law or requested by a law enforcement agent and other sources.

Table 8. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by disposition of patient, according to sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All dispositions	Discharged alive						Status not stated	
		Routine or discharged home	Transferred to						
			Other short-term hospital	Long-term care institution	Other	Died			
Number of discharges in thousands									
All	34,667	26,541	1,514	3,074	1,973	752	*		
Sex									
Male	13,902	10,399	731	1,153	908	383	328		
Female	20,766	16,142	782	1,921	1,065	369	*		
Age									
Under 65 years	21,439	18,707	645	506	915	200	466		
Under 15 years	2,431	2,237	50	13	63	12	*		
15–44 years	10,659	9,684	214	116	383	39	222		
45–64 years	8,349	6,786	381	377	469	148	187		
65 years and over	13,228	7,834	869	2,568	1,058	552	*		
65–74 years	4,900	3,480	317	536	327	135	*		
75–84 years	5,394	3,164	364	1,061	443	232	*		
85 years and over	2,934	1,190	188	971	288	186	*		
Region									
Northeast	7,192	5,269	297	867	527	168	*		
Midwest	7,948	5,847	279	848	499	174	*		
South	12,907	10,088	530	963	630	270	425		
West	6,621	5,337	408	396	316	140	*		
Percent distribution of discharges									
All	100.0	76.6	4.4	8.9	5.7	2.2	*		
Sex									
Male	100.0	74.8	5.3	8.3	6.5	2.8	2.4		
Female	100.0	77.7	3.8	9.3	5.1	1.8	*		
Age									
Under 65 years	100.0	87.3	3.0	2.4	4.3	0.9	2.2		
Under 15 years	100.0	92.0	2.0	0.5	2.6	0.5	*		
15–44 years	100.0	90.9	2.0	1.1	3.6	0.4	2.1		
45–64 years	100.0	81.3	4.6	4.5	5.6	1.8	2.2		
65 years and over	100.0	59.2	6.6	19.4	8.0	4.2	*		
65–74 years	100.0	71.0	6.5	10.9	6.7	2.8	*		
75–84 years	100.0	58.7	6.7	19.7	8.2	4.3	*		
85 years and over	100.0	40.6	6.4	33.1	9.8	6.3	*		
Region									
Northeast	100.0	73.3	4.1	12.1	7.3	2.3	*		
Midwest	100.0	73.6	3.5	10.7	6.3	2.2	*		
South	100.0	78.2	4.1	7.5	4.9	2.1	3.3		
West	100.0	80.6	6.2	6.0	4.8	2.1	*		
Number of days of care in thousands									
All	165,925	107,891	9,175	24,374	13,040	6,586	4,859		
Sex									
Male	72,186	45,713	4,425	10,118	6,056	3,674	2,200		
Female	93,739	62,178	4,750	14,256	6,985	2,912	*		
Age									
Under 65 years	92,775	72,847	3,757	5,386	5,878	2,019	2,888		
Under 15 years	11,371	9,653	450	*	560	*	342		
15–44 years	39,786	33,684	1,230	1,232	2,078	368	1,193		
45–64 years	41,619	29,509	2,077	3,932	3,240	1,508	1,353		
65 years and over	73,149	35,044	5,418	18,988	7,162	4,567	*		
65–74 years	26,068	15,360	1,858	4,506	2,414	1,319	*		
75–84 years	30,228	14,149	2,309	7,963	3,014	2,018	*		
85 years and over	16,853	5,535	1,251	6,519	1,734	1,230	*		

See footnotes at end of table.

Table 8. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by disposition of patient, according to sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All dispositions	Routine or discharged home	Discharged alive				Status not stated	
			Transferred to					
			Other short-term hospital	Long-term care institution	Other	Died		
Number of days of care in thousands								
Northeast	38,730	24,106	1,756	7,241	3,596	1,649	*	
Midwest	33,741	20,875	1,226	5,790	3,028	1,259	*	
South	62,797	42,042	3,314	7,953	4,479	2,213	2,795	
West	30,656	20,867	2,879	3,389	1,937	1,464	119	
Percent distribution of days of care								
All	100.0	65.0	5.5	14.7	7.9	4.0	2.9	
Sex								
Male	100.0	63.3	6.1	14.0	8.4	5.1	3.0	
Female	100.0	66.3	5.1	15.2	7.5	3.1	*	
Age								
Under 65 years	100.0	78.5	4.0	5.8	6.3	2.2	3.1	
Under 15 years	100.0	84.9	4.0	*	4.9	1.3	*	
15–44 years	100.0	84.7	3.1	3.1	5.2	0.9	3.0	
45–64 years	100.0	70.9	5.0	9.4	7.8	3.6	3.3	
65 years and over	100.0	47.9	7.4	26.0	9.8	6.2	*	
65–74 years	100.0	58.9	7.1	17.3	9.3	5.1	*	
75–84 years	100.0	46.8	7.6	26.3	10.0	6.7	*	
85 years and over	100.0	32.8	7.4	38.7	10.3	7.3	*	
Region								
Northeast	100.0	62.2	4.5	18.7	9.3	4.3	*	
Midwest	100.0	61.9	3.6	17.2	9.0	3.7	*	
South	100.0	66.9	5.3	12.7	7.1	3.5	4.5	
West	100.0	68.1	9.4	11.1	6.3	4.8	0.4	
Average length of stay								
All	4.8	4.1	6.1	7.9	6.6	8.8	6.0	
Sex								
Male	5.2	4.4	6.1	8.8	6.7	9.6	6.7	
Female	4.5	3.9	6.1	7.4	6.6	7.9	5.5	
Age								
Under 65 years	4.3	3.9	5.8	10.6	6.4	10.1	6.2	
Under 15 years	4.7	4.3	9.0	*	8.9	11.8	*	
15–44 years	3.7	3.5	5.7	10.7	5.4	9.4	5.4	
45–64 years	5.0	4.3	5.4	10.4	6.9	10.2	7.2	
65 years and over	5.5	4.5	6.2	7.4	6.8	8.3	5.7	
65–74 years	5.3	4.4	5.9	8.4	7.4	9.8	5.8	
75–84 years	5.6	4.5	6.3	7.5	6.8	8.7	5.9	
85 years and over	5.7	4.7	6.6	6.7	6.0	6.6	5.3	
Region								
Northeast	5.4	4.6	5.9	8.4	6.8	9.8	5.9	
Midwest	4.2	3.6	4.4	6.8	6.1	7.2	5.2	
South	4.9	4.2	6.3	8.3	7.1	8.2	6.6	
West	4.6	3.9	7.1	8.6	6.1	10.5	5.1	

* Figure does not meet standards of reliability or precision.

Table 9. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 10,000 population ¹	Number in thousands	Rate per 10,000 population ¹	
All conditions ²	34,667	1174.4	165,925	5,620.9	4.8
Heart disease	391–392.0,393–398,402,404,410–416,420–429	4,230	143.3	18,898	640.2
Acute myocardial infarction410	683	23.1	3,777	127.9
Coronary atherosclerosis414.0	980	33.2	3,422	115.9
Other ischemic heart disease411–413,414.1–414.9	165	5.6	392	13.3
Cardiac dysrhythmias427	773	26.2	2,593	87.8
Congestive heart failure428.0,428.2–428.4	1,079	36.6	5,506	186.5
Females with deliveriesV27	4,038	136.8	10,425	353.2
Psychoses290–299	1,747	59.2	13,265	449.4
Schizophrenic disorders295	351	11.9	3,709	125.7
Major depressive disorder296.2–296.3	470	15.9	3,251	110.1
Pneumonia480–486	1,368	46.3	7,184	243.4
Malignant neoplasms140–208,230–234	1,196	40.5	8,099	274.4
Malignant neoplasm of large intestine and rectum153–154,197.5	148	5.0	1,222	41.4
Malignant neoplasm of trachea, bronchus, and lung162,176.4,197.0,197.3	154	5.2	1,147	38.9
Fractures, all sites ³800–829	1,009	34.2	5,351	181.3
Fracture of neck of femur ³820	302	10.2	1,904	64.5
Cerebrovascular disease430–438	895	30.3	4,648	157.4
Certain complications of surgical and medical care996–999	863	29.2	5,200	176.1
Osteoarthritis and allied disorders715	761	25.8	2,852	96.6
Diabetes mellitus250	585	19.8	2,774	94.0
Chronic bronchitis491	569	19.3	2,852	96.6
Cellulitis and abscess681–682	551	18.7	2,592	87.8
Volume depletion276.5	500	16.9	1,694	57.4
Septicemia038	490	16.6	4,034	136.6
Asthma493	489	16.6	1,616	54.7
Urinary tract infection599.0	481	16.3	2,177	73.7
Benign neoplasms210–229	368	12.5	1,281	43.4
Benign neoplasm of uterus218–219	202	6.8	578	19.6
Cholelithiasis574	325	11.0	1,383	46.8
Anemias280–285	325	11.0	1,432	48.5
Appendicitis540–543	321	10.9	982	33.3
Noninfectious enteritis and colitis555–558	318	10.8	1,451	49.2
Intervertebral disc disorder722	313	10.6	904	30.6
Diverticula of intestine562	307	10.4	1,614	54.7
Intestinal obstruction560	304	10.3	1,956	66.3
Essential hypertension401	301	10.2	678	23.0
Poisonings960–989	258	8.8	771	26.1
Acute pancreatitis577.0	237	8.0	1,197	40.6
Acute bronchitis and bronchiolitis466	230	7.8	735	24.9
Calculus of kidney and ureter592	184	6.2	425	14.4

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²Includes data for diagnostic conditions not shown in table.

³Excludes fractures coded as 733.1, pathologic fracture.

Table 10. Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	65 years and over	65–74 years	75–84 years	85 years and over
Discharges in thousands				
All conditions ¹	13,228	4,900	5,394	2,934
Heart disease	391–392,0,393–398,402,404,410–416,420–429	2,707	980	1,099
Acute myocardial infarction	410	430	156	164
Coronary atherosclerosis	414.0	546	278	215
Cardiac dysrhythmias	427	537	177	228
Congestive heart failure	428.0,428.2–428.4	800	223	336
Pneumonia	480–486	814	245	306
Cerebrovascular disease	430–438	625	214	271
Malignant neoplasms	140–208,230–234	608	289	237
Fractures, all sites ²	800–829	526	119	211
Fracture of neck of femur ²	820	266	44	107
Osteoarthritis and allied disorders	715	478	265	181
Chronic bronchitis	491	386	164	177
Urinary tract infection	599.0	344	74	147
Certain complications of surgical and medical care	996–999	343	173	129
Septicemia	038	334	103	137
Volume depletion	276.5	227	64	97
Psychoses	290–299	208	94	75
Diabetes mellitus	250	191	93	72
Cellulitis and abscess	681–682	169	56	76
Diverticula of intestine	562	164	55	74
Intestinal obstruction	560	164	54	70
Anemias	280–285	137	42	59
Cholelithiasis	574	124	54	52
Essential hypertension	401	121	45	48
Asthma	493	112	53	40
Noninfectious enteritis and colitis	555–558	108	37	53
Discharge rate per 10,000 population ³				
All conditions ¹	3,595.6	2,628.9	4,131.7	5,758.1
Heart disease	391–392,0,393–398,402,404,410–416,420–429	735.8	525.9	842.1
Acute myocardial infarction	410	116.8	83.5	125.3
Coronary atherosclerosis	414.0	148.4	149.4	164.6
Cardiac dysrhythmias	427	146.1	95.0	174.4
Congestive heart failure	428.0,428.2–428.4	217.4	119.8	257.6
Pneumonia	480–486	221.3	131.4	234.2
Cerebrovascular disease	430–438	169.8	114.6	207.8
Malignant neoplasms	140–208,230–234	165.3	154.9	181.7
Fractures, all sites ²	800–829	142.9	63.8	161.4
Fracture of neck of femur ²	820	72.3	23.7	82.1
Osteoarthritis and allied disorders	715	129.8	142.2	139.0
Chronic bronchitis	491	104.9	87.8	135.5
Urinary tract infection	599.0	93.5	39.6	112.5
Certain complications of surgical and medical care	996–999	93.3	92.9	99.0
Septicemia	038	90.8	55.5	104.7
Volume depletion	276.5	61.8	34.3	74.0
Psychoses	290–299	56.7	50.2	57.8
Diabetes mellitus	250	52.0	49.7	55.2
Cellulitis and abscess	681–682	45.9	30.2	57.9
Diverticula of intestine	562	44.6	29.7	57.0
Intestinal obstruction	560	44.6	29.0	53.2
Anemias	280–285	37.2	22.5	45.2
Cholelithiasis	574	33.7	29.2	39.6
Essential hypertension	401	33.0	24.2	36.9
Asthma	493	30.5	28.6	30.8
Noninfectious enteritis and colitis	555–558	29.4	19.8	40.8

See footnotes at end of table.

Table 10. Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	65 years and over	65–74 years	75–84 years	85 years and over
Average length of stay in days				
All conditions ¹	5.5	5.3	5.6	5.7
Heart disease 391–392,0,393–398,402,404,410–416,420–429	4.7	4.4	4.7	4.9
Acute myocardial infarction 410	6.0	5.8	6.2	5.8
Coronary atherosclerosis 414.0	3.5	3.5	3.7	3.3
Cardiac dysrhythmias 427	3.6	3.3	3.4	4.2
Congestive heart failure 428.0,428.2–428.4	5.1	4.9	5.1	5.3
Pneumonia 480–486	5.7	5.4	5.8	5.9
Cerebrovascular disease 430–438	4.9	4.5	5.1	5.3
Malignant neoplasms 140–208,230–234	6.9	6.7	7.3	6.8
Fractures, all sites ² 800–829	5.7	5.5	5.7	5.7
Fracture of neck of femur ² 820	6.4	7.1	6.7	5.8
Osteoarthritis and allied disorders 715	3.8	3.7	3.9	4.1
Chronic bronchitis 491	5.4	4.9	5.9	5.4
Urinary tract infection 599.0	4.7	4.4	4.6	5.0
Certain complications of surgical and medical care 996–999	6.4	6.6	6.2	6.2
Septicemia 038	7.8	8.1	8.0	7.1
Volume depletion 276.5	4.2	3.9	4.1	4.7
Psychoses 290–299	8.6	9.4	7.8	8.5
Diabetes mellitus 250	5.3	5.2	5.6	5.4
Cellulitis and abscess 681–682	5.3	4.9	5.5	5.6
Diverticula of intestine 562	5.8	5.7	5.5	6.3
Intestinal obstruction 560	6.9	6.9	7.3	6.3
Anemias 280–285	4.0	3.5	3.9	4.6
Cholelithiasis 574	4.8	4.2	4.9	6.1
Essential hypertension 401	2.6	2.3	2.7	2.8
Asthma 493	4.7	4.4	4.9	5.0
Noninfectious enteritis and colitis 555–558	5.1	5.1	4.8	5.9

¹Includes data for diagnostic conditions not shown in table.

²Excludes fractures coded as 733.1, pathologic fracture.

³Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

Table 11. Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15–44 years	45–64 years	65 years and over
Number in thousands					
All conditions	34,667	2,431	10,659	8,349	13,228
Infectious and parasitic diseases	001–139	1,045	142	176	244
Septicemia	038	490	6	38	112
Neoplasms	140–239	1,615	27	270	626
Malignant neoplasms	140–208,230–234	1,196	20	118	449
Malignant neoplasm of large intestine and rectum	153–154,197.5	148	*	5	47
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	154	*	7	54
Benign neoplasms	210–229	368	4	141	164
Benign neoplasm of uterus	218–219	202	*	103	96
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	1,708	203	359	494
Diabetes mellitus	250	585	27	157	209
Volume depletion	276.5	500	148	49	76
Diseases of the blood and blood-forming organs	280–289	455	57	115	100
Anemias	280–285	325	28	93	67
Mental disorders	290–319	2,398	124	1,349	680
Psychoses	290–299	1,747	*	970	494
Schizophrenic disorders	295	351	*	196	130
Major depressive disorder	296.2–296.3	470	*	264	137
Diseases of the nervous system and sense organs	320–389	565	84	138	125
Diseases of the circulatory system	390–459	6,159	29	436	1,813
Essential hypertension	401	301	*	42	137
Heart disease	391–392,0,393–398,402,404,410–416,420–429	4,230	15	266	1,242
Acute myocardial infarction	410	683	*	38	215
Coronary atherosclerosis	414.0	980	*	40	393
Other ischemic heart disease	411–413,414.1–414.9	165	*	22	68
Cardiac dysrhythmias	427	773	5	49	182
Congestive heart failure	428.0,428.2–428.4	1,079	*	42	235
Cerebrovascular disease	430–438	895	*	47	221
Diseases of the respiratory system	460–519	3,746	629	370	825
Acute bronchitis and bronchiolitis	466	230	155	13	18
Pneumonia	480–486	1,368	193	110	251
Chronic bronchitis	491	569	*	16	166
Asthma	493	489	159	98	119
Diseases of the digestive system	520–579	3,422	216	832	1,052
Appendicitis	540–543	321	78	162	60
Noninfectious enteritis and colitis	555–558	318	41	94	75
Intestinal obstruction	560	304	11	40	89
Diverticula of intestine	562	307	*	37	106
Cholelithiasis	574	325	*	106	95
Acute pancreatitis	577.0	237	*	83	88
Diseases of the genitourinary system	580–629	1,983	91	535	496
Calculus of kidney and ureter	592	184	*	75	68
Urinary tract infection	599.0	481	28	43	66
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	535	*	533	*
Diseases of the skin and subcutaneous tissue	680–709	751	*	188	225
Cellulitis and abscess	681–682	551	62	144	176
Diseases of the musculoskeletal system and connective tissue	710–739	1,945	40	304	686
Osteoarthritis and allied disorders	715	761	*	17	266
Intervertebral disc disorders	722	313	*	111	140
Congenital anomalies	740–759	182	129	27	18
Certain conditions originating in the perinatal period	760–779	201	201	*	*
Symptoms, signs, and ill-defined conditions	780–799	198	47	62	51
Injury and poisoning	800–999	2,826	224	807	704
Fractures, all sites ²	800–829	1,009	67	235	181
Fracture of neck of femur ²	820	302	*	8	27
Poisonings	960–989	258	15	154	66
Certain complications of surgical and medical care	996–999	863	43	174	303
Supplementary classifications	V01–V84	4,933	77	4,158	210
Females with deliveries	V27	4,038	*6	4,025	*7

* Figure does not meet standards of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 12. Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*.]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15–44 years	45–64 years	65 years and over
Rate per 10,000 population ¹					
All conditions	1,174.4	400.5	853.3	1,147.0	3,595.6
Infectious and parasitic diseases	001–139	35.4	23.4	14.1	33.6
Septicemia	038	16.6	1.0	3.0	15.4
Neoplasms	140–239	54.7	4.5	21.6	86.0
Malignant neoplasms	140–208,230–234	40.5	3.3	9.5	61.7
Malignant neoplasm of large intestine and rectum	153–154,197.5	5.0	*	0.4	6.5
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	5.2	*	0.6	7.4
Benign neoplasms	210–229	12.5	0.7	11.3	22.5
Benign neoplasm of uterus	218–219	6.8	*	8.3	13.2
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	57.9	33.5	28.8	67.8
Diabetes mellitus	250	19.8	4.5	12.6	28.7
Volume depletion	276.5	16.9	24.3	3.9	10.4
Diseases of the blood and blood-forming organs	280–289	15.4	9.4	9.2	13.7
Anemias	280–285	11.0	4.6	7.5	9.2
Mental disorders	290–319	81.2	20.4	108.0	93.5
Psychoses	290–299	59.2	*	77.6	67.9
Schizophrenic disorders	295	11.9	*	15.7	17.8
Major depressive disorder	296.2–296.3	15.9	*	21.1	18.8
Diseases of the nervous system and sense organs	320–389	19.1	13.9	11.0	17.1
Diseases of the circulatory system	390–459	208.7	4.7	34.9	249.1
Essential hypertension	401	10.2	*	3.3	18.9
Heart disease	391–392,0,393–398,402,404,410–416,420–429	143.3	2.5	21.3	170.6
Acute myocardial infarction	410	23.1	*	3.1	29.5
Coronary atherosclerosis	414.0	33.2	*	3.2	54.0
Other ischemic heart disease	411–413,414.1–414.9	5.6	*	1.8	9.4
Cardiac dysrhythmias	427	26.2	0.8	3.9	25.0
Congestive heart failure	428.0,428.2–428.4	36.6	*	3.3	32.3
Cerebrovascular disease	430–438	30.3	*	3.8	30.4
Diseases of the respiratory system	460–519	126.9	103.7	29.6	113.4
Acute bronchitis and bronchiolitis	466	7.8	25.6	1.1	2.5
Pneumonia	480–486	46.3	31.9	8.8	34.5
Chronic bronchitis	491	19.3	*	1.3	22.8
Asthma	493	16.6	26.2	7.8	16.4
Diseases of the digestive system	520–579	115.9	35.6	66.6	144.6
Appendicitis	540–543	10.9	12.9	13.0	8.2
Noninfectious enteritis and colitis	555–558	10.8	6.7	7.5	10.3
Intestinal obstruction	560	10.3	1.7	3.2	12.2
Diverticula of intestine	562	10.4	*	3.0	14.5
Cholelithiasis	574	11.0	*	8.5	13.0
Acute pancreatitis	577.0	8.0	*	6.6	12.1
Diseases of the genitourinary system	580–629	67.2	15.0	42.9	68.1
Calculus of kidney and ureter	592	6.2	*	6.0	9.4
Urinary tract infection	599.0	16.3	4.6	3.4	9.1
Complications of pregnancy, childbirth, and the puerperium ²	630–677	18.1	*	42.7	*
Diseases of the skin and subcutaneous tissue	680–709	25.4	*	15.1	30.9
Cellulitis and abscess	681–682	18.7	10.3	11.5	24.1
Diseases of the musculoskeletal system and connective tissue	710–739	65.9	6.5	24.3	94.2
Osteoarthritis and allied disorders	715	25.8	*	1.3	36.5
Intervertebral disc disorders	722	10.6	*	8.9	19.2
Congenital anomalies	740–759	6.2	21.3	2.2	2.4
Certain conditions originating in the perinatal period	760–779	6.8	33.1	*	*
Symptoms, signs, and ill-defined conditions	780–799	6.7	7.7	5.0	7.0
Injury and poisoning	800–999	95.7	36.8	64.6	96.8
Fractures, all sites ³	800–829	34.2	11.0	18.8	24.9
Fracture of neck of femur ³	820	10.2	*	0.6	3.8
Poisonings	960–989	8.8	2.4	12.4	9.1
Certain complications of surgical and medical care	996–999	29.2	7.0	14.0	41.6
Supplementary classifications	V01–V84	167.1	12.6	332.9	28.8
Females with deliveries	V27	136.8	*1.0	322.2	*1.0

* Figure does not meet standards of reliability or precision.

... Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

³Excludes fractures coded as 733.1, pathologic fracture.

Table 13. Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15–44 years	45–64 years	65 years and over
Average length of stay in days					
All conditions	4.8	4.7	3.7	5.0	5.5
Infectious and parasitic diseases	6.6	3.5	6.0	7.6	7.3
Septicemia	8.2	*	8.9	9.2	7.8
Neoplasms	140–239	6.0	8.5	4.8	5.5
Malignant neoplasms	140–208,230–234	6.8	9.8	6.9	6.4
Malignant neoplasm of large intestine and rectum	153–154,197.5	8.3	*	8.4	7.3
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	7.4	*	*	6.8
Benign neoplasms	210–229	3.5	5.2	3.0	3.3
Benign neoplasm of uterus	218–219	2.9	*	2.7	3.0
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	4.0	2.5	3.5	4.2
Diabetes mellitus	250	4.7	2.5	3.9	5.1
Volume depletion	276.5	3.4	2.0	2.8	4.0
Diseases of the blood and blood-forming organs	280–289	4.4	3.8	4.9	4.7
Anemias	280–285	4.4	4.3	5.1	4.4
Mental disorders	290–319	6.9	9.8	6.4	7.0
Psychoses	290–299	7.6	9.4	7.0	8.0
Schizophrenic disorders	295	10.6	*	10.2	10.9
Major depressive disorder	296.2–296.3	6.9	7.1	5.8	7.7
Diseases of the nervous system and sense organs	320–389	5.2	4.3	4.4	4.5
Diseases of the circulatory system	390–459	4.6	6.4	4.2	4.3
Essential hypertension	401	2.2	*	2.0	2.1
Heart disease	391–392,0,393–398,402,404,410–416,420–429	4.5	7.2	4.0	4.1
Acute myocardial infarction	410	5.5	*	4.1	4.9
Coronary atherosclerosis	414.0	3.5	*	2.8	3.5
Other ischemic heart disease	411–413,414.1–414.9	2.4	*	1.9	2.2
Cardiac dysrhythmias	427	3.4	4.0	2.5	2.9
Congestive heart failure	428.0,428.2–428.4	5.1	*6.5	4.4	5.2
Cerebrovascular disease	430–438	5.2	*8.1	6.2	5.6
Diseases of the respiratory system	460–519	5.2	3.1	4.4	5.3
Acute bronchitis and bronchiolitis	466	3.2	2.9	3.3	4.4
Pneumonia	480–486	5.3	3.1	5.0	5.5
Chronic bronchitis	491	5.0	*	3.9	4.2
Asthma	493	3.3	2.3	2.7	3.9
Diseases of the digestive system	520–579	4.7	4.0	3.6	4.8
Appendicitis	540–543	3.1	3.8	2.3	3.4
Noninfectious enteritis and colitis	555–558	4.6	2.5	4.9	4.6
Intestinal obstruction	560	6.4	4.7	6.0	5.9
Diverticula of intestine	562	5.3	*	3.7	5.0
Cholelithiasis	574	4.2	*	3.1	4.8
Acute pancreatitis	577.0	5.1	*6.5	4.2	5.4
Diseases of the genitourinary system	580–629	3.8	3.2	2.8	3.5
Calculus of kidney and ureter	592	2.3	*	2.1	2.3
Urinary tract infection	599.0	4.5	3.6	3.6	4.6
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	2.9	*	2.9	*
Diseases of the skin and subcutaneous tissue	680–709	5.0	2.7	4.2	5.6
Cellulitis and abscess	681–682	4.7	3.3	4.1	5.1
Diseases of the musculoskeletal system and connective tissue	710–739	3.9	4.8	3.3	3.6
Osteoarthritis and allied disorders	715	3.7	*	3.6	3.6
Intervertebral disc disorders	722	2.9	*	2.3	2.9
Congenital anomalies	740–759	6.8	7.5	3.6	*
Certain conditions originating in the perinatal period	760–779	10.5	10.5	*	*
Symptoms, signs, and ill-defined conditions	780–799	2.8	2.3	2.5	2.5
Injury and poisoning	800–999	5.2	4.5	4.5	5.4
Fractures, all sites ²	800–829	5.3	3.4	5.0	5.3
Fracture of neck of femur ²	820	6.3	*	3.7	6.4
Poisonings	960–989	3.0	1.9	2.8	3.4
Certain complications of surgical and medical care	996–999	6.0	6.1	5.3	6.0
Supplementary classifications	V01–V84	3.7	5.0	2.7	7.4
Females with deliveries	V27	2.6	*2.3	2.6	*3.3

* Figure does not meet standards of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 14. Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
Number in thousands			
All conditions	34,667	13,902	20,766
Infectious and parasitic diseases	001–139	1,045	498
Septicemia	038	490	240
Neoplasms	140–239	1,615	654
Malignant neoplasms	140–208,230–234	1,196	579
Malignant neoplasm of large intestine and rectum	153–154,197.5	148	71
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	154	79
Benign neoplasms	210–229	368	50
Benign neoplasm of uterus	218–219	202	202
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	1,708	713
Diabetes mellitus	250	585	283
Volume depletion	276.5	500	206
Diseases of the blood and blood-forming organs	280–289	455	174
Anemias	280–285	325	120
Mental disorders	290–319	2,398	1,200
Psychoses	290–299	1,747	832
Schizophrenic disorders	295	351	193
Major depressive disorder	296.2–296.3	470	179
Diseases of the nervous system and sense organs	320–389	565	238
Diseases of the circulatory system	390–459	6,159	3,136
Essential hypertension	401	301	126
Heart disease	391–392.0,393–398,402,404,410–416,420–429	4,230	2,222
Acute myocardial infarction	410	683	406
Coronary atherosclerosis	414.0	980	629
Other ischemic heart disease	411–413,414.1–414.9	165	82
Cardiac dysrhythmias	427	773	365
Congestive heart failure	428.0,428.2–428.4	1,079	491
Cerebrovascular disease	430–438	895	418
Diseases of the respiratory system	460–519	3,746	1,764
Acute bronchitis and bronchiolitis	466	230	117
Pneumonia	480–486	1,368	651
Chronic bronchitis	491	569	261
Asthma	493	489	192
Diseases of the digestive system	520–579	3,422	1,546
Appendicitis	540–543	321	179
Noninfectious enteritis and colitis	555–558	318	119
Intestinal obstruction	560	304	134
Diverticula of intestine	562	307	126
Cholelithiasis	574	325	116
Acute pancreatitis	577.0	237	129
Diseases of the genitourinary system	580–629	1,983	642
Calculus of kidney and ureter	592	184	94
Urinary tract infection	599.0	481	157
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	535	...
Diseases of the skin and subcutaneous tissue	680–709	751	400
Cellulitis and abscess	681–682	551	300
Diseases of the musculoskeletal system and connective tissue	710–739	1,945	831
Osteoarthritis and allied disorders	715	761	292
Intervertebral disc disorders	722	313	153
Congenital anomalies	740–759	182	99
Certain conditions originating in the perinatal period	760–779	201	120
Symptoms, signs, and ill-defined conditions	780–799	198	89
Injury and poisoning	800–999	2,826	1,402
Fractures, all sites ²	800–829	1,009	435
Fracture of neck of femur ²	820	302	86
Poisonings	960–989	258	122
Certain complications of surgical and medical care	996–999	863	412
Supplementary classifications	V01–V84	4,933	396
Females with deliveries	V27	4,038	4,038

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

²Excludes fractures coded as 733.1, pathologic fracture.

Table 15. Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
Rate per 10,000 population ¹			
All conditions	1,174.4	959.0	1,382.2
Infectious and parasitic diseases	001–139	35.4	34.4
Septicemia	038	16.6	16.6
Neoplasms	140–239	54.7	45.1
Malignant neoplasms	140–208,230–234	40.5	40.0
Malignant neoplasm of large intestine and rectum	153–154,197.5	5.0	4.9
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	5.2	5.5
Benign neoplasms	210–229	12.5	3.5
Benign neoplasm of uterus	218–219	6.8	13.4
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	57.9	49.2
Diabetes mellitus	250	19.8	19.5
Volume depletion	276.5	16.9	14.2
Diseases of the blood and blood-forming organs	280–289	15.4	12.0
Anemias	280–285	11.0	8.3
Mental disorders	290–319	81.2	82.8
Psychoses	290–299	59.2	57.4
Schizophrenic disorders	295	11.9	13.3
Major depressive disorder	296.2–296.3	15.9	12.4
Diseases of the nervous system and sense organs	320–389	19.1	16.4
Diseases of the circulatory system	390–459	208.7	216.3
Essential hypertension	401	10.2	8.7
Heart disease	391–392.0,393–398,402,404,410–416,420–429	143.3	153.3
Acute myocardial infarction	410	23.1	28.0
Coronary atherosclerosis	414.0	33.2	43.4
Other ischemic heart disease	411–413,414.1–414.9	5.6	5.7
Cardiac dysrhythmias	427	26.2	25.2
Congestive heart failure	428.0,428.2–428.4	36.6	33.9
Cerebrovascular disease	430–438	30.3	28.9
Diseases of the respiratory system	460–519	126.9	121.7
Acute bronchitis and bronchiolitis	466	7.8	8.1
Pneumonia	480–486	46.3	44.9
Chronic bronchitis	491	19.3	18.0
Asthma	493	16.6	13.3
Diseases of the digestive system	520–579	115.9	106.6
Appendicitis	540–543	10.9	12.4
Noninfectious enteritis and colitis	555–558	10.8	8.2
Intestinal obstruction	560	10.3	9.3
Diverticula of intestine	562	10.4	8.7
Cholelithiasis	574	11.0	8.0
Acute pancreatitis	577.0	8.0	8.9
Diseases of the genitourinary system	580–629	67.2	44.3
Calculus of kidney and ureter	592	6.2	6.5
Urinary tract infection	599.0	16.3	10.8
Complications of pregnancy, childbirth, and the puerperium ²	630–677	18.1	...
Diseases of the skin and subcutaneous tissue	680–709	25.4	27.6
Cellulitis and abscess	681–682	18.7	20.7
Diseases of the musculoskeletal system and connective tissue	710–739	65.9	57.3
Osteoarthritis and allied disorders	715	25.8	20.1
Intervertebral disc disorders	722	10.6	10.6
Congenital anomalies	740–759	6.2	6.8
Certain conditions originating in the perinatal period	760–779	6.8	8.3
Symptoms, signs, and ill-defined conditions	780–799	6.7	6.1
Injury and poisoning	800–999	95.7	96.7
Fractures, all sites ³	800–829	34.2	30.0
Fracture of neck of femur ³	820	10.2	6.0
Poisonings	960–989	8.8	8.4
Certain complications of surgical and medical care	996–999	29.2	28.4
Supplementary classifications	V01–V84	167.1	27.3
Females with deliveries	V27	136.8	...
...			

... Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

³Excludes fractures coded as 733.1, pathologic fracture.

Table 16. Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
	Average length of stay in days		
All conditions	4.8	5.2	4.5
Infectious and parasitic diseases	001–139	6.6	6.8
Septicemia	038	8.2	8.3
Neoplasms	140–239	6.0	6.7
Malignant neoplasms	140–208,230–234	6.8	6.9
Malignant neoplasm of large intestine and rectum	153–154,197.5	8.3	8.5
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	7.4	7.4
Benign neoplasms	210–229	3.5	5.0
Benign neoplasm of uterus	218–219	2.9	2.9
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	4.0	4.2
Diabetes mellitus	250	4.7	4.9
Volume depletion	276.5	3.4	3.2
Diseases of the blood and blood-forming organs	280–289	4.4	4.1
Anemias	280–285	4.4	4.1
Mental disorders	290–319	6.9	7.0
Psychoses	290–299	7.6	7.7
Schizophrenic disorders	295	10.6	10.5
Major depressive disorder	296.2–296.3	6.9	7.2
Diseases of the nervous system and sense organs	320–389	5.2	5.2
Diseases of the circulatory system	390–459	4.6	4.5
Essential hypertension	401	2.2	2.2
Heart disease	391–392.0,393–398,402,404,410–416,420–429	4.5	4.3
Acute myocardial infarction	410	5.5	5.6
Coronary atherosclerosis	414.0	3.5	3.3
Other ischemic heart disease	411–413,414.1–414.9	2.4	2.5
Cardiac dysrhythmias	427	3.4	3.1
Congestive heart failure	428.0,428.2–428.4	5.1	4.9
Cerebrovascular disease	430–438	5.2	5.5
Diseases of the respiratory system	460–519	5.2	5.2
Acute bronchitis and bronchiolitis	466	3.2	3.1
Pneumonia	480–486	5.3	5.2
Chronic bronchitis	491	5.0	5.1
Asthma	493	3.3	2.8
Diseases of the digestive system	520–579	4.7	4.7
Appendicitis	540–543	3.1	3.2
Noninfectious enteritis and colitis	555–558	4.6	4.2
Intestinal obstruction	560	6.4	6.4
Diverticula of intestine	562	5.3	5.1
Cholelithiasis	574	4.2	5.1
Acute pancreatitis	577.0	5.1	4.6
Diseases of the genitourinary system	580–629	3.8	4.3
Calculus of kidney and ureter	592	2.3	1.9
Urinary tract infection	599.0	4.5	4.9
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	2.9	...
Diseases of the skin and subcutaneous tissue	680–709	5.0	4.8
Cellulitis and abscess	681–682	4.7	4.5
Diseases of the musculoskeletal system and connective tissue	710–739	3.9	3.8
Osteoarthritis and allied disorders	715	3.7	3.6
Intervertebral disc disorders	722	2.9	2.7
Congenital anomalies	740–759	6.8	6.5
Certain conditions originating in the perinatal period	760–779	10.5	10.1
Symptoms, signs, and ill-defined conditions	780–799	2.8	2.7
Injury and poisoning	800–999	5.2	5.3
Fractures, all sites ²	800–829	5.3	5.4
Fracture of neck of femur ²	820	6.3	6.8
Poisonings	960–989	3.0	3.3
Certain complications of surgical and medical care	996–999	6.0	6.0
Supplementary classifications	V01–V84	3.7	8.2
Females with deliveries	V27	2.6	2.6

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

²Excludes fractures coded as 733.1, pathologic fracture.

Table 17. Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	Northeast	Midwest	South	West
Number in thousands					
All conditions	34,667	7,192	7,948	12,907	6,621
Infectious and parasitic diseases	001–139	1,045	235	382	202
Septicemia	038	490	97	115	99
Neoplasms	140–239	1,615	348	345	600
Malignant neoplasms	140–208,230–234	1,196	259	259	434
Malignant neoplasm of large intestine and rectum	153–154,197.5	148	30	36	54
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	154	36	38	59
Benign neoplasms	210–229	368	77	74	147
Benign neoplasm of uterus	218–219	202	38	40	35
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	1,708	383	381	645
Diabetes mellitus	250	585	148	119	215
Volume depletion	276.5	500	91	121	217
Diseases of the blood and blood-forming organs	280–289	455	106	98	177
Anemias	280–285	325	80	69	129
Mental disorders	290–319	2,398	665	518	*
Psychoses	290–299	1,747	468	395	213
Schizophrenic disorders	295	351	*	60	147
Major depressive disorder	296.2–296.3	470	104	114	30
Diseases of the nervous system and sense organs	320–389	565	135	124	195
Diseases of the circulatory system	390–459	6,159	1,392	1,465	2,226
Essential hypertension	401	301	68	67	48
Heart disease	391–392,0,393–398,402,404,410–416,420–429	4,230	977	1,021	1,492
Acute myocardial infarction	410	683	171	154	221
Coronary atherosclerosis	414.0	980	204	239	137
Other ischemic heart disease	411–413,414.1–414.9	165	44	36	25
Cardiac dysrhythmias	427	773	185	191	259
Congestive heart failure	428.0,428.2–428.4	1,079	239	275	139
Cerebrovascular disease	430–438	895	188	211	325
Diseases of the respiratory system	460–519	3,746	749	967	1,353
Acute bronchitis and bronchiolitis	466	230	37	50	56
Pneumonia	480–486	1,368	253	375	251
Chronic bronchitis	491	569	116	156	75
Asthma	493	489	122	115	173
Diseases of the digestive system	520–579	3,422	712	797	1,225
Appendicitis	540–543	321	64	70	90
Noninfectious enteritis and colitis	555–558	318	74	78	51
Intestinal obstruction	560	304	57	84	61
Diverticula of intestine	562	307	76	79	102
Cholelithiasis	574	325	68	74	51
Acute pancreatitis	577.0	237	42	56	66
577.0	237	42	56	96	43
Diseases of the genitourinary system	580–629	1,983	376	464	782
Calculus of kidney and ureter	592	184	37	50	20
Urinary tract infection	599.0	481	95	114	86
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	535	77	91	255
Diseases of the skin and subcutaneous tissue	680–709	751	155	169	119
Cellulitis and abscess	681–682	551	122	135	92
Diseases of the musculoskeletal system and connective tissue	710–739	1,945	405	504	631
Osteoarthritis and allied disorders	715	761	160	214	406
Intervertebral disc disorders	722	313	61	79	115
Congenital anomalies	740–759	182	48	*	65
Certain conditions originating in the perinatal period	760–779	201	23	*	69
Symptoms, signs, and ill-defined conditions	780–799	198	52	37	30
Injury and poisoning	800–999	2,826	555	670	1,007
Fractures, all sites ²	800–829	1,009	191	257	594
Fracture of neck of femur ²	820	302	57	76	216
Poisonings	960–989	258	53	63	48
Certain complications of surgical and medical care	996–999	863	185	182	311
Supplementary classifications	V01–V84	4,933	773	1,017	1,911
Females with deliveries	V27	4,038	591	824	1,574
					1,049

* Figure does not meet standards of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

²Excludes fractures coded as 733.1, pathologic fracture.

Table 18. Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	Northeast	Midwest	South	West
Rate per 10,000 population ¹					
All conditions	1,174.4	1,317.6	1,206.2	1,208.3	975.1
Infectious and parasitic diseases	001–139	35.4	43.0	34.2	35.8
Septicemia	038	16.6	17.8	17.5	16.6
Neoplasms	140–239	54.7	63.8	52.3	56.1
Malignant neoplasms	140–208,230–234	40.5	47.5	39.3	40.6
Malignant neoplasm of large intestine and rectum	153–154,197.5	5.0	5.5	5.5	5.0
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	5.2	6.6	5.7	5.6
Benign neoplasms	210–229	12.5	14.1	11.2	13.8
Benign neoplasm of uterus	218–219	6.8	7.0	6.0	8.3
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	57.9	70.2	57.8	60.4
Diabetes mellitus	250	19.8	27.1	18.1	20.1
Volume depletion	276.5	16.9	16.6	18.3	20.3
Diseases of the blood and blood-forming organs	280–289	15.4	19.5	14.8	16.5
Anemias	280–285	11.0	14.7	10.4	12.1
Mental disorders	290–319	81.2	121.9	78.7	*
Psychoses	290–299	59.2	85.7	59.9	*
Schizophrenic disorders	295	11.9	*	9.0	*
Major depressive disorder	296.2–296.3	15.9	19.1	17.3	*
Diseases of the nervous system and sense organs	320–389	19.1	24.8	18.9	18.3
Diseases of the circulatory system	390–459	208.7	255.1	222.3	208.4
Essential hypertension	401	10.2	12.4	10.2	11.1
Heart disease	391–392,0,393–398,402,404,410–416,420–429	143.3	179.0	155.0	139.7
Acute myocardial infarction	410	23.1	31.3	23.4	20.7
Coronary atherosclerosis	414.0	33.2	37.4	36.2	33.2
Other ischemic heart disease	411–413,414.1–414.9	5.6	8.0	5.5	5.6
Cardiac dysrhythmias	427	26.2	33.9	29.0	24.2
Congestive heart failure	428.0,428.2–428.4	36.6	43.8	41.7	38.0
Cerebrovascular disease	430–438	30.3	34.5	32.1	30.5
Diseases of the respiratory system	460–519	126.9	137.3	146.7	126.7
Acute bronchitis and bronchiolitis	466	7.8	6.9	7.6	8.1
Pneumonia	480–486	46.3	46.4	56.9	45.8
Chronic bronchitis	491	19.3	21.3	23.6	20.7
Asthma	493	16.6	22.3	17.5	16.2
Diseases of the digestive system	520–579	115.9	130.5	121.0	114.7
Appendicitis	540–543	10.9	11.6	10.7	9.1
Noninfectious enteritis and colitis	555–558	10.8	13.6	11.9	10.7
Intestinal obstruction	560	10.3	10.4	12.7	9.6
Diverticula of intestine	562	10.4	13.9	12.0	9.5
Cholelithiasis	574	11.0	12.5	11.2	11.0
Acute pancreatitis	577.0	8.0	7.8	8.5	9.0
Diseases of the genitourinary system	580–629	67.2	68.9	70.5	73.2
Calculus of kidney and ureter	592	6.2	6.8	7.5	7.3
Urinary tract infection	599.0	16.3	17.4	17.3	17.4
Complications of pregnancy, childbirth, and the puerperium ²	630–677	18.1	14.2	13.8	23.8
Diseases of the skin and subcutaneous tissue	680–709	25.4	28.5	25.6	28.8
Cellulitis and abscess	681–682	18.7	22.3	20.4	18.9
Diseases of the musculoskeletal system and connective tissue	710–739	65.9	74.1	76.5	59.0
Osteoarthritis and allied disorders	715	25.8	29.4	32.4	21.0
Intervertebral disc disorders	722	10.6	11.1	12.0	10.8
Congenital anomalies	740–759	6.2	8.8	*	6.0
Certain conditions originating in the perinatal period	760–779	6.8	4.1	*	6.1
Symptoms, signs, and ill-defined conditions	780–799	6.7	9.6	5.7	7.3
Injury and poisoning	800–999	95.7	101.7	101.7	94.3
Fractures, all sites ³	800–829	34.2	35.1	39.0	32.3
Fracture of neck of femur ³	820	10.2	10.4	11.5	10.2
Poisonings	960–989	8.8	9.7	9.6	8.8
Certain complications of surgical and medical care	996–999	29.2	33.8	27.6	29.1
Supplementary classifications	V01–V84	167.1	141.7	154.3	178.9
Females with deliveries	V27	136.8	108.3	125.1	147.4
					154.4

* Figure does not meet standards of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

³Excludes fractures coded as 733.1, pathologic fracture.

Table 19. Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	Northeast	Midwest	South	West
Average length of stay in days					
All conditions	4.8	5.4	4.2	4.9	4.6
Infectious and parasitic diseases	001–139	6.6	7.4	5.5	6.7
Septicemia	038	8.2	9.7	6.7	8.3
Neoplasms	140–239	6.0	6.8	5.4	5.9
Malignant neoplasms	140–208,230–234	6.8	7.6	6.1	6.8
Malignant neoplasm of large intestine and rectum	153–154,197.5	8.3	9.2	7.3	8.9
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	7.4	7.8	5.8	7.8
Benign neoplasms	210–229	3.5	4.1	3.0	3.6
Benign neoplasm of uterus	218–219	2.9	3.6	2.4	2.9
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	4.0	4.6	3.8	3.9
Diabetes mellitus	250	4.7	5.5	4.0	4.9
Volume depletion	276.5	3.4	4.0	3.4	3.2
Diseases of the blood and blood-forming organs	280–289	4.4	4.7	3.4	4.5
Anemias	280–285	4.4	4.5	3.4	4.4
Mental disorders	290–319	6.9	8.6	5.4	6.9
Psychoses	290–299	7.6	9.5	6.0	7.5
Schizophrenic disorders	295	10.6	13.0	8.3	9.9
Major depressive disorder	296.2–296.3	6.9	9.2	5.4	6.7
Diseases of the nervous system and sense organs	320–389	5.2	5.7	4.4	5.1
Diseases of the circulatory system	390–459	4.6	4.9	4.1	4.7
Essential hypertension	401	2.2	2.3	2.2	2.3
Heart disease	391–392,0,393–398,402,404,410–416,420–429	4.5	4.8	4.0	4.7
Acute myocardial infarction	410	5.5	5.9	5.0	5.6
Coronary atherosclerosis	414.0	3.5	3.4	3.0	4.0
Other ischemic heart disease	411–413,414.1–414.9	2.4	2.5	2.5	2.3
Cardiac dysrhythmias	427	3.4	3.9	3.0	3.4
Congestive heart failure	428.0,428.2–428.4	5.1	5.6	4.7	5.2
Cerebrovascular disease	430–438	5.2	5.4	4.3	4.9
Diseases of the respiratory system	460–519	5.2	5.7	4.4	5.3
Acute bronchitis and bronchiolitis	466	3.2	3.9	3.1	3.0
Pneumonia	480–486	5.3	5.7	4.6	5.5
Chronic bronchitis	491	5.0	5.4	4.2	4.8
Asthma	493	3.3	3.5	2.7	3.7
Diseases of the digestive system	520–579	4.7	5.0	4.3	5.0
Appendicitis	540–543	3.1	3.4	2.8	3.0
Noninfectious enteritis and colitis	555–558	4.6	4.5	3.7	5.2
Intestinal obstruction	560	6.4	6.7	6.0	6.9
Diverticula of intestine	562	5.3	6.0	4.4	5.6
Cholelithiasis	574	4.2	4.8	3.4	4.1
Acute pancreatitis	577.0	5.1	5.2	4.6	5.4
Diseases of the genitourinary system	580–629	3.8	4.4	3.4	3.7
Calculus of kidney and ureter	592	2.3	2.2	2.1	2.4
Urinary tract infection	599.0	4.5	5.1	4.1	4.6
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	2.9	2.6	3.1	2.9
Diseases of the skin and subcutaneous tissue	680–709	5.0	5.5	4.4	4.8
Cellulitis and abscess	681–682	4.7	5.2	4.0	4.7
Diseases of the musculoskeletal system and connective tissue	710–739	3.9	4.1	3.6	3.9
Osteoarthritis and allied disorders	715	3.7	4.0	3.7	3.5
Intervertebral disc disorders	722	2.9	2.8	2.6	3.1
Congenital anomalies	740–759	6.8	6.8	8.2	5.8
Certain conditions originating in the perinatal period	760–779	10.5	10.2	8.5	11.8
Symptoms, signs, and ill-defined conditions	780–799	2.8	2.7	2.5	2.8
Injury and poisoning	800–999	5.2	5.2	4.5	5.8
Fractures, all sites ²	800–829	5.3	5.1	4.6	5.9
Fracture of neck of femur ²	820	6.3	6.7	5.8	6.6
Poisonings	960–989	3.0	4.1	2.5	2.8
Certain complications of surgical and medical care	996–999	6.0	6.1	5.5	6.5
Supplementary classifications	V01–V84	3.7	4.2	3.6	3.4
Females with deliveries	V27	2.6	2.8	2.5	2.5

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

²Excludes fractures coded as 733.1, pathologic fracture.

Table 20. Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15–44 years	45–64 years	65 years and over
Number in thousands					
All conditions	169,359	7,242	41,680	42,790	77,647
Infectious and parasitic diseases	001–139	4,331	466	901	1,182
Septicemia	038	861	24	89	213
Neoplasms	140–239	4,454	112	613	1,560
Malignant neoplasms	140–208,230–234	3,439	97	288	1,173
Malignant neoplasm of large intestine and rectum	153–154,197.5	258	*	10	84
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	494	*	20	165
Benign neoplasms	210–229	855	9	302	346
Benign neoplasm of uterus	218–219	434	*	217	203
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	20,325	622	2,619	6,270
Diabetes mellitus	250	5,480	40	611	1,881
Volume depletion	276.5	2,558	326	326	475
Diseases of the blood and blood-forming organs	280–289	5,109	186	1,045	1,293
Anemias	280–285	3,831	100	804	913
Mental disorders	290–319	12,695	356	5,330	3,911
Psychoses	290–299	4,154	107	1,486	977
Schizophrenic disorders	295	586	*	263	229
Major depressive disorder	296.2–296.3	653	24	341	196
Diseases of the nervous system and sense organs	320–389	3,942	373	678	912
Diseases of the circulatory system	390–459	34,677	140	2,104	8,997
Essential hypertension	401	9,003	18	710	2,869
Heart disease	391–392.0,393–398,402,404,410–416,420–429	19,747	76	923	4,568
Acute myocardial infarction	410	865	*	47	253
Coronary atherosclerosis	414.0	4,340	*	123	1,195
Other ischemic heart disease	411–413,414.1–414.9	1,988	*	99	642
Cardiac dysrhythmias	427	4,528	25	194	795
Congestive heart failure	428.0,428.2–428.4	4,141	10	127	804
Cerebrovascular disease	430–438	1,832	*	89	412
Diseases of the respiratory system	460–519	12,384	1,212	1,475	2,999
Acute bronchitis and bronchiolitis	466	396	198	37	55
Pneumonia	480–486	2,385	258	216	470
Chronic bronchitis	491	1,202	*	39	322
Asthma	493	1,928	347	521	535
Diseases of the digestive system	520–579	11,108	537	2,232	3,495
Appendicitis	540–543	352	80	179	68
Noninfectious enteritis and colitis	555–558	686	120	162	164
Intestinal obstruction	560	773	26	113	215
Diverticula of intestine	562	634	*	55	181
Cholelithiasis	574	579	*2	169	168
Acute pancreatitis	577.0	342	5	116	125
Diseases of the genitourinary system	580–629	9,073	204	2,065	2,209
Calculus of kidney and ureter	592	324	*	122	113
Urinary tract infection	599.0	2,230	47	238	389
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	10,394	19	10,353	22
Diseases of the skin and subcutaneous tissue	680–709	2,632	218	502	764
Cellulitis and abscess	681–682	1,174	84	266	389
Diseases of the musculoskeletal system and connective tissue	710–739	6,884	101	979	2,089
Osteoarthritis and allied disorders	715	1,809	*	66	500
Intervertebral disc disorders	722	560	*	166	247
Congenital anomalies	740–759	718	422	124	95
Certain conditions originating in the perinatal period	760–779	614	611	*	*
Symptoms, signs, and ill-defined conditions	780–799	8,829	771	1,951	2,522
Injury and poisoning	800–999	7,122	476	2,108	1,824
Fractures, all sites ²	800–829	1,642	95	461	332
Fracture of neck of femur ²	820	338	*	9	32
Poisonings	960–989	466	23	288	112
Certain complications of surgical and medical care	996–999	2,201	86	403	740
Supplementary classifications	V01–V84	14,070	416	6,600	2,645
Females with deliveries	V27	4,038	*6	4,025	*7

* Figure does not meet standards of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 21. Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnoses and ICD-9-CM code	Both sexes	Male	Female
Number in thousands			
All conditions	169,359	69,385	99,974
Infectious and parasitic diseases	001–139	4,331	1,955
Septicemia	038	861	428
Neoplasms	140–239	4,454	1,921
Malignant neoplasms	140–208,230–234	3,439	1,695
Malignant neoplasm of large intestine and rectum	153–154,197.5	258	125
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	494	267
Benign neoplasms	210–229	855	156
Benign neoplasm of uterus	218–219	434	434
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	20,325	8,570
Diabetes mellitus	250	5,480	2,477
Volume depletion	276.5	2,558	1,081
Diseases of the blood and blood-forming organs	280–289	5,109	2,032
Anemias	280–285	3,831	1,418
Mental disorders	290–319	12,695	6,005
Psychoses	290–299	4,154	1,878
Schizophrenic disorders	295	586	308
Major depressive disorder	296.2–296.3	653	253
Diseases of the nervous system and sense organs	320–389	3,942	1,708
Diseases of the circulatory system	390–459	34,677	16,812
Essential hypertension	401	9,003	3,841
Heart disease	391–392.0,393–398,402,404,410–416,420–429	19,747	10,090
Acute myocardial infarction	410	865	499
Coronary atherosclerosis	414.0	4,340	2,528
Other ischemic heart disease	411–413,414.1–414.9	1,988	1,197
Cardiac dysrhythmias	427	4,528	2,253
Congestive heart failure	428.0,428.2–428.4	4,141	1,850
Cerebrovascular disease	430–438	1,832	851
Diseases of the respiratory system	460–519	12,384	5,804
Acute bronchitis and bronchiolitis	466	396	190
Pneumonia	480–486	2,385	1,159
Chronic bronchitis	491	1,202	568
Asthma	493	1,928	653
Diseases of the digestive system	520–579	11,108	4,786
Appendicitis	540–543	352	187
Noninfectious enteritis and colitis	555–558	686	276
Intestinal obstruction	560	773	344
Diverticula of intestine	562	634	238
Cholelithiasis	574	579	213
Acute pancreatitis	577.0	342	181
Diseases of the genitourinary system	580–629	9,073	3,339
Calculus of kidney and ureter	592	324	160
Urinary tract infection	599.0	2,230	643
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	10,394	...
Diseases of the skin and subcutaneous tissue	680–709	2,632	1,323
Cellulitis and abscess	681–682	1,174	624
Diseases of the musculoskeletal system and connective tissue	710–739	6,884	2,457
Osteoarthritis and allied disorders	715	1,809	637
Intervertebral disc disorders	722	560	258
Congenital anomalies	740–759	718	371
Certain conditions originating in the perinatal period	760–779	614	344
Symptoms, signs, and ill-defined conditions	780–799	8,829	3,964
Injury and poisoning	800–999	7,122	3,694
Fractures, all sites ²	800–829	1,642	783
Fracture of neck of femur ²	820	338	96
Poisonings	960–989	466	206
Certain complications of surgical and medical care	996–999	2,201	1,068
Supplementary classifications	V01–V84	14,070	4,298
Females with deliveries	V27	4,038	4,038

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

²Excludes fractures coded as 733.1, pathologic fracture.

Table 22. Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnoses and ICD-9-CM code	United States	Northeast	Midwest	South	West
	Number in thousands				
All conditions	169,359	35,502	41,303	60,850	31,703
Infectious and parasitic diseases	001–139	4,331	936	937	1,617
Septicemia	038	861	178	189	324
Neoplasms	140–239	4,454	992	991	1,642
Malignant neoplasms	140–208,230–234	3,439	770	779	1,250
Malignant neoplasm of large intestine and rectum	153–154,197.5	258	58	61	40
Malignant neoplasm on trachea, bronchus, and lung	162,176.4,197.0,197.3	494	115	123	185
Benign neoplasms	210–229	855	182	174	334
Benign neoplasm of uterus	218–219	434	84	82	191
Endocrine, nutritional and metabolic diseases, and immunity disorders	240–279	20,325	4,338	5,228	7,146
Diabetes mellitus	250	5,480	1,236	1,330	1,992
Volume depletion	276.5	2,558	492	668	955
Diseases of the blood and blood-forming organs	280–289	5,109	1,008	1,221	1,876
Anemias	280–285	3,831	746	936	1,420
Mental disorders	290–319	12,695	3,108	3,229	4,470
Psychoses	290–299	4,154	1,052	1,005	1,563
Schizophrenic disorders	295	586	157	108	255
Major depressive disorder	296.2–296.3	653	141	162	55
Diseases of the nervous system and sense organs	320–389	3,942	849	999	1,354
Diseases of the circulatory system	390–459	34,677	7,822	8,659	12,260
Essential hypertension	401	9,003	2,044	2,211	3,243
Heart disease	391–392.0,393–398,402,404,410–416,420–429	19,747	4,541	5,039	6,843
Acute myocardial infarction	410	865	219	207	274
Coronary atherosclerosis	414.0	4,340	1,011	1,111	1,511
Other ischemic heart disease	411–413,414.1–414.9	1,988	458	487	694
Cardiac dysrhythmias	427	4,528	1,034	1,183	1,502
Congestive heart failure	428.0,428.2–428.4	4,141	882	1,112	1,508
Cerebrovascular disease	430–438	1,832	381	447	644
Diseases of the respiratory system	460–519	12,384	2,543	3,080	4,498
Acute bronchitis and bronchiolitis	466	396	68	94	147
Pneumonia	480–486	2,385	458	623	849
Chronic bronchitis	491	1,202	253	299	470
Asthma	493	1,928	472	459	653
Diseases of the digestive system	520–579	11,108	2,310	2,716	4,005
Appendicitis	540–543	352	69	81	106
Noninfectious enteritis and colitis	555–558	686	151	163	259
Intestinal obstruction	560	773	141	212	276
Diverticula of intestine	562	634	151	172	209
Cholelithiasis	574	579	115	140	204
Acute pancreatitis	577.0	342	64	76	139
Diseases of the genitourinary system	580–629	9,073	1,829	2,125	3,474
Calculus of kidney and ureter	592	324	75	82	128
Urinary tract infection	599.0	2,230	465	535	840
Complications of pregnancy, childbirth, and the puerperium ¹	630–677	10,394	1,539	2,153	4,080
Diseases of the skin and subcutaneous tissue	680–709	2,632	587	594	975
Cellulitis and abscess	681–682	1,174	263	277	428
Diseases of the musculoskeletal system and connective tissue	710–739	6,884	1,457	1,858	2,270
Osteoarthritis and allied disorders	715	1,809	392	510	586
Intervertebral disc disorders	722	560	109	147	204
Congenital anomalies	740–759	718	157	142	237
Certain conditions originating in the perinatal period	760–779	614	80	* ²	215
Symptoms, signs, and ill-defined conditions	780–799	8,829	1,863	2,377	3,010
Injury and poisoning	800–999	7,122	1,363	1,702	2,536
Fractures, all sites ²	800–829	1,642	296	418	580
Fracture of neck of femur ²	820	338	63	83	126
Poisonings	960–989	466	84	129	169
Certain complications of surgical and medical care	996–999	2,201	485	458	782
Supplementary classifications	V01–V84	14,070	2,720	3,167	5,187
Females with deliveries	V27	4,038	591	824	1,574

* Figure does not meet standards of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."

²Excludes fractures coded as 733.1, pathologic fracture.

Table 23. Number and rate of discharges from short-stay hospitals with a human immunodeficiency virus diagnosis, by selected characteristics: United States, selected years, 1990–2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Data are for discharges with at least one of the following *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM) codes: 042–044, 279.19, 795.8 (prior to 1995); 042, V08 (beginning in 1995)]

Characteristic	1990	1995	2000	2001	2002	2003	2004	2005
Number in thousands								
All HIV discharges ¹	146	249	173	185	189	207	204	185
Sex								
Male	114	183	115	115	124	136	132	113
Female	32	66	58	70	65	71	73	72
Age								
Under 25 years	14	17	7	10	10	9	10	8
25–29 years	20	26	11	8	11	13	8	9
30–34 years	38	48	28	28	23	21	15	11
35–39 years	36	59	35	41	35	38	35	26
40–44 years	17	49	34	43	42	38	46	45
45 years and over	21	51	58	56	68	87	90	85
Region								
Northeast	66	95	68	61	57	70	67	71
Midwest	20	27	20	18	20	17	21	18
South	37	87	64	86	81	94	99	79
West	24	41	21	21	31	27	17	17
Rate per 100,000 population ²								
All HIV discharges ¹	58.1	93.9	61.6	65.2	65.8	71.6	69.9	62.6
Sex								
Male	93.4	141.6	83.9	82.8	88.5	95.9	91.7	77.7
Female	24.7	48.4	40.1	48.2	44.1	48.2	48.9	48.1
Age								
Under 25 years	14.8	17.6	7.5	10.1	9.5	8.9	10.2	8.2
25–29 years	95.1	132.2	56.6	41.3	57.5	71.3	38.9	43.3
30–34 years	172.8	216.2	134.9	137.6	113.0	104.6	71.9	56.6
35–39 years	179.0	262.7	154.4	182.9	159.7	180.8	168.3	125.8
40–44 years	94.8	241.4	152.6	187.4	185.2	166.6	201.3	197.6
45 years and over	27.4	60.2	59.8	55.8	66.6	83.1	84.6	77.9
Region								
Northeast	128.8	183.3	126.6	112.3	107.9	129.4	122.1	130.2
Midwest	33.6	43.5	30.8	27.9	31.7	25.9	32.6	26.8
South	42.1	93.1	64.2	84.5	81.7	90.1	94.3	74.3
West	44.1	69.5	33.4	32.6	49.9	40.2	25.7	24.7

¹HIV is human immunodeficiency virus.

²Rates are based on the U.S. Census Bureau estimates of the civilian population. Populations for 1990–1995 were adjusted for the net underenumeration in the 1990 census. Population for 2000–2005 were 2000-based postcensal estimates.

Table 24. Annual average number and rate of discharges from short-stay hospitals with a first-listed injury diagnosis, by Barell injury diagnosis matrix categories: United States, 2003–2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. *International Classification of Diseases, Ninth Revision, Clinical Modification* code numbers for Barell matrix categories are shown in Table III.]

Nature of injury	All body regions	Traumatic brain injury	Other head, face, and neck	Spinal cord	Vertebral column	Torso	Upper extremities	Lower extremities	Other and unspecified
Number in thousands ¹									
All injuries	1,934	200	126	11	99	242	248	680	25
Fracture	1,033	53	38	8	86	99	160	590	*
Dislocation	16	...	*	...	2	*	5	7	—
Sprains and strains	51	...	-	...	10	3	16	20	*2
Internal	232	147	...	4	...	81	*
Open wound	95	...	38	13	25	18	*
Amputation	9	7	*1	...
Blood vessels	8	...	*	*2	3	*1	*
Contusions and superficial injuries	65	...	15	20	6	21	3
Crush	4	...	*	*	*2	*	—
Burns	*	...	*	*	*	*	*
Nerves and spinal cord	3	*	*	*	*2	...	*
Unspecified	32	...	20	6	*	2	*3
System-wide	303
Rate per 10,000 population ²									
All injuries	66.1	6.8	4.3	0.4	3.4	8.3	8.5	23.2	0.9
Fracture	35.3	1.8	1.3	0.3	2.9	3.4	5.5	20.2	*
Dislocation	0.5	...	*	...	0.1	*	0.2	0.2	—
Sprains and strains	1.7	...	-	...	0.3	0.1	0.5	0.7	*0.1
Internal	7.9	5.0	...	0.1	...	2.8	*
Open wound	3.2	...	1.3	0.4	0.8	0.6	*
Amputation	0.3	0.3	*0.0	...
Blood vessels	0.3	...	*	*0.1	0.1	*0.0	*
Contusions and superficial injuries	2.2	...	0.5	0.7	0.2	0.7	0.1
Crush	0.1	...	*	*	*0.1	*	—
Burns	*	...	*	*	*	*	*
Nerves and spinal cord	0.1	*	*	*	*0.1	...	*
Unspecified	1.1	...	0.7	0.2	*	0.1	*0.1
System-wide	10.4

* Figure does not meet standards of reliability or precision.

... Category not applicable.

— Quantity zero.

0.0 Quantity more than zero but less than 0.05.

¹The annual average number of discharges was computed by adding the number of discharges with a first-listed injury diagnosis during the 3-year period and dividing by three.

²Rates were calculated by dividing the annual average number of discharges by the 2004 population. The population is the U.S. Census Bureau 2000-based postcensal estimate of the civilian population.

Table 25. Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnosis: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groups and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

First-listed diagnosis and ICD-9-CM code	All ages	Under 65 years	65 years and over	All ages	Under 65 years	65 years and over
	Number of deaths in thousands			Rate per 100 discharges		
All conditions ¹	752	200	552	2.2	0.9	4.2
Heart disease 391–392,0,393–398,402,404,410–416,420–429	117	18	100	2.8	1.1	3.7
Acute myocardial infarction 410	54	7	47	7.9	2.9	10.9
Cardiac dysrhythmias 427	10	*2	8	1.3	*1.1	1.5
Congestive heart failure 428.0,428.2–428.4	33	*	29	3.0	*	3.6
Septicemia 038	94	22	72	19.3	14.4	21.5
Malignant neoplasms 140–208,230–234	81	31	50	6.8	5.3	8.3
Malignant neoplasm on trachea, bronchus, and lung 162,176.4,197.0,197.3	15	6	9	9.5	9.3	9.7
Cerebrovascular disease 430–438	58	18	40	6.5	6.6	6.4
Respiratory failure 518.81,518.83,518.84	57	12	45	19.3	10.6	25.1
Pneumonia 480–486	45	7	38	3.3	1.3	4.6
Pneumonitis due to solids and liquids 507	27	*	25	15.2	*	18.9
Fractures, all sites ² 800–829	22	*6	16	2.1	*1.2	3.0
Fracture of neck of femur ² 820	11	*	10	3.5	*	3.9
Nephritis, nephrotic syndrome and nephrosis 580–589	20	*3	17	6.5	*3.0	8.3

* Figure does not meet standards of reliability or precision.

¹Includes data for deaths not shown in table.

²Excludes fractures coded as 733.1, pathologic fracture.

Table 26. Number of discharges from short-stay hospitals with and without procedures and percentage with procedures, by selected characteristics: United States, 2005

Characteristic	All discharges	Discharges without procedures	Discharges with procedures		
			Total	With surgical procedures	Total
		Number in thousands			Percent
All patients	34,667	13,342	21,325	15,782	61.5
Age					
Under 15 years	2,431	1,270	1,160	647	47.7
15–44 years	10,659	3,091	7,567	5,936	71.0
45–64 years	8,349	3,127	5,223	3,985	62.6
65 years and over	13,228	5,854	7,375	5,214	55.7
65–74 years	4,900	1,819	3,081	2,323	47.4
75–84 years	5,394	2,432	2,962	2,073	54.9
85 years and over	2,934	1,603	1,331	818	27.9
Sex					
Male	13,902	5,787	8,115	5,807	58.4
Female	20,766	7,555	13,210	9,975	63.6
Region					
Northeast	7,192	2,663	4,529	2,979	63.0
Midwest	7,948	3,388	4,560	3,425	57.4
South	12,907	5,202	7,705	5,930	59.7
West	6,621	2,090	4,531	3,448	52.1

NOTE: All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.

Table 27. Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals, by selected procedure categories: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Number in thousands	Rate per 10,000 population ¹
Surgical procedures ²	26,564	899.9
Cesarean section ³	1,262	42.8
Repair of current obstetric laceration ³	1,259	42.6
Cardiac catheterization	1,209	41.0
Artificial rupture of membranes ³	928	31.4
Balloon angioplasty of coronary artery or coronary atherectomy	645	21.9
Reduction of fracture	621	21.0
Open reduction of fracture with internal fixation	424	14.4
Insertion of coronary artery stent(s)	620	21.0
Endoscopy of small intestine with biopsy	620	21.0
Hysterectomy	575	19.5
Episiotomy with or without forceps or vacuum extraction ³	537	18.2
Total knee replacement	534	18.1
Coronary artery bypass graft ⁴	466	15.8
Oophorectomy and salpingo-oophorectomy	434	14.7
Cholecystectomy	398	13.5
Insertion, replacement, removal, and revision of pacemaker leads or device	384	13.0
Debridement of wound, infection, or burn	358	12.1
Lysis of peritoneal adhesions	348	11.8
Bilateral destruction or occlusion of fallopian tubes	346	11.7
Appendectomy, excluding incidental	341	11.6
Excision or destruction of intervertebral disc	294	9.9
Partial excision of large intestine	266	9.0
Total hip replacement	235	7.9
Endoscopy of large intestine with biopsy	218	7.4
Bronchoscopy with biopsy	191	6.5
Partial excision of bone	178	6.0
Prostatectomy	155	5.3
Cystoscopy with biopsy	12	0.4
Nonsurgical procedures ⁵	18,387	622.9
Arteriography and angiography using contrast material	1,986	67.3
Transfusion of blood and blood components	1,902	64.4
Respiratory therapy	1,160	39.3
Diagnostic ultrasound	884	30.0
Computerized axial tomography	811	27.5
Medical induction of labor ³	616	20.9
Hemodialysis	593	20.1
Insertion of endotracheal tube	506	17.2
Endoscopy of small intestine without biopsy	444	15.1
Physical therapy procedures	406	13.8
Spinal tap	343	11.6
Endoscopy of large intestine without biopsy	338	11.5
Injection or infusion of cancer chemotherapeutic substance	209	7.1
Cystoscopy without biopsy	130	4.4
Bronchoscopy without biopsy	92	3.1

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.

³See Table 38 for rate per 100 deliveries.

⁴See Table 29 for comparison of number of procedures with number of discharges with procedures.

⁵Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.

Table 28. Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	65 years and over	65–74 years	75–84 years	85 years and over
Number in thousands				
Surgical procedures ¹	8,858	4,118	3,500	1,240
Cardiac catheterization	37.21–37.23	605	319	241
Total knee replacement	81.54	331	195	121
Endoscopy of small intestine with biopsy	45.14,45.16	328	107	144
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02,36.05	326	173	129
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7–37.8	325	74	159
Insertion of coronary artery stent(s)	36.06–36.07	311	167	121
Coronary artery bypass graft ²	36.1	264	155	101
Reduction of fracture	76.7,79.0–79.3	254	73	95
Open reduction of fracture with internal fixation	79.3	184	59	65
Partial excision of large intestine	45.7	142	67	53
Total hip replacement	81.51	139	69	58
Cholecystectomy	51.2	136	64	57
Debridement of wound, infection, or burn	86.22,86.28	119	58	39
Prostatectomy	60.2–60.6	102	56	35
Nonsurgical procedures ³	7,236	2,875	2,974	1,387
Transfusion of blood and blood components	99.0	1,144	388	485
Arteriography and angiography using contrast material	88.4–88.5	997	494	409
Respiratory therapy	93.9,96.7	512	207	207
Diagnostic ultrasound	88.7	441	157	184
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	353	121	152
Physical therapy procedures	93.1–93.3	277	88	117
Hemodialysis	39.95	270	146	101
Insertion of endotracheal tube	96.04	249	99	101
Endoscopy of small intestine without biopsy	45.11–45.13	245	90	99
Endoscopy of large intestine without biopsy	45.21–45.24	209	76	87
Rate per 10,000 population ⁴				
Surgical procedures ¹	2,407.6	2,209.2	2,681.0	2,432.6
Cardiac catheterization	37.21–37.23	164.4	171.2	184.4
Total knee replacement	81.54	90.1	104.5	92.8
Endoscopy of small intestine with biopsy	45.14,45.16	89.1	57.6	110.2
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02,36.05	88.6	92.9	98.9
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7–37.8	88.4	39.5	122.1
Insertion of coronary artery stent(s)	36.06–36.07	84.5	89.8	92.8
Coronary artery bypass graft ²	36.1	71.8	83.3	77.7
Reduction of fracture	76.7,79.0–79.3	68.9	39.1	73.0
Open reduction of fracture with internal fixation	79.3	49.9	31.4	50.1
Partial excision of large intestine	45.7	38.5	36.1	40.2
Total hip replacement	81.51	37.8	36.8	44.4
Cholecystectomy	51.2	36.9	34.4	43.3
Debridement of wound, infection, or burn	86.22,86.28	32.4	31.2	29.5
Prostatectomy	60.2–60.6	27.8	30.1	26.5
Nonsurgical procedures ³	1,966.8	1,542.3	2,278.2	2,721.9
Transfusion of blood and blood components	99.0	311.0	208.0	371.7
Arteriography and angiography using contrast material	88.4–88.5	271.1	265.1	313.3
Respiratory therapy	93.9,96.7	139.1	111.0	158.2
Diagnostic ultrasound	88.7	120.0	84.3	140.9
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	96.0	65.0	116.7
Physical therapy procedures	93.1–93.3	75.4	47.1	89.9
Hemodialysis	39.95	73.5	78.5	77.3
Insertion of endotracheal tube	96.04	67.8	53.2	77.7
Endoscopy of small intestine without biopsy	45.11–45.13	66.5	48.1	76.0
Endoscopy of large intestine without biopsy	45.21–45.24	56.9	40.8	66.4

¹Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.

²See Table 29 for a comparison of number of procedures with number of discharges with procedures.

³Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.

⁴Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

Table 29. Number and rate of discharges from short-stay hospitals with a coronary artery bypass graft and number and rate of procedures, by sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification*]

Sex, age, and region	Discharges with procedures		Procedures	
	Number in thousands	Rate per 10,000 population ¹	Number in thousands	Rate per 10,000 population ¹
Total	261	8.8	466	15.8
Sex				
Male	178	12.3	321	22.2
Female	83	5.5	144	9.6
Age				
Under 15 years	*	*	*	*
15–44 years	8	0.7	15	1.2
45–64 years	102	14.0	187	25.6
65 years and over	151	41.0	264	71.8
65–74 years	88	47.4	155	83.3
75–84 years	58	44.4	101	77.7
85 years and over	*	*	*	*
Region				
Northeast	44	8.1	77	14.2
Midwest	61	9.2	111	16.8
South	98	9.1	174	16.3
West	58	8.6	103	15.1

* Figure does not meet standards of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

NOTE: The number of procedures exceeds the number of discharges because these procedures are often assigned more than one code for a single operation. See "Methods" section.

Table 30. Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15–44 years	45–64 years	65 years and over
Number in thousands					
All procedures	44,950	2,203	14,744	11,910	16,093
Operations on the nervous system	01–05	1,227	202	358	310
Spinal tap03.31	343	131	86	66
Operations on the endocrine system	06–07	104	*2	27	47
Operations on the eye	08–16	74	14	20	16
Operations on the ear	18–20	42	22	9	*4
Operations on the nose, mouth, and pharynx	21–29	302	64	91	72
Operations on the respiratory system	30–34	1,118	64	168	370
Bronchoscopy with or without biopsy	33.21–33.24,33.27	284	18	36	96
Operations on the cardiovascular system	35–39	6,989	200	655	2,524
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02,36.05	645	*	34	285
Insertion of coronary artery stent(s)	36.06–36.07	620	*	34	275
Coronary artery bypass graft ¹	36.1	466	*	15	187
Cardiac catheterization	37.21–37.23	1,209	9	86	509
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7–37.8	384	*	*9	44
Hemodialysis39.95	593	*	76	245
Operations on the hemic and lymphatic system	40–41	343	17	57	131
Operations on the digestive system	42–54	5,580	246	1,244	1,716
Endoscopy of small intestine with or without biopsy45.11–45.14,45.16	1,064	19	155	317
Endoscopy of large intestine with or without biopsy	45.21–45.25	556	7	77	144
Partial excision of large intestine45.7	266	*	32	89
Appendectomy, excluding incidental	47.0	341	80	170	65
Cholecystectomy	51.2	398	*	133	128
Lysis of peritoneal adhesions	54.5	348	6	133	114
Operations on the urinary system	55–59	999	36	236	314
Cystoscopy with or without biopsy	57.31–57.33	142	*	28	44
Operations on the male genital organs	60–64	233	23	15	64
Prostatectomy	60.2–60.6	155	*	*	51
Operations on the female genital organs	65–71	1,967	10	1,145	613
Oophorectomy and salpingo-oophorectomy	65.3–65.6	434	*	183	204
Bilateral destruction or occlusion of fallopian tubes	66.2–66.3	346	*	344	*
Hysterectomy	68.3–68.7,68.9	575	*	289	240
Obstetrical procedures	72–75	6,858	*11	6,838	*
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	537	*	535	...
Artificial rupture of membranes	73.0	928	*	926	*
Medical induction of labor	73.4	616	*	614	*
Cesarean section	74.0–74.2,74.4,74.99	1,262	*	1,258	*
Repair of current obstetric laceration	75.5–75.6	1,259	*	1,254	*
Operations on the musculoskeletal system	76–84	4,139	168	893	1,444
Partial excision of bone	76.2–76.3,77.6–77.8	178	10	46	71
Reduction of fracture	76.7,79.0–79.3	621	44	185	138
Open reduction of fracture with internal fixation	79.3	424	17	121	102
Excision or destruction of intervertebral disc	80.5	294	*	101	148
Total hip replacement81.51	235	*	15	80
Total knee replacement81.54	534	*	8	194
Operations on the integumentary system	85–86	1,538	*	476	478
Debridement of wound, infection, or burn86.22,86.28	358	*	101	113
Miscellaneous diagnostic and therapeutic procedures and new technologies ²	87–99.00	13,437	954	2,512	3,792
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	811	38	191	229
Arteriography and angiography using contrast material	88.4–88.5	1,986	14	180	794
Diagnostic ultrasound	88.7	884	*	150	249
Physical therapy procedures	93.1–93.3	406	10	39	81
Respiratory therapy	93.9,96.7	1,160	188	173	288
Insertion of endotracheal tube96.04	506	40	81	136
Transfusion of blood and blood components99.0	1,902	67	220	472
Injection or infusion of cancer chemotherapeutic substance99.25	209	42	39	73

* Figure does not meet standards of reliability or precision.

... Category not applicable.

¹See Table 29 for comparison of number of procedures with number of discharges with procedures.

²Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of tables."

Table 31. Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15–44 years	45–64 years	65 years and over
Rate per 10,000 population ¹					
All procedures	1,522.7	362.9	1,180.4	1,636.2	4,374.4
Operations on the nervous system	01–05	41.6	33.3	28.7	42.6
Spinal tap	03.31	11.6	21.6	6.9	9.1
Operations on the endocrine system	06–07	3.5	*0.4	2.2	6.4
Operations on the eye	08–16	2.5	2.2	1.6	2.2
Operations on the ear	18–20	1.4	3.7	0.7	*0.6
Operations on the nose, mouth, and pharynx	21–29	10.2	10.5	7.3	9.9
Operations on the respiratory system	30–34	37.9	10.5	13.5	50.9
Bronchoscopy with or without biopsy	33.21–33.24,33.27	9.6	3.0	2.9	13.2
Operations on the cardiovascular system	35–39	236.8	33.0	52.5	346.8
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02,36.05	21.9	*	2.7	39.1
Insertion of coronary artery stent(s)	36.06–36.07	21.0	*	2.8	37.7
Coronary artery bypass graft ²	36.1	15.8	*	1.2	25.6
Cardiac catheterization	37.21–37.23	41.0	1.5	6.9	69.9
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7–37.8	13.0	*	*0.7	6.1
Hemodialysis	39.95	20.1	*	6.1	33.7
Operations on the hemic and lymphatic system	40–41	11.6	2.8	4.6	18.8
Operations on the digestive system	42–54	189.0	40.6	99.6	235.8
Endoscopy of small intestine with or without biopsy	45.11–45.14,45.16	36.0	3.2	12.4	43.6
Endoscopy of large intestine with or without biopsy	45.21–45.25	18.8	1.1	6.1	19.8
Partial excision of large intestine	45.7	9.0	*	2.6	12.2
Appendectomy, excluding incidental	47.0	11.6	13.1	13.6	9.0
Cholecystectomy	51.2	13.5	*	10.6	17.6
Lysis of peritoneal adhesions	54.5	11.8	1.0	10.6	15.6
Operations on the urinary system	55–59	33.8	6.0	18.9	43.2
Cystoscopy with or without biopsy	57.31–57.33	4.8	*	2.2	6.0
Operations on the male genital organs	60–64	7.9	3.8	1.2	8.8
Prostatectomy	60.2–60.6	5.3	*	*	7.0
Operations on the female genital organs	65–71	66.6	1.6	91.7	84.2
Oophorectomy and salpingo-oophorectomy	65.3–65.6	14.7	*	14.6	28.0
Bilateral destruction or occlusion of fallopian tubes	66.2–66.3	11.7	*	27.6	*
Hysterectomy	68.3–68.7,68.9	19.5	*	23.1	33.0
Obstetrical procedures	72–75	232.3	*1.9	547.4	*
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	18.2	*	42.8	*
Artificial rupture of membranes	73.0	31.4	*	74.1	*
Medical induction of labor	73.4	20.9	*	49.2	*
Cesarean section	74.0–74.2,74.4,74.99	42.8	*	100.7	*
Repair of current obstetric laceration	75.5–75.6	42.6	*	100.4	*
Operations on the musculoskeletal system	76–84	140.2	27.6	71.5	198.4
Partial excision of bone	76.2–76.3,77.6–77.8	6.0	1.6	3.7	9.8
Reduction of fracture	76.7,79.0–79.3	21.0	7.2	14.8	19.0
Open reduction of fracture with internal fixation	79.3	14.4	2.9	9.7	14.0
Excision or destruction of intervertebral disc	80.5	9.9	*	8.1	20.4
Total hip replacement	81.51	7.9	*	1.2	11.0
Total knee replacement	81.54	18.1	*	0.6	26.7
Operations on the integumentary system	85–86	52.1	*	38.1	65.6
Debridement of wound, infection, or burn	86.22,86.28	12.1	*	8.0	15.5
Miscellaneous diagnostic and therapeutic procedures and new technologies ³	87–99.00	455.2	157.2	201.1	520.9
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	27.5	6.2	15.3	31.5
Arteriography and angiography using contrast material	88.4–88.5	67.3	2.3	14.4	109.1
Diagnostic ultrasound	88.7	30.0	*	12.0	34.2
Physical therapy procedures	93.1–93.3	13.8	1.6	3.1	11.1
Respiratory therapy	93.9,96.7	39.3	31.0	13.8	39.5
Insertion of endotracheal tube	96.04	17.2	6.5	6.5	18.7
Transfusion of blood and blood components	99.0	64.4	11.0	17.6	64.8
Injection or infusion of cancer chemotherapeutic substance	99.25	7.1	6.9	3.1	10.1

* Figure does not meet standards of reliability or precision.

. . . Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²See Table 29 for comparison of rate of procedures with rate of discharges with procedures.

³Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 32. Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code		Both sexes	Male	Female
		Number in thousands		
All procedures		44,950	17,991	26,959
Operations on the nervous system	01–05	1,227	569	658
Spinal tap03.31	343	168	175
Operations on the endocrine system	06–07	104	29	75
Operations on the eye	08–16	74	40	34
Operations on the ear	18–20	42	27	15
Operations on the nose, mouth, and pharynx	21–29	302	181	121
Operations on the respiratory system	30–34	1,118	617	501
Bronchoscopy with or without biopsy	33.21–33.24, 33.27	284	153	131
Operations on the cardiovascular system	35–39	6,989	4,062	2,927
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02, 36.05	645	443	202
Insertion of coronary artery stent(s)	36.06–36.07	620	429	192
Coronary artery bypass graft ¹	36.1	466	321	144
Cardiac catheterization	37.21–37.23	1,209	742	467
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7–37.8	384	200	184
Hemodialysis39.95	593	311	282
Operations on the hemic and lymphatic system	40–41	343	168	175
Operations on the digestive system	42–54	5,580	2,414	3,166
Endoscopy of small intestine with or without biopsy	45.11–45.14, 45.16	1,064	481	583
Endoscopy of large intestine with or without biopsy	45.21–45.25	556	222	334
Partial excision of large intestine45.7	266	123	143
Appendectomy, excluding incidental47.0	341	178	163
Cholecystectomy51.2	398	143	256
Lysis of peritoneal adhesions54.5	348	76	272
Operations on the urinary system55–59	999	466	533
Cystoscopy with or without biopsy57.31–57.33	142	67	74
Operations on the male genital organs60–64	233	233	...
Prostatectomy60.2–60.6	155	155	...
Operations on the female genital organs65–71	1,967	...	1,967
Oophorectomy and salpingo-oophorectomy65.3–65.6	434	...	434
Bilateral destruction or occlusion of fallopian tubes66.2–66.3	346	...	346
Hysterectomy68.3–68.7, 68.9	575	...	575
Obstetrical procedures72–75	6,858	...	6,858
Episiotomy with or without forceps or vacuum extraction72.1, 72.21, 72.31, 72.71, 73.6	537	...	537
Artificial rupture of membranes73.0	928	...	928
Medical induction of labor73.4	616	...	616
Cesarean section74.0–74.2, 74.4, 74.99	1,262	...	1,262
Repair of current obstetric laceration75.5–75.6	1,259	...	1,259
Operations on the musculoskeletal system76–84	4,139	1,972	2,167
Partial excision of bone76.2–76.3, 77.6–77.8	178	99	79
Reduction of fracture76.7, 79.0–79.3	621	293	328
Open reduction of fracture with internal fixation79.3	424	190	235
Excision or destruction of intervertebral disc80.5	294	148	146
Total hip replacement81.51	235	109	126
Total knee replacement81.54	534	184	349
Operations on the integumentary system85–86	1,538	731	807
Debridement of wound, infection, or burn86.22, 86.28	358	205	153
Miscellaneous diagnostic and therapeutic procedures and new technologies ²87–99.00	13,437	6,481	6,956
Computerized axial tomography87.03, 87.41, 87.71, 88.01, 88.38	811	378	434
Arteriography and angiography using contrast material88.4–88.5	1,986	1,125	860
Diagnostic ultrasound88.7	884	411	473
Physical therapy procedures93.1–93.3	406	167	240
Respiratory therapy93.9, 96.7	1,160	606	554
Insertion of endotracheal tube96.04	506	268	238
Transfusion of blood and blood components99.0	1,902	788	1,115
Injection or infusion of cancer chemotherapeutic substance99.25	209	116	93

... Category not applicable.

¹See Table 29 for comparison of number of procedures with number of discharges with procedures.

²Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 33. Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code		Both sexes	Male	Female
		Rate per 10,000 population ¹		
All procedures		1,522.7	1,241.1	1,794.5
Operations on the nervous system	01–05	41.6	39.3	43.8
Spinal tap03.31	11.6	11.6	11.7
Operations on the endocrine system	06–07	3.5	2.0	5.0
Operations on the eye	08–16	2.5	2.8	2.3
Operations on the ear	18–20	1.4	1.8	1.0
Operations on the nose, mouth, and pharynx	21–29	10.2	12.5	8.1
Operations on the respiratory system	30–34	37.9	42.6	33.4
Bronchoscopy with or without biopsy	33.21–33.24,33.27	9.6	10.5	8.7
Operations on the cardiovascular system	35–39	236.8	280.2	194.8
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02,36.05	21.9	30.6	13.4
Insertion of coronary artery stent(s)36.06–36.07	21.0	29.6	12.8
Coronary artery bypass graft ²36.1	15.8	22.2	9.6
Cardiac catheterization37.21–37.23	41.0	51.2	31.1
Insertion, replacement, removal, and revision of pacemaker leads or device37.7–37.8	13.0	13.8	12.2
Hemodialysis39.95	20.1	21.4	18.8
Operations on the hemic and lymphatic system40–41	11.6	11.6	11.6
Operations on the digestive system42–54	189.0	166.5	210.7
Endoscopy of small intestine with or without biopsy45.11–45.14,45.16	36.0	33.2	38.8
Endoscopy of large intestine with or without biopsy45.21–45.25	18.8	15.3	22.2
Partial excision of large intestine45.7	9.0	8.5	9.5
Appendectomy, excluding incidental47.0	11.6	12.3	10.9
Cholecystectomy51.2	13.5	9.8	17.0
Lysis of peritoneal adhesions54.5	11.8	5.2	18.1
Operations on the urinary system55–59	33.8	32.2	35.5
Cystoscopy with or without biopsy57.31–57.33	4.8	4.7	4.9
Operations on the male genital organs60–64	7.9	16.1	...
Prostatectomy60.2–60.6	5.3	10.7	...
Operations on the female genital organs65–71	66.6	...	130.9
Oophorectomy and salpingo-oophorectomy65.3–65.6	14.7	...	28.9
Bilateral destruction or occlusion of fallopian tubes66.2–66.3	11.7	...	23.0
Hysterectomy68.3–68.7,68.9	19.5	...	38.3
Obstetrical procedures72–75	232.3	...	456.5
Episiotomy with or without forceps or vacuum extraction72.1,72.21,72.31,72.71,73.6	18.2	...	35.7
Artificial rupture of membranes73.0	31.4	...	61.8
Medical induction of labor73.4	20.9	...	41.0
Cesarean section74.0–74.2,74.4,74.99	42.8	...	84.0
Repair of current obstetric laceration75.5–75.6	42.6	...	83.8
Operations on the musculoskeletal system76–84	140.2	136.0	144.2
Partial excision of bone76.2–76.3,77.6–77.8	6.0	6.8	5.3
Reduction of fracture76.7,79.0–79.3	21.0	20.2	21.8
Open reduction of fracture with internal fixation79.3	14.4	13.1	15.6
Excision or destruction of intervertebral disc80.5	9.9	10.2	9.7
Total hip replacement81.51	7.9	7.5	8.4
Total knee replacement81.54	18.1	12.7	23.3
Operations on the integumentary system85–86	52.1	50.4	53.7
Debridement of wound, infection, or burn86.22,86.28	12.1	14.2	10.2
Miscellaneous diagnostic and therapeutic procedures and new technologies ³87–99.00	455.2	447.1	463.0
Computerized axial tomography87.03,87.41,87.71,88.01,88.38	27.5	26.1	28.9
Arteriography and angiography using contrast material88.4–88.5	67.3	77.6	57.3
Diagnostic ultrasound88.7	30.0	28.4	31.5
Physical therapy procedures93.1–93.3	13.8	11.5	16.0
Respiratory therapy93.9,96.7	39.3	41.8	36.9
Insertion of endotracheal tube96.04	17.2	18.5	15.8
Transfusion of blood and blood components99.0	64.4	54.3	74.2
Injection or infusion of cancer chemotherapeutic substance99.25	7.1	8.0	6.2

... Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²See Table 29 for comparison of rate of procedures with rate of discharges with procedures.

³Chapter 00, "Procedures and interventions not else where classified," is included in this category; see "Use of Tables."

Table 34. Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	Northeast	Midwest	South	West
Number in thousands					
All procedures	44,950	9,689	9,770	15,962	9,529
Operations on the nervous system	01–05	1,227	253	324	384
Spinal tap	03.31	343	64	86	123
Operations on the endocrine system	06–07	104	22	20	36
Operations on the eye	08–16	74	17	12	25
Operations on the ear	18–20	42	9	* ¹	14
Operations on the nose, mouth, and pharynx	21–29	302	66	42	116
Operations on the respiratory system	30–34	1,118	248	230	410
Bronchoscopy with or without biopsy	33.21–33.24,33.27	284	63	61	94
Operations on the cardiovascular system	35–39	6,989	1,340	1,550	2,608
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02,36.05	645	119	182	213
Insertion of coronary artery stent(s)	36.06–36.07	620	105	176	208
Coronary artery bypass graft ¹	36.1	466	77	111	174
Cardiac catheterization	37.21–37.23	1,209	210	289	468
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7–37.8	384	107	74	114
Hemodialysis	39.95	593	112	99	228
Operations on the hemic and lymphatic system	40–41	343	66	68	125
Operations on the digestive system	42–54	5,580	1,146	1,237	2,014
Endoscopy of small intestine with or without biopsy	45.11–45.14,45.16	1,064	214	237	387
Endoscopy of large intestine with or without biopsy	45.21–45.25	556	122	125	195
Partial excision of large intestine	45.7	266	52	66	95
Appendectomy, excluding incidental	47.0	341	67	79	104
Cholecystectomy	51.2	398	74	96	154
Lysis of peritoneal adhesions	54.5	348	66	82	141
Operations on the urinary system	55–59	999	241	183	399
Cystoscopy with or without biopsy	57.31–57.33	142	39	26	57
Operations on the male genital organs	60–64	233	47	53	77
Prostatectomy	60.2–60.6	155	32	35	46
Operations on the female genital organs	65–71	1,967	276	408	888
Oophorectomy and salpingo-oophorectomy	65.3–65.6	434	58	104	191
Bilateral destruction or occlusion of fallopian tubes	66.2–66.3	346	36	60	173
Hysterectomy	68.3–68.7,68.9	575	85	130	247
Obstetrical procedures	72–75	6,858	1,045	1,563	2,509
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	537	82	100	220
Artificial rupture of membranes	73.0	928	120	275	317
Medical induction of labor	73.4	616	88	144	225
Cesarean section	74.0–74.2,74.4,74.99	1,262	190	232	530
Repair of current obstetric laceration	75.5–75.6	1,259	193	284	435
Operations on the musculoskeletal system	76–84	4,139	782	971	1,472
Partial excision of bone	76.2–76.3,77.6–77.8	178	36	34	67
Reduction of fracture	76.7,79.0–79.3	621	124	146	225
Open reduction of fracture with internal fixation	79.3	424	85	98	157
Excision or destruction of intervertebral disc	80.5	294	53	62	109
Total hip replacement	81.51	235	54	60	61
Total knee replacement	81.54	534	109	153	168
Operations on the integumentary system	85–86	1,538	288	295	691
Debridement of wound, infection, or burn	86.22,86.28	358	67	69	161
Miscellaneous diagnostic and therapeutic procedures and new technologies ²	87–99.00	13,437	3,843	2,805	4,194
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	811	386	*	*
Arteriography and angiography using contrast material	88.4–88.5	1,986	361	488	754
Diagnostic ultrasound	88.7	884	318	*	245
Physical therapy procedures	93.1–93.3	406	123	*	157
Respiratory therapy	93.9,96.7	1,160	306	237	352
Insertion of endotracheal tube	96.04	506	108	102	181
Transfusion of blood and blood components	99.0	1,902	327	403	754
Injection or infusion of cancer chemotherapeutic substance	99.25	209	51	28	91

* Figure does not meet standards of reliability or precision.

¹See Table 29 for comparison of number of procedures with number of discharges with procedures.

²Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 35. Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	Northeast	Midwest	South	West	
Rate per 10,000 population ¹						
All procedures	1,522.7	1,775.1	1,482.7	1,494.3	1,403.4	
Operations on the nervous system	01–05	41.6	46.4	49.1	35.9	39.3
Spinal tap03.31	11.6	11.7	13.0	11.5	10.3
Operations on the endocrine system	06–07	3.5	4.0	3.1	3.4	3.7
Operations on the eye	08–16	2.5	3.2	1.8	2.4	2.9
Operations on the ear	18–20	1.4	1.7	* .	1.3	*
Operations on the nose, mouth, and pharynx	21–29	10.2	12.1	6.4	10.9	11.4
Operations on the respiratory system	30–34	37.9	45.4	34.9	38.3	34.0
Bronchoscopy with or without biopsy	33.21–33.24, 33.27	9.6	11.5	9.3	8.8	9.7
Operations on the cardiovascular system	35–39	236.8	245.4	235.3	244.2	219.5
Balloon angioplasty of coronary artery or coronary atherectomy	36.01–36.02, 36.05	21.9	21.8	27.6	20.0	19.3
Insertion of coronary artery stent(s)	36.06–36.07	21.0	19.2	26.7	19.5	19.3
Coronary artery bypass graft ²	36.1	15.8	14.2	16.8	16.3	15.1
Cardiac catheterization	37.21–37.23	41.0	38.4	43.9	43.8	35.7
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7–37.8	13.0	19.7	11.2	10.7	13.0
Hemodialysis39.95	20.1	20.5	15.1	21.3	22.6
Operations on the hemic and lymphatic system	40–41	11.6	12.0	10.3	11.7	12.4
Operations on the digestive system	42–54	189.0	209.9	187.7	188.6	174.3
Endoscopy of small intestine with or without biopsy	45.11–45.14, 45.16	36.0	39.1	36.0	36.2	33.4
Endoscopy of large intestine with or without biopsy	45.21–45.25	18.8	22.3	19.0	18.3	16.8
Partial excision of large intestine	45.7	9.0	9.6	10.1	8.9	7.7
Appendectomy, excluding incidental	47.0	11.6	12.2	12.0	9.7	13.5
Cholecystectomy	51.2	13.5	13.6	14.5	14.4	10.9
Lysis of peritoneal adhesions	54.5	11.8	12.0	12.5	13.2	8.9
Operations on the urinary system	55–59	33.8	44.2	27.8	37.3	25.8
Cystoscopy with or without biopsy	57.31–57.33	4.8	7.1	4.0	5.4	2.9
Operations on the male genital organs	60–64	7.9	8.7	8.0	7.2	8.3
Prostatectomy	60.2–60.6	5.3	5.8	5.4	4.3	6.2
Operations on the female genital organs	65–71	66.6	50.6	61.9	83.2	58.1
Oophorectomy and salpingo-oophorectomy	65.3–65.6	14.7	10.6	15.8	17.8	12.0
Bilateral destruction or occlusion of fallopian tubes	66.2–66.3	11.7	6.6	9.2	16.1	11.3
Hysterectomy	68.3–68.7, 68.9	19.5	15.5	19.7	23.1	16.7
Obstetrical procedures	72–75	232.3	191.4	237.1	234.9	256.6
Episiotomy with or without forceps or vacuum extraction72.1, 72.21, 72.31, 72.71, 73.6	18.2	15.0	15.1	20.6	19.9
Artificial rupture of membranes	73.0	31.4	22.0	41.7	29.7	31.8
Medical induction of labor	73.4	20.9	16.1	21.9	21.1	23.3
Cesarean section	74.0–74.2, 74.4, 74.99	42.8	34.8	35.3	49.6	45.6
Repair of current obstetric laceration	75.5–75.6	42.6	35.4	43.2	40.7	51.0
Operations on the musculoskeletal system	76–84	140.2	143.2	147.4	137.8	134.6
Partial excision of bone	76.2–76.3, 77.6–77.8	6.0	6.6	5.2	6.2	6.0
Reduction of fracture	76.7, 79.0–79.3	21.0	22.8	22.1	21.1	18.5
Open reduction of fracture with internal fixation	79.3	14.4	15.5	14.9	14.7	12.5
Excision or destruction of intervertebral disc	80.5	9.9	9.8	9.4	10.2	10.2
Total hip replacement81.51	7.9	10.0	9.1	5.7	8.8
Total knee replacement81.54	18.1	19.9	23.3	15.7	15.2
Operations on the integumentary system	85–86	52.1	52.9	44.7	64.7	38.8
Debridement of wound, infection, or burn	86.22, 86.28	12.1	12.2	10.4	15.0	9.2
Miscellaneous diagnostic and therapeutic procedures and new technologies ³	87–99.00	455.2	704.0	425.7	392.6	382.2
Computerized axial tomography	87.03, 87.41, 87.71, 88.01, 88.38	27.5	70.8	* .	* .	20.8
Arteriography and angiography using contrast material	88.4–88.5	67.3	66.1	74.0	70.6	56.4
Diagnostic ultrasound	88.7	30.0	58.3	* .	23.0	25.1
Physical therapy procedures	93.1–93.3	13.8	22.6	* .	14.7	8.6
Respiratory therapy	93.9, 96.7	39.3	56.1	35.9	33.0	39.0
Insertion of endotracheal tube96.04	17.2	19.9	15.5	17.0	16.9
Transfusion of blood and blood components99.0	64.4	59.9	61.2	70.6	61.6
Injection or infusion of cancer chemotherapeutic substance99.25	7.1	9.4	4.2	8.5	5.7

* Figure does not meet standards of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²See Table 29 for comparison of number of procedures with number of discharges with procedures.

³Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 36. Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population ¹	Number in thousands	Percent distribution	Rate per 1,000 population ¹	
All deliveries							
10–54 years	4,038	100.0	43.1	10,425	100.0	111.3	2.6
10–19 years	409	10.1	20.0	1,041	10.0	51.0	2.5
20–24 years	1,058	26.2	104.5	2,618	25.1	258.8	2.5
25–29 years	1,104	27.3	113.2	2,761	26.5	283.0	2.5
30–34 years	899	22.3	90.8	2,422	23.2	244.7	2.7
35–39 years	459	11.4	44.0	1,264	12.1	121.3	2.8
40–54 years	110	2.7	3.3	319	3.1	9.6	2.9
All regions	4,038	100.0	43.1	10,425	100.0	111.3	2.6
Northeast	591	14.6	34.2	1,674	16.1	96.8	2.8
Midwest	824	20.4	39.5	2,066	19.8	99.0	2.5
South	1,574	39.0	46.4	4,107	39.4	121.1	2.6
West	1,049	26.0	48.5	2,578	24.7	119.3	2.5
Vaginal deliveries ²							
10–54 years	2,776	100.0	29.6	6,003	100.0	64.1	2.2
10–19 years	327	11.8	16.0	731	12.2	35.8	2.2
20–24 years	765	27.6	75.7	1,633	27.2	161.4	2.1
25–29 years	779	28.0	79.8	1,651	27.5	169.2	2.1
30–34 years	576	20.8	58.2	1,273	21.2	128.5	2.2
35–39 years	271	9.8	26.0	589	9.8	56.5	2.2
40–54 years	58	2.1	1.7	126	2.1	3.8	2.2
All regions	2,776	100.0	29.6	6,003	100.0	64.1	2.2
Northeast	401	14.5	23.2	935	15.6	54.1	2.3
Midwest	592	21.3	28.4	1,267	21.1	60.7	2.1
South	1,044	37.6	30.8	2,318	38.6	68.4	2.2
West	739	26.6	34.2	1,482	24.7	68.6	2.0
Cesarean deliveries ³							
10–54 years	1,262	100.0	13.5	4,422	100.0	47.2	3.5
10–19 years	81	6.4	4.0	309	7.0	15.2	3.8
20–24 years	292	23.1	28.9	985	22.3	97.3	3.4
25–29 years	326	25.8	33.4	1,110	25.1	113.8	3.4
30–34 years	323	25.6	32.7	1,150	26.0	116.1	3.6
35–39 years	188	14.9	18.0	675	15.3	64.8	3.6
40–54 years	52	4.1	1.6	193	4.4	5.8	3.7
All regions	1,262	100.0	13.5	4,422	100.0	47.2	3.5
Northeast	190	15.1	11.0	739	16.7	42.7	3.9
Midwest	232	18.4	11.1	799	18.1	38.3	3.4
South	530	42.0	15.6	1,788	40.4	52.8	3.4
West	310	24.5	14.3	1,096	24.8	50.7	3.5

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2005.

²Discharges of females with deliveries except those with cesarean deliveries.

³Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 37. Number and percent distribution of females with deliveries discharged from short-stay hospitals, by length of stay and type of delivery: United States, selected years, 1980–2005

[Discharges of inpatients from nonfederal hospitals. Data are for discharges with *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)* code V27, females with deliveries]

Length of stay	1980	1985	1990	1995	2000	2005
Number in thousands						
All deliveries	3,762	3,854	4,025	3,766	3,738	4,038
Less than 1 day	23	34	37	36	16	11
1 day	216	373	623	1,368	551	481
2 days	747	1,037	1,649	1,418	1,863	1,880
3 days	1,205	1,140	900	606	863	1,111
4 days	691	586	468	193	292	368
5–7 days	726	598	282	100	112	145
8 days or more	154	86	67	46	41	43
Vaginal deliveries ¹	3,143	2,977	3,080	2,981	2,882	2,776
Less than 1 day	23	33	35	34	14	*8
1 day	215	371	620	1,362	546	467
2 days	743	1,030	1,624	1,271	1,726	1,653
3 days	1,185	1,054	618	228	500	558
4 days	618	338	106	39	53	51
5–7 days	317	125	55	27	25	23
8 days or more	42	26	23	21	17	15
Cesarean deliveries ²	619	877	945	785	855	1,262
Less than 1 day	*	*	*	*	*	*
1 day	*	*	*	*6	*	*
2 days	*	*	25	147	137	227
3 days	20	86	283	379	363	553
4 days	73	249	361	154	239	317
5–7 days	409	473	227	73	87	122
8 days or more	112	60	44	25	24	28
Percent distribution						
All deliveries	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.6	0.9	0.9	1.0	0.4	0.3
1 day	5.7	9.7	15.5	36.3	14.7	11.9
2 days	19.9	26.9	41.0	37.7	49.9	46.6
3 days	32.0	29.6	22.4	16.1	23.1	27.5
4 days	18.4	15.2	11.6	5.1	7.8	9.1
5–7 days	19.3	15.5	7.0	2.7	3.0	3.6
8 days or more	4.1	2.2	1.7	1.2	1.1	1.1
Vaginal deliveries ¹	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.7	1.1	1.1	1.1	0.5	*0.3
1 day	6.8	12.5	20.1	45.7	19.0	16.8
2 days	23.6	34.6	52.7	42.6	59.9	59.5
3 days	37.7	35.4	20.1	7.6	17.3	20.1
4 days	19.7	11.4	3.4	1.3	1.8	1.9
5–7 days	10.1	4.2	1.8	0.9	0.9	0.8
8 days or more	1.3	0.9	0.7	0.7	0.6	0.5
Cesarean deliveries ²	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	*	*	*	*	*	*
1 day	*	*	*	*0.8	*	*
2 days	*	*	2.6	18.7	16.0	18.0
3 days	3.2	9.8	29.9	48.3	42.4	43.8
4 days	11.8	28.4	38.2	19.6	27.9	25.1
5–7 days	66.1	53.9	24.0	9.3	10.2	9.7
8 days or more	18.1	6.8	4.7	3.2	2.8	2.2
Type of delivery						
Average length of stay in days						
All deliveries	3.8	3.3	2.8	2.1	2.5	2.6
Vaginal deliveries ¹	3.2	2.7	2.3	1.7	2.1	2.2
Cesarean deliveries ²	6.5	5.2	4.5	3.6	3.7	3.5

* Figure does not meet standards of reliability or precision.

¹Discharges of females with deliveries except those with cesarean deliveries.

²Discharges of females with deliveries with ICD-9-CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 38. Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years, 1980–2005

[Discharges of inpatients from nonfederal hospitals. Procedure groupings and code number are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD–9–CM). Data are for selected procedures performed on discharges with code V27, females with deliveries]

Procedure category and ICD–9–CM code	1980	1985	1990	1995	2000	2005
Number of procedures in thousands						
All obstetrical procedures	72–75	3,526	4,239	6,660	6,282	6,145
Cesarean section	74.0–74.2,74.4,74.99	619	877	945	785	855
Repair of current obstetric laceration	75.5–75.6	350	543	789	961	1,134
Manually assisted delivery	73.5	70	193	750	866	898
Artificial rupture of membranes	73.0	120	315	691	752	833
Medical induction of labor	73.4	42	66	225	366	435
Episiotomy	72.1,72.21,72.31,72.71,73.6	2,012	1,820	1,714	1,410	944
Vacuum extraction ¹	72.7	22	66	190	279	245
Forceps delivery ¹	72.0–72.4,72.51,72.53,72.6,73.3	555	376	269	176	117
Rate per 100 deliveries ²						
All obstetrical procedures	72–75	93.7	110.0	165.5	166.8	164.4
Cesarean section	74.0–74.2,74.4,74.99	16.5	22.7	23.5	20.8	22.9
Artificial rupture of membranes	73.0	3.2	8.2	17.2	20.0	22.3
Medical induction of labor	73.4	1.1	1.7	5.6	9.7	11.7
Rate per 100 vaginal deliveries ³						
Repair of current obstetric laceration	75.5–75.6	11.1	18.1	25.5	32.0	39.2
Manually assisted delivery	73.5	2.2	6.5	24.3	29.0	31.1
Episiotomy	72.1,72.21,72.31,72.71,73.6	64.0	61.1	55.6	47.2	32.7
Vacuum extraction ¹	72.7	0.7	2.2	6.1	9.2	8.4
Forceps delivery ¹	72.0–72.4,72.51,72.53,72.6,73.3	17.6	12.5	8.6	5.8	4.0

¹Forceps delivery and vacuum extraction with episiotomy are also included in the episiotomy category.

²Procedures per 100 discharges of females with deliveries.

³Procedures for females with vaginal deliveries per 100 discharges with vaginal deliveries. Vaginal deliveries are all deliveries except those with a cesarean section.

Table 39. Number and rate of females with deliveries discharged from short-stay hospitals, by type of delivery: United States, selected years, 1980–2005

[Discharges of inpatients from nonfederal hospitals. Data are for discharges with *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Type of delivery	1980	1985	1990	1995	2000	2005
Number in thousands						
Females with deliveries	3,762	3,854	4,025	3,766	3,738	4,038
Previous cesarean delivery	192	324	426	413	447	571
No previous cesarean delivery	3,571	3,530	3,600	3,353	3,290	3,468
Vaginal deliveries	3,143	2,977	3,080	2,981	2,882	2,776
Vaginal birth after cesarean delivery (VBAC)	*6	21	87	147	123	62
Other vaginal deliveries	3,137	2,956	2,993	2,835	2,759	2,715
Cesarean deliveries	619	877	945	785	855	1,262
No previous cesarean delivery (primary)	434	574	606	519	531	753
Previous cesarean delivery (repeat)	185	303	339	266	324	509
Rate per 100 deliveries						
Cesarean deliveries per 100 deliveries	16.5	22.7	23.5	20.8	22.9	31.3
Primary cesarean deliveries per 100 deliveries without previous cesarean delivery	12.1	16.3	16.8	15.5	16.1	21.7
Vaginal birth after cesarean deliveries (VBAC) per 100 deliveries with previous cesarean delivery	*3.4	6.6	20.4	35.5	27.6	10.8

* Figure does not meet standards of reliability or precision.

NOTES: Previous cesarean deliveries are discharges with ICD–9–CM diagnostic code 654.2. Cesarean deliveries are discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4 or 74.99.

Table 40. Number and rate of females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Data are for discharges with *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)* code V27, females with deliveries]

Age and region	Cesarean deliveries		
	Total ¹	Primary ²	VBAC ^{3,4}
Number in thousands			
10–54 years	1,262	753	62
10–19 years	81	73	*
20–24 years	292	190	13
25–29 years	326	191	18
30–34 years	323	172	16
35–39 years	188	90	12
40–54 years	52	37	*
All regions	1,262	753	62
Northeast	190	117	10
Midwest	232	136	14
South	530	315	23
West	310	184	15
Rate per 100 deliveries			
10–54 years	31.3	21.7	10.8
10–19 years	19.9	18.3	*
20–24 years	27.6	20.2	10.9
25–29 years	29.5	20.1	11.5
30–34 years	36.0	23.5	9.7
35–39 years	40.9	25.8	10.7
40–54 years	47.4	39.9	*
All regions	31.3	21.7	10.8
Northeast	32.1	23.1	11.7
Midwest	28.2	19.1	13.1
South	33.7	23.6	9.7
West	29.5	20.2	10.3

* Figure does not meet standards of reliability or precision.

¹Discharges with ICD-9-CM procedure codes 74.0–74.2, 74.4, or 74.99. Rate per 100 total deliveries.

²Cesarean deliveries without ICD-9-CM diagnostic code 654.2. Rate per 100 deliveries without previous cesarean delivery.

³VBAC is vaginal birth after cesarean delivery.

⁴Vaginal deliveries with ICD-9-CM diagnostic code 654.2. Rate per 100 deliveries with previous cesarean delivery.

Table 41. Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals, by sex, geographic region, and health status: United States, 2005

[Infants born in nonfederal hospitals]

Sex, region, and health status	Discharges		Average length of stay in days
	Number in thousands	Percent distribution	
All newborn infants	3,999	100.0	3.4
Sex			
Male	2,044	51.1	3.4
Female	1,955	48.9	3.4
Region			
Northeast	739	18.5	3.4
Midwest	806	20.1	2.9
South	1,429	35.7	3.9
West	1,026	25.6	3.1
Health status			
Well ¹	2,201	55.0	2.1
Male	1,078	26.9	2.1
Female	1,123	28.1	2.1
Sick ²	1,798	45.0	5.0
Male	967	24.2	4.9
Female	831	20.8	5.0

¹Without any illness or risk-related diagnoses.²With at least one illness or risk-related diagnosis, *International Classification of Diseases, Ninth Revision, Clinical Modification* codes 001–999.**Table 42. Number and percent distribution of newborn infants discharged from short-stay hospitals, by length of stay, and average length of stay, by health status: United States, selected years, 1980–2005**

[Infants born in nonfederal hospitals]

Length of stay and health status	1980	1985	1990	1995	2000	2005
Length of stay						
All newborn infants	3,824	3,794	3,869	3,631	3,643	3,999
Less than 1 day	69	84	79	87	45	44
1 day	270	428	693	1,523	722	674
2 days	842	1,057	1,624	1,145	1,877	1,967
3 days	1,175	1,034	636	486	546	720
4 days	618	528	447	134	205	286
5–7 days	633	497	232	106	94	122
8 days or more	218	167	157	149	153	185
Number in thousands						
All newborn infants	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	1.8	2.2	2.0	2.4	1.2	1.1
1 day	7.1	11.3	17.9	41.9	19.8	16.9
2 days	22.0	27.9	42.0	31.5	51.5	49.2
3 days	30.7	27.3	16.4	13.4	15.0	18.0
4 days	16.2	13.9	11.6	3.7	5.6	7.2
5–7 days	16.5	13.1	6.0	2.9	2.6	3.1
8 days or more	5.7	4.4	4.1	4.1	4.2	4.6
Percent distribution						
All newborn infants	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	1.8	2.2	2.0	2.4	1.2	1.1
1 day	7.1	11.3	17.9	41.9	19.8	16.9
2 days	22.0	27.9	42.0	31.5	51.5	49.2
3 days	30.7	27.3	16.4	13.4	15.0	18.0
4 days	16.2	13.9	11.6	3.7	5.6	7.2
5–7 days	16.5	13.1	6.0	2.9	2.6	3.1
8 days or more	5.7	4.4	4.1	4.1	4.2	4.6
Health status						
All newborn infants	4.3	3.8	3.3	2.8	3.2	3.4
Well ¹	3.2	2.8	2.3	1.7	2.1	2.1
Sick ²	7.1	5.7	4.8	4.4	5.1	5.0
Average length of stay in days						

¹Without any illness or risk-related diagnoses.²With at least one illness or risk-related diagnosis, *International Classification of Diseases, Ninth Revision, Clinical Modification* codes 001–999.

Table 43. Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2005

[Infants born in nonfederal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*]

Diagnostic category and ICD-9-CM code		Both sexes	Male	Female
Number in thousands				
Sick newborn diagnoses	001–999	3,749	2,051	1,699
Hemolytic disease of fetus or newborn, due to isoimmunization, and other perinatal jaundice	773–774	798	419	379
Hemolytic disease due to ABO isoimmunization	773.1	53	20	32
Neonatal jaundice associated with preterm delivery	774.2	157	85	72
Unspecified fetal and neonatal jaundice	774.6	568	303	264
Respiratory conditions of fetus and newborn	768–770	437	246	191
Respiratory distress syndrome769	87	49	38
Transitory tachypnea of newborn	770.6	134	78	55
Disorders relating to short gestation and low birth weight (prematurity)	765.0–765.1	315	170	145
Congenital anomalies	740–759	311	189	122
Congenital anomalies of circulatory system	745–747	86	46	40
Congenital anomalies of musculoskeletal system	754–756	49	27	22
Congenital anomalies of the integument (skin)757	63	33	30
Disorders relating to long gestation and high birthweight766	258	154	103
Maternal causes of perinatal morbidity and mortality	760–763	151	77	75
Endocrine and metabolic disturbances specific to the fetus and newborn775	131	70	61
Neonatal hypoglycemia	775.6	56	31	25
Conditions involving the integument (skin) and temperature regulation of fetus and newborn778	106	70	36
Infections specific to the perinatal period771	102	52	50
Septicemia (sepsis) of newborn771.81	81	42	39
Birth trauma767	101	59	42
Injuries to scalp	767.1	71	40	31
Feeding problems in newborn	779.3	97	53	44
Slow fetal growth and fetal malnutrition764	64	25	39
Fetal and neonatal hemorrhage772	54	26	28

Table 44. Number and percentage of male newborn infants circumcised during hospitalization, by geographic region: United States, selected years, 1980–2005

[Infants born in nonfederal hospitals with *International Classification of Diseases, Ninth Revision, Clinical Modification* procedure code 64.0]

Region	1980	1985	1990	1995	2000	2005
Number circumcised in thousands						
United States	1,261	1,162	1,169	1,200	1,166	1,172
Northeast	253	221	238	249	256	254
Midwest	403	331	341	317	330	330
South	379	393	396	454	438	422
West	226	218	195	180	143	166
Percentage circumcised						
United States	64.7	59.5	59.0	64.1	62.4	57.3
Northeast	67.4	65.2	62.6	68.3	64.6	66.9
Midwest	75.9	70.5	76.0	79.8	81.4	78.7
South	56.0	56.0	57.1	66.1	63.9	58.7
West	61.8	49.0	42.4	42.6	37.3	31.5

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
All codes	34,667	13,902	20,766	2,431	10,659	8,349	13,228	7,192	7,948	12,907	6,621
001–139	1,045	498	547	142	176	244	483	235	225	382	202
003	*8	*	*	*	*	*	*	*	*	*	*
003.0	*6	*	*	*	*	*	*	*	*	*	*
008	173	66	107	44	23	33	74	49	36	64	24
008.4	78	28	50	*	*7	20	49	28	16	27	*8
008.45	76	27	49	*	*6	19	48	27	16	26	*7
008.6	24	13	11	23	*	*	*	*	*	*5	10
008.61	23	13	10	22	*	*	*	*	*	*5	9
008.8	69	24	45	18	15	12	24	18	15	25	11
009	16	*6	9	*	*7	*	*	*	*	*	*5
009.0	10	*	*7	*	*	*	*	*	*	*	*
011	10	*8	*	*	*	*5	*	*	*	*7	*
011.9	*7	*	*	*	*	*	*	*	*	*	*
033	*7	*	*	*7	—	*	—	*	*	*	*
034	*8	*	*	*	*	*	—	*	*	*	*
034.0	*8	*	*	*	*	*	—	*	*	*	*
038	490	240	249	*6	38	112	334	97	115	178	99
038.0	27	15	12	*	*	*6	17	*5	*7	*9	*6
038.1	67	41	26	*	*7	17	42	12	13	29	13
038.10	*5	*	*	*	*	*	*	*	*	*	*
038.11	49	31	18	*	*5	13	31	*9	*9	22	10
038.19	13	*7	*6	*	*	*	*8	*	*	*	*
038.2	*7	*	*	*	*	*	*	*	*	*	*
038.4	95	38	57	*	*7	25	62	23	30	21	21
038.40	*9	*	*	—	*	*	*6	*	*	*	*
038.42	51	17	34	*	*	12	34	13	18	10	9
038.43	*7	*	*	—	*	*	*	*	*	*	*
038.49	27	13	14	*	*	*8	17	*7	*7	*5	*7
038.8	12	*	9	*	*	*	10	*	*	*	*
038.9	278	137	141	*	18	59	198	51	59	112	56
041	12	*6	*6	*	*	*	*	*	*	*5	*
042	58	35	23	*	32	24	*	20	*6	27	*5
047	41	21	20	19	18	*	*	*6	11	13	10
047.9	36	19	17	15	17	*	*	*6	10	13	*8
053	21	*8	13	*	*	*	16	*	*6	*	*6
053.1	*5	*	*	—	*	*	*	*	*	*	*
053.9	10	*	*8	*	*	*	*8	*	*	*	*
054	*6	*	*	*	*	*	*	*	*	*	*
066	*5	*	*	—	*	*	*	—	*	*	*
070	25	16	*9	*	*7	14	*	*	*	10	*7
070.5	*8	*	*	—	*	*5	*	*	*	*	*
070.54	*6	*	*	—	*	*	*	*	*	*	*
070.7	*6	*	*	—	*	*	*	*	*	*	*
075	10	*5	*	*	*7	*	—	*	*	*	*
079	60	30	29	35	*9	*8	*8	14	13	20	13
079.9	54	27	27	30	*8	*8	*8	13	13	18	10
079.99	54	27	27	30	*8	*8	*8	13	13	18	10
112	24	*8	16	*	*	*	13	*	*	12	*
112.0	*7	*	*5	*	*	*	*	*	*	*	*
112.2	*8	*	*6	—	*	*	*7	*	*	*	*
117	*6	*	*	*	*	*	*	*	*	*	*
135	11	*	*6	*	*	*5	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
140–239	1,615	654	961	27	270	626	692	348	345	600	323
141	*6	*	*	—	*	*	*	*	*	*	*
150	13	10	*	—	*	*6	*7	*	*	*	*
151	28	17	11	—	*	10	17	*	*9	10	*
151.0	11	*8	*	—	*	*	*6	*	*	*	*
151.9	*7	*	*	—	*	*	*5	*	*	*	*
153	110	51	58	*	*	32	74	22	27	40	21
153.0	*7	*	*	—	*	*	*6	*	*	*	*
153.1	11	*7	*	—	*	*	*7	*	*	*	*
153.2	*6	*	*5	—	*	*	*	*	*	*	*
153.3	21	10	11	—	*	*7	13	*	*5	*7	*5
153.4	24	10	14	—	*	*7	16	*5	*6	*8	*
153.6	20	9	11	—	*	*5	14	*	*5	*8	*
153.9	13	*7	*6	—	*	*	*8	*	*	*7	*
154	37	20	17	—	*	14	21	*7	*9	13	*7
154.0	16	9	*6	—	*	*6	*9	*	*	*7	*
154.1	19	10	*9	—	*	*7	11	*	*5	*6	*
155	17	10	*7	*	*	*7	*9	*5	*	*7	*
155.0	12	*8	*	*	*	*6	*5	*	*	*5	*
156	*9	*5	*	—	*	*	*6	*	*	*	*
157	28	14	14	*	*	10	17	*	*8	10	*6
157.0	10	*5	*	—	*	*	*7	*	*	*	*
157.9	9	*	*	—	*	*	*	*	*	*	*
161	11	*8	*	—	*	*7	*	*	*	*	*
162	137	71	66	*	*	46	86	32	34	53	17
162.2	*9	*	*	—	*	*	*5	*	*	*	*
162.3	47	25	23	*	*	17	30	11	11	17	*8
162.5	21	11	10	*	*	*8	13	*	*	11	*
162.8	17	*9	*9	—	*	*	12	*7	*	*	*
162.9	39	21	19	—	*	13	25	*7	14	15	*
171	*8	*5	*	*	*	*	*	*	*	*	*
173	*6	*	*	*	*	*	*	*	*	*	*
174	85	—	85	—	15	39	31	15	20	36	14
174.4	10	—	10	—	*	*	*	*	*	*	*
174.8	24	—	24	—	*	13	*7	*6	*	11	*
174.9	40	—	40	—	*6	18	15	*	11	20	*
180	14	—	14	—	*6	*7	*	*	*	*6	*
180.9	*8	—	*8	—	*	*	*	*	*	*5	*
182	34	—	34	—	*	17	14	*8	10	10	*6
182.0	34	—	34	—	*	17	13	*8	10	10	*6
183	20	—	20	*	*	11	*6	*	*	*8	*
183.0	19	—	19	*	*	10	*6	*	*	*8	*
185	78	78	—	*	*	39	38	14	15	27	22
188	39	29	10	—	*	*6	33	11	9	12	*7
188.8	14	10	*	—	*	*	11	*6	*	*	*
188.9	9	*6	*	—	*	*	*8	*	*	*	*
189	45	28	17	*	*	14	26	10	11	14	9
189.0	41	25	16	*	*	13	23	*9	11	14	*8
191	41	26	15	*	*7	14	15	*9	*8	12	13
191.1	12	*9	*	*	*	*5	*	*	*	*	*
191.2	*8	*6	*	*	*	*	*	*	*	*	*
191.8	*5	*	*	*	*	*	*	*	*	*	*
191.9	*6	*	*	*	*	*	*	*	*	*	*
193	24	*5	19	*	10	11	*	*6	*	9	*6

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
196	14	9	*	*	*	*	*9	*	*	*7	*
196.2	*5	*	*	—	*	*	*	*	*	*	*
197	98	42	56	*	*8	39	50	26	20	34	18
197.0	17	*8	9	*	*	*8	*7	*	*	*6	*
197.2	25	*8	16	—	*	9	14	*7	*7	*7	*
197.6	17	*6	11	—	*	*6	*9	*6	*	*5	*
197.7	29	17	13	*	*	12	14	*7	*	11	*6
198	86	37	49	*	*6	40	40	20	24	28	14
198.3	33	14	19	*	*	16	15	*8	*7	11	*6
198.5	36	17	19	*	*	17	17	*7	14	11	*
198.8	10	*	*6	*	*	*	*6	*	*	*	*
198.89	*9	*	*5	*	*	*	*5	*	*	*	*
199	23	9	14	*	*	9	12	*	*	13	*5
199.0	15	*5	10	*	*	*6	*7	*	*	*8	*
199.1	*8	*	*	*	*	*	*5	*	*	*	*
200	*8	*	*	*	*	*	*	*	*	*	*
200.0	*6	*	*	—	*	*	*	*	*	*	*
201	*6	*	*	*	*	*	*	*	*	*	*
202	32	19	13	*	*	9	18	*7	10	9	*6
202.8	28	16	11	*	*	*8	17	*6	*8	*8	*6
202.80	13	*8	*	—	*	*	*9	*	*	*	*
202.83	*5	*	*	—	*	*	*	*	*	*	*
203	15	10	*5	—	*	*	10	*	*	*	*
203.0	14	9	*5	—	*	*	9	*	*	*	*
203.00	14	9	*	—	*	*	9	*	*	*	*
204	19	10	*8	*7	*	*	*6	*	*	*6	*7
204.0	11	*7	*	*7	*	*	*	*	*	*	*
204.00	*9	*6	*	*	*	*	*	*	*	*	*
204.1	*7	*	*	—	—	*	*5	*	*	*	*
204.10	*6	*	*	—	—	*	*5	*	*	*	*
205	18	11	*8	*	*	*7	*7	*5	*	*7	*
205.0	15	*9	*7	*	*	*6	*6	*	*	*6	*
205.00	15	*8	*7	*	*	*6	*6	*	*	*6	*
210	*5	*	*	*	*	*	*	*	*	*	*
211	48	21	27	*	*	17	26	11	9	18	*9
211.3	34	17	18	—	*	12	19	*8	*6	13	*7
218	201	—	201	—	103	95	*	38	40	89	34
218.0	29	—	29	—	11	18	*	*7	*	13	*
218.1	58	—	58	—	31	26	*	15	*8	26	*9
218.2	16	—	16	—	*9	*7	—	*	*	*7	*
218.9	98	—	98	—	52	45	*	14	22	42	20
220	34	—	34	*	16	13	*5	*6	11	10	*7
225	22	*6	16	*	*	*9	*9	*6	*	*7	*6
225.2	14	*	11	—	*	*	*8	*	*	*	*
226	10	*	*9	—	*	*6	*	*	*	*	*
227	16	*5	11	*	*	*8	*	*	*	*6	*
227.3	*8	*	*	*	*	*	*	*	*	*	*
228	*7	*	*	*	*	*	*	*	*	*	*
228.0	*7	*	*	*	*	*	*	*	*	*	*
230	*6	*	*	—	*	*	*	*	*	*	*
233	23	*	23	—	12	*9	*	*	*	13	*
233.0	*8	—	*8	—	*	*	*	*	*	*	*
233.1	13	—	13	—	10	*	*	*	*	*8	*
235	*7	*	*5	—	*	*	*	*	*	*	*
235.2	*5	*	*	—	*	*	*	*	*	*	*
237	*6	*	*	*	*	*	*	*	*	*	*

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
238	18	9	*8	*	*	*	11	*	*	*8	*
238.7	12	*7	*6	*	*	*	10	*	*	*6	*
239	16	*8	*8	*	*	*5	*8	*	*6	*6	*
239.6	*6	*	*	*	*	*	*	*	*	*	*
240–279	1,708	713	995	203	359	494	652	383	381	645	299
241	17	*	15	—	*	10	*	*	*	*6	*
241.0	*6	*	*5	—	*	*	*	*	*	*	*
241.1	9	*	*8	—	*	*5	*	*	*	*	*
242	11	*	*8	*	*	*	*	*	*	*	*
242.9	*5	*	*	*	*	*	*	*	*	*	*
244	19	*	17	*	*	*	*6	*8	*7	*	*5
244.9	18	*	16	*	*	*6	*8	*7	*	*	*
250	585	283	302	27	157	209	191	148	119	215	102
250.0	163	73	90	10	39	63	51	43	33	58	29
250.00	79	37	42	—	17	34	27	18	15	28	17
250.01	20	*8	12	*7	*6	*	*	*	*5	*	*5
250.02	57	25	32	*	15	24	19	17	12	21	*6
250.03	*8	*	*	*	*	*	*	*	*	*	*
250.1	121	59	63	16	70	26	9	22	33	44	22
250.10	24	10	14	—	11	*7	*	*	*6	*8	*
250.11	57	28	29	13	34	*8	*	9	17	19	12
250.12	16	*8	*8	—	*8	*6	*	*	*	*8	*
250.13	24	13	11	—	18	*	*	*5	*7	9	*
250.2	14	*8	*6	*	*	*	*	*	*	*6	*
250.20	*6	*	*	—	*	*	*	*	*	*	*
250.22	*8	*	*	*	*	*	*	*	*	*	*
250.4	25	15	10	—	*	12	*9	*	*	10	*7
250.40	18	11	*7	—	*	*9	*7	*	*	*7	*6
250.6	68	28	39	—	19	32	17	14	13	27	13
250.60	40	17	23	—	*7	20	13	9	*7	15	*8
250.61	10	*	*7	—	*7	*	*	*	*	*	*
250.62	14	*7	*7	—	*	*9	*	*	*	*6	*
250.7	45	26	19	—	*	19	24	14	*7	16	*7
250.70	35	20	16	—	*	14	22	11	*5	14	*6
250.72	*7	*	*	—	*	*	*	*	*	*	*
250.8	135	68	67	*	15	48	71	43	26	49	18
250.80	96	45	51	*	*5	31	59	32	17	32	14
250.81	11	*7	*	*	*	*	*	*	*	*	*
250.82	24	13	11	—	*	11	9	*	*	13	*
250.9	*7	*	*	*	*	*	*	*	*	*	*
250.92	*6	*	*	*	*	*	*	*	*	*	*
251	16	*7	*8	*	*	*	*8	*	*	*6	*
251.2	11	*6	*5	*	*	*	*	*	*	*	*
252	*6	*	*	*	*	*	*	*	*	*	*
252.0	*5	*	*	*	*	*	*	*	*	*	*
253	20	*8	12	*	*	*	13	*5	*	*8	*
253.6	18	*8	10	*	*	*	13	*	*	*8	*
255	12	*5	*6	*	*	*	*6	*	*	*	*
255.4	9	*	*6	*	*	*	*5	*	*	*	*
263	13	*	*8	*	*	*	*8	*	*	*	*
263.9	10	*	*6	*	*	*	*6	*	*	*	*
272	48	24	24	*	*9	25	14	15	*9	18	*6
272.0	19	9	10	—	*	12	*	*7	*	*7	*
272.4	27	14	13	—	*	13	10	*7	*5	10	*
274	17	11	*6	—	*	*8	*7	*	*	*	*
274.0	11	*7	*	—	*	*5	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
275	20	*7	13	*	*	*	10	*	*	*6	*6
275.4	16	*5	11	*	*	*	*9	*	*	*	*5
275.42	11	*	*8	*	*	*	*7	*	*	*	*
276	740	297	444	156	75	144	365	136	182	302	120
276.0	15	*6	9	*	*	*	10	*	*	*	*
276.1	117	44	73	*	*6	26	82	22	33	39	24
276.2	10	*	*5	*	*	*	*	*	*	*	*
276.5	500	206	294	148	49	76	227	91	121	217	72
276.6	*8	*	*	—	*	*	*	*	*	*	*
276.7	35	15	19	*	*	14	18	*8	*7	11	*9
276.8	50	15	35	*	12	18	18	*9	11	22	*7
277	28	16	12	*9	17	*	*	*7	*8	10	*
277.0	15	*8	*7	*6	*9	*	—	*6	*5	*	*
277.02	12	*7	*5	*	*8	*	—	*5	*	*	*
277.1	*6	*6	*	—	*6	*	—	*	*	*5	*
278	129	31	98	*	66	59	*	36	22	39	32
278.0	120	30	90	*	63	54	*	32	21	37	30
278.00	11	*	*8	*	*7	*	*	*	*	*	*
278.01	109	26	83	*	56	50	*	30	20	34	25
278.1	*7	*	*7	—	*	*	*	*	*	*	*
280–289	455	174	281	57	115	100	183	106	98	177	74
280	100	28	71	*	22	22	54	23	23	37	16
280.0	49	14	34	*	13	10	25	12	13	18	*6
280.9	49	13	36	*	*9	11	28	11	10	18	10
282	83	36	46	19	53	*8	*	25	16	37	*
282.4	*9	*5	*	*	*	*	*	*	*	*	*
282.42	*6	*	*	*	*	*	—	*	*	*	*
282.6	72	30	42	17	48	*7	*	20	15	34	*
282.62	59	25	34	12	41	*5	*	16	11	30	*
283	*7	*	*	*	*	*	*	*	*	*	*
284	30	14	16	*	*7	10	10	*5	*7	12	*5
284.8	28	13	14	*	*6	9	9	*5	*7	11	*
285	102	37	64	*	10	26	64	25	20	39	18
285.1	13	*	*9	*	*	*	9	*	*	*	*
285.2	19	*8	11	*	*	*	14	*6	*	*7	*
285.29	10	*	*6	*	*	*	*8	*	*	*	*
285.9	67	24	43	*	*7	18	40	14	11	28	14
286	19	*7	12	*	*	*6	10	*	*5	*8	*
286.5	*8	*	*5	—	*	*	*	*	*	*	*
287	35	16	19	*8	*8	*7	11	9	*7	10	*9
287.3	15	*6	9	*	*	*	*	*	*	*	*
287.5	14	*7	*7	*	*	*	*	*	*	*	*
288	56	23	33	13	*7	16	20	10	13	21	12
288.0	44	19	25	11	*6	14	14	9	*8	17	10
288.8	11	*	*7	*	*	*	*5	*	*	*	*
289	19	*8	12	*7	*	*	*	*	*	*9	*
289.3	*7	*	*	*	*	*	*	*	*	*	*
290–319	2,398	1,200	1,198	124	1,349	680	246	665	518	1,002	213
290	27	*9	18	—	*	*	26	*9	*8	*9	*
290.2	*5	*	*	—	—	*	*	*	*	*	*
290.4	12	*5	*7	—	*	*	11	*	*	*	*
290.40	*5	*	*	—	—	—	*5	*	*	*	*
291	158	111	48	*	78	70	10	41	63	30	24
291.0	22	17	*	—	*8	12	*	*	*7	*5	*5
291.8	133	91	42	—	68	57	*8	36	55	23	19
291.81	128	88	40	—	65	55	*8	35	54	21	17

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
292	109	66	43	*	69	26	14	43	27	21	18
292.0	66	42	24	*	48	17	*	30	21	*8	*7
292.1	*7	*	*	*	*	*	*	*	*	*	*
292.8	33	18	15	*	16	*6	11	11	*	11	*8
292.81	13	*6	*7	—	*	*	10	*	*	*	*
292.84	19	12	*7	*	14	*	*	*7	*	*5	*5
293	16	*7	9	*	*	*	*8	*	*	*	*
293.0	10	*	*6	*	*	*	*7	*	*	*	*
293.8	*6	*	*	*	*	*	*	*	*	*	*
294	40	19	21	*	*	*	36	18	*9	11	*
294.8	37	17	20	—	*	*	34	17	*8	*9	*
295	351	193	158	*	196	130	23	100	60	165	26
295.3	123	74	49	*	68	47	*7	33	18	65	*7
295.30	60	34	25	*	33	23	*	9	*9	40	*
295.32	22	14	*9	*	12	*9	*	*6	*	12	*
295.34	40	26	14	—	23	14	*	18	*7	13	*
295.6	10	*7	*	—	*	*5	*	*	*	*	*
295.7	171	85	86	*	97	62	11	47	31	81	12
295.70	133	67	66	*	79	48	*7	30	25	69	10
295.72	*9	*	*	*	*6	*	*	*	*	*	*
295.74	27	12	15	*	12	12	*	16	*	*7	*
295.9	40	22	18	*	23	13	*	15	*6	14	*
295.90	29	17	13	*	16	9	*	*9	*5	13	*
295.94	*6	*	*	—	*	*	—	*	*	*	*
296	945	374	570	67	568	237	72	219	209	458	58
296.0	9	*	*	*	*6	*	*	*	*	*7	—
296.04	*5	*	*	*	*	*	*	*	*	*	—
296.2	200	74	126	14	114	52	19	33	42	117	*8
296.20	114	40	74	10	70	27	*7	16	23	69	*6
296.22	14	*6	*9	*	9	*	*	*	*	9	*
296.23	41	17	24	*	24	10	*	9	11	19	*
296.24	29	11	18	*	11	10	*7	*6	*	18	*
296.3	270	105	165	*5	149	84	31	71	72	105	22
296.30	68	31	36	*	40	22	*	23	19	24	*
296.32	19	*6	13	*	12	*	*	*	*5	*7	*
296.33	135	48	88	*	73	44	15	33	37	56	10
296.34	46	19	27	*	23	14	*8	13	11	17	*5
296.4	66	25	40	*	38	21	*6	20	13	29	*
296.40	27	9	18	*	16	*8	*	*	*6	15	*
296.43	*7	*	*	—	*	*	*	*	*	*	*
296.44	29	12	17	*	17	10	*	12	*	11	*
296.5	102	37	65	*	67	27	*5	27	27	44	*5
296.50	63	24	39	*	43	15	*	14	17	30	*
296.53	21	*6	15	*	12	*8	*	*8	*5	*6	*
296.54	16	*7	*9	*	11	*	*	*	*	*7	*
296.6	76	33	43	10	48	16	*	14	10	48	*
296.60	39	17	22	*	26	*8	*	*6	*5	26	*
296.63	14	*6	*8	*	*8	*	*	*	*	10	*
296.64	22	10	12	*	12	*	*	*6	*	12	*
296.7	18	*7	11	*	12	*	*	*5	*	*8	*
296.8	154	62	92	20	100	27	*6	23	34	87	10
296.80	122	52	70	19	80	19	*	14	26	75	*8
296.89	31	10	22	*	20	*8	*	*9	*8	12	*
296.9	49	26	23	10	33	*	*	24	*8	13	*
296.90	48	26	23	10	33	*	*	24	*8	13	*
297	*7	*	*	—	*	*	*	*	*	*	*
298	90	48	42	*	51	20	16	30	12	35	14
298.9	86	46	41	*	50	18	16	29	11	34	13

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
300	79	28	50	*	42	24	11	20	18	30	11
300.0	40	13	26	*	19	13	*7	10	*9	15	*6
300.00	28	9	18	*	12	10	*5	*7	*5	12	*
300.01	*8	*	*6	*	*5	*	*	*	*	*	*
300.1	11	*	*7	*	*6	*	*	*	*	*	*
300.11	*6	*	*	*	*	*	*	*	*	*	*
300.4	17	*	12	*	9	*6	*	*6	*	*6	*
300.9	*7	*	*	*	*	*	*	*	*	*	*
303	123	89	35	*	58	59	*6	28	24	62	10
303.0	32	24	*9	—	12	18	*	*8	12	*8	*
303.00	10	*7	*	—	*	*6	*	*	*5	*	*
303.01	20	16	*	—	*8	11	*	*6	*6	*	*
303.9	91	65	26	*	46	40	*	20	12	54	*6
303.90	48	32	16	*	23	23	*	*	*5	37	*
303.91	42	32	10	—	22	18	*	16	*7	17	*
304	81	49	32	*	60	19	*	20	*8	49	*
304.0	31	17	15	—	22	*8	*	10	*	17	*
304.00	15	*7	*7	—	10	*	*	*	*	12	*
304.01	16	*8	*7	—	11	*	*	10	*	*	*
304.2	18	13	*5	—	14	*	*	*	*	12	*
304.20	9	*6	*	—	*7	*	*	*	*	*7	*
304.21	9	*7	*	—	*7	*	—	*	*	*	—
304.7	*6	*	*	—	*5	*	*	*	*	*	*
304.8	11	*7	*	*	*8	*	—	*	*	10	—
304.80	10	*7	*	*	*7	*	—	*	*	10	—
305	95	56	39	*5	57	30	*	15	22	43	15
305.0	37	24	14	*	21	14	*	*7	12	13	*5
305.00	31	19	12	*	18	11	*	*5	11	10	*
305.01	*6	*	*	—	*	*	*	*	*	*	*
305.1	20	12	*8	—	11	*8	*	*	*6	*6	*
305.6	15	*8	*7	—	10	*	*	*	*	*8	*
305.60	12	*7	*5	—	*8	*	—	*	*	*7	*
305.9	16	*8	*7	*	9	*	*	*	*	11	*
305.90	14	*8	*7	*	*8	*	*	*	*	11	*
307	10	*	*9	*	*	*	*	*	*	*	*
309	64	31	33	*	46	11	*	28	13	17	*7
309.0	17	*9	*8	*	11	*	*	*8	*	*	*
309.2	9	*6	*	*	*7	*	*	*	*	*	*
309.28	*8	*5	*	*	*6	*	*	*	*	*	*
309.4	10	*	*6	*	*7	*	*	*	*	*	*
309.8	20	*6	13	*	13	*	*	*6	*5	*7	*
309.81	19	*6	13	*	12	*	*	*5	*5	*7	*
309.9	*7	*	*	*	*7	*	*	*	*	*	—
311	132	71	61	*9	81	34	*7	61	24	36	10
312	24	17	*7	*6	16	*	*	12	*	*8	*
312.3	20	15	*	*	14	*	—	11	*	*5	*
312.30	11	10	*	*	*8	*	—	*8	*	*	*
312.34	*8	*6	*	*	*6	*	—	*	*	*	*
313	*7	*	*	*	*	—	—	*	*	*	—
313.8	*7	*	*	*	*	—	—	*	*	*	—
313.81	*6	*	*	*	*	—	—	*	*	*	—
314	14	10	*	9	*	*	—	*	*	*6	*
314.0	12	*9	*	*8	*	*	—	*	*	*	*
314.01	12	*9	*	*8	*	*	—	*	*	*	*
320–389	565	238	327	84	138	125	219	135	124	195	110
320	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
322	*7	*	*	*	*	*	*	*	*	*	*
322.9	*6	*	*	*	*	*	*	*	*	*	*
324	*8	*	*	*	*	*	*	*	*	*	*
331	108	46	62	*	*	*8	92	25	29	41	14
331.0	68	23	44	—	*	*	66	14	19	29	*6
331.3	16	10	*6	*	*	*	14	*	*	*5	*
331.4	15	*7	*7	*	*	*	*	*	*	*5	*
331.8	*8	*	*	—	—	*	*7	*	*	*	*
331.82	*8	*	*	—	—	*	*7	*	*	*	*
332	17	*7	10	—	*	*	13	*	*	*	*
332.0	17	*7	9	—	*	*	13	*	*	*	*
333	12	*6	*6	*	*	*	*6	*	*	*	*
337	*5	*	*	*	*	*	*	*	*	*	*
340	26	*5	20	*	12	9	*	*8	*	*9	*5
344	*6	*	*	*	*	*	*	*	*	*	*
345	62	31	32	24	18	15	*6	16	10	18	18
345.1	14	*6	*8	*6	*	*	*	*	*	*	*7
345.10	11	*5	*6	*	*	*	*	*	*	*	*5
345.3	17	9	*7	*8	*	*	*	*	*	*	*
345.5	*8	*	*	*	*	*	*	*	*	*	*
345.9	12	*6	*6	*	*	*	*	*	*	*	*
345.90	*9	*	*	*	*	*	*	*	*	*	*
346	59	13	45	*	34	18	*	13	12	22	12
346.8	11	*	9	*	*5	*	*	*	*	*	*
346.80	*7	*	*6	*	*	*	*	*	*	*	*
346.9	37	*8	29	*	23	11	*	*8	*9	12	*7
346.90	29	*6	23	*	17	*9	*	*7	*7	10	*6
346.91	*8	*	*6	*	*6	*	*	*	*	*	*
348	47	23	24	*6	13	10	18	10	10	16	11
348.0	*5	*	*	*	*	*	*	*	*	*	*
348.3	20	9	10	*	*	*	*	12	*	*	*7
348.30	*8	*	*	*	*	*	*	*	*	*	*
348.31	*8	*	*	—	*	*	*6	*	*	*	*
348.8	*5	*	*	*	*	*	*	*	*	*	*
349	14	*	9	*	*5	*	*	*	*	*6	*
349.0	*5	*	*	*	*	*	*	*	*	*	*
349.8	*7	*	*	*	*	*	*	*	*	*	*
349.82	*5	*	*	*	*	*	*	*	*	*	*
351	*7	*	*	*	*	*	*	*	*	*	*
351.0	*5	*	*	*	*	*	*	*	*	*	*
355	*5	*	*	*	*	*	*	*	*	*	*
357	*9	*	*	*	*	*	*	*	*	*	*
357.0	*6	*	*	*	*	*	*	*	*	*	*
358	*6	*	*	*	*	*	*	*	*	*	*
358.0	*6	*	*	*	*	*	*	*	*	*	*
368	*5	*	*	*	*	*	*	*	*	*	*
376	11	*	*6	*6	*	*	*	*	*	*5	*
376.0	10	*	*5	*6	*	*	*	*	*	*	*
376.01	9	*	*	*6	*	*	*	*	*	*	*
380	*8	*	*	*	*	*	*	*	*	*	*
380.1	*6	*	*	*	*	*	*	*	*	*	*
380.10	*6	*	*	*	*	*	*	*	*	*	*
382	13	*8	*	9	*	*	*	*	*	*5	*
382.9	10	*6	*	*7	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
386	42	12	29	*	*	11	26	12	14	9	*6
386.1	21	*7	13	*	*	*6	13	*6	*	*6	*
386.11	15	*6	*9	*	*	*	10	*	*	*	*
386.3	15	*	11	—	*	*	*8	*	*6	*	*
386.30	14	*	10	—	*	*	*8	*	*6	*	*
390–459	6,159	3,136	3,023	29	436	1,813	3,882	1,392	1,465	2,226	1,076
396	17	*6	11	*	*	*	14	*	*	*6	*
396.2	*6	*	*	—	*	*	*5	*	*	*	*
396.3	*6	*	*	*	*	*	*	*	*	*	*
398	30	10	19	—	*	*	24	*8	*6	12	*
398.9	30	10	19	—	*	*	24	*8	*6	12	*
398.91	29	10	19	—	*	*	24	*8	*6	12	*
401	301	126	175	*	42	137	121	68	67	118	48
401.0	26	9	17	*	*5	10	11	*5	*6	11	*
401.9	274	117	158	*	36	127	110	62	62	107	43
402	57	20	37	—	*	15	38	16	12	23	*6
402.0	*6	*	*	—	*	*	*	*	*	*	*
402.9	51	18	33	—	*	13	35	14	11	21	*5
402.90	*8	*	*5	—	*	*	*5	*	*	*	*
402.91	43	15	28	—	*	10	30	12	10	17	*
403	116	56	60	*	24	46	45	22	16	54	23
403.0	*9	*	*	*	*	*	*	*	*	*	*
403.01	*8	*	*	*	*	*	*	*	*	*	*
403.9	106	51	55	*	21	42	43	20	14	50	22
403.91	104	50	54	*	21	40	42	19	14	48	22
404	25	13	12	—	*	*9	14	*	*6	12	*
404.0	*5	*	*	—	*	*	*	*	*	*	*
404.9	20	11	*9	—	*	*6	12	*	*6	9	*
404.93	17	*9	*8	—	*	*	11	*	*	*8	*
410	683	406	277	—	38	215	430	171	154	221	137
410.0	15	*9	*6	—	*	*	*9	*	*	*7	*
410.01	15	*9	*6	—	*	*	*9	*	*	*7	*
410.1	62	36	26	—	*	26	33	12	13	20	18
410.11	60	36	25	—	*	25	31	12	13	19	17
410.2	11	*7	*	—	*	*	*	*	*	*	*
410.21	11	*7	*	—	*	*	*	*	*	*	*
410.3	*7	*6	*	—	*	*	*	*	*	*	*
410.31	*7	*5	*	—	*	*	*	*	*	*	*
410.4	73	55	18	—	*6	35	32	14	17	22	20
410.41	73	55	18	—	*6	35	32	14	17	22	20
410.5	9	*	*	—	*	*	*6	*	*	*	*
410.51	9	*	*	—	*	*	*6	*	*	*	*
410.7	404	235	168	—	19	113	272	113	85	133	71
410.71	398	232	166	—	19	113	267	111	85	131	71
410.8	17	10	*7	—	*	*7	9	*	*7	*	*
410.81	17	9	*7	—	*	*7	*9	*	*7	*	*
410.9	82	42	40	—	*	18	60	15	23	27	17
410.91	80	41	39	—	*	17	58	15	23	26	17
411	89	42	47	—	*9	41	39	25	18	32	14
411.1	83	39	44	—	*9	38	36	22	18	30	13
412	*6	*	*	—	*	*	*	*	*	*	*
413	44	18	25	—	11	15	18	10	11	15	*7
413.9	40	17	23	—	*8	13	18	10	11	12	*6
414	1,007	647	361	*	41	404	562	211	245	365	186
414.0	980	629	351	*	40	393	546	204	239	354	182
414.00	126	76	51	—	*	43	80	34	26	52	14
414.01	817	524	293	*	36	336	444	165	199	288	164
414.02	25	20	*5	—	*	*9	16	*	*9	*8	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
414.04	*5	*	*	—	*	*	*	*	*	*	*
414.05	*5	*	*	—	*	*	*	*	*	*	—
414.8	20	14	*6	—	*	*8	12	*5	*	*7	*
415	140	57	84	*	24	38	77	31	33	53	23
415.1	140	56	83	*	24	38	77	30	33	53	23
415.11	9	*	*5	—	*	*	*6	*	*	*	*
415.19	130	52	78	*	22	36	71	28	31	50	21
416	12	*	*9	*	*	*	*	*	*	*6	*
416.8	*8	*	*6	*	*	*	*	*	*	*	*
420	17	9	*7	*	*	*	*6	*	*6	*	*
420.9	17	9	*7	*	*	*	*6	*	*6	*	*
420.90	11	*6	*6	*	*	*	*5	*	*	*	*
421	16	11	*	*	*	*7	*5	*	*	*7	*
421.0	15	11	*	*	*	*7	*	*	*	*7	*
423	17	*9	*8	*	*	*8	*5	*	*	*7	*
423.9	13	*7	*6	*	*	*6	*	*	*	*6	*
424	100	47	52	*	10	27	61	26	26	27	20
424.0	43	16	27	*	*5	14	23	12	11	14	*6
424.1	51	27	23	*	*	11	36	13	14	12	11
424.9	*5	*	*	*	*	*	*	*	*	*	*
425	41	23	18	*	*8	14	17	*9	10	13	9
425.4	35	21	14	*	*7	12	15	*8	*9	11	*8
426	56	31	26	*	*	*8	43	18	12	14	13
426.0	30	15	15	—	*	*	27	9	*7	*8	*6
426.1	13	*7	*6	*	*	*	10	*	*	*	*
426.13	*8	*	*	*	*	*	*6	*	*	*	*
427	773	365	408	*5	49	182	537	185	191	259	139
427.0	28	11	17	*	*	*6	18	*5	*7	11	*
427.1	59	42	17	*	*	19	36	14	19	20	*7
427.3	454	206	248	*	21	106	327	107	114	150	83
427.31	411	178	233	*	18	91	302	97	106	134	74
427.32	43	29	15	*	*	15	25	10	*8	16	9
427.4	*8	*6	*	*	*	*	*	*	*	*	*
427.41	*8	*6	*	*	*	*	*	*	*	*	*
427.5	*7	*	*	*	*	*	*	*	*	*	*
427.6	*9	*	*	*	*	*	*	*	*	*	*
427.69	*6	*	*	*	*	*	*	*	*	*	*
427.8	202	89	113	*	16	41	143	51	45	68	38
427.81	69	31	38	—	*	*9	60	18	15	24	12
427.89	133	58	75	*	15	32	83	33	30	44	26
427.9	*5	*	*	*	*	*	*	*	*	*	*
428	1,084	494	590	*	42	236	804	239	275	409	161
428.0	1,007	461	546	*	37	223	744	225	253	379	150
428.2	25	13	12	—	*	*5	18	*	10	9	*
428.20	*9	*5	*	—	*	*	*6	*	*	*	*
428.23	9	*	*	—	*	*	*7	*	*	*	*
428.3	43	14	28	—	*	*7	34	10	10	16	*6
428.30	25	*8	17	—	*	*	19	*7	*6	10	*
428.31	10	*	*6	—	*	*	9	*	*	*	*
428.33	*6	*	*	—	*	*	*	*	*	*	*
429	9	*	*6	*	*	*	*	*	*	*	*
430	26	11	15	*	*6	12	*8	*7	*7	*7	*5
431	68	32	36	*	*	21	41	12	13	27	16
432	34	20	14	*	*	*7	26	*6	*8	10	11
432.1	29	18	11	*	*	*7	22	*	*6	*8	9

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
433	145	85	59	*	*	33	110	28	32	60	25
433.1	112	65	48	*	*	26	85	23	25	45	20
433.10	96	55	41	—	*	21	75	19	21	40	17
433.11	16	10	*6	*	*	*5	11	*	*	*	*
433.3	25	16	*9	—	—	*	21	*	*	12	*
433.30	20	14	*6	—	—	*	17	*	*	10	*
433.31	*5	*	*	—	—	*	*	*	*	*	*
434	363	167	196	*	14	82	267	73	86	134	70
434.0	*6	*	*	—	*	*	*	*	*	*	*
434.01	*6	*	*	—	*	*	*	*	*	*	*
434.1	40	19	21	*	*	*8	30	10	10	10	9
434.11	39	18	21	*	*	*8	29	10	10	10	9
434.9	317	144	173	*	12	71	233	61	75	121	60
434.90	*6	*	*	—	*	*	*	*	*	*	*
434.91	311	140	170	*	12	69	229	59	74	118	59
435	195	77	118	*	13	47	135	46	48	69	32
435.3	*7	*	*	—	*	*	*	*	*	*	*
435.8	10	*6	*	—	*	*	*7	*	*	*	*
435.9	178	69	109	*	11	43	123	42	46	63	27
436	11	*6	*5	—	*	*	*9	*	*	*	*
437	32	12	20	*	*	13	15	*8	9	10	*
437.2	*6	*	*	*	*	*	*	*	*	*	*
437.3	11	*	*7	*	*	*6	*	*	*	*	*
438	21	*8	14	*	*	*	15	*5	*6	*	*6
438.2	*6	*	*	*	*	*	*	*	*	*	*
438.20	*5	*	*	*	*	*	*	*	*	*	*
438.8	9	*5	*	—	*	*	*7	*	*	*	*
438.89	*7	*	*	—	*	*	*	*	*	*	*
440	126	71	55	—	*	33	91	24	33	54	16
440.1	14	*7	*7	—	*	*	10	*	*	*6	*
440.2	105	59	46	—	*	26	78	20	27	46	12
440.20	11	*6	*5	—	—	*	*8	*	*	*7	*
440.21	28	16	11	—	*	*9	19	*	9	11	*
440.22	15	*9	*6	—	*	*	11	*	*	*9	*
440.23	16	*9	*8	—	*	*	13	*	*	*9	*
440.24	32	18	14	—	*	*7	24	*8	*8	10	*5
441	63	43	20	*	*	14	47	14	13	26	10
441.0	13	*8	*	—	*	*	*6	*	*	*6	*
441.01	*7	*	*	—	*	*	*	*	*	*	*
441.2	*7	*	*	*	*	*	*	*	*	*	*
441.3	*6	*	*	—	—	*	*6	*	*	*	*
441.4	36	27	9	—	—	*6	31	*8	*7	17	*
442	10	*8	*	—	*	*	*5	*	*	*	*
443	25	14	11	*	*	*8	15	*	*	12	*
443.9	18	10	*8	*	*	*7	11	*	*	10	*
444	31	16	15	—	*	13	17	*7	*	15	*5
444.0	*5	*	*	—	—	*	*	*	*	*	*
444.2	22	10	11	—	*	*8	13	*5	*	10	*
444.22	19	9	10	—	*	*7	11	*5	*	*9	*
446	17	9	*8	*7	*	*	*	*	*	*	*5
446.1	*7	*5	*	*7	*	*	—	*	*	*	*
447	13	*7	*7	*	*	*5	*7	*	*	*	*
447.1	10	*5	*	—	*	*	*6	*	*	*	*
451	17	*8	9	*	*	*	*	*7	*5	*	*
451.1	*6	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
453	132	56	77	*	17	42	73	33	33	47	19
453.4	103	44	58	*	12	32	59	25	27	34	16
453.40	17	*8	9	—	*	*	11	*	*	*7	*
453.41	55	23	32	—	*6	17	32	16	15	17	*8
453.42	30	13	17	*	*	9	17	*8	*9	10	*
453.8	27	10	16	*	*	10	12	*7	*5	12	*
454	*7	*	*6	—	*	*	*6	*	*5	*	—
455	30	14	15	—	*7	*9	14	*8	*7	*8	*6
455.2	20	11	9	—	*	*6	*9	*6	*	*	*
456	*8	*5	*	—	*	*	*	*	*	*	*
458	109	49	60	*	12	20	77	21	28	44	16
458.0	51	20	31	*	*	*6	43	10	14	21	*6
458.2	18	*7	11	—	*	*	11	*	*	*8	*
458.29	16	*6	10	—	*	*	9	*	*	*6	*
458.9	35	20	15	—	*	9	21	*6	*9	14	*6
459	24	10	13	*	*	*6	15	*6	*8	*7	*
459.0	*6	*	*	—	*	*	*	*	*	*	*
459.8	13	*7	*6	*	*	*	*8	*	*	*	*
459.81	13	*7	*6	—	*	*	*7	*	*	*	*
460–519	3,746	1,764	1,983	629	370	825	1,921	749	967	1,353	676
461	*7	*	*	*	*	*	*	*	*	*	*
462	*7	*	*	*	*	*	*	*	*	*	*
463	10	*6	*	*	*7	*	*	*	*	*	*
464	30	16	14	25	*	*	*	*5	*7	*8	10
464.4	21	12	*9	20	*	*	—	*	*6	*6	*6
465	32	18	13	18	*	*5	*5	*5	*6	12	*8
465.9	30	17	13	17	*	*	*	*5	*6	11	*8
466	230	117	113	155	13	18	43	37	50	86	56
466.0	77	28	49	*8	11	17	41	15	20	29	13
466.1	153	89	64	147	*	*	*	23	30	57	43
466.11	80	48	33	79	*	*	*	10	17	27	27
466.19	73	41	32	68	*	*	*	13	13	30	16
473	18	*6	11	*	*6	*	*	*	*	*7	*
473.9	*8	*	*5	*	*	*	*	*	*	*	*
474	15	9	*6	11	*	*	*	*	*	*7	*
474.0	*6	*	*	*	*	*	*	*	*	*	*
474.1	10	*7	*	*8	*	*	—	*	*	*	*
474.10	*6	*	*	*5	*	—	—	*	*	*	*
475	16	10	*6	*	12	*	*	*	*6	*6	*
478	22	10	12	*8	*	*6	*	*	*6	*7	*
478.1	*5	*	*	*	*	*	*	*	*	*	*
478.2	*8	*	*	*	*	*	*	*	*	*	*
480	32	14	18	24	*	*	*	*	*7	11	10
480.1	13	*	9	13	—	*	*	*	*	*5	*
480.9	16	*8	*8	9	*	*	*	*	*	*5	*
481	35	15	20	*	*6	9	17	*7	11	11	*5
482	101	55	46	*	*9	20	68	16	27	39	19
482.0	*5	*	*	*	*	*	*	*	*	*	*
482.1	21	13	*9	*	*	*	14	*	*7	*8	*
482.2	*6	*	*	*	*	*	*5	*	*	*	*
482.3	*6	*	*	*	*	*	*	*	*	*	*
482.4	34	18	16	*	*	*8	22	*5	*7	13	*9
482.41	32	17	15	*	*	*7	21	*	*6	13	*9
482.8	15	10	*6	*	*	*	12	*	*	*8	*
482.83	*9	*6	*	*	*	*	*7	*	*	*5	*
482.9	14	*6	*8	*	*	*	*7	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
483	10	*	*6	*6	*	*	*	*	*	*	*
483.0	*9	*	*	*5	*	*	*	*	*	*	*
485	30	15	15	14	*	*	11	*	*	21	*
486	1,161	549	612	143	92	213	713	222	325	403	211
487	62	29	33	14	*	*8	36	10	27	17	*8
487.0	20	10	10	*	*	*	13	*	*8	*6	*
487.1	40	18	22	10	*	*5	23	*6	19	11	*
490	26	*7	19	*	*	11	11	*	*	15	*
491	569	261	308	*	16	166	386	116	156	222	75
491.2	566	260	305	*	16	166	384	116	155	220	75
491.20	15	*6	9	—	*	*	12	*	*	*9	*
491.21	460	216	244	*	12	135	312	95	127	176	62
491.22	91	38	52	—	*	28	60	19	24	35	12
492	16	10	*6	*	*	*6	10	*	*	*	*
492.8	15	9	*6	*	*	*5	9	*	*	*	*
493	489	192	296	159	98	119	112	122	115	173	78
493.0	38	21	17	29	*	*	*	*9	*	*9	18
493.00	*6	*	*	*	*	*	*	*	*	*	*
493.01	*7	*	*	*6	*	*	—	*	*	*	*
493.02	25	14	11	19	*	*	*	*6	*	*	13
493.2	138	41	97	*	12	56	70	40	32	46	20
493.20	20	*7	14	*	*	*6	12	*	*6	*	*
493.21	*8	*	*6	*	*	*	*	*	*	*	*
493.22	110	33	77	*	*8	47	55	32	25	39	15
493.9	311	130	181	129	82	60	41	73	81	118	40
493.90	51	22	30	20	10	11	9	12	*8	23	*8
493.91	44	21	23	32	*7	*	*	*8	11	20	*5
493.92	216	87	129	77	64	44	31	53	61	75	26
494	*7	*	*	*	*	*	*	*	*	*	*
496	101	47	53	*	*	42	56	18	28	37	18
507	177	98	79	*6	13	24	134	40	42	60	36
507.0	175	96	79	*6	13	23	133	40	41	59	36
510	11	*7	*	*	*	*5	*	*	*	*	*
510.9	11	*7	*	*	*	*	*	*	*	*	*
511	63	34	29	*	*7	16	40	16	17	20	9
511.0	*6	*	*	—	*	*	*	*	*	*	*
511.8	*8	*	*	*	*	*	*	*	*	*	*
511.9	48	25	23	*	*	11	33	13	12	15	*8
512	41	23	18	*	18	*9	13	9	*9	16	*7
512.1	11	*	*7	*	*	*	*5	*	*	*	*
512.8	27	17	11	*	14	*	*8	*6	*6	11	*
514	*6	*	*	—	*	*	*	*	*	*	*
515	36	21	15	*	*	12	22	*6	11	10	*8
516	17	*9	*9	*	*	*7	*9	*	*	*7	*
516.3	*5	*	*	—	—	*	*	*	*	*	*
516.8	12	*6	*5	*	*	*6	*	*	*	*6	*
518	338	155	183	14	31	95	197	66	73	126	73
518.0	10	*	*5	*	*	*	*	*	*	*	*
518.3	*5	*	*	*	*	*	*	*	*	*	*
518.8	311	144	167	12	26	87	186	60	66	117	67
518.81	248	113	135	9	20	66	153	51	50	92	55
518.82	*7	*	*	*	*	*	*	*	*	*	*
518.84	43	19	24	*	*	16	24	*6	11	17	*8
518.89	*8	*	*	*	*	*	*	*	*	*	*
519	16	*6	9	*6	*	*	*	*5	*	*	*
519.8	*6	*	*	*	*	*	*	*	*	*	*

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
520–579	3,422	1,546	1,876	216	832	1,052	1,322	712	797	1,225	687
522	*6	*	*	*	*	*	*	*	*	*	*
522.5	*5	*	*	*	*	*	*	*	*	*	*
524	13	*	*9	*	11	*	*	*	*	*7	*
524.0	*6	*	*	*	*6	*	—	*	*	*	*
527	13	*8	*	*	*	*	*7	*	*	*6	*
527.2	*8	*	*	*	*	*	*	*	*	*	*
528	13	*7	*6	*	*6	*	*	*	*	*7	*
528.3	*6	*	*	*	*	*	*	*	*	*	*
530	225	105	120	29	40	80	77	52	52	71	50
530.1	36	16	20	*	*7	14	14	*8	*9	12	*8
530.10	*8	*	*	*	*	*	*	*	*	*	*
530.11	13	*5	*7	*	*	*6	*	*	*	*	*
530.19	14	*6	*8	*	*	*5	*5	*	*	*	*
530.2	18	9	*9	—	*	*6	*9	*	*	*	*
530.20	*7	*	*	—	*	*	*	*	*	*	*
530.21	11	*5	*6	—	*	*	*8	*	*	*	*
530.3	*7	*	*	*	*	*	*	*	*	*	*
530.7	17	12	*	*	*	*5	*6	*	*	*8	*
530.8	137	59	78	24	22	50	40	35	31	41	29
530.81	124	52	72	24	21	45	34	32	29	37	25
530.85	*6	*	*	—	*	*	*	*	*	*	*
531	87	37	51	*	*8	28	51	11	20	35	22
531.0	12	*	*8	*	*	*	*8	*	*	*	*
531.00	12	*	*8	*	*	*	*8	*	*	*	*
531.4	55	25	30	*	*	13	37	*6	12	22	14
531.40	53	25	29	*	*	13	36	*6	12	22	14
531.9	12	*5	*7	—	*	*7	*	*	*	*	*
531.90	12	*	*7	—	*	*7	*	*	*	*	*
532	58	34	24	*	*7	20	30	10	11	25	12
532.0	10	*6	*	—	*	*	*	*	*	*	*
532.00	10	*6	*	—	*	*	*	*	*	*	*
532.4	29	20	9	*	*	*8	18	*	*	13	*7
532.40	29	20	9	*	*	*8	18	*	*	13	*7
532.9	*8	*	*6	—	*	*5	*	*	*	*5	*
532.90	*7	*	*5	—	*	*	*	*	*	*	*
533	16	*5	11	—	*	*	*8	*	*5	*7	*
533.4	*7	*	*	—	*	*	*5	*	*	*	*
533.40	*7	*	*	—	*	*	*5	*	*	*	*
534	*6	*	*	*	*	*	*	*	*	*	*
535	149	65	84	*6	30	46	67	39	30	55	25
535.0	29	16	13	*	*9	*6	11	10	*7	*9	*
535.00	21	11	11	*	*8	*	*6	10	*	*5	*
535.01	*8	*5	*	*	*	*	*	*	*	*	*
535.1	18	*7	10	*	*	*	11	*7	*	*6	*
535.10	10	*	*7	*	*	*	*6	*5	*	*	*
535.11	*8	*	*	*	*	*	*	*	*	*	*
535.3	*7	*6	*	—	*	*	*	*	*	*	*
535.4	35	14	21	*	*6	13	16	*7	*7	13	*8
535.40	18	*7	11	*	*	*7	*5	*	*	*6	*
535.41	18	*7	10	—	*	*5	10	*	*	*7	*
535.5	53	18	35	*	10	17	24	11	11	23	*9
535.50	34	11	23	*	*8	12	12	*7	*7	14	*5
535.51	19	*7	12	*	*	*5	12	*	*	*9	*
535.6	*7	*	*	*	*	*	*	*	*	*	*
536	53	19	34	*8	11	15	20	11	*9	21	13
536.2	17	*7	10	*	*	*	*	*	*	*8	*
536.3	9	*	*7	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
536.4	16	*6	10	*	*	*	10	*5	*	*	*
536.42	*7	*	*	*	*	*	*	*	*	*	*
536.8	*9	*	*6	*	*	*	*	*	*	*	*
537	28	16	13	*	*	*9	15	*6	*	10	9
537.0	*8	*5	*	*	*	*	*	*	*	*	*
537.8	17	9	*8	—	*	*	11	*	*	*	*6
537.83	10	*5	*	—	*	*	*7	*	*	*	*
540	301	172	130	74	151	55	21	60	64	90	87
540.0	61	37	24	18	22	16	*6	*8	16	18	19
540.1	35	22	13	12	12	*7	*8	*7	11	9	
540.9	205	112	92	44	117	32	12	43	40	61	59
541	11	*	*7	*	*6	*	*	*	*	*5	*
543	*6	*	*	*	*	*	*	*	*	*	*
550	37	31	*6	*6	*	*7	19	10	*7	13	*6
550.1	22	18	*	*	*	*	13	*5	*5	*7	*
550.10	14	11	*	*	*	*	*8	*	*	*6	*
550.11	*7	*6	*	*	*	*	*	*	*	*	*
550.9	14	12	*	*	*	*	*6	*	*	*5	*
550.90	*8	*7	*	*	*	*	*	*	*	*	*
552	69	23	46	*	10	25	33	18	16	23	11
552.1	9	*5	*	—	*	*	*	*	*	*	*
552.2	44	14	30	—	*7	17	20	12	*8	17	*7
552.20	11	*	*8	—	*	*	*	*	*	*	*
552.21	28	9	19	—	*	10	14	*7	*6	10	*
552.29	*5	*	*	—	*	*	*	*	*	*	*
552.8	*6	*	*	*	*	*	*	*	*	*	*
553	80	29	52	*	15	36	29	17	18	32	14
553.2	59	23	37	*	11	28	20	12	16	20	11
553.20	11	*	*8	*	*	*	*	*	*	*6	*
553.21	45	18	26	*	*9	22	14	10	12	13	10
553.3	17	*	13	*	*	*6	*8	*	*	10	*
555	56	23	33	*	35	12	*7	13	15	18	10
555.0	13	*6	*7	*	9	*	*	*	*	*	*
555.1	11	*	*6	*	*7	*	*	*	*	*	*
555.2	*8	*	*	*	*	*	*	*	*	*	*
555.9	25	11	15	*	15	*	*	*	*6	10	*
556	29	15	14	*	15	*6	*6	*7	*6	*9	*7
556.6	*9	*5	*	*	*	*	*	*	*	*	*
556.9	11	*6	*	*	*5	*	*	*	*	*	*
557	49	12	37	*	*	12	34	11	11	17	10
557.0	20	*6	15	*	*	*	15	*6	*	*6	*
557.9	24	*	19	—	*	*6	16	*	*5	9	*5
558	184	68	116	36	41	45	61	44	46	71	23
558.9	180	67	112	36	41	44	60	44	43	70	23
560	304	134	169	11	40	89	164	57	84	102	61
560.0	*6	*	*	*	*	*	*	*	*	*	*
560.1	57	31	26	*	*7	16	32	10	19	20	*8
560.2	11	*5	*6	*	*	*	*6	*	*	*	*
560.3	29	11	19	*	*	*6	17	*	*	15	*5
560.39	28	11	18	*	*	*6	16	*	*	14	*5
560.8	98	39	59	*	11	32	52	18	25	33	21
560.81	72	30	42	*	*8	23	39	14	17	25	16
560.89	26	*8	17	*	*	9	13	*	*9	*8	*
560.9	104	44	59	*	14	32	57	22	29	29	23
562	307	126	181	—	37	106	164	76	79	102	51
562.1	304	125	180	—	37	105	162	75	79	100	50
562.10	26	10	16	—	*	11	13	*7	*	*8	*7
562.11	190	74	116	—	33	77	80	52	55	56	27

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
562.12	72	35	38	—	*	12	59	13	13	31	15
562.13	16	*6	10	—	*	*	11	*	*7	*	*
564	66	23	43	*9	16	16	25	15	15	24	11
564.0	42	16	27	*8	*8	9	17	*8	10	16	*8
564.00	32	12	20	*6	*5	*7	15	*5	*9	13	*6
564.09	10	*	*6	*	*	*	*	*	*	*	*
564.1	14	*	11	*	*7	*	*	*	*	*5	*
564.8	*6	*	*	*	*	*	*	*	*	*	*
564.89	*6	*	*	*	*	*	*	*	*	*	*
566	21	15	*6	*	*8	*8	*	*	*	10	*
567	20	9	11	*	*	*9	*6	*	*6	*6	*
567.2	14	*8	*6	*	*	*8	*	*	*	*	*
568	11	*	10	*	*5	*	*	*	*	*	*
568.0	*7	*	*7	*	*	*	*	*	*	*	*
569	100	43	57	*	12	24	62	19	22	41	18
569.1	*7	*	*6	*	*	*	*	*	*	*	*
569.3	24	12	12	*	*	*	17	*	*	13	*
569.4	*9	*6	*	—	*	*	*6	*	*	*	*
569.6	17	*7	10	*	*	*6	*8	*	*	*7	*
569.69	12	*6	*6	*	*	*	*7	*	*	*	*
569.8	39	16	23	*	*	9	25	*8	10	14	*7
569.81	*5	*	*	*	*	*	*	*	*	*	*
569.82	*8	*	*6	*	*	*	*	*	*	*	*
569.83	*8	*	*	*	*	*	*	*	*	*	*
569.85	15	*8	*7	—	*	*	13	*	*	*	*
570	9	*	*6	*	*	*	*	*	*	*5	*
571	112	68	45	*	18	69	24	17	15	46	34
571.1	13	*7	*6	—	*6	*7	*	*	*	*	*
571.2	48	33	14	—	*8	34	*6	*7	*6	19	16
571.5	38	23	15	*	*	21	15	*	*5	16	12
572	46	28	19	*	*7	24	15	10	*8	15	13
572.2	36	21	15	—	*	19	12	*6	*6	12	11
573	9	*	*	*	*	*	*	*	*	*	*
573.3	*6	*	*	*	*	*	*	*	*	*	*
574	325	116	210	*	106	95	124	68	74	118	66
574.0	120	46	74	*	39	37	43	29	26	42	22
574.00	114	44	70	*	37	37	41	28	26	40	20
574.01	*5	*	*	—	*	*	*	*	*	*	*
574.1	97	33	64	*	31	30	37	21	22	36	18
574.10	95	32	63	*	30	30	35	21	22	35	17
574.2	42	14	28	*	11	12	18	*7	11	15	*8
574.20	40	12	27	*	10	11	18	*6	11	15	*8
574.5	21	*7	14	*	*7	*	*9	*	*5	*7	*
574.50	12	*	*7	*	*	*	*	*	*	*	*
574.51	*9	*	*6	*	*	*	*	*	*	*	*
574.6	11	*	*7	*	*	*	*	*	*	*	*
574.60	*7	*	*5	*	*	*	*	*	*	*	*
574.7	13	*	10	—	*6	*	*	*	*	*	*
574.70	*8	*	*7	—	*	*	*	*	*	*	*
574.8	9	*	*	*	*	*	*	*	*	*	*
574.81	*5	*	*	—	*	*	*	*	*	*	*
575	81	30	51	*	31	28	21	14	23	31	13
575.0	26	12	14	*	*7	10	*9	*5	*9	*7	*5
575.1	45	15	30	*	20	14	11	*7	12	20	*7
575.10	*7	*	*6	*	*	*	*	*	*	*	*
575.11	28	*6	21	*	15	*7	*6	*	*8	13	*
575.12	10	*7	*	—	*	*	*	*	*	*	*
575.8	*5	*	*	*	*	*	*	*	*	*	*

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
576	27	10	17	*	*5	*9	13	*5	*6	9	*7
576.1	*8	*	*	*	*	*	*6	*	*	*	*
576.2	*8	*	*	*	*	*	*	*	*	*	*
576.8	10	*	*7	*	*	*	*	*	*	*	*
577	275	148	126	*	96	105	71	52	60	115	49
577.0	237	129	108	*	83	88	63	42	56	96	43
577.1	25	12	12	*	10	13	*	*6	*	14	*
577.2	*7	*	*	*	*	*	*	*	*	*	*
578	193	93	100	*5	18	38	132	41	60	61	31
578.0	20	10	10	*	*	*6	*9	*7	*5	*5	*
578.1	37	16	21	*	*6	*9	21	10	*7	15	*5
578.9	135	67	69	*	*8	23	103	24	48	41	23
579	*7	*	*	*	*	*	*	*	*	*	*
580–629	1,983	642	1,341	91	535	496	861	376	464	782	361
581	*5	*	*	*	*	*	*	*	*	*	*
584	270	145	125	*	18	65	186	66	62	95	46
584.5	14	*7	*6	*	*	*6	*6	*	*	*	*
584.9	253	135	118	*	15	59	178	62	58	90	42
585	16	*9	*7	*	*	*	9	*6	*	*	*
586	*7	*	*	*	—	*	*6	*	*	*	*
590	143	23	120	29	50	28	36	24	33	55	32
590.1	66	10	56	12	24	15	15	12	15	23	16
590.10	66	10	56	12	24	15	15	12	15	23	16
590.8	73	13	61	17	23	12	21	11	17	30	15
590.80	73	13	61	17	23	12	21	11	17	30	15
591	9	*	*6	*	*	*	*	*	*	*	*
592	184	94	90	*	75	68	37	37	50	77	20
592.0	57	25	32	*	22	18	16	12	16	22	*6
592.1	126	69	57	*	52	50	21	24	33	55	14
593	52	23	28	*9	10	10	23	10	10	22	*9
593.4	10	*	*6	*	*	*	*	*	*	*	*
593.7	*8	*	*7	*8	*	—	*	*	*	*	*
593.70	*7	*	*6	*6	*	—	—	*	*	*	*
593.8	*9	*	*5	*	*	*	*6	*	*	*	*
593.81	*6	*	*	—	*	*	*	*	*	*	*
593.9	19	12	*7	*	*	*6	11	*	*	11	*
595	17	*7	10	*	*	*	12	*	*6	*5	*
595.0	*6	*	*	*	*	*	*	*	*	*	*
595.9	*6	*	*	*	*	*	*	*	*	*	*
596	23	16	*7	*	*	*5	14	*	*	10	*5
596.0	*6	*6	—	—	—	*	*5	*	*	*	*
598	*6	*5	*	*	*	*	*	*	*	*	*
599	522	181	341	29	47	76	370	103	124	200	96
599.0	481	157	324	28	43	66	344	95	114	186	86
599.7	29	20	*9	*	*	*	20	*6	*8	10	*5
599.8	*7	*	*6	—	*	*	*	*	*	*	*
600	80	80	—	—	*	15	64	18	21	22	20
600.0	73	73	—	—	*	14	59	16	19	19	19
600.00	16	16	—	—	—	*	13	*	*	*	*
600.01	57	57	—	—	*	11	46	12	14	16	15
601	*6	*6	—	—	*	*	*	*	*	*	*
604	12	12	—	*	*	*	*	*	*	*	*
604.9	11	11	—	*	*	*	*	*	*	*	*
604.90	*6	*6	—	*	*	*	*	*	*	*	*
604.99	*5	*5	—	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
607	*6	*6	—	*	*	*	*	*	*	*	*
608	11	11	—	*	*5	*	*	*	*	*	*
608.4	*6	*6	—	*	*	*	*	*	*	*	*
611	28	*	27	*	15	10	*	*9	*	14	*
611.0	11	*	10	*	*5	*	*	*	*	*	*
611.1	*9	*	*8	*	*6	*	*	*	*	*	*
611.7	*6	*	*6	—	*	*	*	*	*	*	*
614	45	—	45	*	35	*8	*	*8	*8	23	*6
614.1	*7	—	*7	—	*	*	*	*	*	*5	*
614.2	*8	—	*8	—	*6	*	*	*	*	*	*
614.6	9	—	9	—	*8	*	*	*	*	*	*
614.9	11	—	11	*	10	*	*	*	*	*5	*
616	11	—	11	*	*	*	*	*	*	*	*
617	61	—	61	—	48	12	*	*8	17	25	11
617.0	35	—	35	—	27	*8	*	*	10	15	*5
617.1	11	—	11	—	*9	*	—	*	*	*	*
617.3	*7	—	*7	—	*6	*	—	*	*	*	*
618	141	—	141	*	20	68	53	20	31	58	31
618.0	28	—	28	—	*	13	13	*	*7	9	*7
618.01	15	—	15	—	*	*6	*8	*	*	*	*5
618.04	*8	—	*8	—	*	*	*	*	*	*	*
618.1	14	—	14	*	*	*7	*	*	*	*6	*
618.2	22	—	22	—	*	11	*6	*	*	*8	*7
618.3	14	—	14	—	*	*6	*7	*	*	*	*
618.4	19	—	19	—	*	11	*	*	*	*8	*
618.5	33	—	33	—	*	14	18	*	*7	17	*6
618.8	*8	—	*8	—	*	*	*	*	*	*	*
618.89	*7	—	*7	—	*	*	*	*	*	*	*
620	66	—	66	*	48	13	*	11	18	27	10
620.0	*6	—	*6	—	*5	*	—	*	*	*	*
620.1	9	—	9	*	*8	*	—	*	*	*	*
620.2	38	—	38	*	26	*9	*	*6	10	17	*6
620.8	*7	—	*7	*	*5	*	*	*	*	*	*
621	17	—	17	—	*7	*8	*	*	*	*8	*
621.3	*8	—	*8	—	*	*	*	*	*	*	*
622	*9	—	*9	—	*	*	*	*	*	*	*
622.1	*8	—	*8	—	*	*	*	*	*	*	*
625	69	—	69	*	39	21	*7	10	15	25	18
625.3	12	—	12	*	11	*	—	*	*	*	*
625.6	27	—	27	—	*7	15	*5	*	*8	*8	*8
625.9	22	—	22	*	17	*	*	*	*	10	*
626	121	—	121	*	82	39	*	12	30	57	22
626.2	94	—	94	*	64	30	—	*8	24	44	17
626.8	22	—	22	*	14	*8	*	*	*5	9	*
627	12	—	12	—	*	*5	*	*	*	*	*
627.1	9	—	9	—	*	*	*	*	*	*	*
630–677	535	—	535	*	533	*	—	77	91	255	111
632	21	—	21	*	21	*	—	*	*	12	*
633	33	—	33	—	33	—	—	*	*6	15	*7
633.1	26	—	26	—	26	—	—	*	*	12	*6
633.10	26	—	26	—	26	—	—	*	*	12	*6
634	17	—	17	*	17	*	—	*	*	9	*
634.9	10	—	10	*	10	*	—	*	*	*7	*
634.91	*8	—	*8	*	*8	*	—	*	*	*5	*
640	*8	—	*8	—	*8	—	—	*	*	*	*
641	15	—	15	*	15	—	—	*	*	*	*
641.1	*6	—	*6	—	*6	—	—	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
641.13	*6	—	*6	—	*6	—	—	*	*	*	*
641.9	*5	—	*5	—	*5	—	—	*	*	*	*
641.93	*5	—	*5	—	*5	—	—	*	*	*	*
642	33	—	33	—	33	—	—	*	*5	17	*7
642.3	12	—	12	—	12	—	—	*	*	*5	*
642.33	*9	—	*9	—	*9	—	—	*	*	*	*
642.4	*7	—	*7	—	*7	—	—	*	*	*	*
642.43	*6	—	*6	—	*6	—	—	*	*	*	*
643	33	—	33	*	33	—	—	*	*6	16	*5
643.0	14	—	14	—	14	—	—	*	*	*8	*
643.03	14	—	14	—	14	—	—	*	*	*8	*
643.1	11	—	11	*	11	—	—	*	*	*6	*
643.13	11	—	11	*	11	—	—	*	*	*6	*
643.2	*6	—	*6	—	*6	—	—	*	*	*	*
643.23	*6	—	*6	—	*6	—	—	*	*	*	*
644	109	—	109	*	109	*	—	13	21	50	24
644.0	95	—	95	*	95	*	—	12	19	43	21
644.03	95	—	95	*	95	*	—	12	19	43	21
644.1	14	—	14	—	14	—	—	*	*	*7	*
644.13	14	—	14	—	14	—	—	*	*	*7	*
646	69	—	69	*	69	—	—	*6	10	39	14
646.6	45	—	45	*	45	—	—	*	*5	26	10
646.63	42	—	42	*	42	—	—	*	*5	25	*8
646.8	18	—	18	—	18	—	—	*	*	10	*
646.83	17	—	17	—	17	—	—	*	*	10	*
647	*6	—	*6	—	*6	—	—	*	*	*	*
648	94	—	94	*	94	*	—	16	17	43	18
648.0	9	—	9	*	9	—	—	*	*	*6	*
648.03	9	—	9	*	9	—	—	*	*	*6	*
648.4	13	—	13	*	13	—	—	*	*	*7	*
648.43	9	—	9	*	*9	—	—	*	*	*6	*
648.9	56	—	56	—	56	*	—	11	10	23	13
648.93	52	—	52	—	52	*	—	10	*9	20	13
654	15	—	15	—	15	*	—	*	*	*9	*
654.5	*8	—	*8	—	*8	*	—	*	*	*	*
654.53	*8	—	*8	—	*8	*	—	*	*	*	*
655	*5	—	*5	—	*5	—	—	*	*	*	*
658	*8	—	*8	—	*8	—	—	*	*	*	*
659	*7	—	*7	*	*7	—	—	*	*	*	*
666	*5	—	*5	*	*5	—	—	*	*	*	*
670	*9	—	*9	*	*9	—	—	*	*	*	*
670.0	*9	—	*9	*	*9	—	—	*	*	*	*
670.04	*9	—	*9	*	*9	—	—	*	*	*	*
674	11	—	11	—	11	*	—	*	*	*	*
674.3	*6	—	*6	—	*6	*	—	*	*	*	*
674.34	*6	—	*6	—	*6	*	—	*	*	*	*
680–709	751	400	351	111	188	225	226	155	169	308	119
681	32	17	15	*	*9	13	*8	*8	11	11	*
681.0	19	10	9	*	*8	*	*	*5	*	*7	*
681.00	16	*9	*8	*	*7	*	*	*	*	*6	*
681.1	13	*7	*6	*	*	*8	*	*	*7	*	*
681.10	12	*7	*5	*	*	*7	*	*	*7	*	*
682	518	283	235	60	135	163	161	114	124	191	89
682.0	40	20	20	11	14	10	*6	*6	*6	16	12
682.1	10	*5	*	*	*	*	*	*	*	*	*
682.2	46	24	22	*6	14	16	11	*8	11	19	*9
682.3	50	30	21	*	22	13	12	10	13	20	*7

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
682.4	36	21	16	*	14	9	10	*8	10	12	*6
682.5	24	10	14	*9	*6	*6	*	*	*5	13	*
682.6	261	147	114	19	51	94	97	63	63	90	45
682.7	47	24	22	*	10	12	21	12	12	17	*6
685	*6	*	*	*	*	*	*	*	*	*	*
692	*7	*	*6	*	*	*	*	*	*	*	*
693	*8	*	*	*	*	*	*	*	*	*	*
693.0	*7	*	*	*	*	*	*	*	*	*	*
695	*8	*	*	*	*	*	*	*	*	*	*
695.1	*7	*	*	*	*	*	*	*	*	*	*
701	*7	*	*5	*	*	*	*	*	*	*	—
707	78	39	39	*	12	24	42	15	17	32	14
707.0	47	22	25	*	9	12	26	*8	10	21	*8
707.03	19	*8	11	*	*	*	11	*	*	9	*
707.04	*7	*	*	—	*	*	*	*	*	*	*
707.05	*5	*	*	—	*	*	*	*	*	*	*
707.07	*6	*	*	—	*	*	*	*	*	*	*
707.09	*6	*	*	—	*	*	*	*	*	*	*
707.1	29	16	13	*	*	11	15	*6	*7	9	*6
707.15	9	*5	*	—	*	*	*5	*	*	*	*
707.19	*5	*	*	*	*	*	*	*	*	*	*
708	*6	*	*	*	*	*	*	*	*	*	*
709	51	31	20	34	14	*	*	*	*	50	—
709.2	49	30	20	34	14	*	*	*	*	49	—
710–739	1,945	831	1,114	40	304	686	916	405	504	631	406
710	30	*5	25	*	13	*9	*6	*6	*7	12	*5
710.0	23	*	20	*	12	*6	*	*	*6	*8	*
711	24	14	10	*	*7	*6	*8	*8	*	*8	*
711.0	23	14	10	*	*7	*6	*8	*8	*	*7	*
711.06	12	*8	*	*	*	*	*	*	*	*	*
714	19	*7	12	*	*	*6	*9	*6	*	*6	*
714.0	14	*	10	—	*	*	*8	*	*	*	*
715	761	292	469	*	17	266	478	160	214	224	163
715.1	56	27	30	*	*	15	41	15	*7	19	15
715.15	16	9	*7	—	*	*5	10	*	*	*	*5
715.16	36	16	21	—	*	9	27	10	*6	11	10
715.2	*7	*	*	*	*	*	*	*	*	*	*
715.3	420	163	258	*	10	149	261	103	118	110	89
715.31	14	*5	*9	—	*	*	10	*	*	*5	*5
715.35	115	55	60	—	*	42	69	30	34	25	26
715.36	286	100	186	*	*	101	179	70	82	78	56
715.8	*8	*	*6	—	*	*7	*	*	*	*	*
715.89	*8	*	*6	—	*	*7	*	*	*	*	*
715.9	267	96	171	—	*	97	166	39	84	90	54
715.90	10	*	*6	—	*	*	*6	*	*	*	*
715.91	*7	*	*	—	*	*	*	*	*	*	*
715.95	66	29	37	—	*	20	43	13	22	14	17
715.96	181	59	122	—	*	69	111	24	57	69	31
716	14	*5	9	*	*	*5	*6	*	*	*	*
716.1	*6	*	*	—	*	*	*	*	*	*	*
716.9	*7	*	*5	*	*	*	*	*	*	*	*
718	13	*7	*6	*	*7	*	*	*	*	*	*
719	42	16	26	*	*7	12	21	*8	13	11	11
719.4	28	12	16	*	*6	*8	13	*5	9	*7	*8
719.41	*7	*	*	*	*	*	*	*	*	*	*
719.45	12	*	*8	*	*	*	*7	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
721	109	57	51	*	15	53	41	16	26	38	28
721.0	31	15	15	—	*6	20	*	*	*5	12	10
721.1	24	16	*8	—	*	12	10	*5	*	*8	*7
721.3	46	20	26	—	*7	16	23	*6	16	13	10
721.4	*6	*	*	—	*	*	*	*	*	*	*
722	313	153	159	*	111	140	61	61	79	115	57
722.0	53	26	27	—	23	26	*	11	12	23	*8
722.1	141	78	63	*	59	58	25	30	41	46	24
722.10	139	78	61	*	58	57	24	30	40	45	24
722.4	10	*	*6	—	*	*6	*	*	*	*	*
722.5	65	25	40	*	13	31	20	9	17	25	14
722.52	63	24	39	—	13	30	20	9	16	23	14
722.7	21	11	10	—	*7	*9	*5	*	*	11	*
722.71	14	*7	*7	—	*	*6	*	*	*	*7	*
722.73	*6	*	*	—	*	*	*	*	*	*	*
722.8	*6	*	*	—	*	*	*	*	*	*	*
722.83	*6	*	*	—	*	*	*	*	*	*	*
722.9	14	*	9	—	*	*6	*	*	*	*5	*
722.93	9	*	*6	—	*	*	*	*	*	*	*
723	37	19	18	*	*8	20	*9	*6	*9	12	11
723.0	16	10	*6	*	*	10	*	*	*	*5	*
723.1	12	*6	*6	*	*	*6	*	*	*	*	*
723.4	*8	*	*6	—	*	*	*	*	*	*	*
724	150	63	87	*	21	42	86	37	41	47	25
724.0	84	38	45	—	*	20	60	23	20	27	14
724.02	81	37	44	—	*	20	59	22	19	26	14
724.2	34	15	19	*	9	11	14	*5	12	9	*8
724.4	*8	*	*	—	*	*	*	*	*	*	*
724.5	15	*	11	*	*	*	*	*6	*	*	*5
726	30	15	15	*	*5	10	14	*5	*6	10	9
726.1	*8	*	*	—	*	*	*	*	*	*	*
726.10	*8	*	*	—	*	*	*	*	*	*	*
726.3	*6	*	*	—	*	*	*	*	*	*	*
726.33	*5	*	*	—	*	*	*	*	*	*	*
726.6	*6	*	*	*	*	*	*	*	*	*	*
727	20	*9	12	*	*	*5	9	*	*	10	*
727.0	*7	*	*	*	*	*	*	*	*	*	*
727.6	*7	*	*	—	*	*	*	*	*	*	*
728	38	24	14	*	12	9	15	10	9	13	*6
728.8	35	22	13	*	11	*9	14	*8	*9	13	*
728.88	23	16	*7	*	*8	*6	9	*6	*6	*8	*
728.89	*5	*	*	*	*	*	*	*	*	*	*
729	55	19	36	*	15	18	20	10	15	21	9
729.1	12	*	*9	*	*5	*	*	*	*	*	*
729.5	21	*8	13	*	*6	*9	*5	*	*7	*7	*
729.8	16	*7	10	*	*	*	9	*	*5	*7	*
729.89	12	*5	*7	*	*	*	*8	*	*	*5	*
730	61	42	19	*	11	21	24	10	19	19	13
730.0	12	*8	*	*	*	*	*	*	*	*	*
730.07	*5	*	*	*	*	*	*	*	*	*	*
730.1	10	*7	*	*	*	*	*	*	*	*	*
730.2	38	26	12	*	*6	13	17	*5	13	13	*7
730.26	*6	*	*	*	*	*	*	*	*	*	*
730.27	18	11	*7	*	*	*6	10	*	*7	*6	*
730.28	*8	*5	*	*	*	*	*	*	*	*	*
733	151	51	100	*	23	38	86	34	33	56	27
733.1	70	18	51	*	*	10	57	18	18	23	11
733.13	54	14	40	*	*	*6	47	13	13	19	*8
733.14	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
733.4	27	12	15	*	*7	10	10	*	*	11	*7
733.42	20	10	10	*	*6	*7	*7	*	*	*8	*6
733.6	22	*5	16	*	*6	10	*5	*6	*	10	*
733.8	23	13	10	*	*7	*7	*8	*	*	*9	*6
733.82	19	11	*9	*	*6	*6	*7	*	*	*8	*
736	*8	*	*	*	*	*	*	*	*	*	*
737	16	*	12	9	*	*	*	*	*5	*	*
737.3	14	*	11	*8	*	*	*	*	*	*	*
737.30	10	*	*8	*6	*	*	*	*	*	*	*
738	33	13	20	*	9	13	11	*5	*5	9	13
738.4	29	11	18	—	*7	11	11	*	*	*8	12
740–759	182	99	83	129	27	18	*8	48	29	65	40
742	*8	*	*	*	*	*	*	*	*	*	*
744	*6	*	*	*	*	*	—	*	*	*	*
745	24	13	11	19	*	*	*	*9	*	*8	*
745.4	*6	*	*	*	*	*	*	*	*	*	*
745.5	*7	*	*	*	*	*	*	*	*	*	*
746	13	*7	*6	*8	*	*	*	*	*	*6	*
747	22	9	13	12	*6	*	*	*7	*	*6	*
747.8	*8	*	*6	*	*	*	*	*	*	*	*
747.81	*7	*	*	*	*	*	*	*	*	*	*
748	*6	*	*	*5	*	*	*	*	*	*	*
749	15	9	*6	14	*	—	—	*	*	*	*7
749.2	*9	*6	*	*8	*	—	—	*	*	*	*6
750	15	13	*	15	*	*	—	*	*	*	*
750.5	13	12	*	13	—	—	—	*	*	*	*
751	12	*5	*6	10	*	*	*	*	*	*	*
753	11	*6	*5	*5	*	*	*	*	*	*	*
753.2	*6	*	*	*	*	*	*	*	*	*	*
754	*8	*	*	*6	*	*	—	*	*	*	*
756	18	*9	*9	*9	*	*	*	*	*	*7	*
756.1	*8	*	*	*	*	*	*	*	*	*	*
756.12	*7	*	*	*	*	*	*	*	*	*	*
757	*6	*	*	*	*	*	—	*	*	*6	*
757.3	*6	*	*	*	*	—	—	—	*	*5	*
757.32	*5	*	*	*	*	—	—	—	—	*5	*
759	*7	*	*	*5	*	*	*	*	*	*	*
760–779	201	120	81	201	*	*	*	23	45	65	69
765	27	15	12	27	—	—	—	*	*	12	*8
765.1	20	11	*9	20	—	—	—	*	*	9	*6
765.18	*6	*	*	*6	—	—	—	*	*	*	*
769	18	11	*6	18	—	—	—	*	*8	*	*
770	38	25	13	38	—	—	—	*	*9	11	13
770.6	*8	*7	*	*8	—	—	—	*	*	*	*
770.8	21	12	9	21	—	—	—	*	*	*6	*7
770.81	*6	*	*	*6	—	—	—	*	*	*	*
770.89	10	*	*	10	—	—	—	*	*	*	*
771	14	*8	*7	14	—	—	—	*	*	*	*7
771.8	12	*7	*	12	—	—	—	*	*	*	*6
773	*6	*	*	*6	—	—	—	*	*	*	*
774	58	36	22	58	—	—	*	*7	11	19	21
774.2	*5	*	*	*5	—	—	—	*	*	*	*
774.6	48	30	18	48	—	—	*	*	9	17	17

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
778	*6	*	*	*6	—	—	—	*	*	*	*
778.4	*6	*	*	*6	—	—	—	*	*	*	*
779	20	10	9	20	—	—	—	*	*7	*5	*6
779.3	*6	*	*	*6	—	—	—	*	*	*	*
779.8	10	*6	*	10	—	—	—	*	*	*	*
779.89	10	*5	*	10	—	—	—	*	*	*	*
780–799	198	89	109	47	62	51	38	52	37	78	30
780	60	29	30	28	15	*7	10	16	12	22	10
780.2	11	*	*6	*	*	*	*	*	*	*	*
780.3	26	11	15	14	*7	*	*	*6	*7	*9	*
780.39	22	10	12	10	*7	*	*	*5	*6	*7	*
780.6	13	*7	*6	12	*	*	*	*	*	*5	*
786	51	23	28	*	12	25	10	16	*7	21	*7
786.0	*6	*	*	*	*	*	*	*	*	*	*
786.5	42	20	22	*	12	22	*8	15	*5	17	*6
786.50	20	10	10	*	*6	11	*	*6	*	9	*
786.59	20	9	11	—	*5	10	*	*8	*	*7	*
787	*6	*	*	*	*	*	*	*	*	*	*
789	27	10	16	*8	14	*	*	*7	*	10	*5
789.0	26	10	16	*8	13	*	*	*6	*	9	*5
789.00	*9	*	*	*	*	*	*	*	*	*	*
789.03	*6	*	*5	*	*	*	*	*	*	*	*
799	39	17	22	*	12	12	13	10	*7	17	*5
799.9	38	16	21	*	12	12	12	10	*6	16	*5
800–999	2,826	1,402	1,424	224	807	704	1,091	555	670	1,007	594
800	11	*8	*	*6	*	*	*	*	*	*	*
801	33	25	*7	*6	18	*	*	*	*7	10	12
801.0	11	9	*	*	*5	*	*	*	*	*5	*
801.01	*5	*	*	*	*	*	*	*	*	*	*
801.2	13	10	*	*	*8	*	*	*	*	*	*6
802	31	22	*9	*	18	*	*	*5	*8	12	*6
802.2	11	9	*	*	*7	*	*	*	*	*6	*
802.4	*6	*	*	*	*	*	*	*	*	*	*
805	94	38	55	*	27	16	49	17	27	28	21
805.0	22	11	11	*	*8	*	*9	*	*6	*7	*
805.02	*8	*	*	*	*	*	*	*	*	*	*
805.2	22	*8	13	*	*6	*	11	*	*8	*5	*
805.4	43	17	26	*	12	*7	25	*6	12	13	11
805.6	*7	*	*5	*	*	*	*	*	*	*	*
806	*6	*	*	*	*	*	*	*	*	*	*
807	45	26	19	*	*8	11	26	*8	18	11	*8
807.0	36	21	15	*	*	9	22	*6	16	*8	*6
807.01	*8	*	*	—	*	*	*	*	*	*	*
807.02	*7	*	*	—	*	*	*	*5	*	*	*
807.04	*6	*	*	—	*	*	*	*5	*	*	*
807.2	*7	*	*	*	*	*	*	*	*	*	*
808	59	18	41	*	9	*8	40	13	15	19	12
808.0	14	*8	*6	*	*	*	*5	*	*	*	*
808.2	34	*	30	*	*	*	29	*8	10	9	*7
808.4	*7	*	*	*	*	*	*	*	*	*	*
810	*7	*	*	*	*	*	*	*	*	*	*
810.0	*7	*	*	*	*	*	*	*	*	*	*
812	78	28	50	13	10	19	35	16	19	25	17
812.0	39	10	29	*	*	11	24	*9	12	11	*7
812.00	10	*	*7	*	*	*	*7	*	*	*	*
812.01	11	*	*9	*	*	*	*7	*	*	*	*
812.09	13	*	*9	*	*	*	*8	*	*	*6	*
812.2	13	*5	*8	*	*	*	*6	*	*	*	*

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
812.21	11	*	*6	*	*	*	*	*	*	*	*
812.4	22	10	12	10	*	*	*	*	*	*8	*6
812.41	13	*6	*7	*8	*	*	*	*	*	*	*
813	60	27	33	*7	16	14	23	13	11	23	13
813.0	11	*	*8	*	*	*	*6	*	*	*	*
813.2	*6	*	*	*	*	*	*	*	*	*	*
813.23	*5	*	*	*	*	*	*	*	*	*	*
813.4	29	13	16	*	*6	*7	13	*6	*5	11	*6
813.41	*8	*	*	*	*	*	*	*	*	*	*
813.42	10	*	*6	*	*	*	*	*	*	*	*
813.44	*9	*	*	*	*	*	*	*	*	*	*
813.5	*7	*	*	*	*	*	*	*	*	*	*
816	*7	*6	*	*	*	*	*	*	*	*	*
816.1	*5	*	*	*	*	*	*	*	*	*	*
820	302	86	216	*	*8	27	266	57	76	109	60
820.0	89	25	64	*	*	10	77	13	26	33	17
820.00	*7	*	*6	—	*	*	*6	*	*	*	*
820.03	*6	*	*	*	*	*	*	*	*	*	*
820.09	71	19	52	*	*	*7	63	12	19	26	14
820.2	150	40	110	*	*	14	133	33	34	55	28
820.20	11	*	*6	*	*	*	*9	*	*	*6	*
820.21	126	30	97	*	*	9	115	27	29	43	27
820.22	13	*6	*7	*	*	*	*9	*	*	*6	*
820.8	62	20	42	*	*	*	56	11	16	21	15
821	57	29	28	13	16	*8	21	*9	16	23	10
821.0	34	19	15	10	12	*	9	*	10	14	*6
821.00	*6	*	*	*	*	*	*	*	*	*	*
821.01	28	15	13	*9	11	*	*7	*	*9	12	*
821.2	20	*8	12	*	*	*	10	*	*5	*7	*
821.20	*6	*	*	*	*	*	*	*	*	*	*
821.23	*8	*	*6	*	*	*	*5	*	*	*	*
822	12	*	*7	—	*	*	*5	*	*	*5	*
822.0	11	*	*7	—	*	*	*	*	*	*	*
823	70	42	28	*6	31	22	10	12	16	27	15
823.0	25	13	12	*	*8	11	*6	*	*5	*9	*6
823.00	18	10	*8	*	*7	*8	*	*	*	*7	*
823.02	*6	*	*	*	*	*	*	*	*	*	*
823.2	16	11	*5	*	10	*	*	*	*	*6	*
823.22	11	*7	*	*	*7	*	*	*	*	*	*
823.3	10	*6	*	*	*6	*	*	*	*	*	*
823.32	*6	*	*	*	*	*	*	*	*	*	*
823.8	11	*5	*5	*	*	*	*	*	*	*	*
823.82	*7	*	*	*	*	*	*	*	*	*	*
824	103	45	58	*	38	30	30	26	25	29	23
824.2	*9	*	*5	*	*	*	*	*	*	*	*
824.4	24	9	15	*	*8	*8	*9	*7	*5	*7	*5
824.6	29	*7	22	*	*7	10	12	*7	9	*	*8
824.8	23	14	10	*	11	*5	*	*7	*5	*6	*
824.9	*5	*	*	*	*	*	*	*	*	*	*
825	20	11	*9	*	10	*5	*	*	*5	*7	*
825.0	*8	*	*	*	*	*	*	*	*	*	*
825.2	*7	*	*	*	*	*	*	*	*	*	*
840	16	10	*6	—	*	9	*6	*	*	*7	*
840.4	12	*7	*	—	*	*7	*	*	*	*5	*
844	12	*8	*	*	*	*	*	*	*	*	*
847	11	*6	*	*	*	*	*	*	*	*	*
847.2	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
850	47	29	19	*6	21	9	12	11	12	15	*9
850.0	10	*	*7	*	*	*	*	*	*	*	*
850.1	16	11	*5	*	*7	*	*	*	*	*5	*
850.11	16	11	*5	*	*7	*	*	*	*	*5	*
850.5	17	12	*5	*	*9	*	*	*	*	*6	*
851	25	17	*9	*	9	*6	10	*	*	10	*8
851.8	18	12	*6	*	*7	*	*7	*	*	*8	*6
851.86	*5	*	*	*	*	*	*	*	*	*	*
852	57	32	25	*	*7	11	36	9	14	20	13
852.0	18	10	*8	*	*	*5	9	*	*	*5	*5
852.01	*5	*	*	*	*	*	*	*	*	*	*
852.2	38	21	17	*	*	*6	26	*6	10	14	*8
852.20	10	*7	*	*	*	*	*7	*	*	*	*
852.21	14	*8	*6	*	*	*	11	*	*	*6	*
852.22	*7	*	*	*	*	*	*	*	*	*	*
853	11	*	*6	*	*	*	*6	*	*	*	*
853.0	11	*	*6	*	*	*	*6	*	*	*	*
860	28	20	*9	*	13	10	*6	*5	*7	*9	*8
860.0	14	9	*	*	*	*	*	*	*	*	*
860.4	*7	*	*	*	*	*	*	*	*	*	*
861	*8	*6	*	*	*	*	*	*	*	*	*
861.2	*6	*	*	*	*	*	*	*	*	*	*
861.21	*6	*	*	*	*	*	*	*	*	*	*
863	10	*8	*	*	*8	*	*	*	*	*	*
864	*9	*7	*	*	*5	*	*	*	*	*	*
864.0	*6	*	*	*	*	*	*	*	*	*	*
865	19	14	*5	*	*8	*	*	*	*	*5	*7
865.0	18	13	*5	*	*8	*	*	*	*	*	*7
871	*6	*	*	*	*	*	*	*	*	*	*
873	24	17	*8	*	11	*5	*	*6	*	*8	*6
873.0	*6	*	*	*	*	*	*	*	*	*	*
873.4	10	*6	*	*	*6	*	*	*	*	*	*
879	*5	*	*	*	*	*	*	*	*	*	*
881	10	*7	*	*	*8	*	*	*	*	*	*
882	*7	*5	*	*	*6	*	*	*	*	*	*
883	*8	*8	*	*	*6	*	*	*	*	*	*
886	*6	*6	*	*	*	*	*	*	*	*	*
886.0	*6	*5	*	*	*	*	*	*	*	*	*
891	*8	*	*	*	*	*	*	*	*	*	*
891.1	*5	*	*	*	*	*	*	*	*	*	*
892	*7	*	*	*	*	*	*	*	*	*	*
920	9	*	*7	*	*	*	*6	*	*	*	*
922	17	*8	*8	*	*	*	*9	*	*7	*	*
922.1	9	*	*	*	*	*	*5	*	*	*	*
924	23	*7	16	—	*	*	18	*	*6	*7	*
924.0	13	*	10	—	*	*	11	*	*	*	*
924.01	11	*	*8	—	*	*	10	*	*	*	*
924.1	*6	*	*	—	*	*	*	*	*	*	*
935	*7	*	*	*	*	*	*	*	*	*	*
935.1	*6	*	*	*	*	*	*	*	*	*	*
941	12	9	*	*	*	*	*	*	*	*8	*
941.2	9	*7	*	*	*	*	*	*	*	*5	*
941.29	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
942	13	*8	*	*7	*	*	*	*	*	*9	—
942.2	9	*6	*	*6	*	*	*	*	*	*6	—
943	11	*8	*	*	*6	*	*	*	*	*6	*
943.2	*7	*6	*	*	*	*	*	*	*	*	*
944	*7	*5	*	*	*	*	*	*	*	*	*
945	22	17	*	*6	*7	*	*	*	*	16	*
945.2	*8	*6	*	*	*	*	*	*	*	*7	*
945.3	12	9	*	*	*	*	*	*	*	*8	*
948	*8	*5	*	*6	*	*	*	*	—	*6	—
949	*7	*6	*	*7	*	—	*	—	—	*7	—
949.0	*7	*6	*	*7	*	—	—	—	—	*7	—
958	10	*6	*	*	*	*	*	*	*	*	*
958.3	*5	*	*	*	*	*	*	*	*	*	*
959	36	18	19	*5	12	*6	13	*6	12	*8	11
959.0	21	9	11	*	*5	*	*7	*	*6	*	*8
959.01	19	*8	11	*	*	*	*7	*	*6	*	*7
959.1	*8	*	*	*	*	*	*	*	*	*	*
962	*8	*5	*	*	*	*	*	*	*	*6	*
962.3	*8	*5	*	*	*	*	*	*	*	*6	*
963	*5	*	*	*	*	*	*	*	*	*	*
963.0	*5	*	*	*	*	*	—	*	*	*	*
965	63	23	40	*	40	18	*	13	18	20	13
965.0	24	11	13	*	11	11	*	*5	*	*7	*7
965.09	11	*	*8	*	*	*6	*	*	*	*	*
965.1	*6	*	*	*	*5	*	*	*	*	*	*
965.4	25	*8	17	*	19	*	*	*6	*5	10	*
966	11	*6	*	*	*8	*	*	*	*	*	*
966.3	*9	*	*	*	*7	*	*	*	*	*	*
967	13	*	*9	—	*7	*	*	*	*	*7	*
967.8	*8	*	*6	—	*	*	*	*	*	*5	*
969	77	33	43	*	52	20	*	12	16	29	19
969.0	21	*8	13	*	15	*	*	*	*	*7	*7
969.3	9	*	*	*	*7	*	*	*	*	*	*
969.4	33	13	20	*	22	9	*	*6	*9	14	*
969.7	*7	*	*	*	*	*	*	*	*	*	*
970	18	13	*6	*	13	*5	*	*8	*	*6	*
970.8	18	12	*5	*	12	*	*	*8	*	*6	*
972	9	*	*	*	*	*	*	*	*	*	*
972.6	*5	*	*	*	*	*	*	*	*	*	*
975	*8	*	*5	*	*5	*	*	*	*	*	*
975.2	*6	*	*	*	*	*	*	*	*	*	*
977	*9	*	*5	*	*6	*	*	*	*	*	*
980	*7	*6	*	*	*	*	*	*	*	*	*
989	10	*6	*	*	*	*	*	*	*	*6	*
989.5	*8	*5	*	*	*	*	*	*	*	*5	*
995	39	14	25	*6	*9	12	12	*7	15	11	*6
995.1	11	*	*8	*	*	*	*	*	*	*	*
995.9	12	*6	*7	*	*	*6	*	*	*7	*	*
995.91	9	*	*6	*	*	*5	*	*	*6	*	*
996	487	240	247	27	80	167	211	97	95	191	103
996.0	25	17	*8	*	*	*5	17	*6	*7	*7	*6
996.01	14	*8	*5	*	*	*	10	*	*	*	*
996.04	9	*8	*	*	*	*	*	*6	*	*	*
996.1	10	*6	*	*	*	*	*	*	*	*	*
996.2	21	10	11	*7	9	*	*	*	*5	*8	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
996.4	91	36	55	*	*8	30	52	20	21	35	14
996.5	*8	*	*5	*	*	*	*	*	*	*	*
996.6	175	88	87	11	30	61	73	36	36	66	37
996.61	10	*7	*	*	*	*	*	*	*	*	*
996.62	76	36	40	*8	17	28	23	15	16	28	17
996.63	*6	*	*	*	*	*	*	*	*	*	*
996.64	19	11	*8	—	*	*	15	*	*	10	*
996.66	28	16	12	*	*	10	17	*	9	10	*
996.67	13	*6	*7	*	*	*5	*	*	*	*	*
996.68	*6	*	*	*	*	*	*	*	*	*	*
996.69	12	*5	*7	*	*	*	*	*	*	*5	*
996.7	126	64	62	*	17	50	57	22	20	50	34
996.72	14	9	*	*	*	*7	*6	*	*	*6	*
996.73	42	21	21	*	*6	20	16	*	*6	15	17
996.74	25	12	13	*	*	*9	13	*7	*	10	*
996.77	18	10	*8	—	*	*	13	*	*	*7	*5
996.78	13	*6	*8	*	*	*	*	*	*	*5	*
996.8	27	15	13	*	11	10	*	*5	*	18	*
996.81	18	10	*8	*	*9	*6	*	*	*	11	*
997	112	56	56	*	22	41	45	28	28	34	23
997.0	*7	*	*	*	*	*	*	*	*	*	*
997.1	10	*6	*	*	*	*	*5	*	*	*	*
997.2	9	*5	*	—	*	*	*	*	*	*	*
997.3	11	*	*6	*	*	*	*	*	*	*	*
997.4	41	16	25	*	14	12	13	11	11	12	*7
997.5	11	*8	*	*	*	*5	*	*	*	*	*
997.6	19	12	*7	*	*	*7	10	*5	*	*6	*
997.62	13	9	*	—	*	*5	*6	*	*	*	*
997.69	*5	*	*	*	*	*	*	*	*	*	*
998	257	113	144	10	70	93	84	57	57	85	58
998.1	64	29	36	*	18	17	27	18	16	19	11
998.11	35	19	16	*	*7	*8	18	12	*6	11	*7
998.12	25	*9	16	*	9	*7	*8	*5	*9	*7	*
998.2	*9	*	*6	*	*	*	*	*	*	*	*
998.3	20	*7	12	*	*7	*7	*	*	*	*7	*7
998.32	15	*6	*9	*	*	*6	*	*	*	*6	*
998.5	136	61	75	*5	37	54	40	30	31	47	28
998.51	*6	*	*	*	*	*	*	*	*	*	*
998.59	130	60	70	*	34	52	39	29	29	45	27
998.6	*7	*	*	*	*	*	*	*	*	*	*
998.8	20	*7	12	*	*	*9	*5	*	*	*6	*5
998.83	*6	*	*	*	*	*	*	*	*	*	*
998.89	14	*6	*8	*	*	*6	*	*	*	*	*
999	*7	*	*	*	*	*	*	*	*	*	*
V01–V83	4,933	396	4,538	76	4,158	210	489	773	1,017	1,911	1,232
V10	*7	*	*	—	*	*	*	*	*	*	*
V12	*6	*	*	*	*	*	*	*	*	*	*
V12.5	*5	*	*	*	*	*	*	*	*	*	*
V15	*8	*6	*	*	*	*	*	*	*	*	*
V15.8	*7	*5	*	*	*	*	*	*	*	*	*
V17	*5	*	*	*	*	*	*	*	*	*	*
V27	4,038	—	4,038	*6	4,025	*7	—	591	824	1,574	1,049
V27.0	3,817	—	3,817	*6	3,805	*6	—	544	771	1,500	1,002
V27.1	17	—	17	—	17	—	—	*	*	*6	*6
V27.2	60	—	60	—	59	*	—	11	*6	29	13
V27.9	142	—	142	*	141	*	—	34	43	38	27
V30	*8	*	*5	*8	—	—	—	*	*	*	*
V30.1	*8	*	*5	*8	—	—	—	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
V42	*7	*	*	*	*	*	*	*	*	*	*
V45	22	*8	13	*	*7	*8	*5	*	*7	*5	*6
V45.7	*6	*	*6	*	*	*	*	*	*	*	*
V45.8	11	*7	*	*	*	*	*	*	*	*	*
V45.89	*6	*	*	*	*	*	*	*	*	*	*
V50	*5	*	*	*	*	*	*	*	*	*	*
V53	21	13	*8	*	*	*	16	*	*5	*5	*7
V53.3	20	12	*7	*	*	*	16	*	*5	*	*6
V53.31	13	*8	*6	—	*	*	12	*	*	*	*
V53.32	*6	*	*	*	*	*	*	*	*	*	*
V54	20	*6	14	*	*	*	14	*6	*	*	*8
V54.1	*8	*	*6	*	*	*	*5	*	*	*	*
V54.8	*9	*	*5	*	*	*	*7	*	*	*	*
V54.81	*7	*	*	—	—	*	*6	*	—	*	*
V55	35	18	18	*	*7	14	11	*9	*7	11	*9
V55.2	10	*	*6	*	*	*5	*	*	*	*	*
V55.3	18	9	*9	*	*	*7	*7	*	*	*7	*
V56	*6	*	*	*	*	*	*	*	*	*	*
V56.0	*6	*	*	*	*	*	*	*	*	*	*
V57	492	197	295	*	31	90	368	92	122	178	101
V57.1	86	38	49	*	*6	12	68	18	*	43	21
V57.8	403	159	244	*	25	77	299	73	117	134	79
V57.89	403	159	244	*	25	77	299	73	117	134	79
V58	173	94	79	38	30	56	49	43	30	72	28
V58.0	*7	*	*	*	*	*	*	*	*	*	*
V58.1	140	80	59	35	26	49	30	36	20	62	23
V58.4	*6	*	*	*	*	*	*	*	*	*	*
V58.7	*5	*	*	*	*	*	*	*	*	*	*
V58.8	11	*	*7	*	*	*	*8	*	*7	*	*
V58.89	9	*	*5	*	*	*	*7	*	*7	*	*
V59	*8	*	*	—	*	*	*	*	*	*	*
V59.4	*6	*	*	—	*	*	*	*	*	*	*
V70	14	9	*5	*	10	*	*	*	*	*6	*
V70.7	13	*9	*	*	*9	*	*	*	*	*	*
V71	11	*6	*5	*	*	*	*5	*	*	*5	*
V71.8	*6	*	*	*	*	*	*	*	*	*	*
V71.89	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see "Methods" section.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
All codes	165,925	72,186	93,739	11,371	39,786	41,619	73,149	38,730	33,741	62,797	30,656
001–139	6,932	3,366	3,566	492	1,052	1,858	3,529	1,731	1,234	2,557	1,411
003	*41	*	*	*	*	*	*	*	*	*	*
003.0	*20	*	*	*	*	*	*	*	*	*	*
008	782	291	491	114	74	153	440	260	134	295	93
008.4	537	196	341	*	*32	121	368	204	88	199	*46
008.45	527	193	334	*	*29	119	363	202	87	196	*42
008.6	61	37	24	58	*	*	*	*	*10	23	*19
008.61	59	36	23	57	*	*	*	*	*10	23	*18
008.8	175	55	120	37	40	30	67	45	35	67	28
009	52	*22	31	*	*18	*	*	*	*	*	*20
009.0	34	*	*21	*	*	*	*	*	*	*	*
011	114	*92	*	*	*	*31	*	*	*	*45	*
011.9	*67	*	*	*	*	*	*	*	*	*	*
033	*38	*	*	*38	—	*	—	*	*	*	*
034	*18	*	*	*	*	*	—	*	*	*	*
034.0	*16	*	*	*	*	*	—	*	*	*	*
038	4,034	1,997	2,037	*72	333	1,032	2,597	941	777	1,473	843
038.0	297	154	143	*	*	*72	166	*63	*64	*98	*72
038.1	767	457	310	*	*77	260	407	146	125	330	165
038.10	*59	*	*	*	*	*	*	*	*	*	*
038.11	564	350	213	*	*65	206	290	*108	*85	256	116
038.19	144	*73	*71	*	*	*	*83	*	*	*	*
038.2	*57	*	*	*	*	*	*	*	*	*	*
038.4	823	366	457	*	*59	265	492	229	199	181	215
038.40	*68	*	*	—	*	*	*48	*	*	*	*
038.42	365	124	241	*	*	104	231	107	116	75	66
038.43	*80	*	*	—	*	*	*	*	*	*	*
038.49	297	164	133	*	*	*127	150	*81	*55	*62	*99
038.8	125	*	89	*	*	*	89	*	*	*	*
038.9	1,928	943	985	*	122	397	1,388	440	348	787	353
041	79	*41	*38	*	*	*	*	*	*	*40	*
042	538	305	233	*	287	240	*	191	*42	261	*45
047	116	58	59	49	50	*	*	*20	27	37	32
047.9	102	54	49	39	48	*	*	*19	25	34	*24
053	85	*31	54	*	*	*	61	*	*22	*	*25
053.1	*22	*	*	—	*	*	*	*	*	*	*
053.9	39	*	*31	*	*	*	*29	*	*	*	*
054	*66	*	*	*	*	*	*	*	*	*	*
066	*39	*	*	—	*	*	*	—	*	*	*
070	96	61	*35	*	*39	44	*	*	*	47	*24
070.5	*22	*	*	—	*	*13	*	*	*	*	*
070.54	*16	*	*	—	*	*	*	*	*	*	*
070.7	*18	*	*	—	*	*	*	*	*	*	*
075	34	*14	*	*	*26	*	—	*	*	*	*
079	150	79	71	82	*23	*20	*25	35	26	49	39
079.9	123	59	64	59	*20	*19	*25	33	26	42	22
079.99	123	59	64	59	*20	*19	*25	33	26	42	22
112	143	*42	100	*	*	*	95	*	*	56	*
112.0	*24	*	*19	*	*	*	*	*	*	*	*
112.2	*55	*	*46	—	*	*	*48	*	*	*	*
117	*75	*	*	*	*	*	*	*	*	*	*
135	41	*	*25	*	*	*20	*	*	*	*	*
140–239	9,627	4,392	5,236	232	1,292	3,461	4,641	2,366	1,859	3,536	1,866

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
141	*30	*	*	—	*	*	*	*	*	*	*
150	137	101	*	—	*	*65	*71	*	*	*	*
151	267	161	106	—	*	109	149	*	*74	106	*
151.0	96	*75	*	—	*	*	*47	*	*	*	*
151.9	*56	*	*	—	*	*	*39	*	*	*	*
153	919	449	470	*	*	242	648	208	210	350	152
153.0	*78	*	*	—	*	*	*74	*	*	*	*
153.1	98	*64	*	—	*	*	*65	*	*	*	*
153.2	*50	*	*40	—	*	*	*	*	*	*	*
153.3	169	76	93	—	*	*50	110	*	*42	*54	*43
153.4	191	87	104	—	*	*57	127	*55	*32	*80	*
153.6	181	93	88	—	*	*40	132	*	*42	*75	*
153.9	81	*34	*47	—	*	*	*56	*	*	*54	*
154	288	156	132	—	*	100	175	*65	*51	124	*48
154.0	121	71	*50	—	*	*45	*70	*	*	*56	*
154.1	151	80	*71	—	*	*49	96	*	*26	*62	*
155	108	59	*49	*	*	*43	*55	*35	*	*41	*
155.0	73	*47	*	*	*	*30	*34	*	*	*28	*
156	*85	*47	*	—	*	*	*76	*	*	*	*
157	211	79	133	*	*	89	117	*	*69	77	*35
157.0	87	*42	*	—	*	*	*57	*	*	*	*
157.9	57	*	*	—	*	*	*	*	*	*	*
161	120	*94	*	—	*	*64	*	*	*	*	*
162	1,003	539	464	*	*	332	615	252	195	434	121
162.2	*66	*	*	—	*	*	*34	*	*	*	*
162.3	368	199	169	*	*	125	226	75	76	158	*60
162.5	171	86	85	*	*	*70	97	*	*	104	*
162.8	160	*86	*74	—	*	*	108	*83	*	*	*
162.9	215	113	102	—	*	59	136	*41	61	98	*
171	*48	*26	*	*	*	*	*	*	*	*	*
173	*18	*	*	*	*	*	*	*	*	*	*
174	265	—	265	—	42	101	122	63	54	120	28
174.4	19	—	19	—	*	*	*	*	*	*	*
174.8	95	—	95	—	*	38	*48	*43	*	35	*
174.9	123	—	123	—	*25	46	52	*	33	70	*
180	79	—	79	—	*22	*50	*	*	*	*32	*
180.9	*43	—	*43	—	*	*	*	*	*	*22	*
182	122	—	122	—	*	61	51	*29	32	39	*22
182.0	122	—	122	—	*	60	50	*28	32	39	*22
183	144	—	144	*	*	86	*41	*	*	*66	*
183.0	138	—	138	*	*	83	*39	*	*	*63	*
185	278	278	—	*	*	122	152	61	50	97	71
188	261	209	53	—	*	*84	177	107	48	65	*41
188.8	80	54	*	—	—	*	73	*30	*	*	*
188.9	42	*26	*	—	*	*	*35	*	*	*	*
189	236	136	100	*	*	76	138	41	54	90	51
189.0	217	120	97	*	*	71	123	*36	52	87	*43
191	247	157	90	*	*49	67	95	*53	*32	79	84
191.1	76	*57	*	*	*	*25	*	*	*	*	*
191.2	*42	*32	*	*	*	*	*	*	*	*	*
191.8	*22	*	*	*	*	*	*	*	*	*	*
191.9	*39	*	*	*	*	*	*	*	*	*	*
193	62	*17	45	*	20	34	*	*13	*	28	*13
196	73	42	*	*	*	*	*41	*	*	*34	*
196.2	*29	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
197	735	279	456	*	*91	266	376	206	127	243	160
197.0	136	*39	97	*	*	*37	*39	*	*	*29	*
197.2	189	*63	126	—	*	63	120	*59	*46	*45	*
197.6	159	*61	99	—	*	*73	*77	*63	*	*60	*
197.7	154	80	74	*	*	58	81	*36	*	65	*32
198	680	255	425	*	*103	311	262	155	235	197	94
198.3	282	92	190	*	*	101	98	*52	*126	61	*43
198.5	289	126	163	*	*	162	116	*70	91	95	*
198.8	59	*	*38	*	*	*	*32	*	*	*	*
198.89	*54	*	*34	*	*	*	*29	*	*	*	*
199	170	61	109	*	*	51	98	*	*	110	*25
199.0	100	*40	60	*	*	*39	*41	*	*	*65	*
199.1	*70	*	*	*	*	*	*57	*	*	*	*
200	*77	*	*	*	*	*	*	*	*	*	*
200.0	*62	*	*	—	*	*	*	*	*	*	*
201	*43	*	*	*	*	*	*	*	*	*	*
202	303	172	130	*	*	70	188	*77	106	66	*53
202.8	258	140	119	*	*	*62	170	*61	*88	*58	*52
202.80	114	*80	*	—	*	*	*82	*	*	*	*
202.83	*57	*	*	—	*	*	*	*	*	*	*
203	127	79	*49	—	*	*	76	*	*	*	*
203.0	124	76	*48	—	*	*	73	*	*	*	*
203.00	123	76	*	—	*	*	73	*	*	*	*
204	172	80	*92	*59	*	*	*25	*	*	*48	*74
204.0	144	*62	*	*59	*	*	*	*	*	*	*
204.00	*91	*56	*	*	*	*	*	*	*	*	*
204.1	*27	*	*	—	—	*	*15	*	*	*	*
204.10	*26	*	*	—	—	*	*15	*	*	*	*
205	270	138	*132	*	*	*124	*72	*89	*	*85	*
205.0	250	*123	*127	*	*	*115	*65	*	*	*79	*
205.00	237	*113	*124	*	*	*110	*65	*	*	*76	*
210	*7	*	*	*	*	*	*	*	*	*	*
211	277	121	155	*	*	98	152	82	48	104	*43
211.3	193	93	100	—	*	75	106	*55	*31	73	*34
218	576	—	576	—	282	288	*	138	96	254	87
218.0	71	—	71	—	29	42	*	*17	*	32	*
218.1	152	—	152	—	82	68	*	44	*20	66	*22
218.2	47	—	47	—	*29	*18	—	*	*	*26	*
218.9	306	—	306	—	142	161	*	70	52	130	53
220	96	—	96	*	41	34	*21	*15	34	29	*18
225	117	*35	82	*	*	*39	*47	*31	*	*34	*36
225.2	70	*	54	—	*	*	*38	*	*	*	*
226	16	*	*14	—	*	*10	*	*	*	*	*
227	57	*19	39	*	*	*26	*	*	*	*24	*
227.3	*33	*	*	*	*	*	*	*	*	*	*
228	*37	*	*	*	*	*	*	*	*	*	*
228.0	*31	*	*	*	*	*	*	*	*	*	*
230	*60	*	*	—	*	*	*	*	*	*	*
233	52	*	49	—	24	*20	*	*	*	30	*
233.0	*14	—	*14	—	*	*	*	*	*	*	*
233.1	29	—	29	—	21	*	*	*	*	*19	*
235	*41	*	*28	—	*	*	*	*	*	*	*
235.2	*32	*	*	—	*	*	*	*	*	*	*
237	*27	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
238	112	66	*46	*	*	*	56	*	*	*47	*
238.7	88	*53	*35	*	*	*	47	*	*	*37	*
239	49	*28	*21	*	*	*12	*28	*	*16	*12	*
239.6	*19	*	*	*	*	*	*	*	*	*	*
240–279	6,765	2,976	3,789	518	1,257	2,052	2,938	1,756	1,439	2,504	1,065
241	26	*	23	—	*	15	*	*	*	*11	*
241.0	*8	*	*7	—	*	*	*	*	*	*	*
241.1	15	*	*13	—	*	*10	*	*	*	*	*
242	34	*	*22	*	*	*	*	*	*	*	*
242.9	*15	*	*	*	*	*	*	*	*	*	*
244	46	*	40	*	*	*15	*20	*19	*	*11	*
244.9	42	*	36	*	*	*12	*19	*17	*	*	*
250	2,774	1,394	1,380	69	613	1,069	1,023	819	483	1,058	414
250.0	463	209	255	24	93	170	176	124	93	178	70
250.00	193	91	102	*	38	82	72	47	33	77	36
250.01	41	*17	24	*15	*12	*	*	*	*10	*	*10
250.02	206	92	114	*	39	75	92	59	47	77	*23
250.03	*23	*	*	*	*	*	*	*	*	*	*
250.1	435	203	232	38	228	116	53	88	112	162	73
250.10	108	44	64	*	41	*37	*	*	*23	*33	*
250.11	169	84	85	31	101	*32	*	23	54	60	32
250.12	71	*31	*40	—	*26	*27	*	*	*	*34	*
250.13	88	44	44	*	59	*	*	*23	*24	35	*
250.2	80	*51	*30	*	*	*	*	*	*	*26	*
250.20	*41	*	*	—	*	*	*	*	*	*	*
250.22	*36	*	*	*	*	*	*	*	*	*	*
250.4	172	102	69	—	*	84	*66	*	*	80	*53
250.40	135	80	*55	—	*	*67	*61	*	*	*64	*49
250.6	398	178	221	—	96	191	111	95	75	169	59
250.60	236	102	134	—	*29	116	91	61	*46	92	*37
250.61	58	*	*44	—	*44	*	*	*	*	*	*
250.62	85	*54	*31	—	*	*60	*	*	*	*36	*
250.7	459	255	204	—	*	206	238	157	*64	181	*58
250.70	345	194	152	—	*	141	200	108	*46	146	*46
250.72	*93	*	*	—	*	*	*	*	*	*	*
250.8	704	370	334	*	127	263	313	282	107	240	76
250.80	433	221	212	*	*29	156	247	163	66	144	60
250.81	106	*51	*	*	*	*	*	*	*	*	*
250.82	146	87	59	—	*	75	54	*	*	75	*
250.9	*25	*	*	*	*	*	*	*	*	*	*
250.92	*22	*	*	*	*	*	*	*	*	*	*
251	77	*51	*26	*	*	*	*46	*	*	*45	*
251.2	57	*43	*15	*	*	*	*	*	*	*	*
252	*29	*	*	*	*	*	*	*	*	*	*
252.0	*23	*	*	*	*	*	*	*	*	*	*
253	125	*48	77	*	*	*	93	*48	*	*43	*
253.6	115	*46	69	*	*	*	90	*	*	*40	*
255	63	*20	*44	*	*	*	*26	*	*	*	*
255.4	51	*	*38	*	*	*	*23	*	*	*	*
263	87	*	*43	*	*	*	*51	*	*	*	*
263.9	69	*	*37	*	*	*	*37	*	*	*	*
272	85	39	46	*	*14	39	28	26	*18	28	*13
272.0	33	16	17	—	*	17	*	*13	*	*10	*
272.4	43	22	21	—	*	19	17	*11	*8	17	*
274	65	42	*23	—	*	*29	*32	*	*	*	*
274.0	44	*25	*	—	*	*20	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
275	85	*30	55	*	*	*	50	*	*	*32	*27
275.4	71	*21	50	*	*	*	*43	*	*	*	*24
275.42	56	*	*40	*	*	*	*36	*	*	*	*
276	2,571	1,010	1,561	319	214	564	1,474	562	615	983	411
276.0	79	*33	46	*	*	*	61	*	*	*	*
276.1	415	162	253	*	*24	81	303	90	104	134	87
276.2	41	*	*29	*	*	*	*	*	*	*	*
276.5	1,694	655	1,039	296	138	302	958	364	407	686	237
276.6	*26	*	*	—	*	*	*	*	*	*	*
276.7	123	62	61	*	*	57	58	*30	*27	28	*37
276.8	165	61	104	*	28	78	57	*38	24	78	*24
277	198	112	85	*79	107	*	*	*55	*76	45	*
277.0	133	*71	*62	*59	*73	*	—	*40	*61	*	*
277.02	109	*60	*49	*	*69	*	—	*38	*	*	*
277.1	*26	*22	*	—	*25	*	—	*	*	*16	*
278	392	115	278	*	187	191	*	100	94	109	90
278.0	363	101	263	*	176	174	*	94	92	98	79
278.00	24	*	*19	*	*16	*	*	*	*	*	*
278.01	339	95	243	*	160	168	*	89	87	92	70
278.1	*14	*	*14	—	*	*	*	*	*	*	*
280–289	2,006	713	1,293	219	560	465	763	502	335	789	380
280	362	90	272	*	60	72	226	89	72	124	77
280.0	195	51	143	*	39	26	130	40	42	61	*51
280.9	162	37	125	*	*21	44	94	48	30	61	24
282	421	179	242	72	291	*53	*	143	61	197	*
282.4	*50	*33	*	*	*	*	*	*	*	*	*
282.42	*42	*	*	*	*	*	—	*	*	*	*
282.6	365	142	223	59	258	*47	*	110	59	186	*
282.62	329	129	200	47	241	*41	*	101	48	173	*
283	*49	*	*	*	*	*	*	*	*	*	*
284	219	78	141	*	*92	50	45	*30	*31	59	*99
284.8	195	76	120	*	*91	46	43	*30	*31	52	*
285	369	117	253	*	22	115	229	91	58	155	65
285.1	45	*	*27	*	*	*	33	*	*	*	*
285.2	101	*27	74	*	*	*	59	*27	*	*52	*
285.29	47	*	*26	*	*	*	*42	*	*	*	*
285.9	215	69	146	*	*17	63	132	44	34	87	50
286	69	*28	41	*	*	*18	40	*	*12	*36	*
286.5	*22	*	*15	—	*	*	*	*	*	*	*
287	138	55	83	*17	*33	*26	61	47	*26	33	*31
287.3	64	*19	45	*	*	*	*	*	*	*	*
287.5	49	*27	*22	*	*	*	*	*	*	*	*
288	277	102	175	56	*34	93	94	50	53	118	57
288.0	225	88	137	51	*29	81	63	47	*32	94	51
288.8	44	*	*34	*	*	*	*27	*	*	*	*
289	90	*39	52	*20	*	*	*	*	*	*35	*
289.3	*20	*	*	*	*	*	*	*	*	*	*
290–319	16,652	8,434	8,218	1,210	8,696	4,763	1,982	5,739	2,809	6,942	1,162
290	215	*69	146	—	*	*	205	*53	*73	*78	*
290.2	*54	*	*	—	—	*	*	*	*	*	*
290.4	94	*35	*59	—	*	*	86	*	*	*	*
290.40	*43	*	*	—	—	—	*43	*	*	*	*
291	745	523	222	*	338	351	56	196	297	150	102
291.0	119	101	*	—	*53	60	*	*	*31	*25	*23
291.8	603	406	197	—	281	282	*40	153	261	111	77
291.81	585	397	188	—	273	272	*40	151	258	105	71

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
292	518	268	250	*	343	102	72	252	99	84	82
292.0	316	162	154	*	244	64	*	173	78	*27	*38
292.1	*33	*	*	*	*	*	*	*	*	*	*
292.8	159	85	74	*	82	*25	52	68	*	42	*37
292.81	51	*26	*25	—	*	*	43	*	*	*	*
292.84	97	54	*43	*	76	*	*	*51	*	*20	*24
293	71	*31	40	*	*	*	*36	*	*	*	*
293.0	36	*	*22	*	*	*	*27	*	*	*	*
293.8	*35	*	*	*	*	*	*	*	*	*	*
294	296	158	138	*	*	*	251	159	*45	76	*
294.8	258	125	133	—	*	*	242	153	*40	*49	*
295	3,709	2,024	1,686	*	2,007	1,416	272	1,297	494	1,638	280
295.3	1,316	773	543	*	694	522	*93	438	152	644	*82
295.30	594	328	265	*	304	248	*	126	*72	380	*
295.32	255	154	*101	*	132	*111	*	*77	*	130	*
295.34	459	285	174	—	256	158	*	232	*60	133	*
295.6	112	*79	*	—	*	*50	*	*	*	*	*
295.7	1,778	849	929	*	951	674	146	616	246	821	95
295.70	1,304	647	657	*	723	489	*89	352	200	675	77
295.72	*102	*	*	*	*60	*	*	*	*	*	*
295.74	358	137	221	*	164	150	*	252	*	*61	*
295.9	384	226	158	*	237	120	*	162	*38	118	*
295.90	238	142	96	*	138	73	*	*83	*34	110	*
295.94	*71	*	*	—	*	*	—	*	*	*	*
296	6,875	2,880	3,994	607	3,664	1,856	748	2,104	1,238	3,168	364
296.0	83	*	*	*	*56	*	*	*	*	*61	—
296.04	*55	*	*	*	*	*	*	*	*	*	—
296.2	1,316	478	837	103	640	374	199	243	220	783	*70
296.20	679	235	444	69	365	180	*66	111	122	397	*49
296.22	102	*36	*65	*	55	*	*	*	*	76	*
296.23	278	118	160	*	137	72	*	67	49	145	*
296.24	248	84	164	*	78	85	*72	*52	*	158	*
296.3	1,936	810	1,126	*39	882	680	336	717	393	697	128
296.30	500	263	237	*	234	216	*	234	125	131	*
296.32	97	*27	70	*	60	*	*	*	*19	*38	*
296.33	912	328	584	*	418	303	169	295	172	389	56
296.34	414	187	227	*	162	137	*105	168	76	134	*36
296.4	655	268	387	*	386	191	*64	259	106	259	*
296.40	232	99	132	*	122	*81	*	*	*37	131	*
296.43	*66	*	*	—	*	*	*	*	*	*	*
296.44	335	124	212	*	213	87	*	172	*	105	*
296.5	750	297	452	*	466	217	*54	210	194	299	*46
296.50	436	178	258	*	285	116	*	109	118	198	*
296.53	148	*41	107	*	71	*60	*	*55	*31	*50	*
296.54	151	*76	*75	*	100	*	*	*	*	*49	*
296.6	528	240	288	68	299	138	*	112	64	333	*
296.60	265	121	143	*	158	*72	*	*42	*31	182	*
296.63	87	*37	*50	*	*46	*	*	*	*	61	*
296.64	168	77	91	*	88	*	*	*55	*	87	*
296.7	142	*59	83	*	94	*	*	*44	*	*61	*
296.8	1,045	458	587	169	628	196	*52	204	182	601	58
296.80	860	400	459	160	525	138	*	128	144	539	*48
296.89	181	56	125	*	100	*56	*	*74	*38	60	*
296.9	415	216	199	173	212	*	*	295	*40	70	*
296.90	410	212	198	171	211	*	*	295	*39	66	*
297	*54	*	*	—	*	*	*	*	*	*	*
298	707	393	315	*	402	142	140	299	71	265	73
298.9	684	386	298	*	399	128	133	289	63	262	70
300	260	90	170	*	124	71	42	80	52	93	34
300.0	94	32	62	*	43	30	*17	28	*19	37	*11

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
300.00	56	17	39	*	23	20	*11	*17	*9	25	*
300.01	*19	*	*13	*	*12	*	*	*	*	*	*
300.1	48	*	*31	*	*24	*	*	*	*	*	*
300.11	*25	*	*	*	*	*	*	*	*	*	*
300.4	69	*	50	*	37	*17	*	*22	*	*24	*
300.9	*22	*	*	*	*	*	*	*	*	*	*
303	596	431	165	*	299	253	*42	166	105	291	35
303.0	86	62	*24	—	33	49	*	*25	27	*20	*
303.00	25	*16	*	—	*	*15	*	*	*10	*	*
303.01	57	43	*	—	*21	31	*	*21	*15	*	*
303.9	509	369	140	*	267	205	*	140	78	271	*20
303.90	211	141	70	*	96	103	*	*	*11	170	*
303.91	291	221	70	—	164	102	*	119	*67	99	*
304	561	361	200	*	415	133	*	133	*58	353	*
304.0	180	95	85	—	121	*51	*	50	*	106	*
304.00	84	*44	*40	—	57	*	*	*	*	71	*
304.01	90	*45	*45	—	63	*	*	46	*	*	*
304.2	145	106	*39	—	112	*	*	*	*	83	*
304.20	67	*51	*	—	*54	*	*	*	*	*40	*
304.21	77	*54	*	—	*57	*	—	*	*	*	—
304.7	*44	*	*	—	*40	*	*	*	*	*	*
304.8	67	*44	*	*	*45	*	—	*	*	61	—
304.80	60	*41	*	*	*39	*	—	*	*	58	—
305	255	159	96	*23	137	87	*	42	36	135	43
305.0	96	67	29	*	40	49	*	*22	17	33	*24
305.00	81	57	25	*	35	42	*	*18	14	26	*
305.01	*14	*	*	—	*	*	*	*	*	*	*
305.1	29	17	*12	—	16	*11	*	*	*9	*9	*
305.6	34	*20	*14	—	22	*	*	*	*	*20	*
305.60	27	*16	*10	—	*16	*	—	*	*	*16	*
305.9	67	*37	*31	*	40	*	*	*	*	53	*
305.90	59	*33	*26	*	*32	*	*	*	*	47	*
307	80	*	*75	*	*	*	*	*	*	*	*
309	336	177	158	*	218	37	*	150	49	116	*20
309.0	61	*37	*24	*	45	*	*	*26	*	*	*
309.2	33	*24	*	*	*25	*	*	*	*	*	*
309.28	*31	*22	*	*	*23	*	*	*	*	*	*
309.4	42	*	*18	*	*32	*	*	*	*	*	*
309.8	164	*71	93	*	85	*	*	*54	*33	*73	*
309.81	160	*69	91	*	81	*	*	*50	*33	*73	*
309.9	*28	*	*	*	*25	*	*	*	*	*	—
311	723	380	342	*90	410	189	*33	381	90	211	41
312	280	226	*54	*95	153	*	*	193	*	*73	*
312.3	233	196	*	*	127	*	—	165	*	*55	*
312.30	98	87	*	*	*52	*	—	*70	*	*	*
312.34	*133	*109	*	*	*75	*	—	*	*	*	*
313	*61	*	*	*	*	—	—	*	*	*	—
313.8	*61	*	*	*	*	—	—	*	*	*	—
313.81	*41	*	*	*	*	—	—	*	*	*	—
314	128	100	*	103	*	*	—	*	*	*54	*
314.0	113	*87	*	*93	*	*	—	*	*	*	*
314.01	113	*87	*	*93	*	*	—	*	*	*	*
320–389	2,933	1,249	1,684	366	609	563	1,396	776	553	988	616
320	*77	*	*	*	*	*	*	*	*	*	*
322	*39	*	*	*	*	*	*	*	*	*	*
322.9	*32	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
324	*80	*	*	*	*	*	*	*	*	*	*
331	853	301	552	*	*	*60	726	206	173	279	196
331.0	598	180	418	—	*	*	577	125	119	226	*128
331.3	51	32	*19	*	*	*	46	*	*	*15	*
331.4	103	*53	*49	*	*	*	*	*	*	*29	*
331.8	*93	*	*	—	—	*	*77	*	*	*	*
331.82	*93	*	*	—	—	*	*76	*	*	*	*
332	69	*33	36	—	*	*	59	*	*	*	*
332.0	68	*33	35	—	*	*	59	*	*	*	*
333	45	*20	*25	*	*	*	*28	*	*	*	*
337	*27	*	*	*	*	*	*	*	*	*	*
340	139	*26	113	*	65	46	*	*53	*	*36	*34
344	*30	*	*	*	*	*	*	*	*	*	*
345	251	130	121	92	68	62	*29	78	50	62	61
345.1	49	*24	*25	*26	*	*	*	*	*	*	*26
345.10	27	*12	*15	*	*	*	*	*	*	*	*12
345.3	77	48	*29	*36	*	*	*	*	*	*	*
345.5	*35	*	*	*	*	*	*	*	*	*	*
345.9	43	*21	*21	*	*	*	*	*	*	*	*
345.90	*33	*	*	*	*	*	*	*	*	*	*
346	194	36	158	*	121	56	*	44	47	60	42
346.8	39	*	34	*	*20	*	*	*	*	*	*
346.80	*21	*	*18	*	*	*	*	*	*	*	*
346.9	117	*21	97	*	80	30	*	*27	*33	34	*23
346.90	88	*17	71	*	59	*23	*	*21	*24	25	*18
346.91	*30	*	*25	*	*21	*	*	*	*	*	*
348	297	171	127	*17	71	64	145	78	44	124	51
348.0	*20	*	*	*	*	*	*	*	*	*	*
348.3	146	70	76	*	*	*	110	*	*	*60	*
348.30	*65	*	*	*	*	*	*	*	*	*	*
348.31	*55	*	*	—	*	*	*42	*	*	*	*
348.8	*21	*	*	*	*	*	*	*	*	*	*
349	61	*	43	*	*16	*	*	*	*	*	*23
349.0	*17	*	*	*	*	*	*	*	*	*	*
349.8	*36	*	*	*	*	*	*	*	*	*	*
349.82	*27	*	*	*	*	*	*	*	*	*	*
351	*17	*	*	*	*	*	*	*	*	*	*
351.0	*13	*	*	*	*	*	*	*	*	*	*
355	*39	*	*	*	*	*	*	*	*	*	*
357	*91	*	*	*	*	*	*	*	*	*	*
357.0	*61	*	*	*	*	*	*	*	*	*	*
358	*36	*	*	*	*	*	*	*	*	*	*
358.0	*25	*	*	*	*	*	*	*	*	*	*
368	*14	*	*	*	*	*	*	*	*	*	*
376	36	*	*18	*20	*	*	*	*	*	*	*17
376.0	33	*	*16	*19	*	*	*	*	*	*	*
376.01	30	*	*	*19	*	*	*	*	*	*	*
380	*24	*	*	*	*	*	*	*	*	*	*
380.1	*17	*	*	*	*	*	*	*	*	*	*
380.10	*16	*	*	*	*	*	*	*	*	*	*
382	34	*23	*	20	*	*	*	*	*	*	*18
382.9	21	*12	*	*14	*	*	*	*	*	*	*
386	115	26	89	*	*	24	83	33	45	23	*15
386.1	49	*14	35	*	*	*12	34	*17	*	*13	*
386.11	35	*11	*24	*	*	*	26	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
386.3	39	*	29	—	*	*	*23	*	*14	*	*
386.30	36	*	27	—	*	*	*22	*	*14	*	*
390–459	28,199	14,014	14,185	183	1,819	7,868	18,330	6,789	5,992	10,537	4,881
396	144	*53	91	*	*	*	117	*	*	*60	*
396.2	*58	*	*	—	*	*	*51	*	*	*	*
396.3	*41	*	*	*	*	*	*	*	*	*	*
398	163	53	110	—	*	*	136	*46	*30	63	*
398.9	163	53	110	—	*	*	136	*46	*30	63	*
398.91	162	53	110	—	*	*	136	*46	*30	63	*
401	678	282	397	*	82	283	311	157	150	274	97
401.0	74	25	49	*	*13	28	33	*15	*16	29	*
401.9	602	256	346	*	68	255	277	140	134	246	82
402	279	74	205	—	*	58	206	101	47	115	*16
402.0	*23	*	*	—	*	*	*	*	*	*	*
402.9	254	66	188	—	*	48	195	93	44	105	*13
402.90	*19	*	*13	—	*	*	*12	*	*	*	*
402.91	235	60	176	—	*	42	182	87	41	97	*
403	564	256	307	*	112	237	203	135	76	273	80
403.0	*63	*	*	*	*	*	*	*	*	*	*
403.01	*60	*	*	*	*	*	*	*	*	*	*
403.9	493	223	270	*	92	203	189	122	66	232	73
403.91	479	218	261	*	92	192	186	115	66	225	73
404	202	87	115	—	*	*86	99	*	*40	108	*
404.0	*54	*	*	—	*	*	*	*	*	*	*
404.9	148	71	*77	—	*	*50	91	*	*36	74	*
404.93	130	*58	*73	—	*	*	85	*	*	*67	*
410	3,777	2,255	1,521	—	158	1,054	2,565	1,010	768	1,237	761
410.0	100	*64	*36	—	*	*	*54	*	*	*63	*
410.01	99	*64	*35	—	*	*	*54	*	*	*63	*
410.1	331	186	146	—	*	120	198	82	57	110	82
410.11	313	175	137	—	*	115	185	82	57	100	74
410.2	58	*32	*	—	*	*	*	*	*	*	*
410.21	57	*32	*	—	*	*	*	*	*	*	*
410.3	*41	*30	*	—	*	*	*	*	*	*	*
410.31	*38	*26	*	—	*	*	*	*	*	*	*
410.4	333	223	110	—	*14	160	159	64	76	118	75
410.41	332	223	109	—	*14	160	158	64	76	117	75
410.5	67	*	*	—	*	*	*54	*	*	*	*
410.51	67	*	*	—	*	*	*54	*	*	*	*
410.7	2,334	1,399	935	—	83	555	1,695	670	463	719	482
410.71	2,299	1,383	916	—	83	554	1,662	650	462	706	481
410.8	75	41	*33	—	*	*25	45	*	*27	*	*
410.81	71	38	*33	—	*	*25	*42	*	*26	*	*
410.9	431	241	189	—	*	98	306	88	100	175	68
410.91	421	236	185	—	*	96	299	85	99	170	68
411	189	84	105	—	*16	76	97	54	42	62	30
411.1	176	78	99	—	*16	71	89	47	41	59	29
412	*12	*	*	—	*	*	*	*	*	*	*
413	96	41	55	—	18	35	43	25	18	33	*20
413.9	83	34	50	—	*13	27	43	25	18	27	*13
414	3,517	2,157	1,360	*	119	1,406	1,990	716	737	1,461	603
414.0	3,422	2,089	1,334	*	114	1,372	1,934	690	710	1,426	596
414.00	336	201	135	—	*	113	217	105	61	130	40
414.01	2,986	1,811	1,175	*	106	1,223	1,656	570	615	1,252	549
414.02	60	46	*14	—	*	*19	41	*	*21	*24	*
414.04	*15	*	*	—	*	*	*	*	*	*	*
414.05	*21	*	*	—	*	*	*	*	*	*	—
414.8	64	45	*18	—	*	*29	34	*19	*	*25	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
415	831	336	495	*	134	201	475	172	202	342	115
415.1	809	317	492	*	134	200	472	170	184	341	115
415.11	46	*	*20	—	*	*	*31	*	*	*	*
415.19	763	291	472	*	124	194	441	164	177	317	106
416	138	*	*115	*	*	*	*	*	*	*100	*
416.8	*47	*	*37	*	*	*	*	*	*	*	*
420	79	38	*41	*	*	*	*38	*	*16	*	*
420.9	79	38	*41	*	*	*	*38	*	*16	*	*
420.90	53	*24	*28	*	*	*	*29	*	*	*	*
421	199	128	*	*	*	*78	*60	*	*	*92	*
421.0	185	128	*	*	*	*77	*	*	*	*92	*
423	95	*48	*47	*	*	*41	*35	*	*	*42	*
423.9	59	*29	*30	*	*	*29	*	*	*	*31	*
424	662	343	318	*	60	127	456	185	153	209	115
424.0	281	125	156	*	*46	60	169	87	56	109	*29
424.1	347	194	153	*	*	61	274	85	85	96	80
424.9	*30	*	*	*	*	*	*	*	*	*	*
425	131	74	57	*	*21	41	54	*24	21	57	30
425.4	101	64	38	*	*16	35	38	*21	*15	49	*17
426	180	97	83	*	*	*20	150	58	40	47	35
426.0	110	58	52	—	*	*	98	34	*25	*29	*22
426.1	40	*20	*20	*	*	*	33	*	*	*	*
426.13	*23	*	*	*	*	*	*18	*	*	*	*
427	2,593	1,125	1,468	*20	120	525	1,927	725	563	877	428
427.0	61	24	37	*	*	*12	44	*16	*13	25	*
427.1	207	143	64	*	*	64	132	49	61	72	*24
427.3	1,613	662	951	*	55	325	1,232	462	357	547	249
427.31	1,475	574	901	*	46	274	1,156	423	336	491	226
427.32	138	88	50	*	*	51	76	39	*21	56	23
427.4	*35	*27	*	*	*	*	*	*	*	*	*
427.41	*34	*26	*	*	*	*	*	*	*	*	*
427.5	*37	*	*	*	*	*	*	*	*	*	*
427.6	*20	*	*	*	*	*	*	*	*	*	*
427.69	*13	*	*	*	*	*	*	*	*	*	*
427.8	605	243	361	*	36	94	462	183	118	195	108
427.81	247	105	142	—	*	*28	216	79	43	80	45
427.89	358	138	220	*	32	66	247	104	75	115	64
427.9	*15	*	*	*	*	*	*	*	*	*	*
428	5,525	2,434	3,091	*	181	1,230	4,098	1,329	1,294	2,129	774
428.0	5,071	2,232	2,839	*	154	1,142	3,760	1,248	1,171	1,938	714
428.2	150	81	69	—	*	*29	105	*	58	63	*
428.20	*64	*40	*	—	*	*	*46	*	*	*	*
428.23	57	*	*	—	*	*	*41	*	*	*	*
428.3	264	94	170	—	*	*54	200	63	58	107	*36
428.30	135	*39	96	—	*	*	107	*34	*31	61	*
428.31	70	*	*31	—	*	*	63	*	*	*	*
428.33	*50	*	*	—	*	*	*	*	*	*	*
429	30	*	*16	*	*	*	*	*	*	*	*
430	248	96	152	*	*60	112	*74	*74	*50	*88	*35
431	560	273	288	*	*	237	269	98	67	198	197
432	297	199	98	*	*	*123	167	*45	*58	59	135
432.1	268	182	86	*	*	*117	146	*	*49	*50	129
433	453	262	191	*	*	120	324	99	100	191	64
433.1	343	192	151	*	*	92	248	69	76	144	53
433.10	239	131	108	—	*	49	189	49	42	111	37
433.11	104	61	*43	*	*	*42	59	*	*	*	*
433.3	76	49	*27	—	—	*	65	*	*	34	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
433.30	47	32	*15	—	—	*	42	*	*	24	*
433.31	*29	*	*	—	—	*	*	*	*	*	*
434	2,052	998	1,054	*	90	434	1,526	445	453	769	384
434.0	*39	*	*	—	*	*	*	*	*	*	*
434.01	*38	*	*	—	*	*	*	*	*	*	*
434.1	235	111	124	*	*	*58	166	58	57	68	53
434.11	232	109	123	*	*	*57	164	56	56	68	52
434.9	1,778	858	920	*	78	366	1,332	378	391	686	323
434.90	*27	*	*	—	*	*	*	*	*	*	*
434.91	1,751	844	908	*	78	356	1,316	373	385	672	321
435	569	218	351	*	29	115	425	166	121	199	83
435.3	*22	*	*	—	*	*	*	*	*	*	*
435.8	32	*21	*	—	*	*	*22	*	*	*	*
435.9	511	186	325	*	24	103	383	146	114	180	71
436	48	*24	*25	—	*	*	*33	*	*	*	*
437	145	69	76	*	*	72	48	*33	37	54	*
437.2	*30	*	*	*	*	*	*	*	*	*	*
437.3	36	*	*25	*	*	*19	*	*	*	*	*
438	276	*146	130	*	*	*	224	*37	*19	*	*194
438.2	*186	*	*	*	*	*	*	*	*	*	*
438.20	*178	*	*	*	*	*	*	*	*	*	*
438.8	70	*50	*	—	*	*	*37	*	*	*	*
438.89	*56	*	*	—	*	*	*	*	*	*	*
440	845	451	394	—	*	234	603	176	240	340	89
440.1	26	*14	*12	—	*	*	19	*	*	*11	*
440.2	776	415	361	—	*	204	566	167	225	306	77
440.20	63	*38	*25	—	—	*	*54	*	*	*39	*
440.21	112	59	53	—	*	*30	81	*	31	53	*
440.22	98	*59	*39	—	*	*	78	*	*	*46	*
440.23	99	*55	*44	—	*	*	80	*	*	*51	*
440.24	371	201	170	—	*	*126	243	*100	*122	113	*36
441	486	313	173	*	*	170	290	134	80	202	71
441.0	161	*88	*	—	*	*	*49	*	*	*77	*
441.01	*76	*	*	—	*	*	*	*	*	*	*
441.2	*42	*	*	*	*	*	*	*	*	*	*
441.3	*50	*	*	—	—	*	*42	*	*	*	*
441.4	219	169	50	—	—	*52	167	*59	*44	94	*
442	52	*43	*	—	*	*	*30	*	*	*	*
443	144	63	82	*	*	*56	74	*	*	84	*
443.9	115	52	*63	*	*	*52	62	*	*	65	*
444	184	100	84	—	*	70	106	*44	*	77	*29
444.0	*32	*	*	—	—	*	*	*	*	*	*
444.2	128	65	63	—	*	*43	77	*37	*	56	*
444.22	115	60	55	—	*	*39	69	*36	*	*48	*
446	91	42	*49	*26	*	*	*	*	*	*	*27
446.1	*26	*19	*	*25	*	*	—	*	*	*	*
447	63	*30	*32	*	*	*25	*33	*	*	*	*
447.1	33	*14	*	—	*	*	*20	*	*	*	*
451	68	*37	31	*	*	*	*33	*25	*	*	*
451.1	*35	*	*	—	*	*	*	*	*	*	*
453	652	258	395	*	88	190	372	183	148	233	88
453.4	491	191	301	*	56	148	287	131	119	166	75
453.40	74	*32	42	—	*	*	51	*	*	*34	*
453.41	280	106	174	—	*34	89	156	91	65	83	*41
453.42	137	53	85	*	*	39	80	*30	*37	49	*
453.8	135	51	84	*	*	41	74	*38	*29	59	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
454	*74	*	*71	—	*	*	*72	*	*50	*	—
455	93	42	51	—	*17	*25	51	*30	*18	*24	*22
455.2	62	31	31	—	*	*17	*35	*24	*	*	*
456	*33	*22	*	—	*	*	*	*	*	*	*
458	441	201	240	*	42	91	305	97	100	193	51
458.0	212	82	130	*	*	*20	177	44	48	102	*19
458.2	65	*23	42	—	*	*	38	*	*	*30	*
458.29	53	*19	34	—	*	*	29	*	*	*23	*
458.9	149	90	59	—	*	58	80	*35	*33	57	*23
459	141	78	63	*	*	*37	86	*51	*35	*38	*
459.0	*29	*	*	—	*	*	*	*	*	*	*
459.8	91	*57	*34	*	*	*	*60	*	*	*	*
459.81	88	*56	*33	—	*	*	*58	*	*	*	*
460–519	19,468	9,179	10,289	1,937	1,613	4,391	11,526	4,296	4,285	7,129	3,758
461	*19	*	*	*	*	*	*	*	*	*	*
462	*18	*	*	*	*	*	*	*	*	*	*
463	22	*14	*	*	*11	*	*	*	*	*	*
464	71	38	33	59	*	*	*	*14	*12	*17	28
464.4	36	24	*12	35	*	*	—	*	*9	*11	*9
465	71	39	32	37	*	*12	*16	*15	*15	27	*14
465.9	67	36	31	33	*	*	*	*15	*15	23	*14
466	735	360	375	456	44	80	154	145	153	268	168
466.0	282	94	188	*23	35	78	147	77	68	96	42
466.1	452	266	187	433	*	*	*	68	86	172	126
466.11	263	157	106	259	*	*	*	29	57	88	89
466.19	190	109	81	174	*	*	*	39	28	85	37
473	59	*23	37	*	*20	*	*	*	*	*23	*
473.9	*25	*	*18	*	*	*	*	*	*	*	*
474	21	14	*8	16	*	*	*	*	*	*8	*
474.0	*7	*	*	*	*	*	*	*	*	*	*
474.1	14	*10	*	*12	*	*	—	*	*	*	*
474.10	*9	*	*	*8	*	—	—	*	*	*	*
475	39	26	*13	*	27	*	*	*	*12	*16	*
478	107	37	70	*44	*	*22	*	*	*35	*27	*
478.1	*12	*	*	*	*	*	*	*	*	*	*
478.2	*33	*	*	*	*	*	*	*	*	*	*
480	126	54	72	87	*	*	*	*	*26	40	45
480.1	46	*	32	46	—	*	*	*	*	*17	*
480.9	62	*28	*33	31	*	*	*	*	*	*20	*
481	195	84	111	*	*38	49	98	*36	60	69	*31
482	848	454	394	*	*77	178	561	130	203	339	176
482.0	*45	*	*	*	*	*	*	*	*	*	*
482.1	198	108	*91	*	*	*	122	*	*56	*72	*
482.2	*40	*	*	*	*	*	*37	*	*	*	*
482.3	*33	*	*	*	*	*	*	*	*	*	*
482.4	363	197	167	*	*	*93	231	*53	*68	146	*96
482.41	346	183	163	*	*	*83	225	*	*65	137	*94
482.8	106	62	*44	*	*	*	78	*	*	*54	*
482.83	*56	*34	*	*	*	*	*44	*	*	*31	*
482.9	62	*22	*40	*	*	*	*34	*	*	*	*
483	48	*	*31	*20	*	*	*	*	*	*	*
483.0	*38	*	*	*16	*	*	*	*	*	*	*
485	102	47	55	31	*	*	53	*	*	57	*
486	5,864	2,739	3,125	425	418	1,121	3,900	1,233	1,406	2,152	1,074

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
487	244	115	129	38	*	*25	169	39	101	72	*33
487.0	109	57	53	*	*	*	79	*	*40	*40	*
487.1	131	56	75	24	*	*15	88	*26	59	31	*
490	74	*20	54	*	*	29	34	*	*	42	*
491	2,852	1,328	1,523	*	62	696	2,089	630	657	1,065	499
491.2	2,840	1,324	1,515	*	62	695	2,082	627	656	1,061	496
491.20	66	*31	35	—	*	*	55	*	*	*42	*
491.21	2,336	1,118	1,218	*	50	575	1,711	523	543	847	423
491.22	437	176	262	—	*	111	317	99	100	171	67
492	101	61	*40	*	*	*47	51	*	*	*	*
492.8	85	50	*35	*	*	*39	45	*	*	*	*
493	1,616	533	1,083	362	261	470	524	421	308	636	252
493.0	90	47	43	63	*	*	*	*19	*	*29	37
493.00	*18	*	*	*	*	*	*	*	*	*	*
493.01	*18	*	*	*14	*	*	—	*	*	*	*
493.02	53	29	25	39	*	*	*	*13	*	*	26
493.2	632	171	461	*	35	244	350	183	115	235	99
493.20	75	*22	53	*	*	*19	52	*	*21	*	*
493.21	*46	*	*38	*	*	*	*	*	*	*	*
493.22	511	142	370	*	*28	200	282	148	90	195	78
493.9	891	314	577	296	213	213	170	217	188	370	116
493.90	154	60	94	55	26	32	41	40	*15	71	*27
493.91	131	52	79	79	*19	*	*	*27	23	69	*13
493.92	606	202	404	162	167	152	126	150	150	230	76
494	*37	*	*	*	*	*	*	*	*	*	*
496	358	135	223	*	*	161	190	73	80	157	49
507	1,482	872	610	*76	118	186	1,101	320	282	442	437
507.0	1,468	859	609	*76	118	179	1,095	320	281	437	430
510	142	*85	*	*	*	*	*60	*	*	*	*
510.9	131	*75	*	*	*	*	*	*	*	*	*
511	445	231	214	*	*48	80	311	128	103	166	47
511.0	*18	*	*	—	*	*	*	*	*	*	*
511.8	*93	*	*	*	*	*	*	*	*	*	*
511.9	298	139	159	*	*	51	229	85	76	102	*34
512	209	120	89	*	78	*45	84	42	*44	85	*38
512.1	37	*	*20	*	*	*	*20	*	*	*	*
512.8	161	94	67	*	66	*	*63	*29	*34	66	*
514	*29	*	*	—	*	*	*	*	*	*	*
515	192	114	77	*	*	59	124	*40	55	57	*40
516	139	*80	*60	*	*	*87	*48	*	*	*84	*
516.3	*36	*	*	—	—	*	*	*	*	*	*
516.8	102	*61	*41	*	*	*75	*	*	*	*69	*
518	3,037	1,397	1,640	159	239	849	1,790	773	543	1,090	631
518.0	27	*	*17	*	*	*	*	*	*	*	*
518.3	*27	*	*	*	*	*	*	*	*	*	*
518.8	2,927	1,362	1,565	150	217	825	1,735	751	523	1,044	610
518.81	2,353	1,073	1,280	110	142	656	1,445	656	383	801	513
518.82	*52	*	*	*	*	*	*	*	*	*	*
518.84	426	184	242	*	*	149	225	*67	114	179	*66
518.89	*51	*	*	*	*	*	*	*	*	*	*
519	64	*28	36	*16	*	*	*	*26	*	*	*
519.8	*17	*	*	*	*	*	*	*	*	*	*
520–579	16,217	7,251	8,966	855	2,967	5,077	7,317	3,530	3,449	6,114	3,123
522	*20	*	*	*	*	*	*	*	*	*	*
522.5	*15	*	*	*	*	*	*	*	*	*	*
524	29	*	*19	*	22	*	*	*	*	*16	*
524.0	*13	*	*	*	*12	*	—	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
527	85	*37	*	*	*	*	*64	*	*	*54	*
527.2	*68	*	*	*	*	*	*	*	*	*	*
528	53	*28	*25	*	*22	*	*	*	*	*31	*
528.3	*20	*	*	*	*	*	*	*	*	*	*
530	731	350	381	117	102	227	285	179	155	247	150
530.1	145	54	92	*	*29	53	58	*35	*28	57	*25
530.10	*30	*	*	*	*	*	*	*	*	*	*
530.11	50	*18	*31	*	*	*24	*	*	*	*	*
530.19	57	*21	*36	*	*	*20	*25	*	*	*	*
530.2	72	33	*39	—	*	*25	*34	*	*	*	*
530.20	*33	*	*	—	*	*	*	*	*	*	*
530.21	39	*17	*22	—	*	*	*26	*	*	*	*
530.3	*25	*	*	*	*	*	*	*	*	*	*
530.7	66	48	*	*	*	*24	*31	*	*	*30	*
530.8	357	171	186	106	45	101	104	98	68	110	82
530.81	296	138	158	104	40	76	76	82	59	87	67
530.85	*20	*	*	—	*	*	*	*	*	*	*
531	432	159	274	*	*33	117	279	55	100	163	115
531.0	61	*	*41	*	*	*	*48	*	*	*	*
531.00	61	*	*41	*	*	*	*48	*	*	*	*
531.4	256	103	153	*	*	50	187	*35	66	98	57
531.40	248	101	147	*	*	48	184	*35	63	93	57
531.9	49	*15	*35	—	*	*26	*	*	*	*	*
531.90	45	*	*32	—	*	*25	*	*	*	*	*
532	396	211	184	*	*39	112	243	72	79	157	88
532.0	38	*23	*	—	*	*	*	*	*	*	*
532.00	38	*23	*	—	*	*	*	*	*	*	*
532.4	140	98	42	*	*	*40	87	*	*	70	*21
532.40	139	98	41	*	*	*39	87	*	*	70	*21
532.9	*43	*	*20	—	*	*33	*	*	*	*34	*
532.90	*42	*	*19	—	*	*	*	*	*	*	*
533	75	*24	50	—	*	*	*37	*	*28	*28	*
533.4	*35	*	*	—	*	*	*27	*	*	*	*
533.40	*34	*	*	—	*	*	*26	*	*	*	*
534	*27	*	*	*	*	*	*	*	*	*	*
535	552	228	324	*16	81	162	293	159	91	216	87
535.0	99	53	46	*	*24	*22	48	29	*16	*41	*
535.00	68	32	36	*	*21	*	*32	26	*	*28	*
535.01	*30	*21	*	*	*	*	*	*	*	*	*
535.1	86	*29	57	*	*	*	63	*41	*	*29	*
535.10	52	*	*38	*	*	*	*38	*33	*	*	*
535.11	*34	*	*	*	*	*	*	*	*	*	*
535.3	*28	*25	*	—	*	*	*	*	*	*	*
535.4	133	48	86	*	*20	43	69	*31	*23	48	*31
535.40	60	*20	40	*	*	*24	*21	*	*	*22	*
535.41	74	*27	46	—	*	*18	48	*	*	*26	*
535.5	175	56	119	*	24	53	93	42	33	76	*24
535.50	95	32	63	*	*20	31	39	*27	*18	36	*14
535.51	80	*24	56	*	*	*22	54	*	*	*40	*
535.6	*31	*	*	*	*	*	*	*	*	*	*
536	259	101	158	*35	36	70	118	43	*41	121	53
536.2	54	*22	33	*	*	*	*	*	*	*28	*
536.3	55	*	*35	*	*	*	*	*	*	*	*
536.4	106	*42	64	*	*	*	76	*25	*	*	*
536.42	*38	*	*	*	*	*	*	*	*	*	*
536.8	*38	*	*21	*	*	*	*	*	*	*	*
537	168	79	89	*	*	*57	87	*41	*	55	49
537.0	*59	*35	*	*	*	*	*	*	*	*	*
537.8	79	39	*40	—	*	*	51	*	*	*	*28
537.83	43	*18	*	—	*	*	*31	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
540	936	553	383	294	345	190	107	205	177	276	278
540.0	312	191	121	111	75	73	*53	*45	63	88	117
540.1	241	155	85	98	63	*53	*	*81	*42	69	49
540.9	383	207	176	85	207	65	26	79	73	119	112
541	23	*	*13	*	*13	*	*	*	*	*11	*
543	*14	*	*	*	*	*	*	*	*	*	*
550	127	98	*28	*12	*	*16	89	36	*27	43	*20
550.1	92	72	*	*	*	*	73	*27	*22	*28	*
550.10	61	45	*	*	*	*	*47	*	*	*24	*
550.11	*27	*26	*	*	*	*	*	*	*	*	*
550.9	31	22	*	*	*	*	*13	*	*	*13	*
550.90	*20	*14	*	*	*	*	*	*	*	*	*
552	384	108	277	*	38	127	216	106	73	141	64
552.1	34	*15	*	—	*	*	*	*	*	*	*
552.2	250	66	185	—	*26	90	134	70	*30	112	*38
552.20	57	*	*45	—	*	*	*	*	*	*	*
552.21	161	44	118	—	*	59	91	*41	*22	71	*
552.29	*32	*	*	—	*	*	*	*	*	*	*
552.8	*40	*	*	*	*	*	*	*	*	*	*
553	273	82	191	*	42	106	121	59	55	121	38
553.2	200	66	134	*	32	89	78	44	50	80	26
553.20	31	*	*23	*	*	*	*	*	*	*18	*
553.21	157	52	105	*	*24	75	57	37	42	56	22
553.3	62	*	51	*	*	*14	*40	*	*	36	*
555	379	131	248	*	254	75	*41	84	89	156	50
555.0	73	*36	*37	*	48	*	*	*	*	*	*
555.1	138	*	*107	*	*119	*	*	*	*	*	*
555.2	*51	*	*	*	*	*	*	*	*	*	*
555.9	116	45	72	*	60	*	*	*	*21	47	*
556	172	94	78	*	80	*30	*50	*47	*25	*64	*36
556.6	*62	*39	*	*	*	*	*	*	*	*	*
556.9	55	*29	*	*	*27	*	*	*	*	*	*
557	347	79	268	*	*	76	250	66	60	148	73
557.0	162	*39	123	*	*	*	119	*44	*	*58	*
557.9	155	*	129	—	*	*34	115	*	*28	78	*35
558	553	190	363	76	107	161	209	137	111	230	74
558.9	534	186	348	75	103	154	202	135	104	224	71
560	1,956	864	1,092	50	242	529	1,135	382	503	711	359
560.0	*30	*	*	*	*	*	*	*	*	*	*
560.1	341	233	108	*	*69	80	188	87	85	126	*44
560.2	108	*51	*57	*	*	*	*62	*	*	*	*
560.3	153	59	93	*	*	*19	111	*	*	54	*55
560.39	144	59	85	*	*	*16	108	*	*	50	*54
560.8	864	322	542	*	70	269	496	159	226	336	143
560.81	663	250	413	*	*60	187	391	126	151	270	116
560.89	201	*72	129	*	*	82	105	*	*74	*67	*
560.9	461	175	286	*	59	129	270	97	127	141	95
562	1,614	647	967	—	140	529	945	455	350	571	238
562.1	1,570	623	947	—	140	515	915	449	344	541	236
562.10	149	61	87	—	*	64	74	*49	*	*36	*43
562.11	1,000	369	631	—	126	378	496	319	240	309	132
562.12	331	149	182	—	*	56	273	61	50	162	58
562.13	90	*44	46	—	*	*	72	*	*33	*	*
564	235	69	167	*15	45	72	103	53	55	88	39
564.0	113	37	76	*12	*17	27	58	*21	23	46	*23
564.00	86	27	59	*9	*10	*20	47	*13	*19	36	*18
564.09	26	*	*17	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
564.1	49	*	43	*	*20	*	*	*	*	*21	*
564.8	*50	*	*	*	*	*	*	*	*	*	*
564.89	*50	*	*	*	*	*	*	*	*	*	*
566	78	51	*27	*	*18	*39	*	*	*	33	*
567	127	57	70	*	*	*59	*39	*	*46	*41	*
567.2	98	*52	*46	*	*	*55	*	*	*	*	*
568	46	*	43	*	*16	*	*	*	*	*	*
568.0	*31	*	*29	*	*	*	*	*	*	*	*
569	722	236	486	*	81	217	404	156	145	295	126
569.1	*42	*	*37	*	*	*	*	*	*	*	*
569.3	104	56	48	*	*	*	78	*	*	50	*
569.4	*42	*26	*	—	*	*	*35	*	*	*	*
569.6	181	*51	130	*	*	*50	*85	*	*	*79	*
569.69	108	*32	*76	*	*	*	*70	*	*	*	*
569.8	322	89	233	*	*	120	165	*75	80	127	*40
569.81	*94	*	*	*	*	*	*	*	*	*	*
569.82	*57	*	*48	*	*	*	*	*	*	*	*
569.83	*87	*	*	*	*	*	*	*	*	*	*
569.85	69	*32	*37	—	*	*	60	*	*	*	*
570	58	*	*26	*	*	*	*	*	*	*21	*
571	673	395	278	*	94	441	137	110	83	265	216
571.1	62	*34	*27	—	*28	*30	*	*	*	*	*
571.2	292	211	81	—	*49	209	*34	*52	*33	119	88
571.5	248	128	120	*	*	154	82	*	*30	89	94
572	279	181	97	*	*47	154	76	55	*43	108	72
572.2	222	139	83	—	*	122	66	*28	*37	96	61
573	49	*	*	*	*	*	*	*	*	*	*
573.3	*30	*	*	*	*	*	*	*	*	*	*
574	1,383	586	796	*	326	459	596	330	248	480	324
574.0	469	183	285	*	121	134	214	127	93	151	97
574.00	441	170	271	*	113	131	197	122	90	144	85
574.01	*28	*	*	—	*	*	*	*	*	*	*
574.1	318	122	197	*	85	91	142	69	66	135	48
574.10	307	118	189	*	80	90	136	69	65	128	45
574.2	275	148	127	*	29	166	80	*70	27	80	*99
574.20	232	106	126	*	28	124	80	*27	27	80	*98
574.5	76	*25	51	*	*22	*	*39	*	*17	*28	*
574.50	44	*	*27	*	*	*	*	*	*	*	*
574.51	*32	*	*24	*	*	*	*	*	*	*	*
574.6	62	*	*30	*	*	*	*	*	*	*	*
574.60	*45	*	*24	*	*	*	*	*	*	*	*
574.7	58	*	39	—	*20	*	*	*	*	*	*
574.70	*33	*	*23	—	*	*	*	*	*	*	*
574.8	53	*	*	*	*	*	*	*	*	*	*
574.81	*32	*	*	—	*	*	*	*	*	*	*
575	319	149	170	*	86	102	130	62	76	128	53
575.0	110	60	51	*	*19	37	*55	*30	*30	*34	*16
575.1	168	70	98	*	54	46	67	*25	37	71	*34
575.10	*21	*	*13	*	*	*	*	*	*	*	*
575.11	94	*32	62	*	44	*18	*33	*	*22	40	*
575.12	52	*29	*	—	*	*	*	*	*	*	*
575.8	*19	*	*	*	*	*	*	*	*	*	*
576	139	58	81	*	*22	*44	69	*29	*23	56	*31
576.1	*53	*	*	*	*	*	*37	*	*	*	*
576.2	*43	*	*	*	*	*	*	*	*	*	*
576.8	32	*	*22	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
577	1,420	699	721	*	422	570	409	262	292	639	227
577.0	1,197	589	608	*	352	476	352	219	255	522	201
577.1	125	64	61	*	57	62	*	*24	*	64	*
577.2	*65	*	*	*	*	*	*	*	*	*	*
578	905	479	426	*13	54	153	684	205	330	245	126
578.0	72	36	36	*	*	*23	*34	*22	*18	*16	*
578.1	140	54	86	*	*17	*39	80	43	*25	59	*13
578.9	694	389	305	*	*27	91	570	139	287	171	97
579	*119	*	*	*	*	*	*	*	*	*	*
580–629	7,537	2,775	4,762	295	1,494	1,728	4,021	1,649	1,599	2,896	1,393
581	*21	*	*	*	*	*	*	*	*	*	*
584	1,842	968	874	*	110	473	1,254	528	366	588	360
584.5	161	*103	*58	*	*	*58	*90	*	*	*	*
584.9	1,661	850	811	*	91	409	1,155	477	336	539	308
585	66	*38	*29	*	*	*	40	*37	*	*	*
586	*25	*	*	*	—	*	*22	*	*	*	*
590	499	91	408	87	160	102	150	84	113	191	112
590.1	226	30	197	38	70	52	67	40	53	83	50
590.10	226	30	196	38	70	51	67	40	53	83	50
590.8	254	60	194	49	79	46	80	39	57	97	60
590.80	254	60	194	49	79	46	80	39	57	97	60
591	43	*	*26	*	*	*	*	*	*	*	*
592	425	181	244	*	158	154	107	82	104	185	55
592.0	149	52	97	*	57	47	42	31	39	61	*18
592.1	274	128	146	*	97	107	65	49	65	122	37
593	209	95	115	*26	36	47	100	45	39	98	*28
593.4	41	*	*24	*	*	*	*	*	*	*	*
593.7	*23	*	*20	*21	*	—	*	*	*	*	*
593.70	*17	*	*14	*16	*	—	—	*	*	*	*
593.8	*38	*	*22	*	*	*	*27	*	*	*	*
593.81	*26	*	*	—	*	*	*	*	*	*	*
593.9	82	46	*36	*	*	*30	45	*	*	48	*
595	84	*35	48	*	*	*	59	*	*25	*25	*
595.0	*22	*	*	*	*	*	*	*	*	*	*
595.9	*36	*	*	*	*	*	*	*	*	*	*
596	111	67	*44	*	*	*23	66	*	*	51	*21
596.0	*11	*11	—	—	—	*	*10	*	*	*	*
598	*11	*10	*	*	*	*	*	*	*	*	*
599	2,310	844	1,466	103	170	327	1,710	511	493	908	398
599.0	2,177	760	1,417	100	153	304	1,620	484	464	854	375
599.7	101	66	*35	*	*	*	75	*22	*24	41	*13
599.8	*10	*	*8	—	*	*	*	*	*	*	*
600	219	219	—	—	*	32	188	55	53	55	56
600.0	203	203	—	—	*	29	175	51	47	51	54
600.00	36	36	—	—	—	*	30	*	*	*	*
600.01	167	167	—	—	*	23	144	39	37	45	46
601	*22	*22	—	—	*	*	*	*	*	*	*
604	47	47	—	*	*	*	*	*	*	*	*
604.9	40	40	—	*	*	*	*	*	*	*	*
604.90	*20	*20	—	*	*	*	*	*	*	*	*
604.99	*19	*19	—	*	*	*	*	*	*	*	*
607	*21	*21	—	*	*	*	*	*	*	*	*
608	53	53	—	*	*17	*	*	*	*	*	*
608.4	*42	*42	—	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
611	66	*	65	*	32	23	*	*26	*	27	*
611.0	37	*	37	*	*19	*	*	*	*	*	*
611.1	*10	*	*9	*	*6	*	*	*	*	*	*
611.7	*9	*	*9	—	*	*	*	*	*	*	*
614	140	—	140	*	103	*29	*	*23	*26	72	*18
614.1	*21	—	*21	—	*	*	*	*	*	*17	*
614.2	*36	—	*36	—	*24	*	*	*	*	*	*
614.6	24	—	24	—	*18	*	*	*	*	*	*
614.9	29	—	29	*	25	*	*	*	*	*16	*
616	62	—	62	*	*	*	*	*	*	*	*
617	155	—	155	—	125	29	*	*19	39	67	31
617.0	89	—	89	—	71	*16	*	*	23	41	*13
617.1	31	—	31	—	*24	*	—	*	*	*	*
617.3	*20	—	*20	—	*14	*	—	*	*	*	*
618	324	—	324	*	43	143	136	44	71	141	68
618.0	56	—	56	—	*	22	26	*	*13	20	*16
618.01	30	—	30	—	*	*10	*17	*	*	*	*10
618.04	*15	—	*15	—	*	*	*	*	*	*	*
618.1	29	—	29	*	*	*17	*	*	*	*12	*
618.2	52	—	52	—	*	23	*18	*	*	*13	*17
618.3	35	—	35	—	*	*14	*20	*	*	*	*
618.4	47	—	47	—	*	29	*	*	*	*20	*
618.5	79	—	79	—	*	27	51	*	*14	50	*10
618.8	*15	—	*15	—	*	*	*	*	*	*	*
618.89	*15	—	*15	—	*	*	*	*	*	*	*
620	152	—	152	*	109	30	*	27	40	62	23
620.0	*15	—	*15	—	*11	*	—	*	*	*	*
620.1	23	—	23	*	*19	*	—	*	*	*	*
620.2	85	—	85	*	57	*19	*	*11	20	38	*15
620.8	*17	—	*17	*	*12	*	*	*	*	*	*
621	53	—	53	—	*24	*20	*	*	*	*27	*
621.3	*33	—	*33	—	*	*	*	*	*	*	*
622	*24	—	*24	—	*	*	*	*	*	*	*
622.1	*21	—	*21	—	*	*	*	*	*	*	*
625	135	—	135	*	75	38	*18	19	31	55	30
625.3	22	—	22	*	20	*	—	*	*	*	*
625.6	48	—	48	—	*11	24	*13	*	*14	*16	*13
625.9	47	—	47	*	34	*	*	*	*	22	*
626	273	—	273	*	193	79	*	26	70	132	44
626.2	212	—	212	*	150	61	—	*18	53	105	36
626.8	50	—	50	*	34	*16	*	*	*15	20	*
627	32	—	32	—	*	*12	*	*	*	*	*
627.1	27	—	27	—	*	*	*	*	*	*	*
630–677	1,530	—	1,530	*	1,525	*	—	199	282	745	303
632	32	—	32	*	32	*	—	*	*	17	*
633	67	—	67	—	67	—	—	*	*16	29	*14
633.1	52	—	52	—	52	—	—	*	*	24	*11
633.10	51	—	51	—	51	—	—	*	*	23	*10
634	32	—	32	*	32	*	—	*	*	22	*
634.9	18	—	18	*	18	*	—	*	*	*13	*
634.91	*14	—	*14	*	*14	*	—	*	*	*11	*
640	*14	—	*14	—	*14	—	—	*	*	*	*
641	58	—	58	*	58	—	—	*	*	*	*
641.1	*28	—	*28	—	*28	—	—	*	*	*	*
641.13	*28	—	*28	—	*28	—	—	*	*	*	*
641.9	*16	—	*16	—	*16	—	—	*	*	*	*
641.93	*16	—	*16	—	*16	—	—	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
642	85	—	85	—	85	—	—	*	*17	39	*20
642.3	24	—	24	—	24	—	—	*	*	*10	*
642.33	*17	—	*17	—	*17	—	—	*	*	*	*
642.4	*21	—	*21	—	*21	—	—	*	*	*	*
642.43	*19	—	*19	—	*19	—	—	*	*	*	*
643	74	—	74	*	74	—	—	*	*12	37	*15
643.0	35	—	35	—	35	—	—	*	*	*18	*
643.03	35	—	35	—	35	—	—	*	*	*18	*
643.1	27	—	27	*	27	—	—	*	*	*14	*
643.13	27	—	27	*	27	—	—	*	*	*14	*
643.2	*11	—	*11	—	*11	—	—	*	*	*	*
643.23	*11	—	*11	—	*11	—	—	*	*	*	*
644	338	—	338	*	338	*	—	36	79	148	74
644.0	321	—	321	*	321	*	—	35	78	139	69
644.03	321	—	321	*	321	*	—	35	78	139	69
644.1	16	—	16	—	16	—	—	*	*	*8	*
644.13	16	—	16	—	16	—	—	*	*	*8	*
646	241	—	241	*	241	—	—	*29	25	157	31
646.6	172	—	172	*	171	—	—	*	*14	112	20
646.63	166	—	166	*	166	—	—	*	*14	110	*17
646.8	51	—	51	—	51	—	—	*	*	31	*
646.83	51	—	51	—	51	—	—	*	*	31	*
647	*23	—	*23	—	*23	—	—	*	*	*	*
648	264	—	264	*	262	*	—	34	60	123	47
648.0	31	—	31	*	31	—	—	*	*	*19	*
648.03	31	—	31	*	31	—	—	*	*	*19	*
648.4	56	—	56	*	55	—	—	*	*	*27	*
648.43	31	—	31	*	*30	—	—	*	*	*25	*
648.9	141	—	141	—	141	*	—	22	26	62	31
648.93	130	—	130	—	130	*	—	19	*22	58	31
654	52	—	52	—	52	*	—	*	*	*27	*
654.5	*37	—	*37	—	*37	*	—	*	*	*	*
654.53	*37	—	*37	—	*37	*	—	*	*	*	*
655	*9	—	*9	—	*9	—	—	*	*	*	*
658	*40	—	*40	—	*40	—	—	*	*	*	*
659	*11	—	*11	*	*9	—	—	*	*	*	*
666	*11	—	*11	*	*11	—	—	*	*	*	*
670	*34	—	*34	*	*34	—	—	*	*	*	*
670.0	*34	—	*34	*	*34	—	—	*	*	*	*
670.04	*34	—	*34	*	*34	—	—	*	*	*	*
674	48	—	48	—	48	*	—	*	*	*	*
674.3	*25	—	*25	—	*25	*	—	*	*	*	*
674.34	*25	—	*25	—	*25	*	—	*	*	*	*
680–709	3,721	1,923	1,798	297	797	1,267	1,360	863	746	1,463	649
681	128	65	63	*	*32	49	*40	*27	39	53	*
681.0	73	35	38	*	*29	*	*	*18	*	*37	*
681.00	63	*31	*32	*	*22	*	*	*	*	*34	*
681.1	54	*30	*24	*	*	*34	*	*	*29	*	*
681.10	48	*29	*18	*	*28	*	*	*29	*	*	*
682	2,464	1,291	1,173	197	555	852	861	601	502	889	473
682.0	140	70	69	38	43	37	*22	*19	*22	55	44
682.1	60	*24	*	*	*	*	*	*	*	*	*
682.2	214	98	116	*17	55	84	57	*28	46	89	*51
682.3	218	122	95	*	87	60	59	53	44	90	*30
682.4	144	84	60	*	63	35	38	*27	31	60	*25
682.5	92	35	57	*27	*19	*28	*	*	*21	52	*
682.6	1,372	742	630	69	228	530	545	369	273	463	267
682.7	214	108	106	*	32	52	119	64	53	62	*35

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
685	*19	*	*	*	*	*	*	*	*	*	*
692	*19	*	*11	*	*	*	*	*	*	*	*
693	*34	*	*	*	*	*	*	*	*	*	*
693.0	*30	*	*	*	*	*	*	*	*	*	*
695	*37	*	*	*	*	*	*	*	*	*	*
695.1	*29	*	*	*	*	*	*	*	*	*	*
701	*43	*	*40	*	*	*	*	*	*	*	—
707	748	390	357	*	130	232	386	159	137	326	126
707.0	532	265	267	*	113	156	263	*103	98	246	*85
707.03	223	*128	95	*	*	*	127	*	*	112	*
707.04	*84	*	*	—	*	*	*	*	*	*	*
707.05	*67	*	*	—	*	*	*	*	*	*	*
707.07	*47	*	*	—	*	*	*	*	*	*	*
707.09	*61	*	*	—	*	*	*	*	*	*	*
707.1	203	119	84	*	*	72	116	*53	*37	74	*40
707.15	79	*49	*	—	*	*	*46	*	*	*	*
707.19	*37	*	*	*	*	*	*	*	*	*	*
708	*15	*	*	*	*	*	*	*	*	*	*
709	100	62	38	54	22	*	*	*	*	81	—
709.2	79	42	36	54	22	*	*	*	*	78	—
710–739	7,582	3,139	4,444	188	1,004	2,503	3,887	1,663	1,820	2,470	1,629
710	161	*35	126	*	69	*48	*35	*30	*31	65	*35
710.0	120	*	101	*	63	*31	*	*	*25	*39	*
711	193	109	84	*	*48	*40	*90	*73	*	*50	*
711.0	187	105	82	*	*45	*38	*89	*73	*	*45	*
711.06	72	*50	*	*	*	*	*	*	*	*	*
714	102	*42	60	*	*	*38	*43	*27	*	*37	*
714.0	66	*	45	—	*	*	*33	*	*	*	*
715	2,852	1,061	1,792	*	59	965	1,826	649	791	838	574
715.1	212	99	113	*	*	53	157	62	*27	67	56
715.15	61	37	*24	—	*	*18	42	*	*	*	*20
715.16	135	57	78	—	*	34	100	42	*22	36	36
715.2	*19	*	*	*	*	*	*	*	*	*	*
715.3	1,583	613	970	*	33	550	998	410	426	419	328
715.31	31	*10	*22	—	*	*	21	*	*	*13	*12
715.35	450	218	233	—	*	149	285	122	124	98	106
715.36	1,081	376	705	*	*	385	679	275	297	305	204
715.8	*31	*	*23	—	—	*	*27	*	*	*	*
715.89	*31	*	*23	—	—	*	*27	*	*	*	*
715.9	1,000	329	671	—	*	346	633	159	326	336	179
715.90	36	*	*26	—	*	*	*24	*	*	*	*
715.91	*16	*	*	—	—	*	*	*	*	*	*
715.95	259	103	157	—	*	75	170	50	97	57	56
715.96	676	208	468	—	*	250	422	97	217	255	107
716	43	*15	28	*	*	*16	*18	*	*	*	*
716.1	*17	*	*	—	*	*	*	*	*	*	*
716.9	*19	*	*16	*	*	*	*	*	*	*	*
718	27	*16	*11	*	*13	*	*	*	*	*	*
719	136	47	89	*	*21	23	83	*24	41	33	37
719.4	79	33	46	*	*11	*15	48	*15	20	*22	*23
719.41	*14	*	*	*	*	*	*	*	*	*	*
719.45	43	*	*27	*	*	*	*34	*	*	*	*
721	372	196	176	*	39	154	179	70	80	135	87
721.0	52	23	29	—	*7	31	*	*	*9	20	13
721.1	98	68	*30	—	*	51	44	*21	*	*31	*36
721.3	176	74	102	—	*20	53	103	*26	53	65	32
721.4	*35	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
722	904	407	497	*	253	401	249	168	205	356	176
722.0	87	41	46	—	36	43	*	17	17	40	*13
722.1	363	199	164	*	128	154	81	66	100	128	69
722.10	354	197	157	*	125	149	80	65	99	126	63
722.4	18	*	*12	—	*	*9	*	*	*	*	*
722.5	273	87	186	*	47	127	99	41	58	120	55
722.52	263	85	178	—	47	119	97	41	57	111	54
722.7	61	31	30	—	*20	*21	*19	*	*	31	*
722.71	33	*18	*15	—	*	*10	*	*	*	*15	*
722.73	*20	*	*	—	*	*	*	*	*	*	*
722.8	*19	*	*	—	*	*	*	*	*	*	*
722.83	*17	*	*	—	*	*	*	*	*	*	*
722.9	80	*	48	—	*	*36	*	*	*	*23	*
722.93	68	*	*44	—	*	*	*	*	*	*	*
723	106	56	50	*	*21	58	*27	*24	*24	30	29
723.0	50	35	*15	*	*	32	*	*	*	*14	*
723.1	30	*12	*18	*	*	*15	*	*	*	*	*
723.4	*20	*	*14	—	*	*	*	*	*	*	*
724	496	199	297	*	55	129	310	126	131	151	88
724.0	295	124	171	—	*	65	222	79	75	90	51
724.02	285	119	166	—	*	60	217	76	71	87	51
724.2	115	49	66	*	28	39	48	*15	35	36	*28
724.4	*23	*	*	—	*	*	*	*	*	*	*
724.5	38	*	27	*	*	*	*21	*	*	*12	*
726	93	41	51	*	*14	25	52	*14	*16	38	25
726.1	*14	*	*	—	*	*	*	*	*	*	*
726.10	*13	*	*	—	*	*	*	*	*	*	*
726.3	*20	*	*	—	*	*	*	*	*	*	*
726.33	*20	*	*	—	*	*	*	*	*	*	*
726.6	*24	*	*	*	*	*	*	*	*	*	*
727	59	*25	34	*	*	*14	30	*	*	23	*
727.0	*23	*	*	*	*	*	*	*	*	*	*
727.6	*14	*	*	—	*	*	*	*	*	*	*
728	222	134	88	*	62	60	97	59	47	77	*39
728.8	208	126	82	*	58	*59	90	*54	*44	75	*
728.88	101	67	*35	*	*32	*16	53	*41	*19	*33	*
728.89	*54	*	*	*	*	*	*	*	*	*	*
729	243	51	192	*	51	55	130	34	32	71	107
729.1	28	*	*21	*	*9	*	*	*	*	*	*
729.5	53	*19	34	*	*13	*20	*14	*	*11	*22	*
729.8	119	*19	100	*	*	*	98	*	*13	*20	*
729.89	37	*15	*22	*	*	*	*22	*	*	*17	*
730	569	377	192	*	112	210	220	92	150	207	120
730.0	101	*64	*	*	*	*	*	*	*	*	*
730.07	*40	*	*	*	*	*	*	*	*	*	*
730.1	114	*90	*	*	*	*	*	*	*	*	*
730.2	350	219	131	*	*56	138	147	*52	101	151	*46
730.26	*72	*	*	*	*	*	*	*	*	*	*
730.27	122	72	*50	*	*	*35	68	*	*37	*52	*
730.28	*108	*74	*	*	*	*	*	*	*	*	*
733	668	213	455	*	77	173	398	156	137	249	126
733.1	384	107	277	*	*	72	282	102	85	134	63
733.13	268	69	199	*	*	*43	218	67	47	110	*45
733.14	*53	*	*	*	*	*	*	*	*	*	*
733.4	113	46	67	*	*24	45	44	*	*	38	*36
733.42	86	34	52	*	*17	*37	*32	*	*	*26	*30
733.6	46	*9	38	*	*14	21	*11	*10	*	25	*
733.8	89	41	48	*	*21	*32	*35	*	*	*40	*19
733.82	77	35	*42	*	*18	*29	*29	*	*	*38	*

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
736	*40	*	*	*	*	*	*	*	*	*	*
737	98	*	77	59	*	*	*	*	*	*29	*
737.3	84	*	64	*51	*	*	*	*	*	*	*
737.30	58	*	*50	*36	*	*	*	*	*	*	*
738	129	40	89	*	28	52	48	*20	*17	35	57
738.4	119	37	82	—	*24	48	48	—	—	*31	55
740–759	1,233	643	591	964	99	128	*43	326	242	375	291
742	*43	*	*	*	*	*	*	*	*	*	*
744	*13	*	*	*	*	*	—	—	*	*	*
745	211	136	75	196	*	*	*	*40	*	*75	*
745.4	*26	*	*	*	*	*	*	*	*	*	*
745.5	*33	*	*	*	*	*	*	*	*	*	*
746	104	*55	*48	*86	*	*	*	*	*	*47	*
747	232	102	130	143	*17	*	*	*65	*	*46	*
747.8	*91	*	*75	*	*	*	*	*	*	*	*
747.81	*79	*	*	*	*	*	*	*	*	*	*
748	*87	*	*	*85	*	*	*	*	*	*	*
749	22	13	*9	21	*	—	—	*	*	*	*9
749.2	*12	*8	*	*11	*	—	—	*	*	*	*7
750	54	46	*	54	*	*	—	*	*	*	*
750.5	32	30	*	32	—	—	—	*	*	*	*
751	144	*76	*69	123	*	*	*	*	*	*	*
753	47	*25	*22	*25	*	*	*	*	*	*	*
753.2	*13	*	*	*	*	*	*	*	*	*	*
754	*23	*	*	*16	*	*	—	*	*	*	*
756	130	*47	*83	*90	*	*	*	*	*	*46	*
756.1	*35	*	*	*	*	*	*	*	*	*	*
756.12	*28	*	*	*	*	*	*	*	*	*	*
757	*10	*	*	*	*	*	*	—	*	*	*7
757.3	*7	*	*	*	*	—	—	—	*	*	*5
757.32	*5	*	*	*	*	—	—	—	—	*	*5
759	*48	*	*	*41	*	*	*	*	*	*	*
760–779	2,107	1,219	888	2,106	*	*	*	229	386	765	727
765	851	464	387	851	—	—	—	*	*	378	*294
765.1	447	266	*181	447	—	—	—	*	*	211	*153
765.18	*89	*	*	*89	—	—	—	*	*	*	*
769	316	189	*127	316	—	—	—	*	*74	*	*
770	395	264	131	395	—	—	—	*	*65	110	156
770.6	*88	*76	*	*88	—	—	—	*	*	*	*
770.8	203	124	80	203	—	—	—	*	*	*78	*71
770.81	*56	*	*	*56	—	—	—	*	*	*	*
770.89	78	*	*	78	—	—	—	*	*	*	*
771	72	*39	*33	72	—	—	—	*	*	*	*36
771.8	61	*37	*	61	—	—	—	*	*	*	*30
773	*13	*	*	*13	—	—	—	*	*	*	*
774	112	72	40	112	—	—	*	*14	17	42	39
774.2	*11	*	*	*11	—	—	—	*	*	*	*
774.6	92	61	31	92	—	—	*	*	15	37	30
778	*18	*	*	*18	—	—	—	*	*	*	*
778.4	*15	*	*	*15	—	—	—	*	*	*	*
779	140	74	65	140	—	—	—	*	*40	*46	*37
779.3	*31	*	*	*31	—	—	—	*	*	*	*
779.8	69	*26	*	69	—	—	—	*	*	*	*
779.89	67	*25	*	67	—	—	—	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
780–799	549	236	313	109	152	126	162	143	94	220	91
780	126	60	66	58	30	*14	24	34	24	50	18
780.2	21	*	*13	*	*	*	*	*	*	*	*
780.3	56	22	34	29	*16	*	*	*14	*11	*23	*
780.39	49	21	27	21	*16	*	*	*13	*11	*19	*
780.6	31	*19	*12	25	*	*	*	*	*	*12	*
786	87	37	50	*	20	38	20	24	*13	38	*12
786.0	*14	*	*	*	*	*	*	*	*	*	*
786.5	66	29	38	*	19	31	*15	22	*8	26	*11
786.50	31	15	15	*	*11	16	*	*11	*	14	*
786.59	30	12	18	—	*8	13	*	*10	*	*11	*
787	*12	*	*	*	*	*	*	*	*	*	*
789	46	15	31	*13	24	*	*	*12	*	18	*8
789.0	45	15	30	*13	23	*	*	*11	*	17	*8
789.00	*15	*	*	*	*	*	*	*	*	*	*
789.03	*12	*	*10	*	*	*	*	*	*	*	*
799	224	100	124	*	58	59	89	63	*37	94	*30
799.9	219	98	121	*	58	59	84	62	*35	92	*29
800–999	14,796	7,450	7,345	1,016	3,625	3,817	6,338	2,906	2,991	5,815	3,084
800	52	*41	*	*20	*	*	*	*	*	*	*
801	318	275	*43	*48	161	*	*	*	*41	114	142
801.0	86	79	*	*	*68	*	*	*	*	*58	*
801.01	*23	*	*	*	*	*	*	*	*	*	*
801.2	178	158	*	*	*56	*	*	*	*	*	*115
802	93	71	*22	*	58	*	*	*14	*19	39	*21
802.2	33	29	*	*	*23	*	*	*	*	*18	*
802.4	*17	*	*	*	*	*	*	*	*	*	*
805	471	189	282	*	106	83	279	103	128	145	95
805.0	129	63	65	*	*37	*	*63	*	*34	*44	*
805.02	*52	*	*	*	*	*	*	*	*	*	*
805.2	97	*43	54	*	*27	*	52	*	*35	*20	*
805.4	199	78	121	*	36	*29	133	*37	51	63	47
805.6	*42	*	*36	*	*	*	*	*	*	*	*
806	*104	*	*	*	*	*	*	*	*	*	*
807	203	129	74	*	*51	42	108	*31	63	52	*57
807.0	153	89	64	*	*	38	93	*29	57	*37	*29
807.01	*26	*	*	—	*	*	*	*	*	*	*
807.02	*31	*	*	—	*	*	*21	*	*	*	*
807.04	*28	*	*	—	*	*	*24	*	*	*	*
807.2	*22	*	*	*	*	*	*	*	*	*	*
808	364	109	255	*	68	*93	200	85	66	120	93
808.0	115	*55	*60	*	*	*	*35	*	*	*	*
808.2	158	*	140	*	*	*	138	*45	40	40	*33
808.4	*71	*	*	*	*	*	*	*	*	*	*
810	*29	*	*	*	*	*	*	*	*	*	*
810.0	*29	*	*	*	*	*	*	*	*	*	*
812	272	96	176	21	30	78	144	49	63	98	62
812.0	147	41	106	*	*	33	106	*29	40	45	*33
812.00	32	*	*23	*	*	*	*25	*	*	*	*
812.01	48	*	*40	*	*	*	*39	*	*	*	*
812.09	49	*	*29	*	*	*	*31	*	*	*22	*
812.2	41	*10	*31	*	*	*	*20	*	*	*	*
812.21	32	*	*24	*	*	*	*	*	*	*	*
812.4	66	36	30	14	*	*	*	*	*	*31	*11
812.41	25	*10	*15	*11	*	*	*	*	*	*	*
813	188	87	102	*14	51	55	68	37	31	86	35
813.0	44	*	*36	*	*	*	*16	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
813.2	*21	*	*	*	*	*	*	*	*	*	*
813.23	*18	*	*	*	*	*	*	*	*	*	*
813.4	84	35	49	*	*16	*16	43	*18	*15	36	*15
813.41	*21	*	*	*	*	*	*	*	*	*	*
813.42	30	*	*20	*	*	*	*	*	*	*	*
813.44	*28	*	*	*	*	*	*	*	*	*	*
813.5	*23	*	*	*	*	*	*	*	*	*	*
816	*19	*14	*	*	*	*	*	*	*	*	*
816.1	*13	*	*	*	*	*	*	*	*	*	*
820	1,904	586	1,317	*	*28	176	1,692	379	442	722	360
820.0	571	173	398	*	*	45	517	86	156	229	100
820.00	*37	*	*28	—	*	*	*31	*	*	*	*
820.03	*52	*	*	*	*	*	*	*	*	*	*
820.09	454	120	334	*	*	*35	416	73	122	175	83
820.2	922	252	671	*	*	98	810	222	186	354	160
820.20	80	*	*47	*	*	*	*68	*	*	*38	*
820.21	755	183	572	*	*	58	688	173	162	268	152
820.22	87	*36	*51	*	*	*	*53	*	*	*49	*
820.8	398	150	248	*	*	*	365	71	100	128	99
821	329	171	158	62	99	*46	122	*43	85	146	54
821.0	206	120	86	53	75	*	56	*	55	93	*36
821.00	*40	*	*	*	*	*	*	*	*	*	*
821.01	166	94	71	*49	70	*	*36	*	*43	79	*
821.2	100	*31	69	*	*	*	63	*	*28	*39	*
821.20	*27	*	*	*	*	*	*	*	*	*	*
821.23	*43	*	*32	*	*	*	*32	*	*	*	*
822	48	*	*34	—	*	*	*23	*	*	*24	*
822.0	39	*	*28	—	*	*	*	*	*	*	*
823	441	256	185	*23	210	131	77	65	87	225	65
823.0	127	65	62	*	*27	61	*38	*	*29	*49	*25
823.00	85	45	*39	*	*21	*40	*	*	*	*37	*
823.02	*37	*	*	*	*	*	*	*	*	*	*
823.2	59	36	*23	*	34	*	*	*	*	*	*28
823.22	47	*29	*	*	*29	*	*	*	*	*	*
823.3	85	*35	*	*	*56	*	*	*	*	*	*
823.32	*50	*	*	*	*	*	*	*	*	*	*
823.8	59	*25	*33	*	*	*	*	*	*	*	*
823.82	*41	*	*	*	*	*	*	*	*	*	*
824	387	155	232	*	140	111	128	92	75	146	73
824.2	*66	*	*52	*	*	*	*	*	*	*	*
824.4	84	22	61	*	*14	*24	*46	*28	*17	*21	*17
824.6	80	*17	63	*	*14	24	43	*19	24	*	*20
824.8	74	40	34	*	29	*24	*	*22	*17	*18	*
824.9	*35	*	*	*	*	*	*	*	*	*	*
825	70	41	*29	*	36	*22	*	*	*19	*29	*
825.0	*23	*	*	*	*	*	*	*	*	*	*
825.2	*25	*	*	*	*	*	*	*	*	*	*
840	27	14	*13	—	*	15	*12	*	*	*11	*
840.4	19	*10	*	—	*	*10	*	*	*	*7	*
844	39	*24	*	*	*	*	*	*	*	*	*
847	29	*15	*	*	*	*	*	*	*	*	*
847.2	*18	*	*	*	*	*	*	*	*	*	*
850	177	119	58	*58	46	27	45	34	22	50	*71
850.0	22	*	*16	*	*	*	*	*	*	*	*
850.1	45	28	*17	*	*18	*	*	*	*	*22	*
850.11	45	28	*17	*	*18	*	*	*	*	*22	*
850.5	94	79	*15	*	*18	*	*	*	*	*20	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
851	162	102	*60	*	54	*39	67	*	*	74	*41
851.8	120	70	*50	*	*44	*	*58	*	*	*58	*26
851.86	*28	*	*	*	*	*	*	*	*	*	*
852	412	227	185	*	*43	118	242	63	91	188	71
852.0	89	46	*43	*	*	*31	46	*	*	*30	*22
852.01	*28	*	*	*	*	*	*	*	*	*	*
852.2	318	179	140	*	*	*86	193	*51	64	155	*48
852.20	87	*66	*	*	*	*	*66	*	*	*	*
852.21	119	*66	*53	*	*	*	87	*	*	*65	*
852.22	*55	*	*	*	*	*	*	*	*	*	*
853	86	*	*43	*	*	*	*51	*	*	*	*
853.0	86	*	*43	*	*	*	*51	*	*	*	*
860	175	116	*60	*	78	56	*40	*29	*32	*83	*33
860.0	56	37	*	*	*	*	*	*	*	*	*
860.4	*72	*	*	*	*	*	*	*	*	*	*
861	*42	*37	*	*	*	*	*	*	*	*	*
861.2	*33	*	*	*	*	*	*	*	*	*	*
861.21	*33	*	*	*	*	*	*	*	*	*	*
863	132	*114	*	*	*105	*	*	*	*	*	*
864	*58	*47	*	*	*49	*	*	*	*	*	*
864.0	*30	*	*	*	*	*	*	*	*	*	*
865	101	69	*33	*	*38	*	*	*	*	*29	*37
865.0	100	68	*33	*	*38	*	*	*	*	*	*37
871	*14	*	*	*	*	*	*	*	*	*	*
873	65	38	*27	*	32	*13	*	*17	*	*24	*15
873.0	*15	*	*	*	*	*	*	*	*	*	*
873.4	21	*12	*	*	*12	*	*	*	*	*	*
879	*27	*	*	*	*	*	*	*	*	*	*
881	21	*13	*	*	*17	*	*	*	*	*	*
882	*31	*18	*	*	*26	*	*	*	*	*	*
883	*23	*21	*	*	*16	*	*	*	*	*	*
886	*15	*13	*	*	*	*	*	*	*	*	*
886.0	*13	*12	*	*	*	*	*	*	*	*	*
891	*34	*	*	*	*	*	*	*	*	*	*
891.1	*24	*	*	*	*	*	*	*	*	*	*
892	*27	*	*	*	*	*	*	*	*	*	*
920	23	*	*17	*	*	*	*16	*	*	*	*
922	45	*20	*25	*	*	*	*27	*	*23	*	*
922.1	21	*	*	*	*	*	*13	*	*	*	*
924	82	*17	65	—	*	*	68	*	*19	*24	*
924.0	48	*	39	—	*	*	42	*	*	*	*
924.01	40	*	*34	—	*	*	37	*	*	*	*
924.1	*23	*	*	—	*	*	*	*	*	*	*
935	*10	*	*	*	*	*	*	*	*	*	*
935.1	*8	*	*	*	*	*	*	*	*	*	*
941	73	42	*	*	*	*	*	*	*	*51	*
941.2	33	*30	*	*	*	*	*	*	*	*19	*
941.29	*16	*	*	*	*	*	*	*	*	*	*
942	67	*51	*	*17	*	*	*	*	*	*39	—
942.2	31	*19	*	*15	*	*	*	*	*	*16	—
943	75	*47	*	*	*44	*	*	*	*	*40	*
943.2	*23	*19	*	*	*	*	*	*	*	*	*
944	*25	*17	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
945	152	125	*	*20	*30	*	*	*	*	80	*
945.2	*20	*16	*	*	*	*	*	*	*	*17	*
945.3	120	100	*	*	*	*	*	*	*	*54	*
948	*103	*87	*	*66	*	*	*	*	—	*87	—
949	*98	*78	*	*92	*	—	*	—	—	*98	—
949.0	*98	*78	*	*92	*	—	—	—	—	*98	—
958	105	*43	*	*	*	*	*	*	*	*	*
958.3	*75	*	*	*	*	*	*	*	*	*	*
959	109	40	69	*8	29	*28	44	*16	29	*30	34
959.0	58	19	39	*	*12	*	*31	*	*20	*	*13
959.01	56	*17	39	*	*	*	*31	*	*20	*	*12
959.1	*12	*	*	*	*	*	*	*	*	*	*
962	*24	*13	*	*	*	*	*	*	*	*15	*
962.3	*24	*13	*	*	*	*	*	*	*	*15	*
963	*14	*	*	*	*	*	*	*	*	*	*
963.0	*13	*	*	*	*	*	—	*	*	*	*
965	199	93	106	*	115	64	*	54	41	66	38
965.0	94	55	40	*	42	45	*	*28	*	*27	*24
965.09	40	*	*28	*	*	*28	*	*	*	*	*
965.1	*10	*	*	*	*7	*	*	*	*	*	*
965.4	77	*27	50	*	57	*	*	*23	*11	35	*
966	54	*44	*	*	*44	*	*	*	*	*	*
966.3	*32	*	*	*	*26	*	*	*	*	*	*
967	28	*	*19	—	*13	*	*	*	*	*15	*
967.8	*18	*	*12	—	*	*	*	*	*	*9	*
969	216	100	116	*	117	82	*	38	37	78	63
969.0	59	*29	30	*	35	*	*	*	*	*23	*22
969.3	25	*	*	*	*14	*	*	*	*	*	*
969.4	89	44	45	*	40	42	*	*23	*21	37	*
969.7	*23	*	*	*	*	*	*	*	*	*	*
970	58	41	*17	*	39	*18	*	*32	*	*12	*
970.8	56	40	*16	*	38	*	*	*32	*	*12	*
972	23	*	*	*	*	*	*	*	*	*	*
972.6	*9	*	*	*	*	*	*	*	*	*	*
975	*31	*	*21	*	*18	*	*	*	*	*	*
975.2	*27	*	*	*	*	*	*	*	*	*	*
977	*19	*	*12	*	*13	*	*	*	*	*	*
980	*12	*10	*	*	*	*	*	*	*	*	*
989	16	*9	*	*	*	*	*	*	*	*10	*
989.5	*13	*8	*	*	*	*	*	*	*	*9	*
995	155	74	81	*24	*25	53	53	*25	62	51	*18
995.1	23	*	*14	*	*	*	*	*	*	*	*
995.9	81	*41	*40	*	*	*34	*	*	*39	*	*
995.91	56	*	*35	*	*	*32	*	*	*38	*	*
996	2,985	1,440	1,545	194	442	1,054	1,295	609	520	1,292	564
996.0	76	50	*26	*	*	*21	46	*18	*21	*23	*15
996.01	37	*19	*18	*	*	*	24	*	*	*	*
996.04	25	*22	*	*	*	*	*13	*	*	*	*
996.1	48	*23	*	*	*	*	*	*	*	*	*
996.2	91	36	56	*32	31	*	*	*	*31	*33	*
996.4	406	132	274	*	*39	132	233	94	87	162	64
996.5	*37	*	*23	*	*	*	*	*	*	*	*
996.6	1,559	774	785	108	227	566	657	334	276	618	331
996.61	86	*67	*	*	*	*	*	*	*	*	*
996.62	718	321	397	*58	149	270	242	141	126	279	172
996.63	*63	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
996.64	102	65	*37	—	*	*	79	*	*	50	*
996.66	286	157	129	*	*	106	172	*	81	107	*
996.67	124	*64	*60	*	*	*53	*	*	*	*	*
996.68	*57	*	*	*	*	*	*	*	*	*	*
996.69	93	*42	*52	*	*	*	*	*	*	*43	*
996.7	548	284	264	*	61	192	285	93	70	271	115
996.72	37	25	*	*	*	*16	*15	*	*	*15	*
996.73	161	83	78	*	*19	81	60	*	*19	79	44
996.74	165	92	73	*	*	*50	99	*37	*	94	*
996.77	86	38	*48	—	*	*	70	*	*	*41	*26
996.78	39	*14	*25	*	*	*	*	*	*	*14	*
996.8	210	120	89	*	67	88	*	*36	*	152	*
996.81	121	69	*52	*	*49	*35	*	*	*	83	*
997	647	328	318	*	102	242	283	161	159	199	127
997.0	*52	*	*	*	*	*	*	*	*	*	*
997.1	46	*29	*	*	*	*	*31	*	*	*	*
997.2	35	*21	*	—	*	*	*	*	*	*	*
997.3	66	*	*43	*	*	*	*	*	*	*	*
997.4	244	100	145	*	68	89	80	67	66	63	*49
997.5	46	*33	*	*	*	*23	*	*	*	*	*
997.6	139	92	*47	*	*	*51	72	*40	*	*49	*
997.62	112	79	*	—	*	*42	*58	*	*	*	*
997.69	*27	*	*	*	*	*	*	*	*	*	*
998	1,542	678	864	43	367	526	607	351	324	512	356
998.1	238	85	153	*	65	50	119	63	62	63	51
998.11	98	51	48	*	*14	*20	62	26	*18	29	*25
998.12	126	*32	93	*	46	*24	*55	*34	*39	*29	*
998.2	*75	*	*56	*	*	*	*	*	*	*	*
998.3	114	*50	65	*	*38	*43	*	*	*	*44	*37
998.32	86	*39	*47	*	*	*38	*	*	*	*39	*
998.5	940	421	519	*25	223	329	362	226	204	335	175
998.51	*30	*	*	*	*	*	*	*	*	*	*
998.59	910	416	494	*	216	314	355	220	197	324	169
998.6	*97	*	*	*	*	*	*	*	*	*	*
998.8	70	*28	42	*	*	*28	*19	*	*	*27	*16
998.83	*36	*	*	*	*	*	*	*	*	*	*
998.89	33	*13	*20	*	*	*14	*	*	*	*	*
999	*26	*	*	*	*	*	*	*	*	*	*
V01–V83	18,070	3,227	14,843	379	11,225	1,550	4,916	3,264	3,627	6,951	4,228
V10	*17	*	*	—	*	*	*	*	*	*	*
V12	*15	*	*	*	*	*	*	*	*	*	*
V12.5	*13	*	*	*	*	*	*	*	*	*	*
V15	*15	*11	*	*	*	*	*	*	*	*	*
V15.8	*14	*11	*	*	*	*	*	*	*	*	*
V17	*6	*	*	*	*	*	*	*	*	*	*
V27	10,425	—	10,425	*15	10,387	*24	—	1,674	2,066	4,107	2,578
V27.0	9,605	—	9,605	*14	9,572	*19	—	1,503	1,892	3,809	2,401
V27.1	35	—	35	—	35	—	—	*	*	*13	*10
V27.2	269	—	269	—	267	*	—	47	*21	144	57
V27.9	503	—	503	*	500	*	—	118	140	137	108
V30	*54	*	*38	*54	—	—	—	*	*	*	*
V30.1	*54	*	*38	*54	—	—	—	*	*	*	*
V42	*12	*	*	*	*	*	*	*	*	*	*
V45	52	*20	33	*	*18	*15	*17	*	*20	*13	*10
V45.7	*17	*	*16	*	*	*	*	*	*	*	*
V45.8	24	*16	*	*	*	*	*	*	*	*	*
V45.89	*11	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data from categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
V50	*10	*	*	*	*	*	*	*	*	*	*
V53	39	26	*13	*	*	*	30	*	*9	*10	*10
V53.3	36	23	*13	*	*	*	30	*	*9	*	*10
V53.31	22	*14	*8	—	*	*	20	*	*	*	*
V53.32	*14	*	*	*	*	*	*	*	*	*	*
V54	247	*54	194	*	*	*	230	*65	*	*	*160
V54.1	*154	*	*151	*	*	*	*144	*	*	*	*
V54.8	*88	*	*39	*	*	*	*85	*	*	*	*
V54.81	*62	*	*	—	—	*	*61	*	—	*	*
V55	194	96	98	*	*35	74	71	*50	*41	64	*39
V55.2	60	*	*36	*	*	*25	*	*	*	*	*
V55.3	118	65	*53	*	*	*44	*49	*	*	*40	*
V56	*28	*	*	*	*	*	*	*	*	*	*
V56.0	*17	*	*	*	*	*	*	*	*	*	*
V57	5,723	2,354	3,369	*	440	1,083	4,155	1,146	1,289	2,105	1,183
V57.1	1,060	463	597	*	*100	208	745	286	*	484	249
V57.8	4,619	1,891	2,728	*	340	860	3,379	844	1,248	1,608	919
V57.89	4,617	1,891	2,726	*	340	860	3,378	842	1,248	1,608	919
V58	965	527	438	168	198	266	333	225	148	400	192
V58.0	*52	*	*	*	*	*	*	*	*	*	*
V58.1	743	420	323	150	174	230	190	168	95	336	145
V58.4	*37	*	*	*	*	*	*	*	*	*	*
V58.7	*50	*	*	*	*	*	*	*	*	*	*
V58.8	60	*	*34	*	*	*	*46	*	*43	*	*
V58.89	56	*	*31	*	*	*	*44	*	*42	*	*
V59	*21	*	*	—	*	*	*	*	*	*	*
V59.4	*19	*	*	—	*	*	*	*	*	*	*
V70	36	23	*14	*	20	*	*	*	*	*19	*
V70.7	29	*21	*	*	*13	*	*	*	*	*	*
V71	16	*9	*8	*	*	*	*8	*	*	*8	*
V71.8	*10	*	*	*	*	*	*	*	*	*	*
V71.89	*10	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by an asterisk (*), use according to size of corresponding estimate in Table 45; see "Methods" section.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
All codes	169,359	69,384	99,974	7,242	41,680	42,790	77,647	35,502	41,303	60,850	31,703
001–139	4,331	1,955	2,376	466	901	1,182	1,782	936	937	1,617	840
003	13	*	*8	*	*	*	*	*	*	*7	*
003.0	10	*	*6	*	*	*	*	*	*	*5	*
005	*7	*	*	*	*	*	*	*	*	*	*
005.9	*6	*	*	*	*	*	*	*	*	*	*
008	439	173	266	95	47	78	218	124	92	163	60
008.4	258	101	157	*9	19	56	174	86	56	85	30
008.45	252	98	154	*7	18	55	172	84	55	84	28
008.5	*5	*	154	*	*	*	*	*	*	*	*
008.6	56	28	28	52	*	*	*	*	*6	10	28
008.61	53	27	26	51	*	*	*	*	*6	10	28
008.8	118	42	76	33	26	20	39	31	25	44	19
009	23	10	14	*	*8	*6	*8	*6	*	*7	*7
009.0	15	*5	9	*	*6	*	*	*	*	*	*
011	17	13	*	*	*5	*8	*	*	*	10	*
011.9	12	*9	*	*	*	*5	*	*	*	*8	*
011.90	*8	*7	*	*	*	*	*	*	*	*6	*
031	14	*7	*7	*	*	*	*6	*	*	10	*
031.0	*6	*	*	*	*	*	*	*	*	*	*
031.2	*7	*	*	*	*	*	*	*	*	*	*
033	*9	*	*	*8	*	*	*	*	*	*	*
034	24	12	12	12	10	*	*	*6	*6	*9	*
034.0	23	11	12	11	10	*	*	*6	*6	*8	*
038	861	428	434	24	89	213	534	178	189	324	171
038.0	52	27	25	*	*8	14	28	11	12	18	11
038.1	145	79	66	*5	21	43	76	28	28	62	27
038.10	12	*7	*5	*	*	*	*7	*	*	*5	*
038.11	103	59	44	*	15	30	55	20	18	45	20
038.19	29	13	17	*	*5	*8	14	*5	*8	12	*
038.2	12	*7	*	*	*	*	*5	*	*	*	*
038.3	*6	*	*	*	*	*	*	*	*	*	*
038.4	147	63	84	*	17	39	87	34	40	41	32
038.40	16	*8	*8	*	*	*5	*9	*	*	*6	*
038.42	72	26	46	*	*8	17	45	17	22	19	15
038.43	15	*8	*7	*	*	*	*8	*	*	*5	*
038.49	42	20	21	*	*	12	25	12	11	10	9
038.8	24	11	13	*	*	*6	15	*7	*6	*8	*
038.9	475	238	237	11	39	107	318	92	97	190	95
040	*6	*	*	*	*	*	*	*	*	*	*
041	1,317	542	775	145	260	306	606	245	311	489	272
041.0	183	76	107	15	41	44	83	37	42	67	37
041.00	25	14	11	*	*	*7	11	*	*	12	*
041.01	12	*6	*6	*	*	*	*	*	*	*	*
041.02	36	9	26	*	19	*7	*8	*6	9	13	*7
041.04	82	32	50	*	*6	20	51	18	20	29	14
041.09	25	14	11	*	*	*8	10	*6	*	*8	*6
041.1	380	212	168	52	101	110	117	66	83	147	83
041.10	19	11	*9	*	*	*6	*7	*	*	*7	*
041.11	320	179	141	43	90	91	96	55	68	126	71
041.19	41	23	18	*6	*7	13	14	*7	10	14	10
041.3	84	25	59	*	9	17	52	19	18	33	14
041.4	362	91	272	43	56	57	207	67	100	116	80
041.6	47	15	31	*	*	*6	35	*8	*9	22	*7
041.7	77	39	38	9	10	21	37	16	16	30	16
041.8	157	76	82	15	32	43	67	24	38	65	30
041.84	13	*5	*8	*	*	*	*5	*	*	*6	*
041.85	56	24	32	*6	*8	14	27	*7	16	24	*9

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
041.86	47	27	21	*	10	17	19	*8	*9	20	11
041.89	34	17	16	*5	*8	*9	12	*6	*8	12	*7
041.9	18	*	14	*	*7	*	*	*	*	*6	*
042	125	78	47	*	68	53	*	42	13	55	15
047	44	22	22	20	19	*	*	*7	12	14	11
047.8	*5	*	*	*	*	*	—	*	*	*	*
047.9	39	20	19	16	18	*	*	*7	11	13	*8
052	*5	*	*	*	*	*	*	*	*	*	*
053	66	24	42	*	*	10	52	16	17	18	16
053.1	22	*9	14	*	*	*	20	*5	*8	*	*6
053.19	18	*7	12	*	*	*	17	*	*7	*	*
053.2	*8	*	*	*	*	*	*6	*	*	*	*
053.9	31	10	21	*	*	*6	22	*9	*7	*9	*6
054	67	14	52	*5	42	10	9	9	11	30	17
054.1	25	*	23	*	21	*	*	*	*	11	*6
054.10	16	*	15	—	14	*	*	*	*	*8	*
054.9	28	*6	23	*	18	*5	*	*	*	15	*
057	*6	*	*	*	*	*	*	*	*	*	*
066	*6	*	*	—	*	*	*	*	*	*	*
070	448	268	180	*	134	259	54	102	62	168	116
070.1	*5	*	*	*	*	*	*	*	*	*	*
070.3	51	33	18	*	19	26	*6	12	*7	20	11
070.30	29	19	10	—	10	16	*	*5	*	14	*6
070.32	21	13	*8	*	*9	9	*	*7	*	*6	*
070.4	*6	*	*	—	*	*	*	*	*	*	*
070.44	*6	*	*	—	*	*	*	*	*	*	*
070.5	198	120	79	*	59	114	26	45	22	66	64
070.51	15	*8	*7	—	*	9	*	*6	*	*	*
070.54	182	111	71	—	54	104	24	39	19	62	62
070.7	180	104	76	—	49	111	20	39	31	76	35
070.70	178	103	75	—	49	108	20	39	30	74	34
075	18	9	*9	*	13	*	—	*	*7	*	*
078	25	9	16	*	15	*	*	*	*6	11	*
078.1	11	*	*9	*	*9	*	*	*	*	*5	*
078.10	*7	*	*5	*	*	*	*	*	*	*	*
078.5	*8	*	*	*	*	*	*	*	*	*	*
079	167	73	94	84	36	19	28	35	37	58	36
079.6	*8	*	*	*8	*	*	*	*	*	*	*
079.8	*8	*	*	*6	*	*	*	*	*	*	*
079.89	*8	*	*	*6	*	*	*	*	*	*	*
079.9	145	63	82	69	31	18	27	32	33	51	29
079.99	142	63	79	69	28	18	27	32	33	48	28
088	*6	*	*	*	*	*	*	*	*	*	*
088.8	*6	*	*	*	*	*	*	*	*	*	*
088.81	*6	*	*	*	*	*	*	*	*	*	*
094	*8	*	*	—	*	*	*	*	*	*	*
094.0	*6	*	*	—	*	*	*	*	*	*	*
098	*7	*	*6	*	*6	*	—	*	*	*	*
099	*5	*	*5	*	*5	*	—	*	*	*	*
110	64	34	30	*	11	18	32	18	17	20	*8
110.1	33	16	17	*	*	*7	23	11	9	11	*
110.4	14	*9	*5	*	*	*7	*	*	*5	*	*
110.5	*6	*	*	*	*	*	*	*	*	*	*
112	308	118	190	30	59	87	132	66	78	120	44
112.0	173	70	102	19	31	49	74	38	42	70	22
112.1	19	—	19	*	9	*	*	*	*	*8	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
112.2	34	12	22	*	*	*6	23	*7	*9	11	*7
112.3	14	*	9	*5	*	*	*	*	*	*	*
112.4	9	*5	*	*	*	*	*5	*	*	*	*
112.5	9	*	*	*	*	*	*	*	*	*	*
112.8	41	17	24	*	*7	18	15	10	11	14	*6
112.84	29	13	17	*	*6	12	11	9	*8	*8	*
112.89	10	*	*6	*	*	*5	*	*	*	*	*
112.9	*9	*	*5	*	*	*	*	*	*	*5	*
117	26	10	16	*	*	13	*8	*8	*6	*7	*
117.3	*7	*	*	*	*	*	*	*	*	*	*
117.5	10	*	*7	*	*	*6	*	*	*	*	*
117.9	*8	*	*	*	*	*	*	*	*	*	*
125	*7	*	*7	—	*	*	*	*	*	*6	*
125.9	*5	*	*	—	*	—	*	*	*	*	*
131	11	*	11	*	*9	*	*	*	*	*5	*
131.0	*7	*	*6	*	*5	*	*	*	*	*	*
135	60	19	41	*	13	32	14	20	14	19	*7
136	22	13	*9	*	*8	*6	*7	*	*	13	*
136.3	12	*7	*	—	*7	*	*	*	*	*7	*
136.9	9	*5	*	*	*	*	*6	*	*	*6	*
138	23	*9	14	—	*	*5	17	*	10	*7	*
140–239	4,454	1,921	2,532	112	613	1,560	2,169	992	991	1,642	828
141	12	10	*	—	*	*6	*5	*	*	*	*
146	*6	*	*	—	*	*	*	*	*	*	*
150	35	25	10	—	*	12	21	*8	*7	11	*8
150.5	*7	*	*	—	*	*	*	*	*	*	*
150.8	*6	*	*	—	*	*	*	*	*	*	*
150.9	18	13	*	—	*	*	13	*	*	*7	*
151	42	24	18	—	*	15	24	*7	13	15	*6
151.0	14	10	*	—	*	*6	*7	*	*	*6	*
151.8	*5	*	*	—	*	*	*	*	*	*	*
151.9	14	*7	*7	—	*	*	9	*	*	*6	*
152	*8	*6	*	—	*	*	*	*	*	*	*
153	181	85	96	*	*6	56	118	41	44	68	28
153.0	*9	*	*	—	*	*	*7	*	*	*	*
153.1	12	*8	*	—	*	*	*8	*	*	*	*
153.2	*8	*	*6	—	*	*	*	*	*	*	*
153.3	25	12	13	—	*	*8	15	*5	*7	*8	*5
153.4	28	12	15	—	*	*8	19	*6	*6	10	*6
153.6	23	10	13	—	*	*6	17	*	*7	*8	*5
153.8	*7	*	*	—	*	*	*	*5	*	*	*
153.9	63	30	33	—	*	22	39	14	14	31	*
154	66	39	27	—	*	24	39	14	16	26	10
154.0	27	17	9	—	*	10	15	*6	*6	11	*
154.1	35	20	15	—	*	11	22	*7	9	13	*5
155	36	22	14	*	*	17	16	10	*	15	*6
155.0	26	17	9	*	*	13	11	*9	*	10	*
155.1	*5	*	*	—	*	*	*	*	*	*	*
156	13	*8	*6	—	*	*	9	*	*	*	*
157	61	33	28	*	*	18	39	12	17	19	12
157.0	17	*8	*8	—	*	*	11	*	*	*	*6
157.8	9	*	*	—	*	*	*6	*	*	*	*
157.9	29	16	13	—	*	*8	18	*	10	10	*
158	*8	*	*5	*	*	*	*	*	*	*	*
161	18	13	*	—	*	10	*7	*	*	*6	*6
161.9	*5	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
162	361	201	161	*	*7	117	236	86	91	136	48
162.2	15	*9	*6	—	*	*5	10	*	*	*6	*
162.3	80	45	35	*	*	26	52	19	18	29	13
162.5	33	18	15	*	*	12	20	*7	*5	16	*
162.8	35	21	14	—	*	13	20	16	*	12	*
162.9	194	105	88	*	*	58	131	40	58	72	24
170	24	13	11	11	*7	*	*	*7	*	*5	*8
170.6	*8	*7	*	*6	*	*	*	*	*	*	*7
170.7	*7	*	*	*	*	*	*	*	*	*	*
171	21	14	*7	*	*	*	11	*5	*6	*6	*
171.5	*5	*	*	*	*	*	*	*	*	*	*
172	11	*6	*5	*	*	*	*6	*	*	*	*
173	19	13	*7	*	*	*	14	*	*	*8	*
174	169	—	169	—	25	71	73	32	44	69	24
174.4	10	—	10	—	*	*	*	*	*	*	*
174.8	35	—	35	—	*6	17	11	10	*7	13	*5
174.9	110	—	110	—	14	45	51	17	30	50	12
179	*8	—	*8	—	*	*	*	*	*	*	*
180	25	—	25	—	9	12	*	*7	*	9	*7
180.8	*6	—	*6	—	*	*	*	*	*	*	*
180.9	17	—	17	—	*6	*9	*	*	*	*8	*
182	46	—	46	—	*	23	20	11	12	14	10
182.0	46	—	46	—	*	23	20	11	12	14	10
183	55	—	55	*	*	24	26	12	*8	22	13
183.0	53	—	53	*	*	23	26	11	*7	21	13
184	*6	—	*6	—	*	*	*	*	*	*	*
185	180	180	—	*	*	51	127	44	38	59	40
186	*6	*6	—	*	*6	*	*	*	*	*	*
186.9	*6	*6	—	*	*5	*	*	*	*	*	*
188	70	52	18	*	*	11	57	20	14	23	13
188.2	*5	*	*	—	*	*	*	*	*	*	*
188.8	18	12	*6	*	*	*	14	*8	*	*	*
188.9	31	24	*7	*	*	*	26	*8	*6	12	*
189	67	40	27	*	*	21	40	14	15	27	11
189.0	62	36	26	*	*	19	37	12	14	25	10
191	70	43	27	15	10	22	24	13	13	23	22
191.1	16	11	*	*	*	*7	*5	*	*	*6	*
191.2	10	*8	*	*	*	*	*	*	*	*	*
191.3	*7	*	*	*	*	*	*	*	*	*	*
191.8	*9	*5	*	*	*	*	*	*	*	*	*
191.9	15	10	*	*	*	*	*7	*	*	*	*
192	*6	*	*	*	*	*	*	*	*	*	*
193	30	*6	25	*	11	15	*	*6	*6	12	*7
194	10	*6	*	*7	*	*	*	*	*	*	*
194.0	*6	*	*	*5	*	*	*	*	*	*	*
195	10	*5	*	*	*	*	*	*	*	*	*
196	179	75	104	*	19	75	84	37	37	67	38
196.0	27	14	14	*	*	15	10	*5	*	9	*8
196.1	24	11	13	*	*	10	13	*6	*	*8	*7
196.2	65	34	31	*	*	24	38	14	14	25	12
196.3	36	*	33	*	9	16	11	*7	11	13	*5
196.5	*6	*	*	—	*	*	*	*	*	*	*
196.6	10	*5	*5	*	*	*	*	*	*	*6	*
196.9	*8	*	*	—	*	*	*5	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
197	494	229	265	*	34	184	272	121	118	166	90
197.0	129	64	65	*	13	47	67	28	32	47	22
197.1	*9	*	*6	*	*	*	*5	*	*	*	*
197.2	51	20	31	—	*	18	31	13	13	17	*8
197.4	*9	*	*	—	*	*	*	*	*	*	*
197.5	12	*	10	*	*	*	*8	*	*	*	*
197.6	62	20	42	*	*	24	33	16	14	21	10
197.7	205	107	98	*	12	77	114	53	51	64	38
197.8	15	*7	*9	*	*	*5	9	*	*	*6	*
198	416	191	225	*7	30	163	215	91	104	146	74
198.1	13	*6	*6	—	*	*	*7	*	*	*	*
198.2	*8	*	*6	*	*	*	*	*	*	*	*
198.3	95	40	55	*	*8	41	44	20	27	35	13
198.4	*6	*	*	*	*	*	*	*	*	*	*
198.5	189	94	95	*	12	69	103	40	50	66	33
198.6	*5	—	*5	—	*	*	*	*	*	*	*
198.7	17	12	*5	—	*	*6	10	*	*	*7	*
198.8	77	30	47	*	*5	33	39	16	16	27	18
198.82	11	*	*8	—	*	*6	*	*	*	*	*
198.89	63	27	36	*	*	26	32	13	14	22	15
199	144	61	82	*	9	47	85	26	29	66	23
199.0	51	18	34	*	*	21	27	*7	*	29	11
199.1	92	43	49	*	*6	26	58	19	25	37	11
200	26	18	*8	*	*	*9	11	*	*	11	*6
200.0	17	10	*6	*	*	*5	10	*	*	*7	*
200.00	*7	*	*	*	*	*	*	*	*	*	*
200.2	*6	*	*	*	*	*	*	*	*	*	*
201	26	17	10	*	11	*6	*6	*7	*	*9	*6
201.9	22	13	*9	*	*8	*6	*5	*6	*	*8	*
201.90	18	11	*7	*	*6	*6	*	*	*	*6	*
202	138	70	68	*	13	39	85	30	31	51	25
202.0	*7	*	*	—	*	*	*	*	*	*	*
202.1	*6	*	*	*	*	*	*	*	*	*	*
202.8	118	58	60	*	10	33	75	25	27	45	22
202.80	79	38	41	*	*5	23	51	19	17	30	14
202.81	11	*6	*	*	*	*	*5	*	*	*	*
202.82	*5	*	*	*	*	*	*	*	*	*	*
202.83	10	*5	*5	*	*	*	*7	*	*	*	*
202.88	*8	*	*	—	*	*	*	*	*	*	*
203	67	36	31	*	*	19	47	13	18	25	11
203.0	66	35	31	*	*	18	46	13	18	24	11
203.00	64	35	30	*	*	18	45	12	18	24	10
204	121	65	56	27	11	14	69	26	27	42	26
204.0	43	25	18	27	11	*	*	*8	*6	15	14
204.00	28	17	11	15	*8	*	*	*5	*	12	*6
204.01	15	*8	*6	11	*	*	*	*	*	*	*7
204.1	75	39	37	—	*	10	65	17	21	25	12
204.10	72	36	36	—	*	9	62	16	20	24	11
205	58	34	24	*	10	21	22	13	12	21	12
205.0	45	26	20	*	9	16	16	10	9	15	11
205.00	41	23	18	*	*8	15	16	*9	*8	13	10
205.1	12	*8	*	*	*	*	*6	*	*	*6	*
205.10	11	*7	*	*	*	*	*5	*	*	*6	*
208	*7	*	*	*	*	*	*	*	*	*	*
210	*6	*	*	*	*	*	*	*	*	*	*
210.2	*5	*	*	*	*	*	*	*	*	*	*
211	190	89	101	*	16	58	116	45	41	63	41
211.1	22	9	13	*	*	*6	13	*	*	10	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
211.3	138	69	70	*	*7	43	88	33	28	45	33
211.4	16	*5	11	—	*	*	*8	*	*	*	*
212	*6	*	*	*	*	*	*	*	*	*	*
213	*7	*	*	*	*	*	*	*	*	*	*
214	20	10	10	*	*	*8	*7	*	*	11	*
214.8	*8	*	*5	*	*	*	*	*	*	*	*
216	12	*	9	*	*6	*	*	*	*	*	*
216.5	*6	*	*	*	*	*	*	*	*	*	*
218	432	—	432	—	217	201	14	83	82	191	77
218.0	45	—	45	—	17	28	*	12	*7	18	*7
218.1	109	—	109	—	53	52	*5	27	14	49	19
218.2	56	—	56	—	29	26	*	13	9	26	*8
218.9	222	—	222	—	119	96	*7	31	52	97	42
220	56	—	56	*	26	20	*9	10	15	20	11
223	*8	*	*	*	*	*	*	*	*	*	*
225	37	11	25	*	*6	11	19	10	*7	10	*9
225.2	25	*7	18	*	*	*6	16	*7	*5	*7	*6
226	12	*	11	—	*	*7	*	*	*	*	*
227	29	11	18	*	*7	12	10	*7	*8	*9	*6
227.0	*9	*	*6	*	*	*5	*	*	*	*	*
227.1	*7	*	*5	—	*	*	*	*	*	*	*
227.3	14	*7	*6	*	*6	*	*	*	*	*	*
228	24	12	12	*	*6	*8	*5	*6	*5	*8	*
228.0	23	11	12	*	*6	*8	*5	*6	*5	*7	*
228.04	*7	*	*	*	*	*	*	*	*	*	*
228.09	*7	*	*	*	*	*	*	*	*	*	*
230	*7	*	*	—	*	*	*	*	*	*	*
233	30	*	29	—	14	11	*	*	*	16	*
233.0	11	*	11	—	*	*7	*	*	*	*	*
233.1	14	—	14	—	11	*	*	*	*	*9	*
235	16	*6	11	—	*	*7	*7	*	*	*7	*
235.2	10	*	*8	—	*	*	*5	*	*	*	*
236	*5	*	*	—	*	*	*	*	*	*	*
237	20	*7	13	*	*7	*	*6	*7	*	*	*
237.5	*6	*	*	*	*	*	*	*	*	*	*
237.7	10	*	*8	*	*	*	*	*	*	*	*
237.70	*7	*	*5	*	*	*	*	*	*	*	*
238	85	40	44	*	*7	18	58	21	20	35	9
238.4	17	9	*8	*	*	*7	*8	*	*	*7	*
238.7	57	26	31	*	*	*9	46	16	13	23	*6
239	32	16	16	*	*7	*9	16	*7	9	11	*5
239.6	10	*5	*	*	*	*	*6	*	*	*	*
240–279	20,325	8,570	11,754	622	2,619	6,270	10,813	4,338	5,228	7,146	3,613
240	19	*6	13	*	*	*7	*8	*	*6	*8	*
240.9	19	*6	12	*	*	*7	*8	*	*5	*8	*
241	37	*5	31	*	*7	17	12	10	*8	11	*7
241.0	15	*	13	*	*	*7	*	*	*	*	*
241.1	18	*	16	—	*	*9	*6	*	*	*7	*
242	99	25	75	*	23	28	47	27	22	30	20
242.0	32	*8	24	*	*8	12	12	*8	*7	10	*7
242.00	30	*7	23	*	*7	11	12	*8	*6	*9	*7
242.8	*5	*	*	—	*	*	*	*	*	*	*
242.9	57	15	42	*	14	15	27	17	12	17	11
242.90	53	13	40	*	14	14	25	17	11	15	10

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
244	1,767	356	1,411	*7	207	457	1,096	415	497	549	306
244.0	54	*8	46	*	*7	18	29	13	13	17	11
244.1	12	*	10	—	*	*	*6	*	*	*	*
244.8	11	*	*9	*	*	*	*	*	*	*	*
244.9	1,687	342	1,345	*7	195	430	1,055	396	477	523	291
245	18	*	16	*	*7	*7	*	*6	*	*	*
245.2	13	*	11	*	*	*6	*	*	*	*	*
246	22	*	18	*	*7	*7	*8	*	*	10	*5
246.9	15	*	13	—	*	*	*6	*	*	*8	*
250	5,480	2,477	3,003	40	611	1,881	2,949	1,236	1,330	1,992	923
250.0	4,075	1,799	2,276	21	379	1,353	2,322	919	993	1,511	652
250.00	3,532	1,571	1,960	*8	284	1,162	2,077	805	877	1,281	568
250.01	186	73	113	9	44	57	75	40	43	73	31
250.02	334	144	190	*	45	127	161	68	69	146	52
250.03	23	10	14	*	*5	*7	*8	*6	*	11	*
250.1	150	70	80	16	82	35	17	27	42	54	28
250.10	33	13	20	*	13	10	*8	*7	*7	12	*7
250.11	68	31	37	13	40	11	*	10	22	22	14
250.12	21	10	11	—	*9	*9	*	*	*	9	*
250.13	28	16	12	*	20	*5	*	*5	*9	11	*
250.2	26	13	13	*	*6	*9	10	*7	*	12	*
250.20	13	*7	*6	—	*	*	*7	*	*	*7	*
250.22	10	*	*6	*	*	*	*	*	*	*	*
250.3	*6	*	*	*	*	*	*	*	*	*	*
250.4	283	143	140	*	30	111	142	55	62	87	78
250.40	204	102	102	—	11	76	117	36	41	60	67
250.41	34	18	16	—	11	17	*6	*8	9	11	*5
250.42	35	19	16	*	*	16	15	*9	9	12	*5
250.43	*9	*	*5	—	*	*	*	*	*	*	*
250.5	126	57	69	*	16	51	58	28	36	38	23
250.50	77	35	42	—	*	30	44	17	22	21	17
250.51	21	11	10	—	9	9	*	*6	*5	*8	*
250.52	22	*8	14	*	*	10	10	*	*7	*8	*
250.53	*6	*	*	—	*	*	*	*	*	*	*
250.6	409	191	218	—	55	172	181	90	101	148	70
250.60	275	130	145	—	16	117	142	63	64	96	52
250.61	55	21	35	—	25	17	14	11	16	18	11
250.62	65	33	31	—	*7	34	23	13	17	28	*6
250.63	14	*7	*7	—	*7	*	*	*	*	*7	*
250.7	84	46	38	—	*	33	48	24	18	28	14
250.70	62	35	28	—	*	23	38	19	10	22	11
250.71	*7	*	*	—	*	*	*	*	*	*	*
250.72	13	*7	*6	—	*	*7	*5	*	*	*	*
250.8	282	136	147	*	27	102	152	82	62	95	43
250.80	203	95	107	*	12	68	123	60	45	65	33
250.81	27	15	12	*	*7	10	*9	10	*6	*8	*
250.82	45	21	24	*	*5	21	19	10	9	20	*6
250.83	*7	*	*	*	*	*	*	*	*	*	*
250.9	41	20	20	*	*9	15	17	*	13	16	*8
250.90	*8	*	*	—	*	*	*	*	*	*	*
250.92	29	14	15	*	*5	12	12	*	10	10	*6
251	101	49	52	12	15	28	46	23	23	40	15
251.1	11	*6	*5	*	*	*	*	*	*	*	*
251.2	36	17	19	*8	*7	*7	15	*7	*7	17	*
251.8	52	26	27	*	*7	18	27	13	13	17	*9
252	38	*9	29	*	*	13	19	*8	12	10	*7
252.0	27	*6	20	*	*	11	12	*5	10	*8	*
252.00	17	*	13	*	*	*6	*8	*	*7	*	*
252.01	*8	*	*6	—	*	*	*	*	*	*	*
252.1	9	*	*7	*	*	*	*	*5	*	*	*

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
253	91	39	53	11	10	20	51	19	18	31	23
253.2	15	*7	*7	*	*	*	*8	*	*	*	*
253.5	14	*6	*8	*	*	*	*5	*	*	*7	*
253.6	50	19	31	*	*	12	34	13	9	17	12
255	90	37	53	*	18	32	37	18	25	27	20
255.0	10	*	*8	*	*	*	*	*	*	*	*
255.4	56	25	31	*	10	20	24	12	17	15	12
255.9	12	*	*7	*	*	*	*5	*	*	*	*
256	16	—	16	*	14	*	*	*	*	*7	*
256.4	13	—	13	*	12	*	*	*	*	*5	*
257	*5	*5	—	*	*	*	*	*	*	*	*
259	*6	*	*	*	*	*	*	*	*	*	*
260	16	*5	11	—	*	*	14	*	*	12	*
261	32	17	15	*	*	*8	19	*7	*8	10	*7
262	*7	*	*	*	*	*	*	*	*	*	*
263	424	191	233	*7	45	99	273	78	109	175	62
263.0	*5	*	*	*	*	*	*	*	*	*	*
263.8	19	*8	11	*	*	*	12	*	*	*8	*
263.9	395	178	217	*	43	93	256	69	103	165	59
266	47	18	29	*	*	10	34	14	10	17	*6
266.2	44	16	28	—	*	9	32	14	9	16	*6
269	10	*	*6	*	*	*	*	*	*	*	*
269.9	*5	*	*	*	*	*	*	*	*	*	*
270	22	*8	14	*	*	*7	*8	*5	*	9	*
270.4	*8	*	*	*	*	*	*	*	*	*	*
270.6	*6	*	*	*	*	*	*	*	*	*	*
271	24	12	12	*	*	*8	*8	9	*6	*	*
271.3	20	*8	11	*	*	*7	*7	*8	*6	*	*
272	3,852	1,948	1,903	*7	277	1,394	2,174	939	1,019	1,206	688
272.0	1,277	628	650	*	98	473	706	424	308	348	197
272.1	54	33	21	*	12	27	14	*9	17	18	10
272.2	12	*8	*	—	*	*	*6	*	*	*	*
272.4	2,449	1,251	1,199	*	160	866	1,419	494	671	814	470
272.8	9	*	*	—	*	*	*	*	*	*	*
272.9	41	21	21	—	*	14	23	*	15	15	*7
273	68	32	36	*5	10	23	30	11	21	24	12
273.1	*7	*	*	—	—	*	*5	*	*	*	*
273.8	49	21	28	*	*9	16	19	*7	16	18	*8
274	287	198	89	*	16	89	182	67	74	94	52
274.0	61	41	20	—	*	18	40	14	16	23	*8
274.9	223	155	68	*	13	70	140	53	58	69	43
275	311	128	183	12	59	105	135	49	91	120	52
275.0	14	*9	*	*	*6	*	*	*	*	*6	*
275.2	127	49	78	*	22	48	54	18	40	54	15
275.3	43	19	24	*	11	12	17	*7	12	17	*7
275.4	124	49	74	*	20	41	58	19	37	41	27
275.41	58	23	35	*	12	23	21	*8	18	21	11
275.42	51	20	30	*	*6	16	28	*9	15	16	11
275.49	11	*	*7	*	*	*	*7	*	*	*	*
276	6,038	2,493	3,545	455	785	1,434	3,365	1,089	1,601	2,239	1,109
276.0	168	69	99	10	15	29	115	38	45	59	26
276.1	1,029	432	597	36	84	265	644	158	289	355	227
276.2	341	156	184	36	48	102	155	51	84	138	68
276.3	63	30	33	*	*9	18	31	9	18	27	*9
276.4	25	10	14	*	*	*6	15	*	9	*7	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
276.5	2,558	1,081	1,477	326	326	475	1,432	492	668	955	443
276.6	99	45	54	*	12	31	52	10	27	41	20
276.7	483	233	250	*8	40	130	305	94	125	166	98
276.8	1,135	379	756	24	231	332	549	218	289	447	181
276.9	137	57	80	*7	18	45	67	15	46	43	33
277	80	46	35	18	36	16	10	19	22	25	14
277.0	27	15	12	10	16	*	*	12	*8	*5	*
277.00	*6	*	*	*	*	*	*	*	*	*	*
277.02	18	11	*8	*6	12	*	*	*8	*6	*	*
277.1	*9	*7	*	—	*7	*	*	*	*	*5	*
277.7	23	11	12	*	*6	11	*	*	*8	*8	*
277.8	*8	*	*	*	*	*	*	*	*	*	*
278	1,277	434	843	24	434	557	262	267	292	463	255
278.0	1,253	427	826	24	422	548	260	261	289	451	252
278.00	729	265	464	20	229	305	175	151	171	252	155
278.01	524	162	362	*	193	243	84	111	117	199	97
278.1	16	*	13	—	*8	*7	*	*	*	*7	*
278.8	*7	*	*	*	*	*	*	*	*	*	*
279	26	10	16	*	*5	*9	*8	*	*6	13	*
279.0	11	*	*7	*	*	*	*	*	*	*7	*
280–289	5,109	2,032	3,077	186	1,045	1,293	2,584	1,008	1,221	1,876	1,004
280	789	279	510	18	153	196	422	162	207	281	140
280.0	454	170	283	*	82	113	255	89	120	167	78
280.8	*8	*	*	*	*	*	*	*	*	*	*
280.9	326	103	222	14	69	81	162	71	85	110	60
281	88	35	53	*	*7	20	60	13	27	29	20
281.0	28	*7	21	—	*	*6	21	*	12	10	*
281.1	*7	*	*5	—	*	*	*	*	*	*	*
281.9	49	23	26	*	*	12	32	*6	13	14	16
282	160	57	104	30	95	23	13	46	29	68	18
282.4	31	14	17	*	13	*	*9	12	*	*8	*7
282.42	*7	*	*	*	*5	*	—	*	*	*	*
282.49	22	10	13	*	*7	*	*9	*7	*	*5	*7
282.5	26	*	22	*	20	*	*	*6	*	13	*
282.6	97	36	61	22	61	14	*	25	19	46	*8
282.60	16	*	12	*	*7	*6	*	*	*	*7	*
282.61	*8	*	*6	*	*	*	*	*	*	*	*
282.62	66	26	39	13	46	*7	*	17	12	33	*
283	26	9	17	*	*	9	12	*	*	11	*7
283.0	10	*	*5	*	*	*	*6	*	*	*	*
283.9	11	*	*9	*	*	*	*	*	*	*6	*
284	209	115	94	12	30	64	103	36	49	82	42
284.8	199	111	88	11	29	61	98	33	48	79	39
284.9	*8	*	*6	*	*	*	*	*	*	*	*
285	2,558	923	1,635	38	515	601	1,405	485	620	950	504
285.1	633	228	406	*	94	160	375	105	199	197	131
285.2	284	123	161	*	34	78	168	57	65	109	53
285.21	83	42	41	*	12	27	44	13	21	33	15
285.22	55	24	31	*	*	20	30	10	15	22	*8
285.29	146	57	89	*	19	32	94	34	29	53	30
285.8	41	16	25	*	9	13	18	*8	*7	15	11
285.9	1,597	556	1,041	29	377	349	842	314	348	627	308
286	214	101	113	9	31	66	108	43	41	80	50
286.0	*7	*	*	*	*	*	*	*	*	*	*
286.3	19	*6	13	*	*6	*5	*7	*	*6	*	*
286.4	*8	*	*	*	*	*	*	*	*	*	*
286.5	20	9	11	*	*	*5	15	*	*	9	*
286.6	17	*8	9	*	*	*5	*8	*	*	9	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
286.7	35	17	18	*	*	11	19	10	*5	12	*8
286.9	106	53	53	*	14	35	53	18	19	41	28
287	513	263	250	19	83	149	261	112	119	175	107
287.1	*5	*	*	*	*	*	*	*	*	*	*
287.3	50	21	29	*8	13	9	20	12	12	16	10
287.4	79	43	37	*	12	25	42	20	17	20	22
287.5	368	194	174	9	56	110	193	78	87	130	72
287.9	*5	*	*	*	*	*	*	*	*	*	*
288	408	191	218	43	89	117	159	73	94	152	90
288.0	176	84	92	23	31	54	68	35	34	67	40
288.3	*6	*	*	*	*	*	*	*	*	*	*
288.8	220	102	118	18	57	59	85	36	58	80	46
289	143	60	83	14	39	48	42	34	32	49	28
289.0	10	*6	*	*	*	*6	*	*	*	*	*
289.3	14	*	*9	*7	*	*	*	*	*	*6	*
289.5	*9	*	*5	*	*	*	*	*	*	*	*
289.59	*5	*	*	*	*	*	*	*	*	*	*
289.8	70	28	42	*	23	23	22	20	15	21	15
289.81	41	13	28	*	17	13	10	13	9	12	*8
289.89	25	14	11	*	*6	*8	11	*6	*	*8	*6
289.9	31	14	17	*	*7	11	12	*7	*8	12	*
290–319	12,695	6,005	6,690	356	5,330	3,911	3,098	3,108	3,229	4,470	1,888
290	194	68	126	—	*	*9	184	65	43	59	27
290.0	99	31	68	—	*	*	95	38	20	26	16
290.2	11	*	*8	—	—	*	10	*	*	*	*
290.21	*6	*	*	—	—	*	*6	*	*	*	*
290.3	*9	*	*6	—	—	*	*9	*	*	*	*
290.4	68	29	39	—	*	*	64	18	13	29	*9
290.40	51	23	28	—	*	*	48	13	*8	22	*7
290.41	*9	*	*	—	*	*	*7	*	*	*	*
290.43	*5	*	*	—	—	*	*5	*	*	*	—
291	316	228	88	*	136	148	33	82	99	80	55
291.0	53	43	10	—	18	28	*7	11	14	18	11
291.2	15	13	*	—	*	*7	*7	*	*5	*	*
291.8	241	167	75	—	115	109	18	68	79	56	39
291.81	230	160	70	—	110	102	18	66	76	52	36
291.89	11	*7	*	—	*	*6	*	*	*	*	*
292	263	149	114	*	129	67	65	81	63	74	44
292.0	122	73	49	*	81	37	*	47	33	27	15
292.1	20	11	9	*	*6	*	9	*	*	*7	*
292.12	17	*9	*8	*	*	*	*8	*	*	*6	*
292.8	112	60	52	*	39	23	50	30	24	38	21
292.81	69	34	35	*	*8	14	47	13	19	24	13
292.84	39	24	15	*	28	9	*	15	*	13	*7
292.9	9	*6	*	—	*	*	*	*	*	*	*
293	109	53	56	*	12	23	74	26	32	36	15
293.0	69	33	36	*	*	10	55	15	23	21	10
293.8	28	14	14	*	9	11	*8	*7	*7	12	*
293.83	21	12	9	*	*7	9	*	*6	*5	*9	*
293.9	11	*5	*6	—	*	*	10	*	*	*	*
294	924	346	578	*	18	37	868	218	250	319	137
294.1	352	130	222	*	*	15	333	86	95	115	57
294.10	284	104	180	*	*	12	270	70	74	90	50
294.11	67	26	41	*	*	*	62	15	20	25	*7
294.8	542	202	340	*	*	18	519	126	150	191	75
294.9	27	12	15	*	*8	*	14	*6	*	11	*5
295	586	308	278	*	263	229	92	157	108	255	67
295.3	172	103	69	*	82	67	21	42	26	87	17
295.30	96	55	41	*	42	39	14	15	16	55	10

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
295.32	32	20	12	*	15	13	*	*8	*	17	*
295.34	43	28	15	—	25	15	*	18	*7	14	*
295.6	27	16	11	—	*7	12	*8	*8	*6	*7	*6
295.60	*8	*5	*	—	*	*	*	*	*	*	*
295.62	16	*9	*7	—	*	*7	*7	*	*	*	*
295.7	237	114	123	*	123	91	22	64	47	101	25
295.70	189	92	97	*	100	72	16	43	39	87	20
295.72	15	*7	*8	*	*8	*5	*	*	*	*6	*
295.74	31	14	17	*	15	13	*	17	*	*8	*
295.9	138	67	71	*	44	54	39	39	25	56	18
295.90	122	59	63	*	36	48	38	32	24	52	14
295.92	*9	*	*	—	*	*	*	*	*	*	*
295.94	*7	*	*	—	*	*	—	*	*	*	*
296	1,503	595	908	82	835	419	167	347	361	652	143
296.0	13	*6	*7	*	*7	*	*	*	*	*8	*
296.04	*6	*	*	*	*	*	*	*	*	*5	—
296.2	309	122	188	19	159	85	46	54	69	163	23
296.20	194	73	120	12	104	53	24	33	46	102	13
296.22	21	*8	13	*	11	*	*	*	*	11	*
296.23	51	21	30	*	28	12	*8	11	13	24	*
296.24	40	18	22	*	15	14	9	*7	*6	24	*
296.3	344	131	212	*6	182	111	45	87	93	132	32
296.30	93	41	52	*	51	32	*9	31	27	32	*
296.32	27	9	17	*	16	*7	*	*	*8	11	*
296.33	161	55	106	*	84	54	19	36	45	67	14
296.34	59	25	34	*	29	17	12	15	13	22	*9
296.4	80	32	48	*	46	24	*8	22	18	33	*7
296.40	35	14	20	*	21	*9	*	*6	*8	16	*
296.43	*9	*	*6	—	*	*	*	*	*	*	*
296.44	33	13	20	*	19	10	*	12	*6	12	*
296.5	130	48	83	*	83	35	10	34	34	54	*8
296.50	84	31	53	*	55	21	*6	20	22	38	*
296.53	24	*7	17	*	14	*8	*	*9	*5	*7	*
296.54	19	*8	11	*	12	*5	*	*	*	*8	*
296.6	90	39	51	10	55	21	*	15	12	53	*9
296.60	45	19	26	*	30	10	*	*7	*6	29	*
296.63	15	*6	*9	*	*9	*	*	*	*	10	*
296.64	27	12	15	*	15	*7	*	*6	*	13	*
296.7	49	21	28	*	27	14	*	17	10	17	*5
296.8	396	151	245	25	220	109	43	79	102	166	49
296.80	344	132	211	23	188	93	40	65	92	147	40
296.89	51	17	33	*	31	15	*	14	10	19	*8
296.9	91	45	45	15	55	14	*7	35	21	25	10
296.90	84	43	41	14	52	12	*5	33	20	22	9
296.99	*7	*	*	*	*	*	*	*	*	*	*
297	35	17	18	—	*8	*6	21	14	*7	10	*
297.1	20	11	10	—	*	*	12	9	*5	*	*
297.9	12	*6	*6	—	*	*	*7	*	*	*	*
298	198	92	106	*	74	40	79	51	37	73	38
298.9	192	89	103	*	72	38	77	49	35	71	37
299	27	22	*	14	11	*	*	10	*6	*6	*
299.0	14	11	*	*7	*5	*	*	*	*	*	*
299.00	13	11	*	*7	*	*	*	*	*	*	*
299.8	9	*8	*	*	*	*	*	*	*	*	*
299.80	9	*8	*	*	*	*	*	*	*	*	*
300	1,282	402	880	20	419	422	421	295	372	452	162
300.0	690	221	470	*7	190	224	269	161	190	250	89
300.00	569	187	381	*	141	183	241	138	158	203	71
300.01	76	19	57	*	33	24	17	16	18	27	15
300.02	44	14	30	*	15	17	11	*7	13	20	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
300.1	32	*7	26	*	21	*9	*	*5	12	11	*
300.11	12	*	*8	*	*6	*	*	*	*	*	*
300.14	*8	*	*8	—	*6	*	—	*	*	*	*
300.15	*7	*	*7	*	*6	*	*	*	*	*	*
300.2	24	*9	16	*	13	*7	*	*6	*6	*9	*
300.21	11	*	*7	—	*8	*	*	*	*	*	*
300.3	34	15	19	*	23	*6	*	12	11	*7	*
300.4	398	103	295	*5	112	151	129	95	119	138	45
300.9	98	46	52	*	59	24	11	14	34	37	14
301	284	117	167	*6	182	82	14	94	82	77	32
301.2	*6	*	*	*	*	*	*	*	*	*	*
301.4	*6	*	*	*	*	*	*	*	*	*	*
301.5	9	*	*7	*	*5	*	*	*	*	*7	*
301.50	*8	*	*5	*	*	*	*	*	*	*6	*
301.6	23	*6	16	*	11	10	*	*	*7	11	*
301.7	35	30	*6	*	28	*7	*	*6	13	11	*
301.8	113	25	88	*	83	27	*	34	43	25	11
301.81	*8	*5	*	—	*	*	*	*	*	*	*
301.83	88	13	76	*	68	19	*	25	35	20	*8
301.89	13	*	*8	*	*8	*	*	*6	*	*	*
301.9	84	44	40	*	48	28	*7	44	11	17	11
302	*7	*6	*	*	*	*	*	*	*	*	*
303	740	510	230	*	317	345	77	196	195	210	139
303.0	146	105	42	*	68	69	*9	40	58	19	29
303.00	40	27	13	*	17	19	*	*	17	*8	11
303.01	101	75	26	—	48	48	*6	34	39	12	16
303.9	594	406	188	*	249	275	69	156	137	191	110
303.90	287	193	93	*	112	136	39	49	64	116	58
303.91	242	173	70	—	112	107	24	89	50	61	42
303.92	13	9	*	*	*7	*	*	*5	*	*	*
303.93	52	30	22	*	18	28	*	13	20	11	*8
304	595	336	260	*	396	179	16	225	102	185	83
304.0	201	108	93	*	116	79	*6	83	30	56	32
304.00	71	35	36	*	40	29	*	11	12	34	15
304.01	121	68	53	—	72	47	*	71	14	21	15
304.03	*8	*	*	—	*	*	*	*	*	*	*
304.1	31	15	16	*	17	11	*	13	*	10	*
304.10	14	*6	*8	*	*6	*7	*	*	*	*7	*
304.11	16	*8	*8	—	10	*	*	10	*	*	*
304.2	147	90	57	*	108	38	*	65	20	48	15
304.20	61	36	25	*	44	17	*	11	11	26	13
304.21	77	48	29	—	58	19	*	51	*6	19	*
304.23	*7	*	*	—	*	*	*	*	*	*	*
304.3	56	37	19	*	47	*8	*	15	25	12	*
304.30	23	14	*9	*	18	*	*	*	10	*6	*
304.31	24	17	*7	*	22	*	*	10	10	*	*
304.33	*7	*	*	*	*6	*	*	*	*	*	*
304.4	25	14	11	—	20	*	*	*	*	*	13
304.40	14	9	*	—	11	*	—	*	*	*	*8
304.41	*6	*	*	—	*	*	*	*	*	*	*
304.6	*7	*	*	*	*	*	*	*	*	*	*
304.7	40	23	17	*	28	12	*	20	*6	10	*
304.70	11	*	*6	*	*7	*	*	*	*	*	*
304.71	26	17	10	—	19	*7	*	16	*	*6	*
304.8	43	26	17	*	29	11	*	12	*	24	*
304.80	32	20	12	*	22	*8	*	*5	*	22	*
304.81	*8	*	*	—	*6	*	*	*	*	*	*
304.9	45	20	25	*	27	14	*	14	*6	18	*7
304.90	32	13	18	*	17	11	*	*9	*	13	*6
304.91	9	*	*6	*	*6	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
305	3,466	1,890	1,575	17	1,673	1,318	458	659	873	1,324	610
305.0	542	375	167	*	255	209	76	122	122	194	103
305.00	386	266	120	*	191	140	55	70	100	142	74
305.01	99	73	27	*	39	45	15	29	14	38	19
305.02	23	14	9	*	12	9	*	13	*	*	*
305.03	34	22	11	—	14	15	*5	11	*5	9	*8
305.1	2,190	1,097	1,093	*	923	896	368	381	595	852	362
305.2	158	96	62	*	125	28	*	38	40	56	23
305.20	115	69	46	*	94	17	*	20	34	44	17
305.21	27	17	10	*	20	*7	*	12	*	*8	*5
305.22	*8	*6	*	*	*7	*	—	*	*	*	*
305.23	*7	*	*	—	*	*	—	*	*	*	*
305.4	18	*7	11	—	12	*	*	*	*	*7	*
305.40	14	*6	*8	—	9	*	*	*	*	*6	*
305.5	80	50	30	*	44	33	*	20	21	20	19
305.50	55	35	20	*	31	22	*	10	19	14	12
305.51	18	12	*6	*	*8	*9	*	*7	*	*	*
305.53	*6	*	*	—	*	*	*	*	*	*	*
305.6	255	146	109	*	163	88	*	57	60	105	32
305.60	196	110	86	*	126	67	*	34	52	86	24
305.61	36	23	13	*	21	13	*	15	*	13	*
305.62	*8	*	*	—	*5	*	*	*	*	*	*
305.63	15	*8	*7	—	11	*	*	*	*	*	*
305.7	58	30	29	*	45	13	*	*	*	11	44
305.70	47	23	24	*	37	10	*	*	*	10	35
305.71	*6	*	*	—	*	*	*	*	*	*	*
305.9	160	87	73	*7	102	47	*	32	28	76	24
305.90	130	71	59	*7	81	38	*	22	26	65	18
305.91	10	*7	*	—	*7	*	*	*	*	*5	*
305.93	18	*8	10	—	13	*6	*	*6	*	*6	*5
306	9	*	*6	*	*	*	*	*	*	*	*
307	89	28	60	9	41	17	22	19	28	28	15
307.1	*9	*	*8	*	*6	*	*	*	*	*	*
307.2	*6	*	*	*	*	*	*	*	*	*	*
307.23	*5	*	*	*	*	*	*	*	*	*	*
307.5	21	*	19	*	15	*	*	*6	9	*	*
307.50	14	*	13	*	10	*	*	*	*7	*	*
307.8	15	*	13	*	*6	*5	*	*	*	*7	*
307.81	13	*	12	—	*5	*	*	*	*	*6	*
307.9	33	16	17	*	*8	*6	16	*	10	12	*6
308	16	*7	10	*	*7	*	*	*	*	*6	*
308.9	*7	*	*	*	*	*	*	*	*	*	*
309	272	95	176	21	169	59	23	94	83	61	34
309.0	48	21	27	*	26	10	10	15	13	11	*8
309.2	37	16	20	*	19	*8	*	11	14	*7	*
309.24	*6	*	*	*	*	*	*	*	*	*	*
309.28	28	12	16	*	17	*7	*	*8	13	*	*
309.4	16	*7	9	*	11	*	*	*6	*	*	*
309.8	151	41	109	11	99	35	*5	52	49	34	16
309.81	148	41	107	11	98	34	*5	51	49	33	15
309.9	15	*8	*7	*	11	*	*	*7	*	*5	*
310	40	19	22	*	*7	*7	25	*8	*6	16	10
310.1	*8	*5	*	*	*	*	*	*	*	*	*6
310.2	*7	*	*	*	*	*	*	*	*	*	*
310.9	23	*7	16	—	*	*	18	*	*	13	*
311	1,291	440	851	18	427	428	418	331	359	394	207
312	64	44	20	13	40	*7	*	28	13	21	*
312.3	39	29	11	*6	28	*	*	19	*8	11	*
312.30	23	18	*6	*	17	*	*	13	*	*5	*
312.34	14	10	*	*	10	*	*	*5	*	*5	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
312.8	*7	*	*	*	*	*	*	*	*	*	*
312.9	17	10	*7	*	*6	*	*	*7	*	*6	*
313	35	20	15	22	13	—	*	*8	*8	17	*
313.8	33	19	14	20	12	—	*	*7	*7	17	*
313.81	29	17	12	18	11	—	—	*7	*6	16	*
314	136	87	49	57	69	*8	*	33	39	50	14
314.0	129	82	47	55	65	*8	*	33	39	44	14
314.00	24	15	9	*6	15	*	*	*7	*7	*6	*
314.01	105	67	38	49	50	*5	*	26	32	37	10
314.9	*6	*	*	*	*	—	—	*	*	*6	—
315	59	35	24	37	16	*	*	17	12	12	18
315.3	*7	*	*	*7	*	—	—	*	*	*	*
315.8	21	14	*7	17	*	*	*	*	*	*6	*8
315.9	24	15	9	10	9	*	*	*8	*	*	*7
317	36	19	18	*	19	9	*	14	9	12	*
318	33	18	15	*	14	11	*6	9	12	10	*
318.0	*7	*	*	*	*	*	*	*	*	*	*
318.1	19	11	*8	*	*8	*7	*	*	*6	*7	*
318.2	*7	*	*	*	*	*	*	*	*	*	*
319	83	49	34	12	28	28	14	18	24	28	13
320–389	3,942	1,708	2,234	373	678	912	1,979	849	999	1,354	740
320	13	*6	*8	*	*	*	*	*	*	*6	*
321	9	*	*6	*	*	*6	*	*	*	*	*
321.0	*8	*	*6	*	*	*6	*	*	*	*	*
322	13	*8	*	*	*	*	*	*	*	*	*6
322.9	10	*7	*	*	*	*	*	*	*	*	*
323	12	*6	*6	*	*5	*	*	*	*	*	*
323.9	9	*	*	*	*	*	*	*	*	*	*
324	15	*9	*6	*	*	*7	*	*	*	*	*
324.1	10	*6	*	—	*	*6	*	*	*	*	*
331	568	223	345	22	18	35	492	132	149	205	82
331.0	415	143	272	—	*	10	403	98	111	151	55
331.3	29	18	11	*	*	*	23	*	*8	10	*6
331.4	72	37	35	20	15	17	21	19	14	25	15
331.8	39	19	20	—	*	*	35	*8	12	15	*
331.82	38	18	19	—	*	*	35	*8	12	14	*
331.9	11	*5	*6	*	*	*	*8	*	*	*	*
332	232	117	115	*	*	21	210	60	70	69	32
332.0	231	117	114	*	*	21	209	60	69	69	32
333	123	40	82	*	15	29	76	21	39	45	18
333.1	36	12	24	*	*	*7	27	*5	10	16	*5
333.2	10	*	*5	*	*	*	*	*	*	*	*
333.8	*9	*	*7	*	*	*	*	*	*	*	*
333.82	*8	*	*6	*	*	*	*	*	*	*	*
333.9	57	15	42	*	*7	14	35	9	20	20	*8
333.99	50	13	37	*	*6	13	30	*7	18	18	*7
334	*8	*	*	*	*	*	*	*	*	*	*
335	15	*9	*6	*	*	*	*8	*	*	*5	*
335.2	12	*7	*5	—	*	*	*8	*	*	*	*
335.20	*9	*5	*	—	*	*	*7	*	*	*	*
336	19	11	*8	*	*	*6	*7	*	*	*8	*
336.3	*5	*	*	*	*	*	*	*	*	*	*
336.9	*7	*	*	*	*	*	*	*	*	*	*
337	45	15	30	*	10	20	14	11	13	13	*8
337.1	20	*7	13	—	*	9	*8	*6	*6	*5	*
337.2	16	*	12	*	*6	*8	*	*	*5	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
337.20	*7	*	*6	—	*	*	*	*	*	*	*
337.22	*6	*	*	*	*	*	*	*	*	*	*
337.9	*6	*	*	*	*	*	*	*	*	*	*
340	123	32	90	*	32	60	28	32	39	31	21
341	*6	*	*	*	*	*	*	*	*	*	*
342	151	72	79	*	12	39	95	30	38	54	29
342.8	*7	*	*	*	*	*	*	*	*	*	*
342.9	137	66	71	*	10	36	87	26	34	50	27
342.90	115	57	58	*	9	29	74	19	32	42	22
342.91	11	*	*7	*	*	*	*7	*	*	*	*
342.92	11	*	*6	*	*	*	*6	*	*	*	*
343	81	45	36	31	34	12	*	17	22	26	16
343.2	13	*7	*6	*7	*	*	*	*6	*	*	*
343.9	58	33	26	18	28	10	*	*8	17	22	11
344	147	94	53	*9	51	44	43	23	33	61	30
344.0	54	37	17	*	21	21	10	*7	14	23	10
344.00	48	33	15	*	19	19	*8	*6	13	21	*8
344.1	55	38	16	*	23	14	14	10	9	23	12
344.4	*6	*	*5	*	*	*	*	*	*	*	*
344.40	*5	*	*	*	*	*	*	*	*	*	*
344.6	10	*7	*	*	*	*	*	*	*	*	*
344.60	*6	*	*	—	*	*	*	*	*	*	*
344.8	*6	*	*	*	*	*	*	*	*	*	*
344.89	*5	*	*	*	*	*	*	*	*	*	*
344.9	13	*6	*6	*	*	*	*7	*	*	*	*
345	118	53	65	32	39	30	16	27	25	39	27
345.1	21	*8	13	*7	*	*7	*	*	*	*5	*9
345.10	17	*7	10	*	*	*6	*	*	*	*5	*7
345.3	24	12	11	9	*6	*	*	*	*6	*7	*6
345.4	*6	*	*	*	*	*	*	*	*	*	*
345.5	13	*	9	*	*	*	*	*	*	*	*
345.50	10	*	*8	*	*	*	*	*	*	*	*
345.8	*6	*	*	*	*	*	*	*	*	*	*
345.9	44	21	23	*8	19	12	*	10	11	18	*5
345.90	39	18	20	*7	16	11	*	*9	10	15	*
346	247	51	196	*6	130	90	20	60	55	86	47
346.1	*5	*	*	*	*	*	*	*	*	*	*
346.2	*8	*	*	*	*	*	*	*	*	*	*
346.20	*6	*	*	*	*	*	*	*	*	*	*
346.8	15	*	13	*	*7	*6	*	*	*	*6	*
346.80	11	*	9	*	*6	*	*	*	*	*	*
346.9	215	42	173	*	114	79	18	53	49	73	40
346.90	206	40	166	*	108	76	18	52	47	70	38
346.91	9	*	*7	*	*7	*	*	*	*	*	*
347	*8	*	*5	*	*	*	*	*	*	*	*
347.0	*8	*	*5	*	*	*	*	*	*	*	*
347.00	*8	*	*5	*	*	*	*	*	*	*	*
348	230	116	113	30	44	58	97	42	47	82	58
348.0	11	*6	*6	*	*	*	*	*	*	*	*
348.1	57	35	23	*8	*8	18	23	*6	13	20	18
348.2	11	*	10	*	9	*	*	*	*	*	*
348.3	91	43	48	*6	9	20	56	19	18	36	18
348.30	39	19	20	*	*	*8	23	*6	*7	18	*8
348.31	30	12	18	*	*	*7	21	10	*6	10	*
348.39	22	11	10	*	*	*	12	*	*	*8	*6
348.4	17	*8	*9	*	*7	*5	*	*	*	*	*
348.5	9	*	*5	*	*	*	*	*	*	*	*
348.8	25	16	*9	*7	*6	*6	*5	*	*5	10	*6
348.9	*8	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
349	64	22	42	*	27	16	18	13	18	23	10
349.0	27	*	22	*	22	*	*	*	*7	9	*6
349.2	*5	*	*	*	*	*	*	*	*	*	*
349.8	29	13	16	*	*	*9	16	*6	*9	11	*
349.82	25	11	14	*	*	*7	15	*	*7	10	*
350	21	*	17	—	*	*7	11	*	*6	*7	*
350.1	20	*	17	—	*	*6	11	*	*6	*6	*
351	29	11	18	*	*5	*8	14	*	*6	12	*7
351.0	26	9	17	*	*	*7	13	*	*5	10	*6
353	11	*6	*6	*	*	*	*	*	*	*	*
353.0	*5	*	*	*	*	*	*	*	*	*	*
354	35	14	21	*	10	14	10	*8	9	10	*7
354.0	19	*5	14	*	*6	*8	*	*	*	*7	*
354.2	*6	*	*	*	*	*	*	*	*	*	*
355	62	26	36	*	10	20	32	13	14	23	12
355.8	27	12	16	*	*	*7	17	*7	*8	*8	*5
355.9	27	10	17	*	*5	10	12	*5	*	12	*
356	106	53	53	*	10	33	62	18	25	43	19
356.9	100	50	50	*	*8	32	59	17	24	40	19
357	269	136	133	*	30	110	127	62	65	92	50
357.0	10	*	*6	*	*	*	*	*	*	*	*
357.2	242	123	120	—	24	102	117	55	59	83	45
357.8	*5	*	*	*	*	*	*	*	*	*	*
358	28	13	15	*	*	*5	18	9	*6	11	*
358.0	25	12	13	*	*	*	17	*8	*5	10	*
358.00	21	10	11	*	*	*	16	*7	*	9	*
359	25	14	11	*	*8	*6	*8	*5	*6	9	*5
359.1	10	*6	*	*	*5	*	*	*	*	*	*
361	*5	*	*	*	*	*	*	*	*	*	*
362	170	65	104	*	15	43	108	36	51	52	31
362.0	89	39	51	*	12	35	42	18	27	27	17
362.01	86	37	49	*	10	34	41	17	27	25	16
362.3	*9	*6	*	—	*	*	*	*	*	*	*
362.34	*5	*	*	—	—	*	*	*	*	*	*
362.5	60	17	43	*	*	*	57	12	20	17	12
362.50	59	16	43	*	*	*	56	11	20	17	11
364	*6	*	*	*	*	*	*	*	*	*	*
365	179	67	112	*	*5	28	146	40	51	55	33
365.9	173	64	109	*	*	27	142	38	50	53	31
366	25	*9	17	*	*	*6	17	*6	9	*5	*
366.9	21	*7	14	*	*	*	14	*	*8	*	*
368	63	26	37	*	14	20	25	13	14	20	16
368.1	15	*5	10	*	*	*	*7	*	*	*7	*
368.16	10	*	*7	*	*	*	*7	*	*	*	*
368.2	16	*8	*8	*	*	*6	*6	*	*	*	*6
368.4	14	*5	*9	*	*	*	*8	*	*	*	*
368.46	*8	*	*	—	*	*	*5	*	*	*	*
368.8	10	*	*6	*	*	*	*	*	*	*	*
368.9	*7	*	*	*	*	*	*	*	*	*	*
369	87	42	45	*	15	16	53	20	21	31	16
369.0	27	14	13	*	*5	*	17	*6	*7	10	*
369.00	26	13	13	*	*	*	17	*6	*7	10	*
369.4	23	*7	15	*	*	*	15	*8	*	*6	*
369.6	26	15	11	*	*	*6	15	*	*6	10	*6
369.60	26	15	11	*	*	*6	14	*	*6	10	*6

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
372	53	29	24	14	12	*7	19	*8	13	22	10
372.0	*9	*	*	*	*	*	*	*	*	*	*
372.3	34	18	16	11	*6	*	13	*	*9	14	*7
372.30	31	17	14	10	*	*	13	*	*8	11	*7
372.7	*8	*	*	*	*	*	*	*	*	*	*
372.72	*7	*	*	*	*	*	*	*	*	*	*
373	9	*	*6	*	*	*	*	*	*	*	*
373.1	*6	*	*	*	*	*	*	*	*	*	*
374	12	*5	*7	*	*	*6	*	*	*	*	*
374.3	*7	*	*	*	*	*	*	*	*	*	*
374.30	*7	*	*	*	*	*	*	*	*	*	*
376	18	*8	11	*8	*6	*	*	*	*	*8	*
376.0	14	*6	*8	*7	*	*	*	*	*	*7	*
376.01	13	*6	*7	*7	*	*	*	*	*	*6	*
377	14	*8	*6	*	*6	*	*	*	*	*6	*
377.3	*7	*	*	*	*	*	*	*	*	*	*
378	17	*9	*8	*	*	*6	*	*	*	*6	*
378.5	11	*6	*	*	*	*	*	*	*	*	*
378.54	*5	*	*	*	*	*	*	*	*	*	*
379	21	9	12	*	*5	*6	*5	*	*	9	*
379.9	*8	*	*	*	*	*	*	*	*	*	*
380	30	15	16	*8	*9	*	*9	*8	*7	*8	*7
380.1	17	*8	*9	*5	*	*	*	*	*	*5	*
380.10	16	*8	*9	*5	*	*	*	*	*	*	*
380.4	9	*5	*	*	*	*	*	*	*	*	*
381	19	*8	11	14	*	*	*	*6	*	*	*
381.4	*8	*	*	*	*	*	*	*	*	*	*
382	130	72	58	108	11	*5	*6	17	29	55	29
382.9	123	67	56	102	10	*5	*6	15	28	52	28
383	10	*6	*	*	*	*	*	*	*	*	*
386	78	25	53	*	*8	20	50	21	24	20	13
386.0	17	*5	12	—	*	*6	10	*	*5	*	*
386.00	17	*	12	—	*	*5	10	*	*5	*	*
386.1	35	12	23	*	*	*7	24	11	*8	10	*6
386.11	26	10	16	*	*	*	19	*9	*5	*7	*
386.12	*6	*	*	—	*	*	*	*	*	*	*
386.3	22	*8	14	—	*	*6	13	*6	*9	*	*
386.30	20	*8	13	—	*	*5	12	*6	*8	*	*
388	18	*7	10	*	*6	*	*7	*	*	*7	*
388.3	*7	*	*	*	*	*	*	*	*	*	*
388.30	*6	*	*	*	*	*	*	*	*	*	*
388.7	*7	*	*	*	*	*	*	*	*	*	*
388.70	*7	*	*	*	*	*	*	*	*	*	*
389	102	51	51	*6	14	16	67	22	29	30	21
389.1	*7	*	*	*	*	*	*	*	*	*	*
389.7	*5	*	*	*	*	*	*	*	*	*	*
389.8	*6	*	*	*	*	*	*	*	*	*	*
389.9	82	41	41	*	*9	12	58	16	23	26	17
390–459	34,677	16,812	17,865	140	2,104	8,997	23,436	7,822	8,659	12,260	5,936
394	24	10	13	*	*	*6	16	*9	*	*5	*5
394.0	*7	*	*5	*	*	*	*5	*	*	*	*
394.1	10	*6	*	*	*	*	*7	*5	*	*	*
396	178	71	107	*	*	18	155	48	50	57	23
396.0	*6	*	*	*	*	*	*6	*	*	*	*
396.2	49	20	29	*	*	*	44	13	17	13	*6
396.3	89	33	56	*	*	9	77	26	24	31	9
396.8	17	*6	12	—	*	*	15	*6	*	*6	*
396.9	12	*8	*	—	*	*	10	*	*	*	*

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
397	191	73	118	*	14	34	141	57	45	59	30
397.0	189	73	116	*	14	34	139	57	45	58	30
398	79	29	50	—	*	11	65	25	18	25	12
398.9	79	29	50	—	*	11	65	25	18	25	12
398.91	75	29	46	—	*	10	63	22	17	24	12
401	9,003	3,841	5,161	18	710	2,869	5,406	2,044	2,211	3,243	1,505
401.0	79	31	47	*	11	27	41	15	15	36	13
401.1	90	36	54	*	*9	25	56	72	*	13	*
401.9	8,834	3,774	5,060	17	691	2,818	5,308	1,958	2,191	3,195	1,490
402	289	115	173	—	9	61	218	71	70	112	36
402.0	11	*	*8	—	*	*	*7	*	*	*6	*
402.01	*7	*	*	—	*	*	*	*	*	*	*
402.9	273	110	164	—	*9	57	208	66	68	105	34
402.90	150	63	87	—	*	30	116	37	40	56	17
402.91	123	46	77	—	*	27	92	29	28	49	18
403	1,269	650	618	*	113	383	770	244	256	493	276
403.0	31	14	18	*	11	12	*9	*	*6	18	*
403.01	28	13	16	*	11	11	*7	*	*	17	*
403.1	12	*8	*	—	*	*	*7	10	*	*	*
403.11	12	*7	*	—	*	*	*7	10	*	*	*
403.9	1,225	629	596	*	101	368	754	231	250	473	272
403.90	49	22	27	*	*	10	36	16	*8	14	11
403.91	1,176	607	569	*	98	357	718	215	241	459	261
404	83	42	41	*	*	24	54	17	22	32	12
404.0	*8	*	*	—	*	*	*	*	*	*	*
404.9	75	38	37	*	*	20	50	15	21	28	10
404.91	*7	*	*	—	*	*	*	*	*	*	*
404.92	17	9	*8	*	*	*6	10	*	*	*7	*
404.93	48	24	24	—	*	12	34	*8	15	19	*6
405	9	*	*6	*	*	*	*5	*	*	*	*
405.9	*8	*	*5	*	*	*	*	*	*	*	*
410	865	499	366	*	47	253	563	219	207	274	166
410.0	17	9	*8	—	*	*	11	*	*	*7	*
410.01	16	9	*7	—	*	*	10	*	*	*7	*
410.1	74	43	31	*	*	27	42	15	16	22	21
410.11	63	37	26	—	*	26	34	13	14	20	17
410.12	*9	*	*	—	*	*	*7	*	*	*	*
410.2	13	*9	*	—	*	*	*7	*	*	*	*
410.21	13	*8	*	—	*	*	*7	*	*	*	*
410.3	*8	*6	*	—	*	*	*	*	*	*	*
410.31	*8	*5	*	—	*	*	*	*	*	*	*
410.4	86	64	22	—	*7	39	40	17	20	27	22
410.41	79	59	20	—	*6	36	37	15	19	24	21
410.42	*6	*	*	—	*	*	*	*	*	*	*
410.5	10	*6	*	—	*	*	*6	*	*	*	*
410.51	10	*5	*	—	*	*	*6	*	*	*	*
410.7	508	287	221	—	22	134	352	144	113	163	87
410.71	478	271	207	—	20	129	329	134	108	154	82
410.72	26	14	12	—	*	*	20	*9	*	*7	*5
410.8	24	13	10	*	*	*8	13	*5	11	*	*
410.81	22	13	9	—	*	*8	12	*5	10	*	*
410.9	123	61	62	—	*7	28	89	24	37	40	22
410.91	106	52	55	—	*6	23	77	20	29	36	20
410.92	14	*8	*6	—	*	*	9	*	*7	*	*
411	576	338	238	*	29	225	320	131	125	218	102
411.0	*6	*	*	*	*	*	*	*	*	*	*
411.1	549	322	227	*	28	217	304	121	122	209	97
411.8	21	11	10	—	*	*6	14	10	*	*6	*
411.89	18	*9	9	—	*	*6	12	9	*	*5	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
412	686	430	255	*	36	214	435	163	177	222	124
413	391	208	183	—	23	117	250	89	104	135	63
413.1	*6	*	*	—	*	*	*	*	*	*	*
413.9	385	206	178	—	21	115	248	89	103	131	61
414	4,675	2,749	1,926	*	133	1,280	3,260	1,086	1,192	1,631	766
414.0	4,340	2,528	1,812	*	123	1,195	3,020	1,011	1,111	1,511	706
414.00	1,225	726	498	—	14	256	954	303	334	425	163
414.01	2,993	1,711	1,283	*	107	899	1,986	689	743	1,040	521
414.02	83	68	15	—	29	53	13	26	24	20	
414.04	12	*9	*	—	*	*	*7	*	*	*	*
414.05	25	13	12	—	*	*6	19	*	*	18	*
414.1	17	10	*7	*	*	*5	10	*	*	*	*5
414.10	*7	*	*	*	*	*	*5	*	*	*	*
414.12	*6	*	*	—	*	*	*	*	*	*	*
414.8	272	184	89	*	*7	69	196	64	66	93	49
414.9	47	28	19	—	*	11	34	*8	11	23	*5
415	211	88	123	*	34	60	116	46	51	76	38
415.1	208	87	121	*	34	59	114	46	51	74	38
415.11	20	*9	11	—	*	*5	10	*	*	*6	*
415.19	189	79	110	*	30	54	104	42	46	68	33
416	278	113	165	*7	26	74	171	57	58	105	58
416.0	22	*6	16	*	*	*	14	*8	*	9	*
416.8	200	75	125	*5	21	50	124	37	45	78	42
416.9	55	31	24	*	*	20	33	12	11	18	13
420	28	17	11	*	*8	*9	10	*7	*8	*8	*
420.9	25	15	10	*	*7	*8	*9	*6	*7	*7	*
420.90	19	10	*8	*	*	*6	*7	*5	*	*5	*
420.91	*6	*	*	*	*	*	*	*	*	*	*
421	33	21	12	*	*7	12	14	*9	*5	12	*6
421.0	30	19	11	*	*6	12	12	*8	*5	11	*6
423	59	32	27	*	9	24	22	11	11	25	12
423.8	*6	*	*	—	*	*	*	*	*	*	*
423.9	45	24	20	*	*7	17	17	*7	*7	20	10
424	1,104	433	671	*8	98	205	793	302	268	356	178
424.0	611	214	397	*	82	142	384	162	153	218	78
424.1	406	188	218	*	10	46	347	118	102	104	82
424.2	34	15	20	*	*	*8	24	11	*	11	*8
424.3	*8	*	*	*	*	*	*	*	*	*	*
424.9	44	13	31	*	*	*7	33	10	*8	18	*8
424.90	42	12	30	*	*	*6	32	*9	*8	18	*8
425	696	391	306	*	67	218	408	138	166	263	129
425.1	18	*	13	*	*	*5	11	*	*6	*	*
425.4	630	357	273	*	60	196	372	121	151	243	116
425.5	16	15	*	—	*	10	*	*	*	*	*
425.8	18	*8	10	*	*	*6	10	*	*	10	*
425.9	10	*	*7	*	*	*	*7	*	*	*	*
426	432	239	193	*	26	73	328	93	118	142	79
426.0	73	40	33	*	*	12	58	19	20	20	14
426.1	110	64	46	*	*7	15	88	23	31	39	18
426.10	*6	*	*	—	*	*	*5	*	*	*	*
426.11	61	36	24	*	*	*8	50	10	18	22	11
426.12	*8	*	*	—	*	*	*5	*	*	*	*
426.13	35	21	14	*	*	*	27	*9	10	11	*5
426.2	*7	*	*	—	*	*	*	*	*	*	*
426.3	99	43	57	—	*	19	78	23	26	32	18
426.4	81	51	30	*	*7	16	58	14	25	29	14
426.5	29	20	10	—	*	*	26	*6	*8	11	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
426.52	10	*	*5	—	*	*	9	*	*	*	*
426.53	14	11	*	—	*	*	12	*	*	*6	*
426.7	14	*8	*6	*	*6	*	*	*	*	*	*
426.8	*6	*	*	*	*	*	*	*	*	*	*
426.89	*6	*	*	*	*	*	*	*	*	*	*
426.9	9	*6	*	—	*	*	*8	*	*	*	*
427	4,528	2,253	2,274	25	194	795	3,514	1,034	1,183	1,502	808
427.0	90	36	54	*	*8	24	57	19	21	36	15
427.1	282	189	93	*	16	80	184	62	85	90	45
427.3	3,095	1,511	1,585	*	66	455	2,573	734	803	1,008	550
427.31	2,872	1,376	1,496	*	53	398	2,419	688	748	928	507
427.32	223	134	89	*	12	56	154	46	55	79	43
427.4	43	31	12	*	*	14	25	*8	10	12	13
427.41	42	30	12	*	*	13	25	*8	9	12	13
427.5	95	56	39	*	*8	27	58	16	27	37	16
427.6	90	48	41	*	11	20	58	17	24	33	16
427.61	23	12	11	*	*	*	18	*	*	*9	*5
427.69	65	36	29	*	*9	16	39	12	19	24	10
427.8	764	356	408	16	76	160	512	167	195	257	144
427.81	168	78	90	*	*	20	144	40	42	55	32
427.89	596	277	319	16	73	139	368	128	154	203	112
427.9	65	25	40	*	*	16	44	10	17	30	*9
428	4,165	1,860	2,305	10	128	809	3,218	885	1,117	1,517	647
428.0	3,875	1,743	2,132	10	118	747	2,999	832	1,038	1,411	593
428.1	*6	*	*	—	*	*	*6	*	*	*	*
428.2	72	41	31	*	*	19	50	10	29	25	*8
428.20	36	22	14	*	*	10	25	*6	12	15	*
428.21	*9	*	*	—	*	*	*7	*	*	*	*
428.22	12	*7	*5	—	*	*	*8	*	*7	*	*
428.23	15	*7	*7	—	*	*	10	*	*6	*	*
428.3	177	57	120	—	*6	34	137	38	41	63	35
428.30	130	42	88	—	*	25	101	28	29	50	23
428.31	19	*7	13	—	*	*	17	*	*	*	*6
428.32	15	*	11	—	*	*	11	*	*	*	*5
428.33	12	*	*9	—	*	*	*8	*	*	*	*
428.4	18	*8	9	*	*	*	14	*	*	*9	*
428.40	11	*	*6	*	*	*	*9	*	*	*6	*
428.9	18	*9	9	*	*	*	13	*	*	*7	*6
429	173	76	97	*	17	43	110	43	39	68	23
429.2	55	23	32	—	*	9	45	19	13	18	*5
429.3	74	32	43	*	*7	21	43	13	15	35	13
429.9	26	12	14	*	*	*6	15	*7	*6	*9	*
430	32	13	19	*	*7	14	11	*8	*8	10	*7
431	87	41	46	*	*8	26	52	16	17	32	21
432	49	28	21	*	*	11	33	*8	10	16	15
432.1	39	23	17	*	*	*9	28	*6	*8	13	13
432.9	*8	*	*	*	*	*	*	*	*	*	*
433	288	155	133	*	*	57	226	59	64	113	52
433.1	211	114	97	*	*	42	166	45	46	82	37
433.10	188	100	89	*	*	36	151	40	40	75	33
433.11	23	14	*9	*	*	*6	15	*5	*6	*7	*
433.2	*5	*	*	—	*	*	*	*	*	*	*
433.3	63	34	29	*	*	9	52	11	14	26	13
433.30	56	31	25	*	*	*7	47	9	13	22	11
433.31	*7	*	*	*	*	*	*5	*	*	*	*
434	453	209	244	*	19	103	329	93	108	166	86
434.0	*9	*6	*	*	*	*	*	*	*	*	*
434.01	*7	*	*	*	*	*	*	*	*	*	*
434.1	53	25	28	*	*	12	38	14	12	16	11

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
434.11	51	24	27	*	*	11	37	14	12	15	11
434.9	391	178	213	*	15	88	286	76	94	147	74
434.90	13	*7	*6	*	*	*	10	*	*	*	*
434.91	378	171	207	*	15	85	276	74	91	143	70
435	252	98	154	*	16	53	183	59	62	92	38
435.3	10	*	*7	—	*	*	*7	*	*	*	*
435.8	12	*7	*	—	*	*	*8	*	*	*	*
435.9	227	86	141	*	14	49	165	53	58	84	33
436	20	*9	12	—	*	*	15	*	*5	*7	*
437	151	61	90	*	9	31	109	34	37	59	20
437.0	59	25	34	—	*	*	54	14	16	22	*7
437.1	11	*	*8	*	*	*	*8	*	*	*	*
437.2	13	*5	*8	*	*	*	*6	*	*	*	*
437.3	21	*8	12	*	*	9	*8	*6	*	*8	*
437.7	*7	*	*	—	*	*	*	*	*	*	*
437.9	29	11	19	—	*	*	24	*6	*8	12	*
438	501	237	264	*	21	113	365	101	135	148	117
438.0	16	*8	*7	—	*	*	12	*	*	*	*6
438.1	82	41	41	*	*	21	58	14	23	23	23
438.10	*5	*	*	—	*	*	*	*	*	*	*
438.11	59	27	32	—	*	13	43	11	17	15	16
438.12	*8	*6	*	*	*	*	*6	*	*	*	*
438.19	10	*6	*	—	*	*	*	*	*	*	*
438.2	226	104	121	*	*7	52	166	53	57	70	46
438.20	194	87	107	*	*6	43	145	48	48	62	35
438.21	19	10	9	*	*	*6	13	*	*6	*	*6
438.22	12	*7	*5	*	*	*	*8	*	*	*	*
438.7	*7	*	*	—	*	*	*	*	*	*	*
438.8	149	70	79	*	9	30	110	24	44	43	38
438.82	34	15	18	—	*	*	27	*6	11	*8	10
438.83	12	*	*8	*	*	*	*9	*	*	*	*5
438.84	*8	*	*	—	*	*	*6	*	*	*	*
438.89	92	46	46	*	*5	20	66	14	28	30	20
438.9	10	*	*5	*	*	*	*7	*	*	*	*
440	354	187	167	—	*9	87	258	73	92	139	50
440.0	19	*8	12	—	*	*6	13	*	*6	*6	*
440.1	64	29	36	—	*	13	46	13	16	26	10
440.2	221	126	96	—	*	59	159	41	56	91	33
440.20	59	31	27	—	*	16	42	*7	18	25	10
440.21	48	29	18	—	*	15	31	*8	14	19	*6
440.22	20	10	10	—	*	*	16	*	*	10	*
440.23	25	14	11	—	*	*5	19	*7	*5	10	*
440.24	64	38	26	—	*	17	46	15	14	25	10
440.29	*6	*	*	—	*	*	*	*	*	*	*
440.3	*6	*	*	—	*	*	*	*	*	*	*
440.8	*5	*	*	—	*	*	*	*	*	*	*
440.9	38	18	20	—	*	*	33	14	9	11	*
441	178	113	65	*	*	35	137	39	46	63	29
441.0	21	14	*7	—	*	*6	12	*	*	9	*
441.01	*8	*6	*	—	*	*	*	*	*	*	*
441.02	*8	*	*	—	*	*	*6	*	*	*	*
441.2	23	12	11	*	*	*8	13	*6	*	*6	*7
441.3	*7	*5	*	—	—	*	*7	*	*	*	*
441.4	116	77	39	—	*	19	97	25	31	43	17
441.9	*5	*	*	*	*	*	*	*	*	*	—
442	35	25	10	*	*	10	22	*7	*9	14	*6
442.2	13	11	*	—	*	*	9	*	*	*6	*
442.3	15	10	*	*	*	*	10	*	*	*6	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
443	475	252	222	*	13	119	343	109	126	171	68
443.0	19	*7	12	—	*	*7	*8	*	*6	*5	*
443.8	49	29	20	—	*	19	29	12	12	15	11
443.81	26	14	12	—	*	11	14	*8	*5	*7	*6
443.89	23	15	*8	—	*	*8	14	*	*6	*8	*
443.9	402	215	187	*	*6	92	304	94	107	149	52
444	89	45	44	*	*5	31	52	18	21	36	14
444.0	*9	*	*	*	*	*	*	*	*	*	*
444.2	55	27	28	*	*	20	32	12	11	23	*9
444.21	*6	*	*	*	*	*	*	*	*	*	*
444.22	49	24	25	*	*	17	28	12	9	21	*7
444.8	21	11	11	*	*	*7	13	*	*6	*8	*
444.81	12	*7	*	*	*	*	*7	*	*	*	*
444.89	10	*	*6	*	*	*	*6	*	*	*	*
446	48	17	31	*9	*	9	27	11	11	15	11
446.1	*9	*7	*	*8	*	*	*	*	*	*	*
446.4	*5	*	*	*	*	*	*	*	*	*	*
446.5	21	*	19	—	*	*	20	*5	*6	*7	*
446.6	*5	*	*	*	*	*	*	*	*	*	*
447	58	27	31	*	*6	23	28	15	*9	22	12
447.1	33	16	17	*	*	13	19	9	*5	13	*6
447.6	*8	*	*	*	*	*	*	*	*	*	*
448	*5	*	*	—	*	*	*	*	*	*	*
451	59	28	31	*	13	20	25	15	13	21	10
451.0	11	*6	*	*	*	*	*	*	*	*	*
451.1	13	*7	*6	—	*	*	9	*	*	*	*
451.19	*9	*6	*	—	*	*	*6	*	*	*	*
451.2	*8	*	*	*	*	*	*	*	*	*	*
451.8	25	10	15	*	*7	12	*6	*6	*	12	*
451.82	11	*	*6	*	*	*6	*	*	*	*5	*
451.84	10	*	*6	*	*	*	*	*	*	*	*
452	9	*	*6	*	*	*6	*	*	*	*	*
453	405	181	224	*5	52	125	222	103	95	140	67
453.4	262	118	144	*	29	81	150	67	71	85	39
453.40	70	31	39	*	*8	19	43	14	16	27	12
453.41	110	48	61	*	12	33	63	32	30	32	15
453.42	83	38	44	*	*9	29	44	21	25	25	12
453.8	121	54	67	*	20	38	61	32	21	45	24
453.9	13	*6	*7	*	*	*	*8	*	*	*7	*
454	35	10	25	—	*	10	22	*8	15	*8	*
454.0	*8	*	*5	—	*	*	*6	*	*	*	*
454.2	*6	*	*	—	*	*	*	*	*	*	*
454.9	16	*	12	—	*	*5	*9	*	*7	*	*
455	184	74	110	*	47	49	87	39	44	65	36
455.0	64	25	39	—	13	16	35	15	15	24	10
455.2	34	18	16	—	*6	13	15	10	*8	10	*7
455.3	23	*9	14	—	*7	*	12	*	*9	*7	*
455.5	12	*6	*6	—	*	*	*5	*	*	*	*
455.6	33	*8	25	*	12	*7	13	*	*6	11	11
455.8	12	*6	*7	—	*	*	*	*	*	*6	*
456	79	56	24	*	13	41	24	15	14	29	21
456.0	*6	*	*	—	*	*	*	*	*	*	*
456.1	*9	*7	*	*	*	*	*	*	*	*	*
456.2	46	33	14	*	*7	28	11	10	*6	17	14
456.20	31	23	*8	*	*	19	*8	*6	*	11	10
456.21	15	10	*6	*	*	9	*	*	*	*6	*
456.8	15	11	*	*	*	*7	*6	*	*	*7	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
457	39	16	23	*	*7	12	17	*8	10	13	*7
457.1	24	*6	17	*	*	*7	12	*6	*6	*7	*
457.2	*7	*	*	*	*	*	*	*	*	*	*
457.8	*7	*5	*	*	*	*	*	*	*	*	*
458	575	254	321	10	70	132	363	106	146	222	100
458.0	131	60	72	*	11	22	96	25	36	52	19
458.1	*6	*	*5	—	*	*	*	*	*	*	*
458.2	74	30	43	*	11	18	43	17	16	28	13
458.29	69	29	41	*	10	17	40	15	15	26	13
458.8	32	16	15	*	*	*8	19	*8	*7	*9	*8
458.9	332	147	185	*5	43	81	202	56	83	134	60
459	183	83	100	*	16	50	116	38	51	68	26
459.0	19	*8	11	*	*	*	13	*	*6	*7	*
459.2	15	*6	9	*	*	*6	*6	*	*	*7	*
459.8	131	62	69	*	11	34	86	29	39	45	18
459.81	119	57	62	*	10	32	77	26	36	41	16
459.89	12	*5	*6	*	*	*	*8	*	*	*	*
459.9	14	*6	*9	*	*	*	10	*	*	*8	*
460–519	12,384	5,804	6,580	1,212	1,475	2,999	6,699	2,543	3,080	4,498	2,262
461	48	21	27	*6	20	12	9	11	17	16	*
461.0	9	*	*	*	*	*	*	*	*	*	*
461.2	*7	*	*	*	*	*	*	*	*	*	*
461.9	24	*8	15	*	11	*	*	*	*8	9	*
462	64	27	37	17	24	12	10	9	19	26	10
463	19	12	*8	*	12	*	*	*	*6	*8	*
464	46	25	21	35	*	*	*	9	11	13	14
464.1	12	*	*7	9	*	*	*	*	*	*	*5
464.10	10	*	*6	*8	*	*	*	*	*	*	*
464.4	25	16	9	25	*	*	*	*5	*6	*8	*6
465	178	81	96	65	37	29	46	34	42	67	35
465.9	174	80	95	64	37	29	45	34	41	65	35
466	396	190	206	198	37	55	107	68	94	147	88
466.0	202	78	125	12	34	53	103	41	57	72	32
466.1	194	113	81	186	*	*	*	27	37	75	56
466.11	98	56	42	96	*	*	*	12	19	34	34
466.19	96	57	39	90	*	*	*	15	18	41	22
470	12	*8	*	*	*	*	*	*	*	*5	*
472	11	*5	*6	*	*	*	*6	*	*	*6	*
472.0	*9	*	*6	*	*	*	*	*	*	*	*
473	164	67	96	26	46	48	44	28	41	70	24
473.0	23	10	13	*	*	*7	*7	*	*	10	*5
473.2	13	*6	*7	*	*	*	*	*	*	*7	*
473.3	*5	*	*	*	*	*	*	*	*	*	*
473.8	13	*6	*7	*	*	*	*	*	*	*6	*
473.9	107	43	64	15	31	34	28	21	31	42	13
474	28	17	11	15	*8	*	*	*6	*	11	*6
474.0	*9	*	*	*	*	*	*	*	*	*	*
474.00	*6	*	*	*	*	*	*	*	*	*	*
474.1	18	12	*6	12	*	*	*	*	*	*6	*
474.10	*8	*6	*	*7	*	—	—	*	*	*	*
474.11	*9	*	*	*	*	*	*	*	*	*	*
475	18	11	*7	*	13	*	*	*	*7	*6	*
477	85	31	55	*7	20	29	30	11	25	33	16
477.9	81	30	51	*6	19	28	29	10	24	32	14
478	115	62	53	27	19	37	32	20	30	42	23
478.1	23	10	13	*7	*5	*5	*5	*	*7	10	*
478.2	49	32	17	10	*8	22	10	9	13	17	9

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
478.29	39	27	12	*	*6	20	*9	*7	11	16	*6
478.3	19	*9	10	*	*	*	11	*	*	*5	*6
478.30	*6	*	*	*	*	*	*	*	*	*	*
478.31	*6	*	*	*	*	*	*	*	*	*	*
478.7	12	*	*7	*6	*	*	*	*	*	*	*
478.74	*6	*	*	*	*	*	*	*	*	*	*
480	43	21	22	30	*	*	*7	*	*9	14	15
480.1	16	*6	10	15	—	*	*	*	*	*6	*5
480.9	24	14	11	13	*	*	*7	*	*5	*8	*8
481	51	24	27	*	*9	14	25	11	15	17	*8
482	213	123	90	10	25	46	132	36	51	79	48
482.0	17	11	*6	*	*	*7	9	*	*	*7	*
482.1	42	25	18	*	*6	*8	26	*7	12	13	10
482.2	11	*6	*	*	*	*	*8	*	*	*	*
482.3	14	*7	*6	*	*	*	*8	*	*	*	*
482.30	9	*	*	*	*	*	*5	*	*	*	*
482.4	76	46	30	*	10	17	47	11	16	31	18
482.41	69	42	27	*	*9	15	43	11	14	29	16
482.8	30	18	12	*	*	*6	21	*6	*7	14	*
482.82	*5	*	*	*	*	*	*	*	*	*	*
482.83	20	13	*7	*	*	*	14	*	*	10	*
482.9	23	9	13	*	*	*	11	*6	*6	*6	*
483	15	*7	*8	9	*	*	*	*	*	*5	*
483.0	13	*6	*7	*9	*	*	*	*	*	*	*
484	*8	*	*	*	*	*	*	*	*	*	*
485	36	19	18	15	*	*7	13	*5	*	23	*
486	2,020	963	1,057	189	176	396	1,259	399	541	709	371
487	98	46	52	23	*7	16	52	14	40	30	14
487.0	24	12	12	*	*	*	14	*	*9	*7	*
487.1	72	33	38	18	*	13	37	10	30	22	9
490	116	43	72	*6	21	37	51	19	22	55	20
491	1,202	568	634	*	39	322	839	253	299	470	179
491.2	1,157	549	609	*	30	308	819	248	289	451	169
491.20	88	46	42	—	*	21	64	14	17	43	14
491.21	926	443	484	*	23	246	657	203	235	351	136
491.22	143	60	83	—	*	41	98	30	37	57	19
491.9	38	17	21	*	*7	12	18	*	*9	18	*8
492	223	120	103	*	10	63	148	45	61	83	33
492.0	13	*8	*	*	*	*	*	*	*	*	*
492.8	210	112	98	*	*6	59	144	42	59	79	31
493	1,928	653	1,275	347	521	535	525	472	459	653	343
493.0	76	38	38	50	10	10	*6	17	*7	18	34
493.00	37	17	20	19	*6	*7	*	*8	*	11	13
493.01	*9	*	*	*7	*	*	*	*	*	*	*
493.02	30	17	13	23	*	*	*	*7	*	*	16
493.2	464	157	307	*	35	160	267	111	124	149	80
493.20	258	90	169	*	21	78	158	59	75	74	50
493.21	14	*	12	*	*	*9	*	*	*	*8	*
493.22	191	65	126	*	12	73	105	48	48	67	28
493.8	*6	*	*	*	*	*	*	*	*	*	*
493.81	*5	*	*	*	*	*	*	*	*	*	*
493.9	1,382	457	925	295	472	364	251	343	325	485	229
493.90	1,016	310	706	148	374	295	199	262	226	348	180
493.91	58	27	31	37	12	*7	*	10	13	27	*8
493.92	308	120	188	109	86	63	50	70	86	110	41
494	39	17	22	*	*	*8	27	10	*8	14	*7
494.0	29	12	17	*	*	*6	20	*7	*6	9	*6
494.1	10	*5	*	*	*	*	*7	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
496	2,127	1,084	1,043	*	58	526	1,539	453	551	762	362
500	9	9	*	—	—	*	*9	*7	*	*	*
501	15	14	*	—	—	*	12	*	*	*6	*
507	385	221	164	13	37	65	270	85	85	131	83
507.0	380	218	162	13	36	63	268	85	85	129	82
508	*8	*	*	—	*	*	*	*	*	*	*
508.0	*6	*	*	—	*	*	*	*	*	*	*
510	32	21	11	*	*6	12	12	*7	*6	13	*6
510.9	28	18	10	*	*	10	11	*6	*	11	*5
511	452	220	233	14	42	116	280	99	118	162	73
511.0	28	13	16	*	*6	10	13	*	12	*8	*
511.8	31	14	16	*	*	9	17	*	*8	11	*6
511.9	391	191	200	13	33	96	249	88	98	141	63
512	111	61	50	*	25	29	52	23	24	46	19
512.0	*5	*	*	*	*	*	*	*	*	*	*
512.1	51	26	25	*	*5	16	28	11	11	20	*9
512.8	54	32	23	*	18	11	22	10	11	24	9
513	12	*8	*	*	*	*	*	*	*	*	*
513.0	12	*8	*	*	*	*	*	*	*	*	*
514	36	12	24	*	*	*8	22	*5	9	18	*
515	224	98	125	*	11	47	166	38	65	76	44
516	35	19	17	*	*	11	21	*7	*8	12	*8
516.3	15	*8	*7	*	*	*	12	*	*	*	*
516.8	19	10	9	*	*	*8	9	*	*	*8	*
517	29	11	18	*	*8	13	*7	*7	*8	*8	*5
517.8	22	*8	13	*	*5	10	*6	*6	*	*6	*
518	1,638	808	830	99	202	449	888	305	354	625	354
518.0	402	189	213	35	50	116	200	65	99	148	90
518.1	*6	*	*	*	*	*	*	*	*	*	*
518.3	23	10	13	*	*	*5	15	*	*6	*8	*
518.4	19	*6	13	*	*	*	10	*	*5	*8	*
518.5	116	68	48	*6	21	34	55	20	25	51	20
518.8	1,069	530	538	55	123	286	605	208	217	405	238
518.81	741	373	367	24	83	199	435	147	149	288	156
518.82	84	42	42	18	15	16	36	11	18	34	21
518.83	54	25	29	*	*	19	30	11	11	16	16
518.84	74	32	42	*	*7	22	44	12	18	30	14
518.89	116	58	58	10	15	30	61	27	22	37	30
519	80	39	42	21	*8	20	31	16	27	23	15
519.0	13	*7	*6	*	*	*6	*	*	*	*	*
519.09	*7	*	*	*	*	*	*	*	*	*	*
519.1	28	13	15	*7	*	*7	10	*	11	*9	*
519.8	26	13	14	*8	*	*	13	*	10	*8	*
519.9	*6	*	*	*	*	*	*	*	*	*	*
520–579	11,108	4,786	6,322	537	2,232	3,495	4,843	2,310	2,716	4,005	2,076
521	23	13	10	*	11	*6	*	*7	*	*8	*
521.0	23	13	10	*	11	*6	*	*7	*	*8	*
521.00	19	11	*8	*	9	*6	*	*5	*	*7	*
522	29	14	14	*	16	*7	*	*8	*6	10	*
522.5	23	12	12	*	13	*6	*	*7	*	*8	*
523	14	*	9	*	*	*	*	*	*	*6	*
524	36	9	27	*	26	*5	*	*7	*	16	*8
524.0	13	*	*9	*	11	*	*	*	*	*7	*
524.6	10	*	9	*	*	*	*	*	*	*	*
524.60	*7	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
525	13	*6	*7	*	*8	*	*	*	*	*5	*
525.9	*9	*	*	*	*6	*	*	*	*	*	*
526	9	*	*	*	*	*	*	*	*	*	*
527	24	13	11	*	*	*5	13	*	*6	10	*
527.2	14	*8	*7	*	*	*	*7	*	*	*	*
528	46	24	21	*7	12	14	13	*8	9	20	*8
528.0	19	10	9	*	*	*6	*7	*	*	*7	*
528.3	9	*	*	*	*	*	*	*	*	*	*
528.9	*8	*	*	*	*	*	*	*	*	*	*
530	2,561	1,047	1,515	118	412	863	1,168	583	628	913	437
530.0	*6	*	*	*	*	*	*	*	*	*	*
530.1	198	92	106	*5	41	68	84	46	48	67	37
530.10	63	28	36	*	13	19	29	15	17	22	9
530.11	71	33	38	*	16	25	27	19	13	24	15
530.12	*6	*	*	*	*	*	*	*	*	*	*
530.19	58	29	29	*	11	22	23	11	16	19	11
530.2	32	18	14	*	*6	9	16	*6	*8	*9	9
530.20	17	10	*7	*	*5	*6	*6	*	*	*	*6
530.21	15	*8	*7	—	*	*	10	*	*	*	*
530.3	56	28	28	*	*	15	34	11	16	20	9
530.5	21	*6	15	*	*	*7	11	*	*	*7	*6
530.6	*6	*	*	*	*	*	*	*	*	*	*
530.7	28	20	*9	*	9	*8	9	*	*	12	*8
530.8	2,211	877	1,334	107	345	753	1,006	509	544	792	366
530.81	2,134	836	1,298	106	333	727	969	491	531	767	346
530.82	19	*9	10	*	*	*8	*8	*	*	*6	*
530.85	41	25	16	—	*	15	20	11	*6	13	10
530.89	15	*5	10	*	*5	*	*7	*	*	*6	*5
531	164	69	94	*	18	52	93	24	40	60	40
531.0	14	*5	*8	*	*	*	*9	*	*	*	*
531.00	14	*5	*8	*	*	*	*9	*	*	*	*
531.4	80	36	44	*	*7	20	53	*9	20	31	20
531.40	79	35	43	*	*6	20	52	*9	20	30	20
531.5	*6	*	*	*	*	*	*	*	*	*	*
531.50	*5	*	*	*	*	*	*	*	*	*	*
531.9	54	23	31	—	*7	22	26	10	13	18	14
531.90	53	23	31	—	*7	21	25	9	12	18	14
532	89	52	37	*	11	32	46	17	17	36	20
532.0	12	*8	*	*	*	*	*	*	*	*	*
532.00	11	*8	*	*	*	*	*	*	*	*	*
532.4	40	27	13	*	*	13	23	*7	*7	16	11
532.40	39	27	13	*	*	13	23	*7	*7	16	10
532.9	24	11	13	*	*	10	11	*6	*	10	*
532.90	23	11	13	*	*	9	11	*6	*	10	*
533	110	50	60	*	19	35	56	21	32	37	19
533.4	12	*5	*7	—	*	*	*8	*	*	*	*
533.40	12	*5	*7	—	*	*	*8	*	*	*	*
533.7	*6	*	*	—	*	*	*	*	*	*	*
533.70	*6	*	*	—	*	*	*	*	*	*	*
533.9	87	39	47	*	16	28	43	17	25	30	15
533.90	87	39	47	*	16	27	43	17	25	30	15
534	*8	*	*6	*	*	*	*	*	*	*	*
535	589	270	319	15	116	177	281	135	139	209	107
535.0	51	27	24	*	11	15	21	14	10	20	*6
535.00	39	19	19	*	10	11	14	14	*8	13	*
535.01	12	*8	*	*	*	*	*7	*	*	*7	*
535.1	80	29	52	*	12	22	45	26	11	24	19
535.10	66	21	45	*	10	19	35	23	11	18	14
535.11	15	*8	*7	*	*	*	10	*	*	*6	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
535.3	19	14	*5	—	9	*9	*	*	*7	*6	*
535.30	13	*8	*	—	*7	*	*	*	*6	*	*
535.31	*7	*6	*	—	*	*	*	*	*	*	*
535.4	132	66	67	*	25	42	64	26	32	47	28
535.40	91	45	47	*	18	31	42	17	23	34	18
535.41	41	21	20	*	*7	11	22	*8	9	13	10
535.5	240	94	146	*7	45	73	115	51	59	93	37
535.50	192	73	120	*	40	60	88	40	48	76	29
535.51	48	21	27	*	*6	13	28	11	12	17	*8
535.6	65	39	26	*	13	16	35	15	19	18	12
535.60	54	32	22	*	12	14	28	12	17	15	10
535.61	11	*7	*	*	*	*	*7	*	*	*	*
536	269	94	175	18	63	89	99	44	67	112	46
536.2	36	14	22	*8	10	*7	11	*	*7	15	9
536.3	106	30	76	*	35	45	26	17	18	54	17
536.4	43	18	25	*6	*	*8	26	11	*7	16	10
536.41	16	*7	9	*	*	*	10	*	*	*6	*
536.42	15	*6	*9	*	*	*	*8	*	*	*	*
536.49	11	*5	*6	*	*	*	*7	*	*	*	*
536.8	75	27	48	*	13	25	33	10	34	23	*8
536.9	*7	*	*	*	*	*	*	*	*	*	*
537	87	38	49	*	*7	26	52	17	18	32	20
537.0	23	11	12	*	*	*8	11	*	*5	11	*
537.8	49	21	29	*	*	12	34	11	10	16	13
537.82	*7	*	*	—	*	*	*	*	*	*	*
537.83	22	11	11	—	*	*	18	*	*	*7	*7
537.89	15	*5	10	—	*	*	9	*	*	*5	*
537.9	*5	*	*	—	*	*	*	*	*	*	*
540	312	174	139	74	158	57	23	62	67	94	90
540.0	62	38	25	18	22	16	*6	*8	17	18	19
540.1	37	23	15	12	13	*8	*	9	*7	11	10
540.9	213	113	99	44	123	33	13	45	42	65	61
541	14	*5	9	*	*7	*	*	*	*	*6	*
542	*6	*	*	*	*	*	*	*	*	*	*
543	19	*5	13	*	9	*6	*	*	*7	*	*
543.9	17	*5	12	*	*8	*6	*	*	*7	*	*
550	72	60	11	10	10	17	34	16	16	26	14
550.1	26	22	*	*	*	*6	15	*6	*6	10	*
550.10	16	13	*	*	*	*	10	*	*	*7	*
550.11	*8	*7	*	*	*	*	*5	*	*	*	*
550.9	45	38	*8	*8	*8	12	18	10	10	16	9
550.90	33	27	*6	*	*6	10	14	*8	*7	12	*6
550.92	9	*8	*	*	*	*	*	*	*	*	*
552	99	33	66	*	15	36	48	23	23	34	19
552.1	19	10	*9	*	*	*8	*8	*	*	*5	*5
552.2	60	18	42	*	*9	24	28	15	11	25	10
552.20	16	*	12	—	*	*6	*7	*	*	*7	*
552.21	37	11	26	*	*5	14	18	9	*7	15	*6
552.29	*7	*	*	*	*	*	*	*	*	*	*
552.8	*9	*	*6	*	*	*	*	*	*	*	*
553	531	202	329	10	80	177	264	111	135	210	76
553.1	52	25	27	*	17	16	16	10	14	15	14
553.2	112	42	70	*	19	48	43	26	31	40	16
553.20	29	11	18	*	*	11	15	*5	*9	13	*
553.21	71	25	46	*	15	32	24	17	20	22	13
553.29	12	*5	*6	*	*	*5	*	*	*	*	*
553.3	358	132	226	*6	41	110	201	73	88	152	44
553.8	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
555	125	54	71	*	57	37	28	30	31	44	20
555.0	19	*9	10	*	11	*	*	*5	*6	*	*
555.1	14	*5	*9	*	*8	*	*	*	*	*	*
555.2	*9	*	*	*	*5	*	*	*	*	*	*
555.9	83	36	47	*	33	26	22	19	19	33	12
556	64	31	33	*	22	19	20	16	15	16	18
556.6	10	*6	*	*	*5	*	*	*	*	*	*
556.8	*6	*	*	*	*	*	*	*	*	*	*
556.9	39	19	20	*	9	14	15	*8	12	*8	11
557	87	26	61	*	*7	21	59	19	22	29	17
557.0	41	14	27	*	*	10	27	11	12	11	*8
557.1	*8	*	*6	*	*	*	*6	*	*	*	*
557.9	37	9	28	*	*	*9	25	*5	*8	16	*8
558	411	166	245	113	76	87	134	87	96	169	58
558.1	*7	*	*	—	*	*	*	*	*	*	*
558.9	400	163	237	111	75	84	130	86	91	168	55
560	773	344	430	26	113	215	420	141	212	276	145
560.0	*8	*6	*	*	*	*	*	*	*	*	*
560.1	354	169	185	13	56	99	186	59	99	131	65
560.2	19	*8	11	*	*	*	10	*	*6	*6	*
560.3	77	29	48	*	*7	13	54	15	17	35	10
560.39	72	29	43	*	*6	12	49	14	14	33	10
560.8	159	63	96	*	20	51	84	30	43	53	33
560.81	94	37	56	*	11	28	51	18	22	33	21
560.89	66	26	40	*	*9	23	33	13	20	21	12
560.9	156	69	88	*	24	47	84	32	45	49	30
562	634	238	397	*	55	181	398	151	172	209	102
562.0	10	*	*7	*	*	*	*7	*	*	*	*
562.00	*9	*	*6	—	*	*	*6	*	*	*	*
562.1	624	234	390	*	55	177	391	149	170	205	100
562.10	260	90	170	*	15	65	180	60	70	84	46
562.11	254	95	159	—	39	92	123	68	76	75	35
562.12	92	43	49	—	*	15	76	17	17	40	18
562.13	18	*7	11	—	*	*5	13	*	*8	*6	*
564	664	225	439	47	136	171	311	124	189	242	109
564.0	476	181	294	39	83	112	241	83	135	179	78
564.00	425	162	263	34	75	99	217	69	126	162	69
564.09	50	19	31	*6	*9	13	23	14	9	18	10
564.1	153	26	126	*	45	51	56	34	48	52	19
564.8	23	13	10	*5	*	*	*8	*	*	*5	9
564.81	*8	*6	*	*	*	*	*	*	*	*	*
564.89	15	*7	*8	*	*	*	*6	*	*	*	*
565	10	*5	*	*	*	*	*	*	*	*	*
565.1	*6	*	*	*	*	*	*	*	*	*	*
566	31	19	12	*	11	10	*7	*	*7	13	*6
567	98	47	51	*	22	37	34	23	22	32	21
567.2	64	34	30	*	15	27	19	16	13	22	13
567.8	16	*6	10	*	*	*	*7	*	*	*5	*
567.9	17	*7	9	*	*	*6	*7	*	*	*5	*
568	174	48	126	*	61	61	48	31	46	63	34
568.0	145	42	103	*	46	54	41	27	37	51	30
568.8	28	*6	22	*	15	*6	*7	*	9	11	*
568.81	18	*	14	*	12	*	*	*	*	*8	*
568.89	9	*	*7	*	*	*	*	*	*	*	*
569	281	126	155	*8	43	80	151	61	64	102	54
569.0	9	*	*6	—	*	*	*	*	*	*	*
569.1	9	*	*8	*	*	*	*6	*	*	*	*
569.3	69	34	35	*	*8	19	41	13	14	28	14

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
569.4	26	15	11	*	*7	*	14	*	*9	*9	*
569.41	*7	*	*	—	*	*	*5	*	*	*	*
569.49	16	*9	*7	*	*	*	*7	*	*6	*	*
569.5	30	14	16	*	*	14	12	11	*5	*8	*6
569.6	32	14	18	*	*5	11	14	*6	*6	12	*8
569.62	*8	*	*5	*	*	*	*	*	*	*	*
569.69	21	10	11	*	*	*6	11	*	*	*6	*5
569.8	100	42	57	*	15	24	58	22	25	34	18
569.81	13	*	9	*	*	*5	*	*	*	*6	*
569.82	16	*6	11	*	*	*	*8	*	*	*6	*
569.83	25	12	13	*	*	*6	14	*6	*6	*9	*
569.84	*8	*	*5	—	*	*	*7	*	*	*	*
569.85	23	11	12	—	*	*	20	*	*6	*7	*6
569.89	13	*7	*7	*	*	*	*	*	*	*	*
570	35	20	15	*	11	14	10	*6	*6	12	10
571	492	286	205	*	90	265	133	99	98	168	128
571.1	49	32	17	—	19	27	*	11	13	13	12
571.2	163	116	48	—	27	108	29	35	28	55	46
571.3	24	18	*5	—	9	11	*	*	11	*	*
571.4	17	*	13	*	*	9	*	*	*	*6	*
571.49	13	*	11	*	*	*8	*	*	*	*6	*
571.5	174	95	79	*	13	82	77	33	30	69	41
571.6	11	*	10	*	*	*5	*5	*	*	*	*
571.8	45	16	29	*	17	16	9	*8	10	13	14
572	166	100	65	*	21	89	54	30	27	63	46
572.0	*9	*6	*	*	*	*	*	*	*	*	*
572.2	81	49	32	—	*8	44	29	15	13	29	25
572.3	45	28	16	*	*6	23	14	*7	*6	19	12
572.4	14	*8	*7	—	*	*8	*	*	*	*	*
572.8	17	10	*7	*	*	9	*	*	*	*8	*
573	101	46	55	*	28	33	37	19	24	35	23
573.0	*5	*	*	*	*	*	*	*	*	*	*
573.1	*6	*	*	*	*	*	*	*	*	*	*
573.3	43	16	27	*	14	13	15	*9	9	15	11
573.8	35	17	17	*	*6	12	17	*7	*7	14	*7
573.9	13	*6	*7	*	*	*6	*	*	*	*	*
574	579	213	366	*	169	168	240	115	140	204	120
574.0	142	53	89	*	45	43	53	32	33	50	26
574.00	135	51	84	*	42	42	51	31	33	47	24
574.01	*7	*	*	—	*	*	*	*	*	*	*
574.1	200	71	130	*	62	62	76	43	45	75	37
574.10	194	68	126	*	60	62	72	43	44	72	35
574.11	*6	*	*	—	*	*	*	*	*	*	*
574.2	138	51	87	*	30	40	66	23	36	46	33
574.20	133	48	85	*	29	37	66	21	35	44	32
574.3	*6	*	*	—	*	*	*	*	*	*	*
574.4	*9	*	*	—	*	*	*5	*	*	*	*
574.40	*6	*	*	—	*	*	*	*	*	*	*
574.5	35	13	22	*	10	*9	16	*6	10	12	*7
574.50	18	*7	11	*	*5	*	*9	*	*	*8	*
574.51	17	*6	11	*	*	*	*8	*	*5	*	*
574.6	13	*	*8	*	*	*	*5	*	*	*	*
574.60	*8	*	*5	*	*	*	*	*	*	*	*
574.7	17	*5	12	—	*7	*	*5	*	*	*6	*
574.70	11	*	*7	—	*	*	*	*	*	*	*
574.71	*6	*	*	—	*	*	*	*	*	*	*
574.8	11	*6	*5	*	*	*	*5	*	*	*	*
574.80	*6	*	*	*	*	*	*	*	*	*	*
574.81	*5	*	*	—	*	*	*	*	*	*	*
574.9	9	*	*6	*	*	*	*	*	*	*	*
574.90	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
575	159	59	100	*	56	49	54	28	49	56	26
575.0	35	17	18	*	*9	11	15	*8	12	9	*6
575.1	66	22	44	*	27	21	18	10	19	27	10
575.10	14	*	11	*	*5	*	*	*	*	*	*
575.11	39	10	29	*	19	10	9	*	12	19	*
575.12	14	*9	*	—	*	*7	*	*	*	*	*
575.3	11	*	*7	*	*	*	*6	*	*	*	*
575.8	29	*9	20	*	12	*8	*8	*	10	10	*5
575.9	10	*	*7	—	*	*	*	*	*	*	*
576	103	40	63	*	20	30	50	20	23	37	23
576.1	31	14	17	*	*	*8	19	*7	*	11	*8
576.2	25	10	15	*	*	*7	13	*5	*6	*8	*6
576.8	44	15	29	*	11	14	18	*7	11	17	9
577	466	243	223	*8	156	178	124	92	101	190	84
577.0	342	181	161	*	116	125	97	64	76	139	64
577.1	73	37	35	*	27	36	*8	17	14	32	9
577.2	25	12	13	*	*6	11	*7	*6	*	11	*
577.8	17	*7	10	*	*6	*	*6	*	*5	*	*
577.9	*9	*	*	—	*	*	*7	*	*	*	*
578	464	227	237	11	47	109	298	97	123	156	89
578.0	63	33	30	*	13	22	24	15	12	17	19
578.1	116	56	59	*	15	29	67	28	26	41	21
578.9	286	138	148	*	19	58	206	54	85	98	49
579	52	24	29	11	*9	15	18	10	12	19	11
579.0	13	*	*8	*	*	*	*8	*	*	*	*
579.3	19	*8	11	*6	*	*7	*	*	*5	*8	*
579.8	12	*7	*6	*	*	*	*	*	*	*	*
579.9	*5	*	*	*	*	*	*	*	*	*	*
580–629	9,073	3,339	5,734	204	2,065	2,209	4,595	1,829	2,125	3,474	1,644
580	*7	*	*5	*	*	*	*	*	*	*	*
581	31	16	15	*	*8	12	*7	*6	*6	11	*8
581.8	14	*7	*7	*	*	*7	*	*	*	*	*
581.81	14	*7	*7	*	*	*7	*	*	*	*	*
581.9	13	*6	*7	*	*	*	*	*	*	*6	*
582	15	*7	*8	*	*5	*6	*	*	*	*5	*
582.8	*6	*	*	*	*	*	*	*	*	*	*
583	136	66	70	*	25	51	58	28	31	39	38
583.8	120	59	62	*	19	45	55	27	30	32	32
583.81	114	56	58	*	17	42	53	24	28	31	31
583.89	*6	*	*	*	*	*	*	*	*	*	*
583.9	12	*6	*7	*	*	*	*	*	*	*5	*5
584	1,101	585	517	*5	88	277	732	243	245	406	208
584.5	82	41	41	*	*7	30	44	20	18	30	14
584.8	*7	*	*	*	*	*	*	*	*	*	*
584.9	1,012	538	474	*	79	244	685	221	225	374	192
585	416	235	181	*	35	101	277	104	120	129	63
586	38	17	20	*	*	*	28	*5	16	12	*
588	21	9	12	*	*	*7	*7	*	*5	9	*
588.8	14	*6	*8	*	*	*	*	*	*	*7	*
588.81	10	*5	*	*	*	*	*	*	*	*	*
590	259	45	214	38	104	53	64	43	56	98	62
590.0	*6	*	*6	*	*	*	*	*	*	*	*
590.00	*6	*	*6	*	*	*	*	*	*	*	*
590.1	104	16	88	13	44	23	25	17	24	35	28
590.10	104	16	88	13	44	23	25	17	24	35	28
590.8	144	27	116	24	55	27	37	23	28	60	33
590.80	144	27	116	24	55	27	37	23	28	59	33

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
591	228	97	131	10	71	65	83	49	50	92	36
592	324	160	165	*6	122	113	84	75	82	128	38
592.0	145	67	78	*	53	48	41	35	38	54	17
592.1	175	91	84	*	67	64	42	39	44	72	21
593	811	415	396	26	64	151	569	163	187	306	154
593.2	19	12	*7	*	*	*	12	*	*	*8	*
593.3	14	*6	*8	*	*5	*	*	*	*	*7	*
593.4	25	10	15	*	*5	*8	10	*6	*5	9	*
593.5	14	*6	*7	*	*	*	*	*	*	*5	*
593.7	22	*7	15	18	*	*	*	*	*	*6	*7
593.70	17	*7	10	15	*	*	*	*	*	*	*6
593.8	24	*8	16	*	*8	*	10	*	*5	10	*
593.81	9	*	*5	*	*	*	*6	*	*	*	*
593.89	15	*	11	*	*6	*	*	*	*	*6	*
593.9	691	364	327	*	37	124	527	138	165	259	129
594	15	11	*	*	*	*	12	*	*	*	*
594.1	13	11	*	*	*	*	11	*	*	*	*
595	81	26	55	*	11	16	52	19	19	30	13
595.0	16	*6	10	*	*	*	13	*	*6	*6	*
595.1	13	*	13	*	*	*	*	*	*	*	*
595.2	9	*	*	*	*	*	*6	*	*	*	*
595.8	*8	*5	*	—	*	*	*6	*	*	*	*
595.82	*5	*	*	—	*	*	*	*	*	*	*
595.9	35	9	25	*	*	*7	22	*8	*	14	*8
596	181	108	73	10	25	36	109	38	46	64	33
596.0	37	35	*	*	*	*	32	*7	10	13	*7
596.1	11	*6	*	*	*	*	*7	*	*	*	*
596.3	*7	*5	*	*	*	*	*	*	*	*	*
596.4	10	*6	*	*	*	*	*7	*	*	*6	*
596.5	80	36	44	*8	14	18	40	17	24	24	15
596.51	13	*	11	—	*	*	9	*	*6	*	*
596.54	60	31	29	*8	11	14	28	11	17	18	13
596.59	*5	*	*	*	*	*	*	*	*	*	*
596.7	*6	*5	*	—	—	*	*	*	*	*	*
596.8	25	12	13	*	*6	*6	12	*5	*	10	*
597	*7	*	*	*	*	*	*	*	*	*	*
597.8	*7	*	*	*	*	*	*	*	*	*	*
597.80	*6	*	*	*	*	*	*	*	*	*	*
598	29	22	*7	*	*	*	19	*5	*6	13	*
598.9	22	16	*6	*	*	*	14	*	*	11	*
599	2,602	869	1,733	55	288	475	1,783	542	623	976	460
599.0	2,230	643	1,586	47	238	389	1,556	465	535	840	390
599.6	32	24	*8	*	*	*8	21	*6	*5	12	*8
599.7	311	193	118	*8	45	64	194	67	78	112	53
599.8	18	*	15	*	*	11	*	*	*	*6	*
599.81	*9	—	*9	—	*	*7	*	*	*	*	*
599.82	*5	—	*5	—	*	*	*	*	*	*	*
600	502	502	—	*	*	84	417	136	114	152	101
600.0	461	461	—	*	*	75	385	128	104	135	94
600.00	339	339	—	*	*	58	280	98	77	99	64
600.01	122	122	—	—	*	17	105	30	27	36	30
600.9	35	35	—	—	—	*7	27	*5	*7	16	*6
600.90	24	24	—	—	—	*	19	*	*5	10	*
600.91	11	11	—	—	—	*	*8	*	*	*6	*
601	41	41	—	—	*	11	29	11	*8	18	*
601.0	11	11	—	—	*	*	*8	*	*	*5	*
601.1	11	11	—	—	*	*	*8	*	*	*	*
601.9	17	17	—	—	*	*	12	*	*	*8	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
602	9	9	—	*	*	*	*7	*	*	*	*
603	10	10	—	*	*	*	*	*	*	*	*
603.9	*7	*7	—	*	*	*	*	*	*	*	*
604	19	19	—	*	*5	*7	*6	*	*7	*6	*
604.9	18	18	—	*	*	*7	*6	*	*7	*5	*
604.90	13	13	—	*	*	*	*	*	*5	*	*
604.99	*6	*6	—	*	*	*	*	*	*	*	*
605	*7	*7	—	*	*	*	*	*	*	*	*
607	19	19	—	*	*	*7	*7	*	*	*7	*6
607.8	10	10	—	*	*	*	*	*	*	*	*
607.84	*7	*7	—	—	*	*	*	*	*	*	*
608	31	31	—	*	11	*9	*8	*5	*	12	*9
608.4	12	12	—	*	*	*	*	*	*	*	*
608.8	9	9	—	*	*	*	*	*	*	*	*
610	24	*	23	*	*6	12	*5	*5	*	12	*
610.1	17	—	17	—	*	*8	*	*	*	*9	*
611	68	*5	63	*	29	24	14	19	9	32	*9
611.0	18	*	17	*	9	*5	*	*7	*	*6	*
611.1	15	*	12	*	*8	*6	*	*	*	*7	*
611.7	22	*	21	*	*7	*7	*8	*	*	10	*
611.72	15	*	14	*	*	*5	*7	*	*	*6	*
611.8	10	—	10	—	*	*	*	*	*	*6	*
614	236	—	236	*	166	61	*7	36	51	110	39
614.1	22	—	22	*	10	10	*	*	*	12	*
614.2	11	—	11	*	*8	*	*	*	*	*	*
614.3	*6	—	*6	*	*5	*	*	*	*	*	*
614.4	*7	—	*7	*	*	*	*	*	*	*	*
614.6	165	—	165	*	115	44	*5	22	38	78	26
614.9	20	—	20	*	18	*	*	*	*	*9	*
615	25	—	25	*	23	*	*	*	*5	10	*6
615.9	19	—	19	—	18	*	*	*	*	*8	*
616	109	—	109	*	59	37	*9	23	23	45	17
616.0	72	—	72	—	38	30	*	17	14	30	11
616.1	23	—	23	*	14	*5	*	*	*7	*8	*
616.10	22	—	22	*	14	*5	*	*	*7	*8	*
616.4	*8	—	*8	*	*	*	*	*	*	*	*
617	241	—	241	*	161	76	*	34	60	100	47
617.0	125	—	125	—	72	49	*	21	28	55	21
617.1	39	—	39	*	27	11	*	*	12	12	10
617.2	*8	—	*8	—	*6	*	*	*	*	*	*
617.3	37	—	37	—	28	*8	*	*	*8	17	*9
617.8	*8	—	*8	—	*7	*	—	*	*	*	*
617.9	22	—	22	*	18	*	*	*	*7	*8	*
618	257	—	257	*	48	116	92	38	55	107	56
618.0	78	—	78	—	11	36	31	11	17	31	18
618.01	40	—	40	—	*6	17	17	*6	9	15	9
618.04	29	—	29	—	*	14	11	*	*5	13	*6
618.1	23	—	23	*	*8	10	*	*	*	9	*
618.2	26	—	26	—	*7	13	*6	*	*	10	*8
618.3	16	—	16	—	*	*7	*7	*	*	*	*
618.4	27	—	27	—	*7	14	*6	*	*8	11	*6
618.5	36	—	36	—	*	16	20	*	*8	18	*6
618.6	31	—	31	—	*7	15	*9	*6	*	15	*5
618.8	19	—	19	—	*5	*5	*8	*	*5	*8	*
618.89	19	—	19	—	*5	*5	*8	*	*5	*8	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
619	*8	—	*8	*	*	*	*	*	*	*	*
619.1	*5	—	*5	—	*	*	*	*	*	*	*
620	310	—	310	*8	195	95	12	50	67	137	56
620.0	47	—	47	*	26	17	—	*7	*7	28	*
620.1	50	—	50	*	34	15	—	*9	12	24	*6
620.2	147	—	147	*	99	39	*7	21	34	63	29
620.5	*7	—	*7	*	*	*	*	*	*	*	*
620.8	50	—	50	*	29	17	*	10	11	16	12
620.9	*5	—	*5	—	*	*	*	*	*	*	*
621	73	—	73	*	33	29	10	13	16	31	12
621.0	24	—	24	—	*8	12	*	*5	*6	9	*
621.2	12	—	12	*	*9	*	*	*	*	*7	*
621.3	19	—	19	—	*	11	*	*	*	*7	*
621.30	*7	—	*7	—	*	*	*	*	*	*	*
621.33	*7	—	*7	—	*	*6	*	*	*	*	*
621.8	11	—	11	*	*6	*	*	*	*	*	*
622	36	—	36	*	19	14	*	*8	*7	17	*
622.1	22	—	22	—	13	*8	*	*5	*	12	*
622.10	14	—	14	—	*8	*6	*	*	*	*9	*
622.7	*7	—	*7	—	*	*	*	*	*	*	*
623	26	—	26	*	14	*6	*	*7	*	10	*
623.8	19	—	19	*	10	*	*	*5	*	*6	*
625	345	—	345	*	199	104	39	37	81	157	71
625.0	26	—	26	—	21	*	*	*	*	16	*
625.3	87	—	87	*	66	21	*	*6	19	46	15
625.6	130	—	130	—	39	57	34	17	27	50	35
625.8	12	—	12	*	*5	*	*	*	*	*	*
625.9	82	—	82	*	62	15	*	*9	25	38	10
626	311	—	311	*	205	104	*	40	74	141	55
626.0	*6	—	*6	*	*5	*	—	*	*	*	*
626.2	229	—	229	*	149	79	*	27	57	106	40
626.4	*7	—	*7	*	*6	*	—	*	*	*	*
626.6	*5	—	*5	—	*	*	*	*	*	*	*
626.8	53	—	53	*	32	19	*	*8	11	23	10
626.9	*9	—	*9	—	*6	*	—	*	*	*	*
627	48	—	48	*	*6	26	16	*8	14	19	*7
627.0	*6	—	*6	—	*	*	—	*	*	*	*
627.1	22	—	22	—	*	11	11	*	*6	*8	*
627.2	*7	—	*7	—	*	*	*	*	*	*	*
627.9	*8	—	*8	—	*	*5	*	*	*	*	*
628	*5	—	*5	*	*	*	—	*	*	*	*
630–677	10,394	—	10,394	19	10,353	22	—	1,539	2,153	4,080	2,621
632	22	—	22	*	22	*	—	*	*	12	*
633	34	—	34	—	34	—	—	*5	*6	16	*7
633.1	27	—	27	—	27	—	—	*	*	13	*6
633.10	26	—	26	—	26	—	—	*	*	13	*6
634	17	—	17	*	17	*	—	*	*	9	*
634.9	11	—	11	*	11	*	—	*	*	*7	*
634.91	*8	—	*8	*	*8	*	—	*	*	*5	*
635	*5	—	*5	*	*	*	—	*	*	*	*
639	*8	—	*8	—	*8	—	—	*	*	*	*
640	12	—	12	—	12	—	—	*	*	*7	*
640.0	*6	—	*6	—	*6	—	—	*	*	*	*
640.03	*5	—	*5	—	*5	—	—	*	*	*	*
640.9	*6	—	*6	—	*6	—	—	*	*	*	*
641	98	—	98	*	97	*	—	12	22	35	29
641.0	9	—	9	—	9	—	—	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
641.01	*9	—	*9	—	*9	—	—	*	*	*	*
641.1	26	—	26	—	25	*	—	*	*6	10	*7
641.11	18	—	18	—	17	*	—	*	*5	*7	*
641.13	*8	—	*8	—	*8	—	—	*	*	*	*
641.2	41	—	41	*	41	*	—	*6	10	12	13
641.21	37	—	37	*	37	*	—	*	9	12	12
641.9	16	—	16	—	16	—	—	*	*	*7	*
641.91	9	—	9	—	9	—	—	*	*	*	*
641.93	*7	—	*7	—	*7	—	—	*	*	*	*
642	406	—	406	*	405	*	—	46	75	188	97
642.0	48	—	48	—	48	*	—	*	9	24	10
642.01	41	—	41	—	41	*	—	*	*8	21	*9
642.03	*6	—	*6	—	*6	*	—	*	*	*	*
642.3	170	—	170	*	170	*	—	17	32	75	46
642.31	152	—	152	*	152	*	—	15	29	67	41
642.33	12	—	12	—	12	—	—	*	*	*5	*
642.34	*5	—	*5	—	*5	—	—	*	*	*	*
642.4	92	—	92	*	92	*	—	13	19	38	21
642.41	82	—	82	*	82	*	—	11	18	34	19
642.43	*8	—	*8	—	*8	—	—	*	*	*	*
642.5	38	—	38	*	38	*	—	*	*6	18	9
642.51	35	—	35	*	35	*	—	*	*5	16	*9
642.7	21	—	21	—	21	*	—	*	*	13	*
642.71	15	—	15	—	15	*	—	*	*	*8	*
642.9	27	—	27	—	27	—	—	*	*	15	*5
642.91	23	—	23	—	23	—	—	*	*	13	*
643	49	—	49	*	49	—	—	*6	9	26	*9
643.0	21	—	21	—	21	—	—	*	*	11	*5
643.03	21	—	21	—	21	—	—	*	*	11	*5
643.1	15	—	15	*	15	—	—	*	*	9	*
643.13	15	—	15	*	15	—	—	*	*	9	*
643.2	10	—	10	—	10	—	—	*	*	*5	*
643.23	*8	—	*8	—	*8	—	—	*	*	*	*
644	424	—	424	*	423	*	—	62	77	191	94
644.0	112	—	112	*	112	*	—	14	22	51	24
644.03	112	—	112	*	112	*	—	14	22	51	24
644.1	16	—	16	—	16	—	—	*	*	*8	*
644.13	16	—	16	—	16	—	—	*	*	*8	*
644.2	297	—	297	*	296	*	—	45	53	133	65
644.21	297	—	297	*	296	*	—	45	53	133	65
645	365	—	365	*	364	*	—	69	74	117	105
645.1	356	—	356	*	355	*	—	67	72	115	102
645.11	352	—	352	*	351	*	—	66	71	114	102
645.2	*9	—	*9	—	*9	—	—	*	*	*	*
645.21	*9	—	*9	—	*9	—	—	*	*	*	*
646	246	—	246	*	245	*	—	31	41	113	61
646.1	27	—	27	—	27	*	—	*	*	*7	13
646.11	20	—	20	—	20	*	—	*	*	*6	10
646.13	*5	—	*5	—	*5	—	—	*	*	*	*
646.2	19	—	19	—	19	*	—	*	*	10	*
646.21	*6	—	*6	—	*6	*	—	*	*	*	*
646.23	10	—	10	—	10	—	—	*	*	*6	*
646.3	*8	—	*8	—	*8	—	—	*	*	*	*
646.31	*8	—	*8	—	*8	—	—	*	*	*	*
646.6	117	—	117	*	117	*	—	17	17	59	24
646.61	43	—	43	—	43	*	—	*8	*7	20	*8
646.62	*5	—	*5	—	*5	—	—	*	*	*	*
646.63	63	—	63	*	63	—	—	*7	*8	36	12
646.64	*5	—	*5	—	*5	—	—	*	*	*	*
646.8	67	—	67	*	67	—	—	*7	12	30	18

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
646.81	40	—	40	—	40	—	—	* ⁹	* ⁹	16	12
646.83	23	—	23	*	23	—	—	*	*	13	*
647	102	—	102	—	102	*	—	11	17	53	21
647.6	59	—	59	—	59	*	—	* ⁶	9	30	13
647.61	52	—	52	—	52	*	—	*	* ⁸	27	12
647.8	36	—	36	—	36	—	—	*	* ⁷	20	* ⁶
647.81	22	—	22	—	22	—	—	*	* ⁶	12	*
647.83	11	—	11	—	11	—	—	*	*	* ⁶	*
648	1,865	—	1,865	*	1,858	*	—	292	399	747	426
648.0	48	—	48	*	48	*	—	* ⁶	* ⁷	25	10
648.01	32	—	32	—	32	*	—	*	*	15	* ⁸
648.03	14	—	14	*	14	—	—	*	*	9	*
648.1	73	—	73	*	73	*	—	13	20	25	15
648.11	65	—	65	—	65	*	—	12	19	21	14
648.13	* ⁷	—	* ⁷	*	* ⁷	—	—	*	*	*	*
648.2	319	—	319	*	317	*	—	50	62	132	74
648.21	190	—	190	*	189	*	—	31	33	82	44
648.22	98	—	98	*	97	—	—	15	23	37	23
648.23	21	—	21	—	21	—	—	*	*	9	*
648.24	* ⁹	—	* ⁹	—	* ⁹	—	—	*	*	*	*
648.3	11	—	11	—	11	—	—	*	*	*	*
648.31	* ⁶	—	* ⁶	—	* ⁶	—	—	*	*	*	*
648.4	233	—	233	*	232	*	—	40	65	80	49
648.41	186	—	186	—	185	*	—	31	52	62	42
648.43	36	—	36	*	36	—	—	* ⁶	* ⁹	15	* ⁶
648.44	10	—	10	—	10	—	—	*	*	*	*
648.6	46	—	46	—	46	*	—	* ⁹	10	21	* ⁶
648.61	36	—	36	—	36	*	—	* ⁷	* ⁹	18	*
648.7	16	—	16	—	16	—	—	*	*	9	*
648.71	* ⁸	—	* ⁸	—	* ⁸	—	—	*	*	*	*
648.73	* ⁷	—	* ⁷	—	* ⁷	—	—	*	*	* ⁵	*
648.8	182	—	182	*	181	*	—	30	34	65	53
648.81	165	—	165	*	164	*	—	26	30	58	51
648.83	15	—	15	*	15	*	—	*	*	* ⁶	*
648.9	934	—	934	*	931	*	—	140	200	383	211
648.91	791	—	791	*	788	*	—	113	170	326	182
648.92	19	—	19	—	19	—	—	*	* ⁶	* ⁶	*
648.93	116	—	116	*	116	*	—	21	23	48	24
648.94	* ⁸	—	* ⁸	—	* ⁸	—	—	*	*	*	*
650	310	—	310	*	310	—	—	36	59	126	90
651	72	—	72	*	72	*	—	11	12	34	16
651.0	66	—	66	*	65	*	—	10	* ⁹	33	15
651.01	47	—	47	—	47	*	—	* ⁷	* ⁶	24	10
651.03	19	—	19	*	19	*	—	*	*	* ⁹	*
652	304	—	304	*	303	*	—	50	57	123	75
652.0	* ⁶	—	* ⁶	—	* ⁶	—	—	*	*	*	*
652.01	* ⁶	—	* ⁶	—	* ⁶	—	—	*	*	*	*
652.1	* ⁷	—	* ⁷	—	* ⁷	—	—	*	*	*	*
652.11	* ⁷	—	* ⁷	—	* ⁷	—	—	*	*	*	*
652.2	127	—	127	*	126	*	—	20	23	52	33
652.21	123	—	123	*	122	*	—	19	23	49	32
652.3	21	—	21	—	21	—	—	*	*	* ⁸	* ⁵
652.31	20	—	20	—	20	—	—	*	*	* ⁸	* ⁵
652.4	* ⁷	—	* ⁷	—	* ⁷	—	—	*	*	*	*
652.41	* ⁷	—	* ⁷	—	* ⁷	—	—	*	*	*	*
652.5	58	—	58	—	58	—	—	10	11	24	12
652.51	58	—	58	—	58	—	—	10	11	24	12
652.6	20	—	20	—	19	*	—	* ⁶	*	* ⁷	*
652.61	19	—	19	—	19	*	—	* ⁵	*	* ⁷	*
652.8	54	—	54	—	54	—	—	* ⁹	9	21	15
652.81	54	—	54	—	54	—	—	* ⁹	9	21	15

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
653	105	—	105	—	105	—	—	12	20	45	27
653.4	87	—	87	—	87	—	—	10	18	38	21
653.41	87	—	87	—	87	—	—	10	18	38	21
653.5	11	—	11	—	11	—	—	*	*	*	*5
653.51	10	—	10	—	10	—	—	*	*	*	*5
654	730	—	730	*	728	*	—	104	139	314	173
654.0	13	—	13	—	13	*	—	*	*	*5	*
654.01	12	—	12	—	12	*	—	*	*	*	*
654.1	31	—	31	—	30	*	—	*6	*	13	*7
654.11	27	—	27	—	27	*	—	*5	*	12	*6
654.2	587	—	587	—	586	*	—	83	112	247	145
654.21	571	—	571	—	570	*	—	82	110	238	140
654.23	16	—	16	—	16	*	—	*	*	9	*
654.4	34	—	34	*	33	*	—	*5	*6	15	*7
654.41	23	—	23	*	22	*	—	*	*	*8	*6
654.43	9	—	9	—	*9	*	—	*	*	*7	*
654.5	35	—	35	*	35	*	—	*	*	20	*6
654.51	20	—	20	—	20	*	—	*	*	10	*
654.53	15	—	15	*	15	*	—	*	*	10	*
654.6	20	—	20	—	20	—	—	*	*5	10	*
654.61	15	—	15	—	15	—	—	*	*	*8	*
654.63	*5	—	*5	—	*5	—	—	*	*	*	*
654.9	*8	—	*8	—	*8	*	—	*	*	*	*
654.91	*8	—	*8	—	*7	*	—	*	*	*	*
655	51	—	51	*	51	—	—	*	11	24	12
655.7	22	—	22	*	22	—	—	*	*6	10	*
655.71	17	—	17	*	17	—	—	*	*5	*6	*
655.73	*5	—	*5	—	*5	—	—	*	*	*	*
655.8	19	—	19	—	19	—	—	*	*	*7	*6
655.81	17	—	17	—	17	—	—	*	*	*6	*6
656	379	—	379	*	378	*	—	53	74	163	89
656.1	76	—	76	—	76	*	—	12	15	36	13
656.11	74	—	74	—	74	*	—	12	15	35	12
656.3	17	—	17	—	17	—	—	*	*	10	*
656.31	17	—	17	—	17	—	—	*	*	10	*
656.4	18	—	18	—	18	—	—	*	*	*5	*7
656.41	17	—	17	—	17	—	—	*	*	*5	*7
656.5	76	—	76	*	76	—	—	12	13	36	15
656.51	73	—	73	*	72	—	—	11	12	35	15
656.6	101	—	101	—	100	*	—	12	23	39	26
656.61	99	—	99	—	98	*	—	12	23	39	25
656.7	16	—	16	—	16	—	—	*	*	*5	*
656.71	15	—	15	—	15	—	—	*	*	*	*
656.8	73	—	73	*	73	*	—	10	12	30	22
656.81	73	—	73	*	73	*	—	10	12	29	22
657	25	—	25	—	25	—	—	*6	*	12	*
657.0	25	—	25	—	25	—	—	*6	*	12	*
657.01	22	—	22	—	22	—	—	*	*	10	*
658	361	—	361	*	361	*	—	63	67	136	96
658.0	113	—	113	*	112	*	—	22	18	47	27
658.01	108	—	108	*	107	*	—	21	17	44	25
658.03	*5	—	*5	—	*5	—	—	*	*	*	*
658.1	137	—	137	*	137	*	—	26	28	45	39
658.11	132	—	132	*	132	*	—	26	26	43	37
658.13	*5	—	*5	—	*5	—	—	*	*	*	*
658.2	53	—	53	—	53	*	—	9	14	19	11
658.21	50	—	50	—	50	*	—	*9	14	17	10
658.4	55	—	55	*	55	—	—	*6	*7	23	20
658.41	54	—	54	*	54	—	—	*6	*6	23	19

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
659	1,097	—	1,097	*5	1,086	*	—	168	228	411	289
659.1	56	—	56	—	56	*	—	*7	14	20	15
659.11	49	—	49	—	49	*	—	*6	12	19	12
659.13	*7	—	*7	—	*7	—	—	*	*	*	*
659.2	54	—	54	*	54	*	—	*7	14	19	15
659.21	54	—	54	*	54	*	—	*7	14	18	15
659.4	24	—	24	—	24	—	—	*	*	11	*7
659.41	22	—	22	—	22	—	—	*	*	9	*7
659.5	51	—	51	—	51	*	—	11	9	17	15
659.51	48	—	48	—	48	*	—	11	*8	16	13
659.6	314	—	314	—	309	*	—	53	62	113	86
659.61	300	—	300	—	296	*	—	50	59	106	84
659.63	14	—	14	—	13	*	—	*	*	*6	*
659.7	531	—	531	*	530	*	—	80	113	211	128
659.71	524	—	524	*	523	*	—	78	111	208	126
659.73	*7	—	*7	—	*7	—	—	*	*	*	*
659.8	53	—	53	*	48	*	—	*8	12	20	14
659.81	52	—	52	*	47	*	—	*8	11	19	14
659.9	11	—	11	—	11	—	—	*	*	*	*9
659.91	11	—	11	—	11	—	—	*	*	*	*9
660	194	—	194	—	194	—	—	25	48	68	52
660.0	40	—	40	—	40	—	—	*8	10	14	*8
660.01	40	—	40	—	40	—	—	*8	10	14	*8
660.1	35	—	35	—	35	—	—	*	*7	13	12
660.11	35	—	35	—	35	—	—	*	*7	13	12
660.2	*5	—	*5	—	*5	—	—	*	*	*	*
660.21	*5	—	*5	—	*5	—	—	*	*	*	*
660.3	40	—	40	—	40	—	—	*6	*7	14	12
660.31	40	—	40	—	40	—	—	*6	*7	14	12
660.4	52	—	52	—	52	—	—	*	19	18	11
660.41	52	—	52	—	52	—	—	*	19	18	11
660.6	*8	—	*8	—	*8	—	—	*	*	*	*
660.61	*7	—	*7	—	*7	—	—	*	*	*	*
660.7	11	—	11	—	11	—	—	*	*	*	*
660.71	11	—	11	—	11	—	—	*	*	*	*
661	380	—	380	*	379	*	—	60	78	148	94
661.0	102	—	102	—	102	*	—	13	25	41	24
661.01	102	—	102	—	102	*	—	13	25	40	24
661.1	87	—	87	*	87	*	—	16	15	36	20
661.11	87	—	87	*	87	*	—	16	15	36	20
661.2	91	—	91	—	91	—	—	17	17	39	17
661.21	90	—	90	—	90	—	—	17	17	39	17
661.3	87	—	87	*	86	—	—	10	19	28	30
661.31	87	—	87	*	86	—	—	10	19	28	30
661.4	*6	—	*6	—	*6	—	—	*	*	*	*
661.41	*5	—	*5	—	*5	—	—	*	*	*	*
661.9	*6	—	*6	—	*6	—	—	*	*	*	*
661.91	*6	—	*6	—	*6	—	—	*	*	*	*
662	35	—	35	*	34	*	—	*6	*8	*8	12
662.0	*6	—	*6	—	*6	—	—	*	*	*	*
662.01	*6	—	*6	—	*6	—	—	*	*	*	*
662.1	*6	—	*6	—	*6	*	—	*	*	*	*
662.11	*6	—	*6	—	*5	*	—	*	*	*	*
662.2	23	—	23	*	23	—	—	*	*5	*5	*8
662.21	23	—	23	*	23	—	—	*	*5	*5	*8
663	833	—	833	*	831	*	—	128	185	298	221
663.0	11	—	11	—	11	—	—	*	*	*	*
663.01	10	—	10	—	10	—	—	*	*	*	*
663.1	129	—	129	*	129	—	—	16	31	44	38

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
663.11	129	—	129	*	129	—	—	16	31	44	38
663.2	43	—	43	—	43	*	—	*7	11	14	11
663.21	43	—	43	—	43	*	—	*7	11	14	11
663.3	610	—	610	*	609	*	—	98	130	222	160
663.31	610	—	610	*	608	*	—	98	130	222	160
663.4	12	—	12	—	12	—	—	*	*	*	*
663.41	12	—	12	—	12	—	—	*	*	*	*
663.8	26	—	26	—	26	—	—	*	*8	*9	*6
663.81	26	—	26	—	26	—	—	*	*8	*9	*6
664	1,259	—	1,259	*	1,254	*	—	195	290	431	343
664.0	556	—	556	*	555	*	—	85	123	193	155
664.01	556	—	556	*	554	*	—	85	123	192	155
664.1	560	—	560	*	557	*	—	87	133	185	155
664.11	559	—	559	*	556	*	—	87	133	185	155
664.2	86	—	86	*	86	—	—	15	21	31	19
664.21	86	—	86	*	86	—	—	15	21	31	19
664.3	31	—	31	—	31	—	—	*	*6	12	*9
664.31	31	—	31	—	31	—	—	*	*6	12	*9
664.4	13	—	13	—	13	—	—	*	*	*6	*
664.41	11	—	11	—	11	—	—	*	*	*5	*
664.8	*9	—	*9	—	*9	—	—	*	*	*	*
664.81	*9	—	*9	—	*9	—	—	*	*	*	*
665	176	—	176	*	176	—	—	17	53	65	42
665.3	*6	—	*6	—	*6	—	—	*	*	*	*
665.31	*6	—	*6	—	*6	—	—	*	*	*	*
665.4	53	—	53	*	53	—	—	*	16	19	15
665.41	53	—	53	*	53	—	—	*	16	19	15
665.5	104	—	104	*	103	—	—	11	32	39	22
665.51	102	—	102	*	102	—	—	10	32	38	22
666	120	—	120	*	119	*	—	12	30	39	38
666.0	*7	—	*7	—	*7	*	—	*	*	*	*
666.02	*6	—	*6	—	*6	*	—	*	*	*	*
666.1	93	—	93	*	93	*	—	9	25	27	31
666.12	93	—	93	—	93	*	—	9	25	27	31
666.2	13	—	13	—	13	—	—	*	*	*6	*
666.22	*8	—	*8	—	*8	—	—	*	*	*	*
666.3	*7	—	*7	—	*7	—	—	*	*	*	*
666.32	*6	—	*6	—	*6	—	—	*	*	*	*
667	20	—	20	—	20	—	—	*	*	*5	*6
667.0	12	—	12	—	12	—	—	*	*	*	*
667.02	12	—	12	—	12	—	—	*	*	*	*
667.1	*7	—	*7	—	*7	—	—	*	*	*	*
667.12	*6	—	*6	—	*6	—	—	*	*	*	*
668	15	—	15	—	14	*	—	*	*	*	*
668.8	14	—	14	—	13	*	—	*	*	*	*
668.82	10	—	10	—	*9	*	—	*	*	*	*
669	140	—	140	*	140	*	—	16	26	56	42
669.2	*7	—	*7	—	*7	—	—	*	*	*	*
669.4	*6	—	*6	—	*6	*	—	*	*	*	*
669.5	61	—	61	*	61	—	—	*5	10	35	12
669.51	61	—	61	*	61	—	—	*5	10	35	12
669.7	*7	—	*7	*	*6	—	—	*	*	*	*
669.71	*7	—	*7	*	*6	—	—	*	*	*	*
669.8	58	—	58	*	57	—	—	*7	12	15	24
669.81	52	—	52	*	52	—	—	*6	9	14	22
669.82	*5	—	*5	—	*5	—	—	*	*	*	*
670	23	—	23	*	23	—	—	*6	*	*7	*6
670.0	23	—	23	*	23	—	—	*6	*	*7	*6
670.02	13	—	13	*	13	—	—	*	*	*5	*
670.04	10	—	10	*	10	—	—	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
671	27	—	27	—	27	—	—	*5	*6	*9	*6
671.8	14	—	14	—	14	—	—	*	*	*	*
671.81	*6	—	*6	—	*6	—	—	*	*	*	*
671.82	*6	—	*6	—	*6	—	—	*	*	*	*
672	18	—	18	—	18	—	—	*	*	*8	*
672.0	18	—	18	—	18	—	—	*	*	*8	*
672.02	17	—	17	—	17	—	—	*	*	*8	*
674	46	—	46	—	46	*	—	*6	11	18	12
674.3	18	—	18	—	18	*	—	*	*6	*5	*
674.32	11	—	11	—	11	*	—	*	*	*	*
674.34	*6	—	*6	—	*6	*	—	*	*	*	*
674.8	21	—	21	—	21	—	—	*	*	10	*6
674.82	15	—	15	—	15	—	—	*	*	*8	*
674.84	*6	—	*6	—	*6	—	—	*	*	*	*
676	*6	—	*6	—	*6	—	—	*	*	*	*
680–709	2,632	1,323	1,309	218	502	764	1,149	587	594	975	477
680	12	*8	*	*	*6	*	*	*	*	*5	*
681	73	43	30	*	17	30	21	18	25	23	*9
681.0	31	19	12	*	14	*8	*6	*8	*7	12	*
681.00	27	17	10	*	12	*7	*	*6	*7	10	*
681.1	42	24	18	*	*	21	16	10	17	11	*
681.10	39	24	16	*	*	20	15	9	17	10	*
682	1,101	580	521	79	248	359	415	245	252	406	198
682.0	59	27	32	14	21	14	11	10	10	24	15
682.1	20	11	9	*	*6	*6	*	*	*	*8	*
682.2	131	56	75	10	39	43	39	22	30	53	26
682.3	117	66	52	*5	40	36	36	25	24	47	21
682.4	53	30	23	*	20	14	15	12	13	18	10
682.5	37	16	21	10	11	*8	*7	*	10	18	*
682.6	518	278	240	24	86	181	227	126	125	175	92
682.7	145	86	59	*5	22	49	69	36	35	54	21
682.8	9	*	*	*	*	*	*	*	*	*	*
682.9	11	*5	*6	*	*	*	*6	*	*	*	*
683	*6	*	*	*	*	*	*	*	*	*	*
684	*8	*	*5	*	*	*	*	*	*	*	*
685	*8	*	*	*	*	*	*	*	*	*	*
685.0	*5	*	*	*	*	*	*	*	*	*	*
686	17	*8	*9	*	*	*7	*	*	*	*5	*
686.9	*9	*5	*	*	*	*	*	*	*	*	*
690	*7	*	*	*	*	*	*	*	*	*	*
690.1	*7	*	*	*	*	*	*	*	*	*	*
690.10	*5	*	*	*	*	*	*	*	*	*	*
691	33	16	16	28	*	*	*	*	*6	13	10
691.0	25	13	12	25	*	*	*	*	*	10	*8
691.8	*7	*	*	*	*	*	*	*	*	*	*
692	75	33	42	22	19	14	21	20	20	25	10
692.9	59	31	29	21	12	12	15	16	14	21	*8
693	47	18	29	*	11	14	18	14	9	18	*6
693.0	45	18	27	*	11	13	18	14	*9	18	*
694	*8	*	*	*	*	*	*5	*	*	*	*
695	41	13	28	*	13	10	14	*8	10	11	12
695.1	10	*	*6	*	*	*	*	*	*	*	*
695.3	*7	*	*	—	*	*	*	*	*	*	*
695.4	*5	*	*	—	*	*	*	*	*	*	*
695.8	9	*	*8	*	*	*	*	*	*	*	*
695.89	9	*	*8	*	*	*	*	*	*	*	*
695.9	*6	*	*	*	*	*	*	*	*	*	*

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
696	59	31	29	*	10	28	20	17	15	14	13
696.0	16	*8	*9	*	*	11	*	*	*7	*	*
696.1	42	22	20	*	*8	17	16	14	*9	11	*9
698	38	11	28	*	16	12	*7	*9	11	11	*8
698.8	*6	*	*5	*	*	*	*	*	*	*	*
698.9	28	*9	19	*	11	*9	*5	*6	*8	*9	*6
701	32	*8	24	*5	14	9	*	*9	*6	13	*
701.4	*8	*	*6	*	*	*	*	*	*	*	*
701.8	*8	*	*7	*	*	*	*	*	*	*	*
701.9	*7	*	*5	*	*	*	*	*	*	*	*
702	12	*5	*7	*	*	*	9	*	*	*	*
702.0	*6	*	*	—	—	*	*5	*	*	*	*
702.1	*6	*	*	*	*	*	*	*	*	*	*
702.19	*6	*	*	*	*	*	*	*	*	*	*
703	9	*	*	*	*	*	*5	*	*	*	*
703.0	*6	*	*	*	*	*	*	*	*	*	*
704	15	*	10	*	*7	*	*	*	*	*	*
704.0	*6	*	*	*	*	*	*	*	*	*	*
704.8	*7	*	*	*	*	*	*	*	*	*	*
705	10	*	*6	*	*6	*	*	*	*	*	*
705.8	*8	*	*6	*	*5	*	*	*	*	*	*
705.83	*7	*	*	*	*	*	*	*	*	*	*
706	23	10	13	*	11	*6	*	*5	*	11	*
706.1	13	*	*8	*	*8	*	—	*	*	*	*
706.2	*8	*	*	*	*	*	*	*	*	*5	*
707	878	447	430	*	72	227	575	201	186	316	175
707.0	499	238	261	*	39	99	359	105	93	195	106
707.00	22	*9	14	—	*	*	17	*	*	11	*8
707.03	223	101	122	*	17	46	160	56	37	88	42
707.04	40	24	15	*	*6	*	29	*5	*7	12	16
707.05	72	33	39	*	*6	17	49	15	15	25	18
707.06	13	*7	*6	—	*	*	*7	*	*	*	*
707.07	71	31	40	*	*	*9	59	15	15	31	10
707.09	52	29	23	*	*	12	35	*9	14	20	9
707.1	355	198	158	*	29	120	206	90	88	111	66
707.10	41	17	25	*	*	17	21	11	14	12	*
707.11	*7	*5	*	—	*	*	*	*	*	*	*
707.12	21	9	12	—	*	*6	14	*	*7	*6	*
707.13	25	15	10	—	*	*	19	*	*	*8	*7
707.14	59	29	30	*	*	19	37	14	14	18	13
707.15	150	93	57	*	13	55	82	43	36	49	22
707.19	52	30	22	*	*	17	30	12	12	15	13
707.8	21	11	10	*	*	*7	9	*	*	9	*
708	22	*9	14	*5	*6	*6	*5	*7	*	*8	*
708.0	13	*6	*7	*	*	*	*	*	*	*	*
708.9	*8	*	*5	*	*	*	*	*	*	*	*
709	92	46	46	37	29	16	10	*7	13	67	*6
709.2	63	32	31	35	23	*	*	*	*	*5	55
709.8	11	*5	*5	*	*	*	*	*	*	*	*
709.9	11	*	*6	*	*	*	*	*	*	*6	*
710–739	6,884	2,457	4,427	101	979	2,089	3,716	1,457	1,858	2,270	1,299
710	198	25	173	*	63	71	60	31	41	85	40
710.0	137	15	122	*	54	49	32	20	27	64	27
710.1	26	*	23	*	*	*9	13	*	*8	*7	*6
710.2	15	*	13	—	*	*7	*5	*	*	*7	*
710.3	*7	*	*6	*	*	*	*	*	*	*	*
710.4	*9	*	*6	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
711	56	31	24	*	12	17	22	12	15	19	10
711.0	52	29	23	*	11	16	22	12	14	17	9
711.05	*7	*	*	*	*	*	*	*	*	*	*
711.06	27	17	10	*	*6	*7	13	*6	9	*8	*
711.07	*6	*	*	*	*	*	*	*	*	*	*
712	*9	*	*6	*	*	*	*6	*	*	*	*
712.3	*8	*	*5	*	*	*	*6	*	*	*	*
713	10	*	*6	*	*	*5	*	*	*	*	*
713.5	9	*	*	—	*	*	*	*	*	*	*
714	266	70	196	*	23	85	155	67	61	90	49
714.0	249	64	186	—	19	81	150	63	58	83	45
714.3	*8	*	*5	*	*	*	*	*	*	*	*
714.30	*7	*	*	*	*	*	*	*	*	*	*
715	1,809	637	1,171	*	66	500	1,243	392	510	586	321
715.0	23	*8	14	—	*	*6	14	*8	*	*9	*
715.09	18	*7	12	—	*	*5	11	*7	*	*6	*
715.1	69	32	37	*	*	19	48	19	*8	22	20
715.15	19	11	*8	—	*	*5	13	*6	*	*	*7
715.16	44	19	24	—	*	12	30	11	*6	14	12
715.2	*8	*	*	*	*	*	*	*	*	*	*
715.3	532	205	327	*	15	181	335	130	144	143	114
715.30	*7	*	*	—	*	*	*	*	*	*	*
715.31	26	11	15	—	*	*8	18	*6	*	*8	*8
715.35	138	63	75	—	*6	47	86	35	41	32	30
715.36	348	123	225	*	*8	121	219	84	96	99	70
715.37	*7	*	*	—	*	*	*	*	*	*	*
715.8	59	19	39	—	*	10	46	11	17	22	*8
715.89	58	19	39	—	*	10	46	11	17	22	*8
715.9	1,119	368	751	—	43	279	797	221	335	389	173
715.90	735	231	504	—	30	155	550	153	218	262	102
715.91	17	*8	*9	—	*	*	11	*	*5	*7	*
715.95	94	39	55	—	*5	27	61	19	31	21	23
715.96	240	79	161	—	*	84	152	39	72	88	40
715.97	*8	*5	*	—	*	*	*5	*	*	*	*
715.98	19	*6	14	—	*	*7	11	*8	*	*6	*
716	276	98	178	*	27	98	149	40	82	109	46
716.1	10	*7	*	—	*	*	*	*	*	*	*5
716.5	9	*5	*	*	*	*	*	*	*	*	*
716.9	252	83	169	*	21	90	140	36	77	101	38
716.90	191	64	127	*	15	65	110	25	65	73	27
716.91	10	*	*6	—	*	*	*5	*	*	*	*
716.95	*6	*	*	—	*	*	*	*	*	*	*
716.96	17	*	12	*	*	*7	*7	*	*	*7	*
716.98	*7	*	*	—	*	*	*	*	*	*	*
716.99	13	*	9	—	*	*	*8	*	*	*8	*
717	26	15	11	—	9	10	*7	*8	*	*8	*5
717.7	*8	*5	*	—	*	*	*	*	*	*	*
717.8	*6	*	*	—	*	*	*	*	*	*	*
718	48	24	24	*5	12	13	17	10	10	13	15
718.3	*6	*	*	*	*	*	*	*	*	*	*
718.4	20	10	10	*	*	*5	*9	*	*	*6	*7
718.5	*9	*6	*	*	*	*	*	*	*	*	*
718.56	*6	*	*	*	*	*	*	*	*	*	*
719	270	105	164	*9	54	72	135	57	96	65	51
719.0	24	14	11	*	*	*7	12	*5	9	*5	*
719.06	18	12	*6	*	*	*6	*8	*	*7	*	*
719.1	*8	*	*	*	*	*	*	*	*	*	*
719.16	*6	*	*	*	*	*	*	*	*	*	*
719.4	173	63	110	*	45	51	74	31	52	48	42

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
719.40	*6	*	*	*	*	*	*	*	*	*	*
719.41	48	22	26	*	9	18	20	9	14	13	11
719.45	52	12	39	*	10	*8	32	*8	19	15	10
719.46	43	17	25	*	14	14	14	*8	11	12	11
719.47	10	*	*6	*	*	*	*	*	*	*	*
719.49	*8	*	*	*	*	*	*	*	*	*	*
719.7	50	20	30	*	*	10	38	19	28	*	*
720	17	*9	*8	*	*	10	*	*	*	*6	*
720.0	10	*8	*	—	*	*6	*	*	*	*	*
720.2	*6	*	*5	*	*	*	*	*	*	*	*
721	305	132	174	*	43	110	152	53	88	104	61
721.0	77	33	44	—	13	33	31	11	19	29	18
721.1	33	20	12	—	*	17	14	*7	*	13	*9
721.2	*6	*	*	—	*	*	*	*	*	*	*
721.3	130	51	79	*	18	36	76	22	47	37	24
721.4	10	*6	*	—	*	*	*	*	*	*	*5
721.41	*5	*	*	—	*	*	*	*	*	*	*
721.8	*9	*6	*	*	*	*	*	*	*	*	*
721.9	39	11	27	—	*	13	22	10	12	13	*
721.90	38	11	27	—	*	13	22	10	12	13	*
722	560	258	302	*	166	247	147	109	147	204	101
722.0	69	34	35	—	26	36	*6	15	15	28	11
722.1	193	99	93	*	75	79	39	44	53	64	31
722.10	188	98	91	*	73	77	38	44	52	62	30
722.2	*8	*	*5	—	*	*	*	*	*	*	*
722.4	37	16	20	—	*7	20	10	*	9	13	11
722.5	147	56	91	*	29	59	59	21	48	50	28
722.51	*6	*	*	*	*	*	*	*	*	*	*
722.52	141	54	87	*	29	56	56	21	47	47	27
722.6	17	*6	11	—	*	*5	*8	*	*	*7	*
722.7	30	14	16	—	*8	15	*7	*6	*	16	*
722.71	18	*9	10	—	*	*9	*	*	*	*8	*
722.73	9	*	*	—	*	*	*	*	*	*6	*
722.8	17	9	*7	—	*	10	*	*	*	9	*
722.83	15	*9	*7	—	*	*9	*	*	*	*8	*
722.9	43	19	24	*	9	21	13	10	10	14	*9
722.91	12	*	*8	—	*	*7	*	*	*	*	*
722.93	26	12	15	*	*6	12	*9	*	*8	9	*
723	108	50	58	*	26	44	35	18	27	38	25
723.0	29	17	12	*	*	16	11	*5	*6	10	*8
723.1	49	22	27	*	18	16	13	*8	14	17	10
723.4	18	*7	11	*	*	9	*5	*	*	*7	*
723.5	*7	*	*	*	*	*	*	*	*	*	*
724	642	261	381	*	147	203	288	136	181	211	115
724.0	174	73	100	*	*7	39	127	48	45	50	30
724.00	35	13	22	—	*	*	31	12	11	*5	*6
724.02	136	59	77	*	*6	36	94	35	34	44	23
724.2	205	89	116	*	60	76	68	33	61	69	41
724.3	25	*6	19	—	*7	*6	13	*7	*6	*7	*
724.4	33	15	17	*	*7	13	13	*7	10	13	*
724.5	180	71	110	*	56	63	59	36	48	64	32
724.6	*6	*	*5	*	*	*	*	*	*	*	*
724.8	12	*	*8	—	*5	*	*	*	*5	*	*
725	49	13	36	—	*	*7	42	12	13	15	*9
726	101	45	56	*	13	37	49	21	23	31	26
726.0	*7	*	*	—	*	*	*	*	*	*	*
726.1	29	13	16	—	*	9	17	*6	*8	*6	*8
726.10	26	12	13	—	*	*9	16	*5	*7	*6	*8
726.2	17	*7	10	—	*	9	*7	*	*	*7	*
726.3	11	*9	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
726.33	10	*8	*	*	*	*	*	*	*	*	*
726.5	15	*	14	*	*	*	13	*	*	*8	*
726.6	12	*8	*	*	*	*	*	*	*	*	*
726.65	*6	*	*	*	*	*	*	*	*	*	*
726.9	*5	*	*	*	*	*	*	*	*	*	*
727	68	28	40	*	14	24	25	14	14	26	14
727.0	27	13	14	*	*8	*9	*7	*5	*6	10	*
727.00	13	*6	*7	*	*	*	*	*	*	*	*
727.05	10	*6	*	*	*	*	*	*	*	*	*
727.4	13	*	*8	*	*	*5	*6	*	*	*	*
727.40	11	*	*7	*	*	*	*6	*	*	*	*
727.5	*6	*	*	—	*	*	*	*	*	*	*
727.51	*6	*	*	—	*	*	*	*	*	*	*
727.6	9	*	*	*	*	*	*6	*	*	*5	*
727.61	*6	*	*	—	*	*	*	*	*	*	*
727.8	11	*	*6	*	*	*	*	*	*	*	*
727.81	*6	*	*	*	*	*	*	*	*	*	*
728	214	128	87	*7	53	52	102	47	47	72	48
728.2	*7	*	*	*	*	*	*6	*	*	*	*
728.3	*8	*	*	*	*	*	*	*	*	*5	*
728.8	182	114	68	*	46	45	87	39	40	61	42
728.85	21	11	10	*	*8	*	*7	*	*	*7	*7
728.86	10	*5	*	*	*	*	*	*	*	*	*
728.87	17	10	*8	*	*	*	10	*	*	*	*
728.88	118	78	40	*	28	29	62	26	29	40	24
728.89	14	9	*	*	*	*5	*6	*	*	*5	*
729	410	124	286	11	92	156	151	67	127	134	82
729.1	170	24	145	*	49	83	37	33	50	54	33
729.2	9	*	*6	*	*	*	*5	*	*	*	*
729.3	*5	*	*	*	*	*	*	*	*	*	*
729.5	95	33	63	*6	24	30	35	17	30	30	18
729.6	*6	*	*	*	*	*	*	*	*	*	*
729.8	119	56	63	*	11	36	70	14	38	40	26
729.81	27	11	16	*	*	9	12	*	11	*8	*
729.82	*6	*	*	*	*	*	*	*	*	*	*
729.89	86	43	43	*	*5	25	55	9	25	30	21
730	159	101	58	*7	28	62	61	39	35	48	36
730.0	36	24	12	*	*	16	14	12	*7	10	*6
730.07	20	13	*6	*	*	9	*8	*8	*	*6	*
730.08	*6	*	*	*	*	*	*	*	*	*	*
730.1	30	21	*8	*	*7	11	11	*7	*	9	9
730.15	*6	*	*	*	*	*	*	*	*	*	*
730.17	12	9	*	*	*	*6	*	*	*	*	*
730.2	92	55	36	*	17	35	36	20	22	28	21
730.25	*7	*	*	*	*	*	*	*	*	*	*
730.26	11	*7	*	*	*	*6	*	*	*	*	*
730.27	50	28	22	*	*5	22	22	11	12	17	11
730.28	14	*8	*6	*	*	*	*6	*	*	*	*
731	43	27	16	*	*	21	18	13	*8	14	*8
731.0	9	*	*	—	*	*	*9	*	*	*	*
731.8	34	23	11	—	*	21	10	11	*	12	*6
732	*8	*6	*	*	*	*	*	*	*	*	*
733	974	171	803	11	54	173	736	236	256	310	173
733.0	609	57	552	*	10	78	517	159	163	189	98
733.00	587	54	533	*	10	76	499	151	160	183	93
733.01	*6	*	*6	—	—	*	*6	*	*	*	*
733.09	16	*	12	*	*	*	12	*5	*	*	*
733.1	154	44	110	*	*	26	122	35	39	49	31
733.13	122	34	88	*	*	19	101	28	31	37	26
733.14	10	*	*7	*	*	*	*6	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
733.15	*5	*	*	*	*	*	*	*	*	*	*
733.19	*7	*	*6	*	*	*	*	*	*	*	*
733.2	*6	*	*	*	*	*	*	*	*	*	*
733.4	48	23	25	*	13	16	20	*9	10	17	13
733.42	33	18	15	*	*9	11	13	*6	*7	11	9
733.49	*8	*	*	*	*	*	*	*	*	*	*
733.6	38	9	28	*	11	15	11	9	*8	16	*
733.8	35	19	16	*	9	12	13	*7	*6	13	9
733.81	*8	*	*	*	*	*	*	*	*	*	*
733.82	28	14	14	*	*8	*9	10	*5	*5	11	*7
733.9	83	13	69	*	*6	24	50	15	27	24	17
733.90	69	9	59	*	*	19	44	13	22	20	14
733.99	*8	*	*6	*	*	*	*	*	*	*	*
734	*5	*	*	*	*	*	*	*	*	*	*
735	*8	*	*	—	*	*	*	*	*	*	*
736	55	25	30	*	*7	15	28	11	13	19	11
736.4	*8	*	*6	*	*	*	*6	*	*	*6	*
736.42	*6	*	*	*	*	*	*	*	*	*	*
736.6	*5	*	*	*	*	*	*	*	*	*	*
736.7	29	14	15	*	*	10	13	*7	*8	*8	*
736.79	24	11	13	*	*	*8	12	*6	*7	*7	*
736.8	*7	*	*	*	*	*	*	*	*	*	*
737	93	29	64	16	20	16	40	20	25	29	18
737.1	17	*7	10	*	*	*	11	*	*6	*6	*
737.10	14	*6	*8	*	*	*	11	*	*5	*5	*
737.3	67	19	48	14	18	14	21	17	16	21	13
737.30	55	14	41	10	15	14	17	14	14	19	*9
737.39	9	*	*6	*	*	*	*	*	*	*	*
737.4	*8	*	*5	*	*	*	*8	*	*	*	*
737.41	*6	*	*	—	—	*	*6	*	*	*	*
738	71	26	45	*	15	29	27	12	13	23	22
738.4	59	23	37	—	10	24	25	10	12	18	20
739	26	*5	21	*	17	*	*6	18	*6	*	—
739.2	*6	*	*5	*	*	*	*	*	*	*	—
739.3	14	*	12	*	12	*	*	12	*	*	—
740–759	718	371	347	422	124	95	76	157	142	237	181
741	25	14	11	11	9	*	*	*	*	10	*6
741.0	14	9	*	*8	*	*	—	*	*	*	*
741.00	*9	*6	*	*	*	*	—	*	*	*	*
741.9	11	*	*7	*	*5	*	*	*	*	*	*
741.90	9	*	*6	*	*	*	*	*	*	*	*
742	37	22	15	25	*8	*	*	10	*7	12	*8
742.2	*6	*	*	*5	*	*	—	*	*	*	*
742.3	14	10	*	10	*	*	*	*	*	*	*
742.5	*6	*	*	*	*	*	*	*	*	*	*
742.59	*5	*	*	*	*	*	*	*	*	*	*
744	*8	*6	*	*6	*	*	—	*	*	*	*
745	103	53	50	76	9	*8	10	24	17	33	28
745.1	*7	*	*	*5	*	*	—	*	*	*	*
745.2	*5	*	*	*	*	*	*	*	*	*	*
745.4	28	13	15	22	*	*	*	*6	*	10	*7
745.5	53	28	25	36	*	*	*9	11	10	16	16
746	62	37	25	34	*8	13	*6	15	11	22	14
746.0	*6	*	*	*6	*	*	*	*	*	*	*
746.02	*5	*	*	*	*	*	*	*	*	*	*
746.4	15	10	*	*	*	*7	*	*	*	*	*5
746.8	21	14	*7	12	*	*5	*	*	*	*9	*
746.85	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
746.89	10	*6	*	*7	*	*	*	*	*	*	*
746.9	*7	*	*	*	*	*	*	*	*	*	*
747	73	37	36	48	10	*	9	17	17	23	16
747.0	22	12	10	20	*	*	*	*	*	*6	*7
747.1	*5	*	*	*	*	*	*	*	*	*	*
747.3	12	*5	*6	10	*	*	*	*	*	*	*
747.6	*8	*	*	*	*	*	*	*	*	*	*
747.8	18	9	*9	*8	*5	*	*	*	*7	*6	*
747.81	10	*	*6	*	*5	*	*	*	*	*	*
747.83	*7	*5	*	*7	—	—	—	*	*	*	*
748	23	12	11	22	*	*	*	*5	*	*7	*7
748.3	16	*8	*7	15	—	*	*	*	*	*	*
749	20	11	9	18	*	—	*	*	*	*6	9
749.0	*7	*	*	*6	*	—	*	*	*	*	*
749.00	*6	*	*	*	*	—	*	*	*	*	*
749.2	11	*6	*	10	*	—	—	*	*	*	*7
750	24	17	*7	20	*	*	*	*	*8	*6	*7
750.5	13	12	*	13	—	—	—	*	*	*	*
751	38	17	21	21	*	*6	*6	*9	*7	13	*9
751.3	*7	*	*	*6	*	*	—	*	*	*	*
751.5	11	*	*7	*	*	*	*	*	*	*	*
752	28	12	16	11	12	*	*	*6	*	10	*8
752.1	*8	—	*8	*	*6	*	*	*	*	*	*
752.11	*7	—	*7	*	*6	*	*	*	*	*	*
752.5	*5	*5	—	*	*	*	—	*	*	*	*
752.6	*6	*6	—	*	*	*	*	*	*	*	*
752.61	*5	*5	—	*	*	*	*	*	*	*	*
753	83	38	45	19	18	21	25	16	22	30	14
753.0	*9	*	*	*	*	*	*	*	*	*	*
753.1	43	19	23	*	*7	13	20	*6	15	17	*
753.10	14	*7	*8	*	*	*	*7	*	*	*6	*
753.12	14	*6	*8	*	*	*	*8	*	*7	*	*
753.19	*6	*	*	*	*	*	*	*	*	*	*
753.2	13	*8	*5	*8	*	*	*	*	*	*	*
753.29	*7	*	*	*	*	*	*	*	*	*	*
753.3	*9	*	*6	*	*	*	*	*	*	*	*
754	21	10	11	18	*	*	*	*	*	*7	*7
754.2	*5	*	*	*	*	*	—	*	*	*	*
755	19	*8	12	10	*	*	*	*7	*	*	*6
755.6	13	*	*8	*	*	*	*	*	*	*	*
755.63	9	*	*5	*	*	*	*	*	*	*	*
756	59	29	30	25	13	13	*8	11	16	20	12
756.0	11	*7	*	10	*	*	*	*	*	*	*
756.1	30	12	18	*	*8	11	*8	*6	10	*9	*5
756.12	20	*8	13	*	*	10	*7	*	*7	*6	*
757	11	*	*7	*7	*	*	*	*	*	*7	*
757.3	*8	*	*5	*6	*	*	*	*	*	*6	*
757.32	*6	*	*	*	*	*	*	*	*	-	*5
758	41	24	17	24	*8	*8	*	11	*9	9	12
758.0	29	17	11	15	*	*8	*	*8	*7	*7	*7
759	40	19	21	25	9	*	*	*9	*6	13	12
759.8	27	14	13	20	*5	*	*	*7	*	*7	9
759.89	18	10	*9	15	*	*	*	*6	*	*	*6
760–779	614	344	270	611	*	*	*	80	124	215	194
764	*6	*	*	*6	—	—	—	*	*	*	*
765	109	58	51	109	*	*	*	15	20	40	34
765.0	11	*	*7	11	—	—	—	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
765.03	*6	*	*	*6	—	—	—	*	*	*	*
765.1	45	25	20	45	—	*	*	*	9	17	14
765.17	*6	*	*	*6	—	—	*	*	*	*	*
765.18	13	*6	*6	13	—	—	—	*	*	*6	*
765.19	12	*8	*	12	—	—	—	*	*	*	*
765.2	53	29	25	53	*	*	—	*8	10	18	17
765.26	*7	*	*	*7	—	—	—	*	*	*	*
765.27	14	*7	*7	14	—	—	—	*	*	*7	*
765.28	17	10	*7	17	—	*	—	*	*	*6	*
767	*7	*	*	*7	—	—	—	*	*	*	*
769	34	20	13	34	—	—	—	*	11	10	9
770	104	60	45	103	*	*	*	19	22	32	31
770.2	*8	*5	*	*8	—	—	—	*	*	*	*
770.6	15	10	*	15	—	—	—	*	*	*5	*
770.7	18	10	*8	18	*	—	*	*5	*	*	*
770.8	52	27	24	51	*	*	*	*8	11	17	16
770.81	15	*9	*7	15	—	—	—	*	*	*	*
770.82	*8	*	*	*7	*	*	*	*	*	*	*
770.89	20	11	*9	20	—	*	*	*	*	*7	*5
771	43	25	19	43	*	*	*	*5	*8	16	14
771.8	34	20	13	34	—	—	*	*	*6	12	12
771.81	22	11	11	22	—	—	*	*	*	*8	*7
771.82	*7	*6	*	*7	—	—	—	*	*	*	*
772	13	*6	*7	13	—	*	*	*	*	*7	*
772.1	10	*	*5	9	—	—	*	*	*	*6	*
773	11	*7	*	11	—	—	—	*	*	*	*
773.2	*6	*	*	*6	—	—	—	*	*	*	*
774	113	67	46	113	*	*	*	12	22	41	37
774.2	37	21	16	37	—	—	—	*	*8	15	10
774.6	70	42	28	70	*	*	*	*8	14	24	24
775	39	20	19	39	*	—	—	*	*6	14	13
775.4	*6	*	*	*6	—	—	—	*	*	*	*
775.5	17	*9	*8	17	—	—	—	*	*	*	*7
775.6	*7	*	*	*7	*	—	—	*	*	*	*
776	20	11	*8	20	—	—	—	*	*	10	*5
776.1	*6	*	*	*6	—	—	—	*	*	*	*
776.6	10	*6	*	10	—	—	—	*	*	*	*
777	10	*	*6	10	—	—	—	*	*	*	*
777.8	*5	*	*	*5	—	—	—	*	*	*	*
778	18	11	*7	18	—	—	—	*	*	*	*7
778.4	9	*5	*	9	—	—	—	*	*	*	*
779	73	38	35	73	—	—	*	*8	17	23	25
779.0	*7	*	*	*7	—	—	—	*	*	*	*
779.3	38	19	19	38	—	—	—	*	*6	13	15
779.8	26	14	12	26	—	—	*	*	*	*7	*8
779.81	*7	*	*	*7	—	—	*	*	*	*	*
779.89	16	*8	*8	16	—	—	*	*	*	*5	*5
780–799	8,829	3,964	4,865	771	1,951	2,522	3,585	1,863	2,377	3,010	1,579
780	2,720	1,302	1,418	253	576	766	1,125	620	736	909	455
780.0	125	61	65	*7	33	27	59	23	33	38	31
780.01	14	*7	*7	*	*	*	*7	*	*	*	*
780.02	*8	*	*6	*	*	*	*	*	*	*	*
780.09	101	50	51	*	27	21	48	19	27	29	26
780.1	24	14	10	*	*8	*	10	*6	*6	9	*
780.2	453	200	254	*7	48	108	290	129	108	147	69
780.3	893	440	453	102	247	255	290	214	243	291	145
780.31	22	10	12	21	*	*	*	*	*	*9	*6

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
780.39	871	430	441	81	247	254	289	211	238	282	139
780.4	145	54	91	* ¹	22	40	82	39	42	39	25
780.5	407	235	172	17	100	184	106	93	115	129	70
780.50	*7	*	*	*	*	*	*	*	*	*	*
780.52	53	19	33	*	18	16	18	10	20	14	*8
780.57	338	207	132	15	79	160	84	80	90	110	58
780.6	327	156	171	104	68	74	82	55	81	133	58
780.7	212	88	124	*6	27	43	136	35	71	73	34
780.79	207	87	121	*6	25	41	135	34	70	71	33
780.8	*8	*	*6	*	*	*	*5	*	*5	*	*
780.9	125	52	73	*8	22	30	65	26	31	48	21
780.93	15	*8	*8	*	*5	*	*6	*	*6	*	*
780.99	104	42	62	*	17	26	58	23	24	43	15
781	271	116	155	17	26	59	169	69	66	81	55
781.0	32	13	19	*6	*7	*7	12	*6	10	10	*6
781.2	158	65	93	*	9	33	114	50	33	47	28
781.3	51	25	26	*7	*6	10	28	10	15	14	12
781.9	24	11	13	*	*	*7	12	*	*6	*8	*7
781.94	17	*8	10	—	*	*6	*9	*	*	*	*6
781.99	*7	*	*	*	*	*	*	*	*	*	*
782	232	104	127	28	58	61	85	41	77	71	42
782.0	53	24	29	*	22	19	11	9	15	20	*9
782.1	47	25	22	15	12	*7	13	10	16	12	9
782.3	91	34	58	*	12	25	52	15	35	25	16
782.4	21	13	*8	*	*7	*6	*6	*	*5	*8	*
782.5	*6	*	*	*5	*	*	*	*	*	*	*
782.7	*6	*	*	*	*	*	*	*	*	*	*
783	232	106	126	76	28	28	100	44	58	83	47
783.0	37	15	22	*	*6	*5	21	*5	9	17	*6
783.2	65	31	34	*7	12	16	31	12	16	26	12
783.21	63	31	32	*6	11	15	31	11	16	26	10
783.3	20	10	10	15	*	*	*	*	*	9	*
783.4	52	26	26	47	*	*	*	10	15	16	11
783.40	18	*9	10	15	*	*	*	*	*6	*	*
783.41	31	15	15	29	*	*	*	*	*8	11	*7
783.7	48	19	28	*	*	*	41	11	14	11	12
784	394	154	240	23	113	112	146	79	114	133	68
784.0	208	65	143	13	90	66	39	37	61	79	30
784.2	17	*8	9	*	*	*	*6	*	*	*6	*
784.3	57	23	34	*	*	13	40	13	18	14	12
784.5	53	29	24	*	*5	14	32	12	18	14	9
784.7	47	25	22	*	*7	12	25	13	10	15	*8
785	420	191	229	31	91	114	183	93	107	149	71
785.0	111	49	62	10	40	29	32	18	33	43	18
785.1	44	14	30	*	10	12	21	11	11	14	*7
785.2	41	14	27	10	12	*8	11	*7	13	16	*6
785.4	58	34	25	*	*5	21	31	18	10	21	9
785.5	107	57	50	*	10	28	65	27	26	32	23
785.50	*8	*	*	*	*	*	*5	*	*	*	*
785.51	23	13	10	*	*	*	16	*7	*6	*5	*5
785.52	64	34	31	*	*7	19	36	16	15	21	12
785.59	12	*6	*6	*	*	*	*7	*	*	*	*
785.6	48	20	28	*5	13	13	17	12	11	18	*7
785.9	10	*	*6	*	*	*	*6	*	*	*	*
786	1,570	722	848	81	293	608	587	323	367	585	295
786.0	217	98	118	58	31	52	75	34	56	83	43
786.03	17	*8	*9	14	*	*	*	*	*	*6	*
786.05	80	34	46	*	13	26	39	14	22	29	15
786.06	*7	*	*	*	*	*	*	*	*	*	*
786.07	*9	*	*5	*6	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
786.09	96	44	51	32	14	20	30	15	24	38	18
786.1	*8	*	*	*6	*	*	*	*	*	*	*
786.2	39	16	23	10	*9	*8	12	*	12	15	*7
786.3	58	31	27	*	13	18	27	11	17	20	10
786.5	1,192	536	656	*	232	514	443	260	265	446	221
786.50	464	219	244	*	88	196	178	90	107	185	81
786.51	37	14	24	*	*9	16	12	*6	*6	18	*7
786.52	56	24	32	*	15	19	22	*8	19	20	10
786.59	634	279	356	*	119	283	231	156	132	224	122
786.6	45	27	18	*	*	14	24	10	13	12	*8
786.8	*7	*6	*	*	*	*	*	*	*	*	*
787	791	313	478	111	185	193	302	145	221	279	145
787.0	399	148	251	70	118	105	106	66	115	141	78
787.01	228	79	149	17	75	69	67	37	61	90	41
787.02	85	23	62	*6	24	27	28	14	31	22	18
787.03	86	46	40	46	19	*9	12	14	23	29	19
787.1	*7	*	*	*	*	*	*	*	*	*	*
787.2	131	62	69	*	13	28	86	30	32	46	22
787.3	14	*7	*7	*	*	*	*	*	*	*5	*
787.6	14	*7	*7	*	*	*	*7	*	*	*	*
787.9	225	85	140	32	44	52	97	43	62	82	38
787.91	219	80	139	32	41	49	97	43	61	77	38
787.99	*6	*6	*	*	*	*	*	*	*	*5	*
788	418	202	216	13	70	94	240	94	131	115	77
788.0	10	*	*6	*	*7	*	*	*	*	*	*
788.1	15	*9	*7	*	*	*	*7	*	*7	*	*
788.2	224	137	87	*	36	47	137	53	69	59	42
788.20	185	112	73	*	30	38	115	44	59	50	33
788.21	*6	*	*	—	*	*	*	*	*	*	*
788.29	32	21	11	*	*5	*8	19	*8	*9	*7	*8
788.3	114	32	82	*5	13	32	65	26	34	30	25
788.30	81	26	54	*	*8	21	48	20	26	20	14
788.31	*9	*	*8	*	*	*	*	*	*	*	*
788.33	15	*	15	—	*	*7	*5	*	*	*5	*6
788.39	*6	*	*	*	*	*	*	*	*	*	*
788.4	22	*8	14	*	*	*	13	*	*7	*7	*
788.41	15	*	11	*	*	*	10	*	*5	*	*
788.5	9	*	*7	*	*	*	*	*	*	*	*
788.6	*5	*	*	*	*	*	*	*	*	*	*
788.9	18	*8	10	*	*	*	14	*	*9	*	*
789	592	236	356	40	209	188	155	118	145	204	125
789.0	419	149	270	34	176	109	101	81	105	140	94
789.00	145	56	89	16	57	37	35	30	31	57	27
789.01	47	14	33	*	22	14	9	*6	14	12	14
789.02	12	*5	*7	*	*	*	*	*	*	*	*
789.03	46	16	30	*7	26	*6	*6	*8	14	13	12
789.04	23	*7	16	*	9	*7	*6	*	*8	*5	*
789.05	*6	*	*	*	*	*	*	*	*	*	*
789.06	56	17	39	*	17	21	16	10	12	19	14
789.07	17	*6	11	*	*9	*	*5	*	*	*7	*
789.09	68	25	43	*5	28	16	19	15	17	20	15
789.1	10	*6	*	*	*6	*	*	*	*	*	*
789.2	13	*8	*	*	*	*	*	*	*	*5	*
789.3	18	*	14	*	*5	*5	*7	*	*	9	*
789.30	10	*	*8	*	*	*	*	*	*	*5	*
789.39	*6	*	*	—	*	*	*	*	*	*	*
789.5	128	68	60	*	17	68	40	31	31	42	23
790	481	221	260	37	104	127	214	106	149	152	75
790.2	63	18	45	*	30	14	18	*8	19	21	16
790.22	17	*	16	—	14	*	*	*	*	*8	*
790.29	42	14	27	*	16	11	14	*	13	12	12

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
790.4	21	9	12	*	*7	*6	*7	*	*9	*6	*
790.5	44	23	21	*	*9	15	18	*9	16	11	*7
790.6	128	58	71	*8	25	40	55	27	39	43	19
790.7	129	66	63	20	20	32	58	35	34	42	19
790.9	89	44	45	*	11	18	57	23	30	28	*8
790.92	67	30	37	*	*6	13	46	17	21	21	*7
790.93	*8	*8	—	—	*	*	*6	*	*	*	*
790.99	12	*	*7	*	*	*	*	*	*6	*	*
791	32	12	20	*	13	*8	*7	*7	12	10	*
791.0	13	*7	*6	*	*6	*	*	*	*	*	*
791.9	13	*	10	*	*	*	*	*	*6	*	*
792	61	12	48	*	40	*6	13	13	19	17	12
792.1	22	10	12	*	*	*6	11	*7	*6	*5	*
792.3	35	—	35	*	35	—	—	*6	11	11	*8
793	22	11	11	*	*6	*6	*8	*	*7	*6	*
793.1	11	*6	*5	*	*	*	*6	*	*	*	*
794	178	83	95	*	46	65	62	35	52	62	29
794.3	85	38	47	*	14	33	38	16	30	28	12
794.31	55	22	34	*	10	21	25	11	19	16	*9
794.39	27	15	11	—	*	12	12	*	10	11	*
794.8	80	41	39	*	28	30	20	16	18	31	16
795	40	10	30	*	27	9	*	12	*6	13	10
795.0	9	—	9	—	*8	*	*	*	*	*	*
795.5	20	*6	13	*	13	*	*	*8	*	*	*7
795.7	10	*	*7	*	*5	*	*	*	*	*	*
795.79	10	*	*7	—	*5	*	*	*	*	*	*
796	59	23	36	*	24	14	19	*7	16	23	12
796.0	*6	*	*	*	*	*	*	*	*	*	*
796.2	41	16	25	*	19	11	10	*	12	16	10
799	316	145	171	47	42	63	164	52	94	119	51
799.0	150	72	78	41	16	27	66	20	46	57	27
799.1	14	*6	*8	*	*	*5	*	*	*5	*	*
799.2	*5	*	*	*	*	*	*	*	*	*	*
799.3	71	31	40	—	*	*8	60	12	27	25	*8
799.4	36	16	20	*	*6	10	20	*7	*7	14	*8
799.9	38	16	21	*	12	12	12	10	*6	16	*5
800–999	7,122	3,694	3,427	476	2,108	1,824	2,714	1,363	1,702	2,536	1,521
800	14	11	*	*8	*	*	*	*	*	*	*
801	48	35	12	*8	25	*6	9	*6	11	16	14
801.0	21	16	*	*	10	*	*	*	*	*8	*
801.01	11	*8	*	*	*5	*	*	*	*	*	*
801.1	*6	*	*	*	*	*	*	*	*	*	*
801.2	17	12	*	*	*9	*	*	*	*	*	*7
802	106	78	28	*8	59	18	21	21	27	36	22
802.0	21	15	*6	*	9	*	*7	*5	*7	*	*
802.2	22	18	*	*	16	*	*	*	*6	10	*
802.3	*5	*	*	*	*	*	*	*	*	*	*
802.4	20	14	*6	*	11	*	*	*	*5	*6	*
802.6	13	11	*	*	*7	*	*	*	*	*	*
802.8	22	14	*8	*	10	*	*	*	*	*8	*
803	*6	*	*	*	*	*	*	*	*	*	*
805	211	98	113	*	71	44	93	32	59	75	45
805.0	51	29	22	*	23	13	14	*8	12	21	11
805.02	15	*7	*7	*	*	*7	*	*	*5	*	*
805.04	*6	*	*	—	*	*	*	*	*	*	*
805.06	*6	*	*	*	*	*	*	*	*	*	*
805.07	*9	*6	*	*	*	*	*	*	*	*	*
805.2	51	22	29	*	15	10	25	10	14	18	*9

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
805.4	82	37	45	*	23	15	44	11	24	26	21
805.6	21	*9	13	*	*7	*6	*8	*	*7	*8	*
806	10	*7	*	*	*7	*	*	*	*	*6	*
806.2	*5	*	*	*	*	*	*	*	*	*	*
807	127	77	49	*	32	39	53	24	35	39	29
807.0	111	68	43	*	26	35	47	22	31	33	25
807.00	*8	*	*	*	*	*	*6	*	*	*	*
807.01	23	15	*8	*	*7	*7	9	*5	*6	*7	*
807.02	17	9	*8	*	*	*5	*9	*	*6	*	*
807.03	15	10	*6	*	*	*6	*	*	*5	*	*
807.04	10	*6	*	*	*	*	*7	*	*	*	*
807.05	*7	*	*	—	*	*	*	*	*	*	*
807.09	20	14	*6	*	*6	*7	*6	*	*6	*6	*6
807.2	11	*6	*5	*	*	*	*6	*	*	*	*
808	120	44	75	*	29	22	65	20	30	42	27
808.0	30	18	13	*	12	*8	10	*	*7	12	*7
808.2	69	17	51	*	12	*8	47	12	17	23	17
808.4	11	*7	*	*	*	*5	*	*	*	*	*
808.41	*7	*	*	*	*	*	*	*	*	*	*
808.8	*8	*	*6	*	*	*	*7	*	*	*	*
810	22	15	*7	*	*8	*6	*6	*	*6	*6	*6
810.0	22	15	*7	*	*8	*6	*6	*	*6	*6	*6
810.00	10	*6	*	*	*	*	*	*	*	*	*
810.02	*8	*6	*	*	*	*	*	*	*	*	*
811	13	10	*	*	*	*5	*	*	*	*	*
811.0	13	10	*	*	*	*5	*	*	*	*	*
811.00	*6	*	*	*	*	*	*	*	*	*	*
812	108	36	71	15	15	24	54	22	30	31	25
812.0	61	16	45	*	*5	16	38	13	21	15	13
812.00	16	*	12	*	*	*	11	*	*5	*	*
812.01	18	*	14	*	*	*	12	*	*8	*	*
812.03	*8	*	*6	—	*	*	*	*	*	*	*
812.09	20	*7	13	*	*	*5	11	*	*5	*7	*
812.2	18	*7	12	*	*	*	10	*	*	*5	*6
812.20	*5	*	*	*	*	*	*	*	*	*	*
812.21	13	*6	*8	*	*	*	*5	*	*	*	*
812.4	23	10	12	10	*	*	*5	*	*	*8	*6
812.41	13	*6	*7	*8	*	*	*	*	*	*	*
813	107	49	58	*9	28	27	42	20	28	38	20
813.0	20	*6	13	*	*	*7	*8	*	*6	*6	*
813.01	*7	*	*	*	*	*	*	*	*	*	*
813.2	10	*7	*	*	*6	*	*	*	*	*	*
813.23	*7	*	*	*	*	*	*	*	*	*	*
813.4	56	23	34	*	9	13	30	10	15	21	10
813.41	13	*	9	*	*	*	*9	*	*	*7	*
813.42	23	11	13	*	*	*7	11	*	*6	*8	*
813.44	15	*6	10	*	*	*	*8	*	*	*5	*
813.5	*9	*5	*	*	*	*	*	*	*	*	*
814	*7	*6	*	—	*	*	*	*	*	*	*
814.0	*6	*	*	—	*	*	*	*	*	*	*
815	18	14	*	*	12	*	*	*	*	*8	*
815.0	14	11	*	*	9	*	*	*	*	*7	*
815.00	*6	*	*	*	*	*	*	*	*	*	*
816	19	14	*5	*	10	*	*5	*	*	*8	*
816.0	10	*5	*	*	*5	*	*	*	*	*5	*
816.01	*7	*	*	—	*	*	*	*	*	*	*
816.1	*9	*8	*	*	*5	*	*	*	*	*	*
816.11	*5	*5	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
820	338	96	242	*	9	32	296	63	83	126	67
820.0	98	27	71	*	*	11	85	15	28	36	20
820.00	*7	*	*6	—	*	*	*6	*	*	*	*
820.03	*6	*	*	*	*	*	*5	*	*	*	*
820.09	80	21	59	*	*	*8	70	13	21	29	16
820.2	168	44	123	*	*	17	147	36	37	63	31
820.20	14	*	9	*	*	*	11	*	*	*7	*
820.21	137	31	105	*	*	11	124	29	32	49	27
820.22	17	*8	9	*	*	*	11	*	*	*8	*
820.8	71	23	48	*	*	*	64	12	17	26	16
821	72	36	37	14	22	10	26	11	19	30	12
821.0	42	23	19	10	15	*	12	*6	12	17	*7
821.00	*9	*	*	*	*	*	*	*	*	*	*
821.01	34	18	16	9	13	*	*8	*	10	14	*
821.2	26	10	16	*	*	*6	13	*	*6	10	*
821.20	*7	*	*	*	*	*	*	*	*	*	*
821.23	11	*	*8	*	*	*	*7	*	*	*6	*
822	16	*7	9	*	*	*	*7	*	*	*8	*
822.0	14	*5	*9	*	*	*	*6	*	*	*6	*
823	94	58	36	*7	42	31	15	14	21	37	21
823.0	38	21	18	*	13	16	9	*5	9	15	*9
823.00	27	14	12	*	*9	11	*6	*	*6	11	*6
823.02	*8	*	*	*	*	*	*	*	*	*	*
823.2	22	16	*6	*	13	*6	*	*	*	*9	*6
823.20	*6	*	*	*	*	*	*	*	*	*	*
823.22	14	9	*	*	*8	*	*	*	*	*	*
823.3	11	*7	*	*	*7	*	*	*	*	*5	*
823.32	*7	*6	*	*	*	*	*	*	*	*	*
823.8	13	*7	*6	*	*	*	*	*	*	*	*
823.82	*8	*	*	*	*	*	*	*	*	*	*
823.9	*5	*	*	*	*	*	*	*	*	*	*
824	129	59	70	*6	47	36	40	30	31	39	29
824.0	*9	*	*	*	*	*	*	*	*	*	*
824.2	14	*6	*8	*	*	*	*	*	*	*	*
824.4	28	12	16	*	10	9	9	*8	*6	*8	*7
824.6	30	*8	23	*	*8	10	12	*7	10	*5	*8
824.8	34	19	15	*	13	*8	*9	*9	*8	10	*6
824.9	*6	*5	*	*	*	*	*	*	*	*	*
825	44	24	20	*	22	13	*8	*7	11	17	*9
825.0	13	*7	*5	*	*	*	*	*	*	*	*
825.2	23	11	12	*	12	*	*6	*	*7	*8	*
825.21	*6	*	*	*	*	*	*	*	*	*	*
825.25	11	*	*8	*	*	*	*	*	*	*	*
825.3	*6	*	*	*	*	*	*	—	*	*	*
826	*8	*	*	*	*	*	*	*	*	*	*
826.0	*6	*	*	*	*	*	*	*	*	*	*
831	12	*8	*	*	*	*5	*	*	*	*	*
831.0	12	*8	*	*	*	*5	*	*	*	*	*
836	16	10	*7	*	*	9	*	*	*	*7	*
836.0	*6	*	*	*	*	*	*	*	*	*	*
836.1	*6	*	*	*	*	*	*	*	*	*	*
839	*8	*6	*	*	*5	*	*	*	*	*	*
839.0	*5	*	*	*	*	*	*	*	*	*	*
840	35	18	18	—	*	15	17	*6	10	16	*
840.4	23	11	12	—	*	10	12	*	*6	9	*
843	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
844	22	12	10	*	*8	*6	*7	*	*6	*7	*
844.8	*6	*	*	*	*	*	*	*	*	*	*
844.9	*8	*	*	*	*	*	*	*	*	*	*
845	23	11	12	*	9	*6	*7	*7	*6	*7	*
845.0	22	11	11	*	9	*5	*7	*7	*5	*6	*
845.00	10	*	*6	*	*	*	*	*	*	*	*
845.03	*5	*	*	*	*	*	*	*	*	*	*
846	*5	*	*	—	*	*	*	*	*	*	*
847	35	16	19	*	12	*8	14	10	9	10	*6
847.0	18	*9	*9	*	*7	*	*5	*5	*	*5	*
847.2	11	*	*6	*	*	*	*6	*	*	*	*
850	85	53	33	*9	40	19	18	19	20	26	20
850.0	15	*5	10	*	*	*	*5	*	*	*	*
850.1	26	18	*8	*	12	*6	*	*6	*6	*8	*5
850.11	25	18	*7	*	12	*6	*	*6	*6	*7	*5
850.5	34	25	9	*	19	*8	*	*6	*7	12	*9
850.9	10	*	*5	*	*	*	*	*	*	*	*
851	35	24	11	*	15	*8	11	*	*5	14	11
851.8	25	17	*9	*	11	*5	*8	*	*	12	*8
851.81	*5	*	*	*	*	*	*	*	*	*	*
851.82	*6	*	*	*	*	*	*	*	*	*	*
851.86	*7	*5	*	*	*	*	*	*	*	*	*
852	74	43	31	*	10	16	44	11	17	26	20
852.0	26	17	10	*	*	*8	13	*	*6	*8	9
852.00	*7	*5	*	*	*	*	*	*	*	*	*
852.01	*6	*	*	*	*	*	*	*	*	*	*
852.02	*7	*	*	*	*	*	*	*	*	*	*
852.2	46	25	20	*	*5	*8	31	*7	11	17	11
852.20	12	*9	*	*	*	*	*8	*	*	*	*
852.21	18	9	*9	*	*	*	14	*	*	*7	*
852.22	*8	*	*5	*	*	*	*5	*	*	*	*
852.26	*5	*	*	—	*	*	*	*	*	*	*
853	16	*8	*8	*	*	*	*9	*	*	*	*5
853.0	16	*8	*8	*	*	*	*9	*	*	*	*5
853.01	*5	*	*	*	*	*	*	*	*	*	*
854	*8	*5	*	*	*	*	*	*	*	*	*
854.0	*8	*5	*	*	*	*	*	*	*	*	*
860	72	52	20	*	40	18	12	11	14	29	18
860.0	40	29	10	*	20	10	*7	*6	*8	16	10
860.2	*8	*6	*	—	*	*	*	*	*	*	*
860.4	15	*9	*6	*	*8	*	*	*	*	*6	*
860.5	*6	*5	*	—	*5	*	*	*	*	*	*
861	39	28	11	*	19	*9	*8	*5	10	13	10
861.2	33	25	*9	*	16	*7	*6	*	9	11	*9
861.21	32	23	*8	*	15	*7	*6	*	9	11	*7
862	11	*9	*	*	*5	*	*	*	*	*	*
862.2	*7	*5	*	*	*	*	*	*	*	*	*
862.29	*6	*	*	*	*	*	*	*	*	*	*
863	25	19	*5	*	16	*	*	*	*	12	*8
863.3	*7	*5	*	*	*5	*	*	*	*	*	*
864	24	16	*8	*	15	*	*	*	*	12	*5
864.0	19	12	*7	*	11	*	*	*	*	9	*
864.05	*6	*	*	*	*	*	*	*	*	*	*
864.1	*5	*	*	*	*	*	*	*	*	*	*
865	38	28	11	*6	19	*8	*	*5	*9	11	13
865.0	36	26	11	*6	17	*8	*	*	*8	10	13
865.00	10	*8	*	*	*6	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
865.01	*6	*	*	*	*	*	*	*	*	*	*
865.03	*6	*	*	*	*	*	*	*	*	*	*
866	11	9	*	*	*7	*	*	*	*	*	*
866.0	10	*8	*	*	*6	*	*	*	*	*	*
867	19	17	*	*	*	*	12	*	*	*8	*
867.0	16	14	*	*	*	*	11	*	*	*7	*
868	15	10	*	*	*9	*	*	*	*	*6	*
868.0	11	*6	*	*	*6	*	*	*	*	*	*
870	13	10	*	*	*5	*	*	*	*	*5	*
870.8	*5	*	*	*	*	*	*	*	*	*	*
871	*7	*	*	*	*	*	*	*	*	*	*
872	*6	*	*	*	*	*	*	*	*	*	*
873	174	110	64	15	74	32	54	33	40	55	46
873.0	53	27	26	*	18	*8	25	11	13	12	17
873.2	*8	*6	*	*	*	*	*	*	*	*	*
873.20	*8	*6	*	*	*	*	*	*	*	*	*
873.4	80	52	28	*7	37	15	21	17	20	25	17
873.40	*7	*	*	*	*	*	*	*	*	*	*
873.41	*5	*	*	*	*	*	*	*	*	*	*
873.42	41	25	15	*	17	*7	13	9	10	13	*8
873.43	15	10	*	*	*7	*	*	*	*	*5	*
873.44	*8	*6	*	*	*	*	*	*	*	*	*
873.5	*9	*8	*	*	*5	*	*	*	*	*	*
873.6	13	10	*	*	*6	*	*	*	*	*	*
873.63	*6	*	*	*	*	*	*	*	*	*	*
874	*7	*6	*	*	*	*	*	*	*	*	*
874.8	*5	*	*	*	*	*	*	*	*	*	*
875	11	10	*	—	*9	*	*	*	*	*	*
875.0	10	9	*	—	*9	*	*	*	*	*	*
879	16	11	*	*	9	*	*	*	*	*7	*
879.2	*7	*5	*	*	*	*	*	*	*	*	*
880	*7	*7	*	*	*6	*	*	*	*	*	*
880.0	*6	*6	*	*	*5	*	*	*	*	*	*
881	49	23	25	*	29	*7	9	*8	18	14	9
881.0	35	15	20	*	18	*6	*9	*6	14	*9	*6
881.00	11	*6	*	*	*6	*	*	*	*	*	*
881.01	11	*	*6	*	*	*	*6	*	*	*	*
881.02	13	*	10	*	9	*	*	*	*	*7	*
881.1	*6	*	*	*	*	*	*	*	*	*	*
881.2	*8	*5	*	*	*7	*	*	*	*	*	*
882	26	15	11	*	16	*	*	*	*7	9	*5
882.0	10	*6	*	*	*5	*	*	*	*	*	*
882.1	12	*7	*6	*	*8	*	*	*	*	*	*
883	30	23	*7	*	18	*9	*	*	*8	10	*7
883.0	11	9	*	*	*	*	*	*	*	*	*
883.1	9	*6	*	*	*6	*	*	*	*	*	*
883.2	10	*8	*	*	*7	*	*	*	*	*	*
886	*7	*6	*	*	*	*	*	*	*	*	*
886.0	*6	*6	*	*	*	*	*	*	*	*	*
890	10	*7	*	*	*7	*	*	*	*	*	*
890.0	*6	*	*	*	*	*	*	*	*	*	*
891	34	19	15	*	14	*6	9	*5	*8	14	*7
891.0	17	9	*8	*	*8	*	*	*	*	*7	*
891.1	14	*7	*7	*	*	*	*	*	*	*7	*
892	15	10	*	*	*5	*6	*	*	*5	*6	*
892.1	*9	*7	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
893	*7	*5	*	*	*	*	*	*	*	*	*
893.0	*6	*5	*	*	*	*	*	*	*	*	*
903	11	*7	*	*	*6	*	*	*	*5	*	*
904	*6	*6	*	*	*	*	—	*	*	*	*
905	44	24	20	*	17	14	11	11	*8	13	12
905.0	*8	*	*	*	*	*	*	*	*	*	*
905.1	*8	*	*	*	*	*	*	*	*	*	*
905.4	16	11	*5	*	*8	*	*	*	*	*5	*
906	18	13	*6	*	*9	*	*	*	*	9	*
907	57	45	12	*	30	16	9	9	*8	17	23
907.0	27	21	*7	*	12	*9	*5	*	*	*	13
907.2	27	22	*5	*	17	*5	*	*	*	11	9
908	16	9	*7	*	*6	*6	*	*	*	*6	*
908.9	11	*7	*	*	*	*	*	*	*	*	*
909	13	*6	*7	*	*	*	*	*	*	*	*
909.2	*7	*	*	*	*	*	*	*	*	*	*
910	35	21	15	*6	14	*5	10	*6	12	10	*8
910.0	34	20	14	*	14	*	10	*6	11	*9	*8
911	17	12	*5	*	*8	*	*	*	*	*	*6
911.0	13	9	*	*	*6	*	*	*	*	*	*
912	*8	*5	*	*	*	*	*	*	*	*	*
912.0	*7	*5	*	*	*	*	*	*	*	*	*
913	25	11	14	*	14	*	*5	*	12	*5	*
913.0	20	*9	11	*	10	*	*	*	9	*	*
914	*8	*6	*	*	*	*	*	*	*	*	*
914.0	*5	*	*	*	*	*	*	*	*	*	*
916	39	23	16	*8	15	*8	*9	*5	9	12	13
916.0	25	14	11	*	12	*5	*6	*	*7	*	9
916.5	*8	*5	*	*	*	*	*	*	*	*	*5
917	*6	*	*	*	*	*	*	*	*	*	*
918	*8	*	*	*	*	*	*	*	*	*	*
918.1	*6	*	*	*	*	*	*	*	*	*	*
919	13	*9	*	*	*6	*	*	*	*	*	*
919.0	11	*8	*	*	*5	*	*	*	*	*	*
920	76	34	42	10	18	13	36	16	24	22	13
921	20	11	*9	*	*8	*	*6	*	*6	*6	*
921.2	*5	*	*	*	*	*	*	*	*	*	*
922	68	36	32	*	17	15	32	12	26	15	15
922.1	32	17	15	*	*7	*7	18	*	15	*7	*7
922.2	17	*9	*8	*	*	*	*6	*	*6	*5	*
922.3	15	*8	*7	*	*	*	*6	*	*	*	*
922.31	10	*6	*	*	*	*	*	*	*	*	*
923	39	15	24	*	*8	*7	22	*	16	11	*7
923.0	13	*6	*7	*	*	*	*7	*	*7	*	*
923.00	11	*5	*6	*	*	*	*6	*	*6	*	*
923.1	15	*6	*9	*	*	*	*8	*	*	*6	*
923.11	11	*	*7	*	*	*	*6	*	*	*	*
923.2	*7	*	*	*	*	*	*	*	*	*	*
924	98	35	63	*	18	19	58	16	33	27	21
924.0	33	9	24	*	*	*	26	*	14	*9	*6
924.00	10	*	*7	*	*	*	*6	*	*5	*	*
924.01	23	*6	17	*	*	*	20	*	*9	*6	*
924.1	31	10	21	*	*	*7	19	*6	10	*8	*7
924.10	12	*	*8	*	*	*	9	*	*	*	*
924.11	19	*5	13	*	*	*5	10	*	*7	*	*
924.8	19	11	*8	*	*6	*5	*6	*	*	*7	*5

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
933	10	*	*6	*	*	*	*5	*	*	*	*
933.1	*9	*	*5	*	*	*	*	*	*	*	*
934	19	11	*8	*	*	*	12	*	*	*8	*7
934.8	9	*7	*	*	*	*	*6	*	*	*	*
935	11	*7	*	*	*	*	*	*	*	*	*
935.1	*8	*	*	*	*	*	*	*	*	*	*
941	26	20	*7	*8	11	*5	*	*	*5	16	*
941.2	20	15	*	*7	9	*	*	*	*	12	*
941.29	9	*6	*	*	*5	*	*	*	*	*7	*
942	30	20	10	11	12	*5	*	*	*7	17	*
942.2	23	15	*8	9	*9	*	*	*	*5	13	*
942.22	*6	*5	*	*	*	*	*	*	*	*	*
942.29	*6	*	*	*	*	*	—	*	*	*	—
942.3	*6	*	*	*	*	*	*	*	*	*	*
943	27	19	*7	9	10	*5	*	*	*5	15	*
943.2	19	15	*	*8	*7	*	*	*	*	11	*
943.21	*6	*	*	*	*	*	*	*	*	*	*
943.29	*6	*5	*	*	*	*	*	*	—	*6	—
943.3	*5	*	*	*	*	*	*	*	*	*	—
944	27	18	*9	*5	14	*	*	*	*8	13	*
944.2	16	12	*	*	*7	*	*	*	*	*8	*
944.28	*5	*	*	*	*	*	*	*	*	*	—
944.3	*6	*	*	*	*	*	*	*	*	*	—
945	37	29	*9	11	14	*8	*	*	*6	24	*
945.2	16	13	*	*7	*6	*	*	*	*	12	*
945.3	17	12	*	*	*7	*	*	*	*	11	*
948	61	44	17	22	22	12	*5	*8	*9	40	*
948.0	40	31	*9	13	14	*9	*	*6	*7	23	*
948.00	40	31	*9	13	14	*9	*	*6	*7	23	*
948.1	15	9	*6	*7	*6	*	*	*	*	13	—
948.10	11	*6	*	*	*	*	*	*	*	*9	—
949	*7	*6	*	*7	*	—	*	—	—	*7	—
949.0	*7	*6	*	*7	*	—	—	—	—	*7	—
952	*8	*7	*	*	*	*	*	*	*	*	*
952.0	*5	*5	*	*	*	*	*	*	*	*	*
955	14	11	*	*	*9	*	*	*	*	*	*
956	*6	*	*	—	*	*	*	*	*	*	*
958	37	23	15	*	16	12	*7	10	*8	12	*8
958.3	10	*	*6	*	*	*	*	*	*	*	*
958.4	*7	*	*	*	*	*	*	*	*	*	*
958.7	10	*6	*	*	*	*	*	*	*	*	*
958.8	*7	*5	*	*	*	*	*	*	*	*	*
959	89	45	44	10	33	18	28	17	24	22	27
959.0	47	23	24	*8	15	9	15	*8	13	10	16
959.01	41	19	23	*7	12	*8	15	*6	13	9	13
959.09	*6	*	*	*	*	*	*	*	*	*	*
959.1	21	11	11	*	9	*	*7	*	*7	*5	*5
959.11	*7	*	*	*	*	*	*	*	*	*	*
959.12	*7	*	*	*	*	*	*	*	*	*	*
959.19	*7	*	*	*	*	*	*	*	*	*	*
959.7	*7	*	*	*	*	*	*	*	*	*	*
962	11	*6	*	*	*	*5	*	*	*	*7	*
962.3	10	*6	*	*	*	*	*	*	*	*6	*
963	12	*6	*6	*	*8	*	*	*	*	*	*
963.0	11	*6	*5	*	*8	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
964	*6	*	*	*	*	*	*	*	*	*	*
964.2	*5	*	*	—	*	*	*	*	*	*	*
965	108	40	68	*	72	25	*7	19	29	37	22
965.0	43	20	23	*	24	15	*	*8	10	14	11
965.00	11	*6	*5	—	*7	*	*	*	*	*5	*
965.01	*5	*	*	*	*	*	*	*	*	*	*
965.02	*6	*	*	*	*	*	*	*	*	*	*
965.09	20	*6	14	*	10	*7	*	*	*6	*6	*6
965.1	*9	*	*5	*	*7	*	*	*	*	*	*
965.4	37	12	25	*	27	*6	*	*8	*9	14	*6
965.6	11	*	*8	*	*9	*	*	*	*	*	*
965.61	11	*	*8	*	*9	*	*	*	*	*	*
966	20	*9	11	*	15	*	*	*	*8	*6	*
966.3	17	*6	11	*	14	*	*	*	*7	*5	*
967	21	*6	15	*	10	*8	*	*	*	10	*
967.8	13	*	9	*	*7	*	*	*	*	*6	*
967.9	*5	*	*	—	*	*	*	*	*	*	*
968	*7	*	*	*	*	*	*	*	*	*	*
969	134	57	78	*	95	32	*	21	34	51	28
969.0	38	17	21	*	29	*8	*	*6	*9	15	*9
969.3	15	*7	*8	*	11	*	*	*	*	*	*
969.4	56	19	37	*	37	16	*	9	15	23	9
969.7	12	*8	*	*	10	*	*	*	*	*	*
969.8	*6	*	*	*	*	*	*	*	*	*	*
970	29	19	10	*	22	*7	*	10	*6	10	*
970.8	26	18	*8	*	19	*6	*	9	*6	*8	*
972	16	*8	*8	*	*	*	*	*5	*5	*	*
972.6	*8	*5	*	*	*	*	*	*	*	*	*
975	13	*	10	*	10	*	*	*	*	*	*
975.2	*8	*	*6	*	*6	*	*	*	*	*	*
977	14	*7	*7	*	*9	*	*	*	*	*6	*
977.8	*8	*	*	*	*	*	*	*	*	*	*
977.9	*6	*	*	*	*	*	*	*	*	*	*
980	25	14	11	*	16	*7	*	*	10	*7	*
980.0	19	10	*9	*	13	*	*	*	*6	*6	*
987	9	*	*5	*	*	*	*	*	*	*	*
987.8	*5	*	*	*	*	*	*	*	*	*	*
989	19	11	*9	*5	*8	*	*	*	*	11	*
989.5	14	10	*	*	*5	*	*	*	*	*9	*
991	13	*8	*6	*	*	*	*	*5	*	*	*
991.6	11	*6	*	*	*	*	*	*	*	*	*
992	*7	*	*	*	*	*	*	*	*	*	*
994	*7	*	*	*	*	*	*	*	*	*	*
995	601	281	320	31	79	158	333	124	139	224	114
995.0	*6	*	*	*	*	*	*	*	*	*	*
995.1	17	*7	11	*	*	*6	*5	*	*5	*7	*
995.2	35	17	18	*	*	10	20	*6	*8	15	*5
995.3	*7	*	*	*	*	*	*	*	*	*	*
995.5	*8	*	*	*8	*	*	—	*	*	*	*
995.8	*8	*	*7	*	*	*	*	*	*	*	*
995.9	518	246	272	17	62	134	304	109	117	193	98
995.90	*7	*	*	*	*	*	*5	*	*	*	*
995.91	345	157	188	12	46	84	203	73	92	121	60
995.92	159	82	77	*	14	48	93	35	23	69	32
996	826	416	410	43	139	280	364	169	160	319	179
996.0	41	25	16	*	*	*7	29	11	11	9	10

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
996.01	23	12	11	*	*	*	19	*6	*	*5	*7
996.04	12	10	*	*	*	*	*7	*	*	*	*
996.1	24	13	11	*	*	*7	*9	*5	*	10	*
996.2	26	12	13	*8	11	*	*	*	*6	9	*6
996.3	*9	*5	*	*	*	*	*	*	*	*	*
996.4	115	45	70	*	11	38	65	25	28	44	19
996.5	22	10	11	*	*	9	*7	*	*	*7	*5
996.59	15	*8	*7	*	*	*5	*5	*	*	*	*
996.6	274	137	137	17	47	92	119	58	55	103	59
996.61	13	*8	*5	*	*	*	*6	*	*	*	*
996.62	133	65	69	13	28	45	47	28	27	51	28
996.63	*7	*	*	*	*	*	*	*	*	*	*
996.64	29	16	13	—	*	*	23	*6	*	13	*6
996.65	*7	*	*	—	*	*	*	*	*	*	*
996.66	39	21	18	*	*	13	24	*6	12	13	*8
996.67	18	9	*9	*	*	*7	*6	*	*	*7	*
996.68	*8	*	*	*	*	*	*	*	*	*	*
996.69	19	*8	11	*	*	*7	*8	*5	*	*8	*
996.7	253	132	121	*6	34	91	123	47	43	99	64
996.71	*5	*	*	*	*	*	*	*	*	*	*
996.72	45	30	15	*	*	19	23	11	*8	17	9
996.73	69	34	35	*	*9	29	30	*9	10	24	26
996.74	53	25	28	*	*	18	28	14	*9	23	*8
996.76	21	14	*7	*	*5	*	11	*	*	*8	*5
996.77	22	11	11	—	*	*6	15	*	*	*8	*6
996.78	22	*8	13	*	*7	*8	*7	*	*	10	*
996.79	10	*	*6	*	*	*	*	*	*	*5	*
996.8	62	34	27	*	23	29	*6	12	*6	35	*8
996.81	40	22	17	*	15	18	*	*8	*6	22	*
996.82	*7	*	*	*	*	*	*	*	*	*	*
996.85	*6	*	*	*	*	*	—	*	*	*	*
997	548	278	270	17	83	166	282	135	119	181	113
997.0	30	14	16	*	*6	10	14	*6	*8	9	*7
997.02	15	*8	*7	*	*	*	*9	*	*	*	*
997.09	12	*5	*7	*	*	*6	*	*	*	*	*
997.1	111	60	51	*	9	29	70	30	24	33	25
997.2	23	10	13	*	*	*8	12	*6	*6	*6	*
997.3	99	46	53	*5	13	33	48	23	20	36	20
997.4	175	82	93	*6	42	51	76	39	40	61	35
997.5	66	40	26	*	*8	19	38	18	12	20	15
997.6	33	20	13	*	*	11	18	9	*6	12	*5
997.62	22	15	*7	—	*	*9	12	*5	*	*8	*
997.69	10	*	*	*	*	*	*6	*	*	*	*
997.9	*7	*	*	*	*	*	*	*	*	*	*
998	784	357	428	24	171	281	308	170	171	268	174
998.0	*8	*	*5	*	*	*	*5	*	*	*	*
998.1	256	127	130	*7	48	84	118	61	55	83	57
998.11	131	71	60	*5	21	45	61	31	22	45	33
998.12	110	49	60	*	23	34	52	25	29	34	22
998.13	15	*6	*9	*	*	*5	*	*	*	*	*
998.2	95	35	60	*	23	30	39	20	20	32	23
998.3	63	33	30	*	14	27	21	*9	12	25	18
998.31	15	*7	*8	*	*7	*	*	*	*	*	*
998.32	49	27	22	*	*7	22	18	*6	9	20	13
998.5	227	102	125	*8	53	86	79	48	53	77	49
998.51	9	*	*8	*	*	*	*	*	*	*	*
998.59	218	100	118	*8	50	83	77	47	49	75	47
998.6	19	11	*8	*	*5	*8	*6	*6	*	*6	*5
998.8	106	41	64	*	25	41	37	25	25	39	18
998.83	19	*8	11	*	*	*9	*	*	*	9	*
998.89	83	32	51	*	20	31	30	19	20	29	15
998.9	*6	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
999	43	18	25	*	9	13	18	11	*8	15	*8
999.2	15	*7	*8	*	*	*5	*6	*	*	*6	*
999.3	9	*	*	*	*	*	*	*	*	*	*
999.8	10	*	*7	*	*	*	*5	*	*	*	*
999.9	*6	*	*	*	*	*	*	*	*	*	*
V01-V83	14,069	4,298	9,771	416	6,600	2,645	4,409	2,719	3,167	5,187	2,997
V01	*7	*	*	*	*	*	*	*	*	*	*
V02	546	23	523	*	516	22	*6	102	108	213	123
V02.5	511	*	506	*	502	*	*	80	106	206	119
V02.51	498	*	498	*	496	*	—	77	105	200	116
V02.59	11	*	*6	*	*	*	*	*	*	*6	*
V02.6	33	18	14	*	12	17	*	22	*	*6	*
V02.61	*6	*	*	—	*	*	*	*	*	*	*
V02.62	26	15	10	*	*8	15	*	19	*	*	*
V03	20	10	10	*	*	*7	9	*	*	*	*8
V03.8	19	9	10	*	*	*6	9	*	*	*	*8
V03.82	19	9	10	*	*	*6	9	*	*	*	*8
V04	25	*	21	*	14	*	*6	*9	*	*9	*
V04.3	10	—	10	—	10	*	—	*	*	*	*
V04.8	15	*	10	*	*	*	*6	*6	*	*	*
V04.81	14	*	10	*	*	*	*6	*6	*	*	*
V05	10	*	*7	*5	*	*	*	*	*	*	*
V05.3	*8	*	*5	*5	*	*	*	*	*	*	*
V06	14	*	9	*	9	*	*	*	*	*	*
V06.4	*8	—	*8	—	*8	—	—	*	*	*	*
V07	52	*	47	*	40	*6	*	11	*9	18	14
V07.2	35	*	35	*	34	*	*	*8	*7	*8	11
V07.4	*7	—	*7	*	*	*	*	*	*	*	*
V07.8	*7	*	*	*	*	*	*	*	*	*	*
V08	60	35	25	*	30	27	*	29	*	25	*
V09	166	94	72	26	51	39	49	31	34	69	33
V09.0	159	92	67	26	51	38	44	28	32	67	31
V10	936	407	529	*	60	263	610	242	225	292	177
V10.0	198	111	88	*	*8	58	131	54	44	66	34
V10.02	14	*7	*8	—	*	*	10	*	*	*	*5
V10.03	*5	*	*	—	*	*	*	*	*	*	*
V10.04	10	*7	*	—	*	*	*6	*	*	*	*
V10.05	132	70	62	—	*	33	95	40	29	46	17
V10.06	19	12	*7	—	*	*9	9	*	*6	*6	*
V10.09	9	*5	*	—	*	*	*	*	*	*	*
V10.1	63	32	31	—	*	14	48	16	20	21	*6
V10.11	63	32	31	—	*	14	48	16	20	21	*6
V10.2	13	*9	*	—	*	*	*8	*	*	*6	*
V10.21	11	*8	*	—	*	*	*8	*	*	*6	*
V10.3	239	*	236	—	15	71	154	62	51	79	47
V10.4	212	130	82	*	14	51	147	54	58	57	42
V10.41	27	—	27	—	*6	14	*7	*6	*8	10	*
V10.42	28	—	28	—	*	9	17	*8	*9	*7	*
V10.43	22	—	22	—	*	9	10	*	*6	*6	*5
V10.46	122	122	—	*	*	14	108	32	34	30	26
V10.47	*6	*6	—	—	*	*	*	*	*	*	*
V10.5	76	52	23	*	*	17	55	20	15	28	13
V10.51	42	29	13	—	*	*8	34	12	*8	14	*7
V10.52	25	18	*8	*	*	*7	16	*7	*5	10	*
V10.7	25	13	12	*	*5	*7	13	*6	*9	*7	*
V10.72	*6	*	*	*	*	*	*	*	*	*	*
V10.79	19	10	*9	*	*	*	12	*	*7	*5	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V10.8	103	53	50	*	11	38	51	26	25	24	28
V10.82d	25	13	12	—	*	11	11	*7	*8	*	*6
V10.83	36	22	14	—	*	*9	27	*8	*9	10	10
V10.85	*9	*	*	*	*	*	*	*	*	*	*
V10.87	16	*	13	—	*	*8	*	*	*	*	*
V10.89	11	*5	*5	*	*	*	*	*	*	*	*
V11	24	13	10	—	*9	12	*	*	*6	*7	*7
V11.3	12	*9	*	—	*	*8	*	*	*	*	*
V11.8	*5	*	*	—	*	*	*	*	*	*	*
V12	785	332	452	12	105	196	472	179	205	274	127
V12.0	39	17	22	*	16	11	11	*	9	15	10
V12.01	12	*7	*	*	*	*	*5	*	*	*	*
V12.09	22	*7	15	*	13	*5	*	*	*6	*9	*
V12.2	*8	*	*5	*	*	*	*	*	*	*	*
V12.4	10	*	*6	*	*	*	*	*	*	*	*
V12.49	*5	*	*	*	*	*	*	*	*	*	*
V12.5	589	239	351	*	58	142	385	144	156	200	90
V12.50	*6	*	*	*	*	*	*	*	*	*	*
V12.51	197	67	130	*	36	58	102	47	59	63	29
V12.59	382	167	215	*	21	80	277	93	95	136	58
V12.6	14	*7	*7	*	*	*	*	*	*	*5	*
V12.7	120	62	58	*	20	34	64	21	32	47	21
V12.71	64	33	31	*	11	18	34	*9	14	28	13
V12.72	32	19	13	—	*	*8	20	*6	12	11	*
V12.79	23	10	13	*	*	*8	9	*5	*7	*7	*
V13	101	44	58	13	34	29	26	16	33	30	22
V13.0	64	33	31	*	20	23	19	11	22	18	13
V13.01	52	30	22	*	17	20	14	*8	19	15	11
V13.09	11	*	*7	*	*	*	*	*	*	*	*
V13.2	9	—	9	—	*8	*	*	*	*	*	*
V13.29	*7	—	*7	—	*6	*	*	*	*	*	*
V13.6	*7	*	*	*	*	*	*	*	*	*	*
V13.69	*6	*	*	*	*	*	—	*	*	*	*
V13.7	*7	*	*	*7	*	—	*	—	*	*	*
V13.8	*8	*	*6	*	*	*	*	*	*	*	*
V14	173	47	126	*9	71	47	46	19	30	99	25
V14.0	87	26	61	*	34	27	23	10	17	44	16
V14.1	10	*	*8	*	*5	*	*	*	*	*	*
V14.2	25	*	21	*	10	*7	*8	*	*	16	*
V14.3	*6	*	*	*	*	*	*	*	*	*	*
V14.5	13	*	9	*	*6	*	*	*	*	*8	*
V14.6	13	*	10	*	*7	*	*	*	*	10	*
V14.8	16	*	11	*	*6	*	*	*	*	10	*
V15	998	536	462	37	237	374	350	202	224	356	216
V15.0	36	12	24	10	13	10	*	*9	*8	16	*
V15.05	*5	*	*	*	*	*	*	*	*	*	*
V15.07	*6	*	*5	*	*	*	*	*	*	*	*
V15.08	*7	*	*	—	*	*	*	*	*	*	*
V15.09	*9	*	*6	*	*	*	*	*	*	*	*
V15.1	23	11	12	12	*6	*	*	*8	*	*7	*
V15.3	23	11	13	*	*	11	*9	*	*5	*7	*7
V15.4	20	*7	13	*	10	*	*	*	*	*9	*
V15.41	15	*5	10	*	*9	*	*	*	*	*7	*
V15.5	28	15	13	*	13	*9	*	*8	*5	10	*
V15.8	863	478	385	*6	191	339	328	169	197	305	192
V15.81	245	137	108	*	109	89	42	54	67	86	36
V15.82	596	328	267	*	77	244	275	108	126	213	148
V15.89	18	10	*9	*	*5	*	*8	*6	*	*	*7

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V16	87	27	60	*	31	34	19	10	22	35	19
V16.0	22	*8	13	*	*7	*7	*8	*	*7	*7	*
V16.1	*8	*	*	*	*	*	*	*	*	*	*
V16.3	23	*	21	*	10	*8	*	*	*	13	*
V16.4	16	*6	10	*	*7	*6	*	*	*	*5	*
V16.41	*6	—	*6	*	*	*	*	*	*	*	*
V16.42	*6	*6	—	—	*	*	*	*	*	*	*
V16.8	*6	*	*	*	*	*	*	*	*	*	*
V16.9	*6	*	*	*	*	*	*	*	*	*	*
V17	223	123	100	18	61	103	41	42	60	84	38
V17.1	12	*6	*6	*	*	*5	*	*	*	*5	*
V17.3	138	79	59	*	34	75	29	32	32	50	24
V17.4	50	26	25	*	17	20	10	*6	16	20	*8
V17.5	12	*7	*5	11	*	*	*	*	*6	*	*
V18	55	25	30	*8	22	18	*6	*5	15	19	15
V18.0	47	22	25	*6	19	17	*	*	11	18	14
V19	*8	*	*	*	*	*	*	*	*	*	*
V19.8	*5	*	*	*	*	*	*	*	*	*	*
V21	12	*8	*	11	*	*	—	*	*	*	*6
V21.3	11	*8	*	11	*	*	—	*	*	*	*6
V21.30	*5	*	*	*5	*	*	—	*	*	*	*
V22	13	—	13	*	13	—	—	*	*	*8	*
V22.1	*5	—	*5	*	*5	—	—	*	*	*	*
V23	153	—	153	*	151	*	—	15	23	76	40
V23.2	16	—	16	—	16	—	—	*	*	12	*
V23.4	16	—	16	—	16	*	—	*	*	*6	*
V23.41	*9	—	*9	—	*9	*	—	*	*	*	*
V23.49	*7	—	*7	—	*7	—	—	*	*	*	*
V23.5	*9	—	*9	—	*9	—	—	*	*	*	*
V23.7	97	—	97	*	96	*	—	*7	15	46	29
V23.8	*6	—	*6	*	*5	*	—	*	*	*	*
V25	313	*	312	—	310	*	—	35	54	156	67
V25.2	307	*	306	—	304	*	—	35	54	151	67
V26	16	*	15	—	11	*	*	*	*	10	*
V26.5	16	*	15	—	11	*	*	*	*	9	*
V26.51	15	—	15	—	10	*	*	*	*	*9	*
V27	4,038	—	4,038	*6	4,025	*7	—	591	824	1,574	1,049
V27.0	3,817	—	3,817	*6	3,805	*6	—	544	771	1,500	1,002
V27.1	17	—	17	—	17	—	—	*	*	*6	*6
V27.2	60	—	60	—	59	*	—	11	*6	29	13
V27.9	142	—	142	*	141	*	—	34	43	38	27
V29	18	10	*8	18	—	—	—	*	*	10	*
V29.0	17	10	*7	17	—	—	—	*	*	9	*
V30	*8	*	*5	*8	—	—	—	*	*	*	*
V30.1	*8	*	*5	*8	—	—	—	*	*	*	*
V42	138	81	57	17	31	47	43	29	27	51	31
V42.0	48	28	21	*	18	21	*8	*7	11	19	11
V42.1	*8	*6	*	*	*	*	*	*	*	*	*
V42.2	31	18	13	—	*	*	27	*7	*6	*9	9
V42.7	23	12	12	*6	*	12	*	*6	*	9	*
V42.8	24	15	9	*8	*7	*7	*	*7	*	*8	*
V42.81	*8	*5	*	*	*	*	*	*	*	*	*
V42.82	*6	*	*	*	*	*	*	*	*	*	*
V42.83	*6	*	*	*	*	*	*	*	*	*	*
V43	477	195	282	*	20	127	329	120	120	153	85
V43.3	123	72	51	*	11	32	79	30	25	44	24
V43.6	346	121	225	*	*9	91	246	89	94	106	57
V43.64	148	55	93	*	*6	34	107	37	42	43	26
V43.65	190	65	125	*	*	55	132	50	50	61	28

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V44	184	91	94	46	33	40	66	51	42	50	42
V44.0	32	19	13	12	*6	*8	*6	*7	*6	11	*8
V44.1	78	36	42	29	12	10	27	24	17	20	18
V44.2	21	11	10	*	*6	*8	*6	*	*	*7	*7
V44.3	40	17	23	*	*5	11	22	11	12	9	*7
V44.6	*7	*	*	*	*	*	*	*	*	*	*
V45	1,842	1,004	838	43	205	533	1,060	413	471	639	319
V45.0	295	157	138	*	14	44	235	85	79	95	36
V45.01	236	115	121	*	9	27	199	70	60	77	29
V45.02	52	37	15	*	*	16	32	14	16	17	*5
V45.09	*5	*	*	—	*	*	*	*	*	*	*
V45.1	118	64	53	*	21	45	50	18	30	42	28
V45.2	34	21	13	18	*8	*	*	*7	*7	11	*8
V45.3	49	10	39	*	21	24	*	*9	14	16	10
V45.4	54	24	30	*	11	25	17	*8	13	20	14
V45.6	10	*5	*	*	*	*	*9	*	*	*	*
V45.61	9	*	*	*	*	*	*9	*	*	*	*
V45.7	203	55	148	*	47	76	77	30	55	79	39
V45.71	37	*	36	—	*7	14	16	*5	*9	14	*9
V45.72	20	11	9	*	*5	*6	*7	*	*6	*6	*
V45.73	26	14	12	*	*6	*7	13	*	*6	12	*
V45.76	11	*	*7	*	*	*	*6	*	*	*	*
V45.77	59	*	54	—	13	28	17	*6	18	24	11
V45.79	42	15	27	*	14	13	14	*8	11	14	*9
V45.8	1,079	668	412	18	83	315	663	254	269	374	182
V45.81	517	348	168	—	*8	126	383	139	120	185	73
V45.82	371	236	134	—	18	137	215	80	102	119	70
V45.89	189	82	107	18	55	51	65	35	47	68	39
V46	85	37	48	*8	*7	19	53	17	22	27	19
V46.1	22	11	11	*	*	*	*9	*5	*7	*6	*
V46.11	22	11	11	*	*	*	*9	*5	*7	*6	*
V46.2	64	26	37	*	*	15	44	12	15	21	16
V49	82	45	37	*	12	33	36	20	25	28	9
V49.6	*5	*	*	—	*	*	*	*	*	*	*
V49.7	65	38	28	*	*9	23	33	16	18	24	*7
V49.72	*7	*	*	*	*	*	*	*	*	*	*
V49.75	31	15	16	*	*	12	14	*6	10	11	*
V49.76	20	14	*6	*	*	*6	12	*6	*	*9	*
V49.8	11	*	*9	*	*	*8	*	*	*	*	*
V49.81	*6	—	*6	—	*	*	*	*	*	*	*
V50	13	*	*8	*	*	*	*	*	*	*5	*
V50.4	*6	*	*6	—	*	*	*	*	*	*	*
V53	31	17	13	*	*	*5	23	*6	*9	*8	*8
V53.3	25	15	10	*	*	*	21	*	*8	*5	*7
V53.31	17	*9	*8	—	*	*	16	*	*6	*	*
V53.32	*7	*6	*	*	*	*	*	*	*	*	*
V54	155	50	106	*	18	28	106	31	38	43	43
V54.0	*9	*	*6	*	*	*	*	*	*	*	*
V54.01	*6	*	*	*	*	*	*	*	*	*	*
V54.1	75	22	53	*	13	11	51	11	20	21	22
V54.11	*9	*	*7	—	*	*	*8	*	*	*	*
V54.13	22	*6	15	*	*	*	19	*	*6	*8	*
V54.15	*5	*	*	*	*	*	*	*	*	*	*
V54.16	*8	*	*5	*	*	*	*	*	*	*	*
V54.17	*7	*	*5	—	*	*	*	*	*	*	*
V54.19	20	*7	13	*	*5	*	10	*	*5	*5	*6
V54.8	68	23	45	*	*	15	50	18	16	19	16
V54.81	52	18	34	—	*	12	40	15	11	14	12
V54.89	16	*5	10	*	*	*	10	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V55	48	24	24	*6	*9	18	15	11	10	17	9
V55.1	*6	*	*	*	*	*	*	*	*	*	*
V55.2	12	*	*8	*	*	*6	*	*	*	*	*
V55.3	21	11	10	*	*	*8	*8	*	*6	*8	*
V56	116	62	54	*	22	56	38	17	10	47	41
V56.0	109	60	50	*	18	54	37	17	10	43	40
V56.8	*6	*	*	*	*	*	*	*	*	*	*
V57	567	229	338	*	46	105	411	116	136	207	107
V57.1	118	51	67	*	11	19	86	27	14	54	23
V57.2	22	*7	15	*	*	*	13	*6	*5	10	*
V57.21	22	*7	15	*	*	*	13	*6	*5	10	*
V57.8	423	170	253	*	31	81	308	83	117	141	82
V57.89	423	170	253	*	31	81	308	83	117	141	82
V58	977	467	510	51	160	300	466	213	239	345	179
V58.0	20	11	*9	*	*	*7	*8	*	*	10	*6
V58.1	159	91	68	37	29	56	36	38	21	72	27
V58.4	17	*5	12	*	*	*	9	*6	*	*6	*
V58.49	11	*	*8	*	*	*	*6	*	*	*	*
V58.6	735	339	396	9	118	224	384	160	196	246	134
V58.61	257	114	144	*	20	63	174	55	68	84	51
V58.62	*6	*	*	*	*	*	*	*	*	*	*
V58.63	14	*8	*6	—	*	*6	*6	*	*	*8	*
V58.65	48	20	27	*	9	16	20	10	17	11	10
V58.66	111	65	47	*	*7	41	63	26	34	32	20
V58.67	246	107	139	*	69	80	94	58	58	88	43
V58.69	49	23	26	*	*8	15	23	*8	12	22	*7
V58.7	17	*6	11	*	*	*	11	*	*	*	*5
V58.73	*6	*	*	*	*	*	*	*	*	*	*
V58.78	*6	*	*	—	*	*	*	*	*	*	*
V58.8	24	12	12	*	*	*	15	*	12	*7	*
V58.83	9	*6	*	—	*	*	*7	*	*	*	*
V58.89	12	*	*8	*	*	*	*8	*	*7	*	*
V59	9	*	*6	*	*	*	*	*	*	*	*
V59.4	*7	*	*	*	*	*	*	*	*	*	*
V60	62	41	21	*	32	22	*7	15	14	18	14
V60.0	42	31	12	*	24	16	*	11	*7	12	11
V60.2	10	*6	*	*	*5	*	*	*	*	*	*
V61	51	15	36	*9	34	*7	*	*8	20	15	*8
V61.0	9	*	*	*	*6	*	*	*	*	*	*
V61.1	*8	*	*6	—	*6	*	*	*	*	*	*
V61.10	*8	*	*5	—	*6	*	*	*	*	*	*
V61.2	14	*	11	*6	*8	*	—	*	*5	*	*
V61.20	11	*	*9	*	*7	*	—	*	*	*	*
V61.5	9	—	9	—	9	—	—	*	*	*	*
V61.8	*6	*	*	*	*	*	*	*	*	*	*
V62	72	41	31	*	46	19	*	23	22	19	*8
V62.0	12	*8	*	—	*6	*6	—	*	*	*	*
V62.8	46	24	22	*	31	10	*	15	17	11	*
V62.81	10	*	*6	*	*8	*	*	*	*7	*	—
V62.89	32	17	15	*	22	*7	*	12	9	*9	*
V64	128	54	74	*	40	50	34	27	28	47	26
V64.1	23	9	14	*	*	9	9	*7	*6	*6	*
V64.2	19	*8	11	*	*7	*6	*7	*	*	*8	*
V64.3	19	*9	10	*	*	9	*	*	*	10	*
V64.4	65	27	38	*	24	26	14	12	17	23	13
V64.41	64	26	38	*	24	25	14	12	17	23	12
V65	38	16	22	12	15	*7	*	*5	*6	21	*5
V65.2	10	*	*5	*	*7	*	*	*	*	*	*
V65.3	24	11	14	11	*6	*	*	*	*	15	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V66	41	21	20	*	*	*6	31	*5	13	15	*7
V66.7	37	18	19	*	*	*	30	*	12	14	*6
V70	27	16	12	*	14	*6	*5	*5	*	11	*7
V70.7	25	15	10	*	12	*5	*5	*	*	10	*6
V71	15	*8	*8	*	*5	*	*6	*	*	*8	*
V71.4	*5	*	*	*	*	*	*	*	*	*	*
V71.8	*7	*	*	*	*	*	*5	*	*	*6	*
V71.89	*7	*	*	*	*	*	*5	*	*	*6	*
V72	12	*	*8	*	*	*	*6	*	*	*	*
V72.8	11	*	*7	*	*	*	*6	*	*	*	*

* Figure does not meet standards of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see "Methods" section.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
All codes	44,950	17,991	26,959	2,203	14,744	11,910	16,093	9,689	9,770	15,962	9,529
00	194	115	79	*	15	65	110	48	44	74	28
00.1	68	35	33	*	*	20	41	18	14	29	*7
00.13	26	14	12	—	*	*7	18	*	*7	13	*
00.14	*7	*	*	—	*	*	*	*	*	*	—
00.17	26	15	11	*	*	*6	16	*8	*	10	*
00.2	22	16	*6	—	*	10	11	*5	*	*8	*
00.24	19	14	*5	—	*	*8	*9	*	*	*8	*
00.3	12	*8	*	*	*	*	*	*5	*	*	*
00.39	*6	*5	*	*	*	*	*	*	*	*	*
00.5	53	33	19	*	*	15	36	10	17	16	10
00.51	39	25	14	—	*	12	26	*8	11	13	*7
00.52	*6	*	*	—	*	*	*	*	*	*	*
00.6	24	15	9	*	*	*6	17	*6	*5	10	*
00.61	13	*8	*	*	*	*	*9	*	*	*5	*
00.63	10	*6	*	—	*	*	*7	*	*	*	*
00.9	15	*8	*7	*	*	*9	*	*	*	*8	*
00.93	10	*5	*	*	*	*6	*	*	*	*5	*
01-05	1,227	569	658	202	358	310	357	253	324	384	267
01	158	98	60	22	35	46	55	31	31	47	50
01.0	11	*6	*	*7	*	*	*	*	*	*	*
01.02	*8	*	*	*6	*	*	*	*	*	*	*
01.1	30	20	9	*	*8	11	*9	*5	*8	9	*7
01.13	10	*6	*	*	*	*	*	*	*	*	*
01.14	*6	*	*	*	*	*	*	*	*	*	*
01.18	13	9	*	*	*5	*	*	*	*6	*	*
01.2	20	12	*8	*	*5	*	*7	*	*	*5	*8
01.24	12	*7	*	*	*	*	*	*	*	*	*
01.3	40	28	12	*	*	12	20	*6	*8	11	14
01.31	32	23	*9	*	*	10	17	*	*7	9	12
01.39	*8	*	*	*	*	*	*	*	*	*	*
01.5	56	31	25	*5	16	17	17	14	9	16	17
01.51	*8	*	*6	—	*	*	*	*	*	*	*
01.59	47	28	19	*5	13	14	15	12	*8	13	14
02	132	68	63	30	36	34	32	27	29	41	34
02.0	22	11	10	*5	*6	*7	*	*6	*	*7	*6
02.06	10	*	*6	*	*	*	*	*	*	*	*
02.1	11	*6	*6	*	*	*	*	*	*	*	*
02.12	10	*	*6	*	*	*	*	*	*	*	*
02.2	30	14	15	*	*8	9	*8	*	*7	9	*9
02.3	30	17	13	*7	*7	*	12	*5	*7	10	*8
02.34	25	14	11	*5	*5	*	11	*	*	*9	*7
02.4	25	14	10	10	*8	*	*	*	*7	*7	*7
02.42	20	10	9	*8	*8	*	*	*	*5	*7	*
02.9	15	*6	*9	*	*	*7	*	*	*	*	*
02.93	*6	*	*	*	*	*	*	*	*	*	*
02.94	*7	*	*	*	*	*	*	*	*	*	*
03	853	361	492	149	262	199	244	169	244	276	164
03.0	208	103	104	*	25	72	109	45	43	70	50
03.09	202	101	102	*	23	69	108	43	42	69	48
03.3	345	168	177	132	86	67	61	64	86	124	71
03.31	343	168	175	131	86	66	60	64	86	123	70
03.4	10	*	*6	*	*	*	*	*	*	*	*
03.5	31	15	16	*	9	10	*9	*	*5	14	*7
03.53	13	*8	*5	*	*6	*	*	*	*	*7	*
03.59	16	*7	9	*	*	*6	*5	*	*	*7	*
03.6	*5	*	*	*	*	*	*	*	*	*	*
03.9	250	66	184	10	138	43	60	52	106	61	31
03.90	84	22	62	*	48	15	19	21	30	22	10
03.91	91	15	76	*	62	12	16	18	51	14	*7

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
03.92	52	23	29	*7	11	11	24	9	17	17	*9
03.95	17	*	14	—	14	*	*	*	*6	*7	*
04	79	39	41	*	24	28	26	26	17	19	17
04.0	*7	*	*	*	*	*	*	*	*	*	*
04.3	*5	*	*	*	*	*	*	*	*	*	*
04.4	19	9	9	*	*8	*9	*	*	*	*6	*6
04.43	*8	*	*	*	*	*	*	*	*	*	*
04.49	*7	*	*	*	*	*	*	*	*	*	*
04.8	38	15	23	*	*6	13	20	18	10	*7	*
04.81	36	14	22	*	*5	13	19	17	*9	*6	*
05	*5	*	*	*	*	*	*	*	*	*	*
06–07	104	29	75	*	27	47	27	22	20	36	25
06	81	16	65	*	20	38	22	18	17	26	21
06.1	*6	*	*	*	*	*	*	*	*	*	*
06.2	23	*	18	*	*5	11	*6	*	*	*8	*6
06.3	*9	*	*7	*	*	*	*	*	*	*	*
06.39	*8	*	*6	*	*	*	*	*	*	*	*
06.4	25	*	21	*	*8	13	*	*6	*	*8	*7
06.8	9	*	*7	—	*	*	*	*	*	*	*
06.89	*9	*	*7	—	*	*	*	*	*	*	*
07	22	12	10	*	*7	*8	*6	*	*	10	*
07.2	*8	*	*	*	*	*	*	*	*	*	*
07.22	*7	*	*	*	*	*	*	*	*	*	*
07.6	*8	*	*	*	*	*	*	*	*	*	*
08–16	74	40	34	14	20	16	25	17	12	25	19
08	30	18	12	*5	11	*7	*7	*7	*6	10	*7
08.8	17	11	*6	*	*8	*	*	*	*	*	*
08.81	14	10	*	*	*7	*	*	*	*	*	*
13	*7	*	*	*	*	*	*	*	*	*	*
14	16	*9	*7	*	*	*	*7	*	*	*6	*5
14.7	*6	*	*	*	*	*	*	*	*	*	*
16	*8	*	*5	*	*	*	*	*	*	*	*
18–20	42	27	15	22	9	*	*5	9	*9	14	10
18	18	13	*	*7	*	*	*	*	*	*8	*6
18.2	*5	*	*	*	*	*	*	*	*	*	*
18.29	*5	*	*	*	*	*	*	*	*	*	*
20	21	12	9	15	*	*	*	*6	*6	*5	*
20.0	16	*8	*7	14	*	*	*	*	*	*	*
20.01	13	*7	*6	11	*	*	*	*	*	*	*
21–29	302	181	121	64	91	72	75	66	42	116	77
21	67	38	29	*6	15	19	27	15	12	26	14
21.0	32	14	18	*	*	*9	20	*9	*6	10	*7
21.01	14	*5	*8	*	*	*	*8	*	*	*	*
21.02	*6	*	*	—	*	*	*	*	*	*	*
21.03	11	*5	*5	*	*	*	*7	*	*	*	*
21.6	*6	*	*	*	*	*	*	*	*	*	*
21.8	18	13	*5	*	*7	*6	*	*	*	*7	*
21.81	*6	*	*	*	*	*	*	*	*	*	*
21.88	*6	*	*	*	*	*	*	*	*	*	*
22	30	17	14	*5	10	*8	*8	*7	*	13	*6
22.6	15	*7	*8	*	*	*	*	*	*	*6	*
22.62	*5	*	*	*	*	*	*	*	*	*	*
22.63	*8	*5	*	*	*	*	*	*	*	*	*
23	26	14	12	*5	11	*5	*	*7	*	*9	*6
23.0	12	*6	*6	*	*5	*	*	*	*	*	*
23.09	11	*5	*6	*	*5	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
23.1	13	*7	*6	*	*5	*	*	*	*	*	*
23.19	12	*6	*6	*	*	*	*	*	*	*	*
24	*8	*	*	*	*	*	*	*	*	*	*
25	13	10	*	*	*	*	*5	*	*	*	*
25.2	*5	*	*	*	*	*	*	*	*	*	*
26	11	*5	*6	*	*	*5	*	*	*	*	*
26.3	*6	*	*	*	*	*	*	*	*	*	*
27	70	43	28	22	23	10	15	11	*9	31	20
27.0	11	*	*7	*	*6	*	*	*	*	*6	*
27.4	*6	*5	*	*	*	*	*	*	*	*	*
27.5	29	16	13	9	*9	*	*9	*	*	11	11
27.51	12	*8	*	*	*6	*	*	*	*	*5	*
27.54	*7	*	*	*7	*	—	—	*	*	*	*
27.6	18	13	*	*9	*5	*	*	*	*	*7	*5
27.62	*7	*	*	*6	*	—	—	*	*	*	*
27.69	*8	*7	*	*	*	*	*	*	*	*	*
28	44	26	18	19	19	*	*	11	*8	17	*8
28.0	12	*7	*5	*	*7	*	*	*	*	*	*
28.2	14	9	*5	*	10	*	*	*	*	*5	*
28.3	12	*8	*	10	*	*	—	*	*	*5	*
29	32	25	*7	*	*6	15	*7	*7	*	10	13
29.1	*8	*5	*	*	*	*	*	*	*	*	*
29.11	*7	*	*	*	*	*	*	*	*	*	*
29.3	*7	*6	*	*	*	*	*	*	*	*	*
29.4	11	*9	*	*	*	*6	*	*	*	*	*
30–34	1,118	617	501	64	168	370	516	248	230	410	231
30	10	*6	*	*	*	*7	*	*	*	*	*
31	168	97	72	24	30	51	62	41	31	62	35
31.1	88	53	35	*	19	30	34	16	16	39	17
31.2	14	*8	*6	*	*	*	*7	*	*	*	*5
31.29	14	*8	*6	*	*	*	*7	*	*	*	*5
31.4	54	29	25	15	*7	14	19	17	11	16	10
31.42	47	24	22	15	*6	10	16	15	10	13	*9
31.43	*5	*	*	—	*	*	*	*	*	*	*
31.7	*5	*	*	*	*	*	*	*	*	*	*
32	76	41	35	*	11	28	35	18	18	26	14
32.2	37	21	16	*	*8	15	14	10	10	11	*6
32.29	34	18	16	*	*6	14	12	*9	*9	11	*5
32.3	*7	*	*	*	*	*	*	*	*	*	*
32.4	27	12	15	*	*	9	15	*6	*5	11	*
33	357	194	163	20	47	122	167	71	74	125	86
33.1	*5	*	*	—	*	*	*	—	*	*	*
33.2	341	184	156	19	42	117	163	70	70	118	84
33.21	*5	*	*	*	*	*	*	*	*	*	*
33.22	49	28	22	*	*5	15	27	11	11	14	14
33.23	38	22	16	*7	*5	12	13	10	*8	12	*8
33.24	147	76	71	*7	20	48	73	31	29	50	37
33.26	36	19	17	*	*5	11	19	*	*6	13	13
33.27	44	24	20	*	*	20	20	10	11	18	*6
33.28	20	12	*8	*	*	10	*8	*	*	9	*5
33.3	*6	*	*	*	*	*	*	*	*	*	*
34	508	280	229	16	80	162	250	117	106	194	92
34.0	190	110	80	*9	47	61	72	44	38	74	33
34.03	10	*8	*	*	*	*6	*	*	*	*	*
34.04	133	76	57	*7	33	40	53	35	27	51	20
34.09	40	22	18	*	11	13	14	*7	*8	15	10
34.2	56	29	26	*	*7	20	27	15	13	20	*8
34.21	22	13	9	*	*	*9	*9	*7	*6	*7	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
34.22	15	*7	*8	—	*	*	*8	*	*	*	*
34.24	9	*6	*	—	*	*	*6	*	*	*	*
34.25	*5	*	*	*	*	*	*	*	*	*	*
34.45	26	19	*7	*	*	11	*9	*6	*	13	*
34.51	21	15	*6	*	*	9	*6	*	*	10	*
34.6	*8	*5	*	*	*	*	*	*	*	*	*
34.7	11	*8	*	*	*	*	*	*	*	*	*
34.79	*7	*	*	*	*	*	*	*	*	*	*
34.9	206	102	105	*	14	60	131	46	46	76	39
34.91	183	92	91	*	12	51	119	39	39	68	37
34.92	22	*9	13	*	*	*8	12	*6	*8	*	*
35–39	6,989	4,062	2,927	200	655	2,524	3,609	1,340	1,550	2,608	1,491
35	147	81	66	26	13	39	68	39	29	47	32
35.1	25	12	13	*	*	10	10	*6	*5	10	*
35.12	16	9	*7	*	*	*6	*8	*	*	*5	*
35.	81	45	36	*	*6	25	48	18	17	24	22
35.21	30	17	13	*	*	*	23	*8	*7	*6	9
35.22	26	16	10	—	*	12	13	*	*5	*9	*7
35.23	10	*	*5	*	*	*	*7	*	*	*	*
35.24	12	*5	*7	*	*	*5	*	*	*	*	*
35.3	*7	*	*	*	*	*	*	*	*	*	*
35.33	*6	*	*	*	*	*	*	*	*	*	*
35.5	*5	*	*	*	*	*	*	*	*	*	*
35.6	*7	*	*	*	*	*	*	*	*	*	*
35.7	*9	*6	*	*	*	*	*	*	*	*	*
35.71	*7	*	*	*	*	*	*	*	*	*	*
35.9	*8	*	*	*5	*	*	*	*	*	*	*
36	1,742	1,200	542	*	84	751	905	305	471	600	366
36.0	1,271	874	397	*	69	563	639	226	358	425	262
36.01	547	376	171	*	31	239	277	102	157	183	105
36.02	12	*8	*	—	*	*6	*6	*	*	*	*
36.05	86	59	27	—	*	40	43	15	24	26	22
36.06	86	61	25	*	*	33	48	15	19	30	22
36.07	534	368	167	*	29	241	263	90	157	178	109
36.1	466	321	144	*	15	187	264	77	111	174	103
36.11	45	27	18	—	*	15	28	*9	12	15	9
36.12	93	60	33	—	*	38	53	16	17	35	25
36.13	70	50	20	*	*	31	38	11	22	27	11
36.14	39	31	*7	—	*	13	23	*	*8	14	12
36.15	206	142	64	*	*6	83	116	34	50	77	45
36.16	*8	*7	*	—	*	*	*	*	*	*	*
37	1,936	1,155	781	20	136	676	1,103	395	449	700	391
37.0	13	*7	*7	*	*	*5	*5	*	*	*6	*
37.1	15	10	*6	*	*	*6	*6	*	*	*6	*
37.12	14	*9	*5	*	*	*6	*5	*	*	*5	*
37.2	1,322	808	513	11	104	548	659	238	316	509	259
37.21	19	10	9	*	*	*8	*8	*5	*	*8	*
37.22	1,071	670	401	*	77	464	530	175	263	420	212
37.23	119	63	57	*8	*7	37	67	29	23	40	27
37.26	69	41	28	*	*9	25	34	19	16	26	*8
37.27	24	14	10	*	*6	*8	9	*6	*6	*8	*
37.29	*8	*	*	*	*	*	*	*5	*	*	*
37.3	65	36	29	*	10	19	35	16	16	23	10
37.33	16	9	*7	*	*	*	9	*	*	*7	*
37.34	46	25	21	*	*8	13	24	12	12	15	*8
37.6	42	26	16	—	*	18	22	*8	11	13	10
37.61	40	25	15	—	*	16	22	*7	11	13	9
37.7	203	107	97	*	*5	25	170	57	39	59	49
37.71	29	14	15	*	*	*	27	*9	*6	10	*5
37.72	118	60	58	*	*	14	101	34	24	34	26

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
37.75	*8	*5	*	*	*	*	*7	*	*	*	*
37.76	*9	*	*	*	*	*	*7	*	*	*	*
37.78	22	13	*9	*	*	*	17	*	*	*5	*8
37.78	180	93	87	*	*	19	155	51	35	55	40
37.81	14	*7	*7	—	*	*	13	*	*	*6	*
37.82	14	*6	*8	*	*	*	13	*	*	*	*
37.83	123	63	59	*	*	14	105	36	23	37	26
37.87	16	*8	*7	*	*	*	13	*	*	*	*
37.89	*6	*	*	*	*	*	*	*	*	*	*
37.9	92	68	24	*	*8	35	50	19	28	29	16
37.94	70	50	19	*	*5	28	37	15	21	21	12
37.98	9	*7	*	*	*	*	*6	*	*	*	*
37.99	*8	*6	*	*	*	*	*	*	*	*	*
38	1,807	873	933	123	271	553	860	350	334	743	381
38.0	27	15	12	*	*	*9	16	*7	*6	10	*5
38.08	17	9	*8	*	*	*6	10	*	*	*6	*
38.1	127	75	52	—	*	32	95	23	23	55	26
38.12	103	62	42	—	*	21	82	18	20	46	20
38.18	19	12	*7	—	*	*8	10	*	*	*8	*
38.2	*9	*	*5	—	*	*	*7	*	*	*	*
38.21	*6	*	*	—	*	*	*5	*	*	*	*
38.4	35	25	9	*	*	9	21	*8	*6	12	*8
38.44	16	11	*	—	*	*	13	*	*	*8	*
38.45	11	*8	*	*	*	*	*	*	*	*	*
38.6	*8	*	*	*	*	*	*	*	*	*	*
38.7	85	38	47	*	9	25	51	25	13	35	12
38.8	26	16	10	11	*5	*6	*	*5	*6	*8	*6
38.85	12	*6	*6	10	*	*	*	*	*	*	*
38.9	1,485	693	792	110	247	467	660	275	276	613	321
38.91	96	57	40	17	13	26	40	13	18	42	23
38.92	9	*7	*	9	—	*	*	*	*	*	*
38.93	1,183	528	655	77	201	374	531	225	223	489	247
38.94	*8	*	*	*	*	*	*	*	*	*	*
38.95	156	85	72	*	29	57	70	27	28	62	39
38.98	26	11	15	*	*	*7	16	*9	*	14	*
38.99	*5	*	*	*	*	*	*	*	*	*	*
39	1,357	753	604	30	150	505	672	251	267	518	322
39.2	118	71	47	*	*8	45	64	23	18	52	26
39.25	17	11	*6	*	*	*7	9	*	*	*7	*
39.27	34	19	15	*	*5	15	14	*7	*	13	10
39.29	63	39	24	—	*	22	39	12	10	29	11
39.3	25	11	15	*	13	*	*8	*	10	*6	*7
39.31	16	*7	10	*	*9	*	*5	*	*7	*	*
39.32	*7	*	*	*	*	*	*	*	*	*	*
39.4	66	37	29	*	*7	27	31	*8	10	26	21
39.42	18	11	*7	—	*	9	*6	*	*	*7	*8
39.43	*6	*	*	—	*	*	*	*	*	*	*
39.49	39	21	18	*	*	15	21	*6	*7	15	11
39.5	178	91	87	*	15	53	105	34	42	71	31
39.50	136	69	68	*	*8	39	87	28	34	53	22
39.51	*9	*	*7	—	*	*	*	*	*	*	*
39.52	9	*5	*	—	*	*	*	*	*	*	*
39.56	*6	*	*	*	*	*	*	*	*	*	*
39.57	*5	*	*	*	*	*	*	*	*	*	*
39.59	*7	*	*	*	*	*	*	*	*	*	*
39.6	245	156	89	16	17	95	117	41	56	84	64
39.61	223	143	80	13	15	89	107	39	53	79	53
39.62	*7	*	*	*	*	*	*	—	*	*	*7
39.63	*6	*	*	*	*	*	*	*	*	*	*
39.64	*6	*	*	*	*	*	*	*	*	*	*
39.7	44	30	14	*	*	11	29	10	10	17	*8

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
39.71	21	16	*5	—	—	*18	18	*	*6	*8	*
39.72	13	*7	*6	*	*	*	*	*	*	*6	*
39.79	11	*8	*	*	*	*	*7	*	*	*	*
39.9	674	354	321	*	84	268	319	131	120	260	163
39.90	59	31	28	*	*	16	41	15	16	23	*6
39.93	*7	*	*	*	*	*	*	*	*	*	*
39.95	593	311	282	*	76	245	270	112	99	228	154
39.98	13	*7	*5	*	*	*	*	*	*	*5	*
40–41	343	168	175	17	57	137	131	66	68	125	84
40	227	105	123	*	34	97	92	45	48	78	57
40.1	57	23	34	*	*8	22	26	14	14	19	10
40.11	54	23	32	*	*8	20	25	14	13	19	9
40.2	66	20	46	*	12	28	26	11	14	28	13
40.23	20	*	19	—	*	10	*7	*	*	*9	*
40.24	*9	*	*	*	*	*	*	*	*	*	*
40.29	33	13	21	*	*5	12	15	*6	*8	15	*
40.3	67	39	28	*	9	32	25	12	13	21	20
40.4	18	11	*7	*	*	*7	9	*	*	*	*8
40.41	17	11	*6	*	*	*6	*9	*	*	*	*7
40.5	17	10	*7	—	*	*8	*6	*	*	*	*6
40.53	*5	*	*	—	*	*	*	*	*	*	*
40.59	*9	*5	*	—	*	*	*	*	*	*	*
41	115	63	52	13	23	40	40	21	20	47	27
41.0	*8	*	*	*	*	*	*	*	*	*	*
41.3	80	45	35	10	13	27	30	14	15	34	17
41.31	80	44	35	10	13	27	30	14	15	34	17
41.5	22	13	10	*	*	*8	*8	*	*	*8	*7
42–54	5,580	2,414	3,166	246	1,244	1,716	2,373	1,146	1,237	2,014	1,183
42	107	66	42	*7	12	37	52	24	26	34	24
42.2	37	23	13	*5	*	12	16	11	*9	11	*6
42.23	29	20	9	*	*	9	13	10	*6	*9	*
42.24	*7	*	*	*	*	*	*	*	*	*	*
42.3	28	20	*9	*	*	16	*7	*	*5	*8	11
42.33	27	19	*8	*	*	15	*7	*	*	*7	11
42.4	*5	*	*	*	*	*	*	*	*	*	*
42.9	30	14	16	*	*	*5	23	*5	*9	11	*
42.92	30	14	16	*	*	*	23	*5	*9	11	*
43	233	118	115	28	16	50	139	48	37	97	51
43.1	171	86	86	13	11	31	116	36	25	73	38
43.11	137	70	67	*	*6	24	102	29	21	57	29
43.19	34	16	18	9	*	*7	13	*6	*	16	*9
43.3	15	13	*	14	—	*	*	*	*	*	*
43.4	22	*9	13	*	*	*9	11	*	*	9	*
43.41	15	*6	9	—	*	*	*9	*	*	*8	*
43.42	*6	*	*	*	*	*	*	*	*	*	*
43.7	*9	*	*6	—	*	*	*	*	*	*	*
43.9	*8	*	*	—	*	*	*	*	*	*	*
43.99	*8	*	*	—	*	*	*	*	*	*	*
44	305	124	182	18	95	111	82	60	57	110	78
44.0	*5	*	*	*	*	*	*	*	*	*	*
44.1	25	*9	17	*	*5	11	*9	*	*6	12	*
44.13	13	*	9	*	*	*5	*	*	*	*6	*
44.14	11	*	*7	*	*	*	*5	*	*	*	*
44.2	13	*8	*5	*	*6	*	*	*	*	*5	*
44.22	*7	*	*	—	*	*	*	*	*	*	*
44.29	*6	*	*	*	*	*	*	*	*	*	*
44.3	105	28	77	*	48	46	10	23	21	32	29
44.31	*6	*	*5	—	*	*	*	*	*	*	*
44.32	*8	*6	*	*	*	*	*	*	*	*	*
44.38	57	11	46	—	29	27	*	10	10	18	19

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
44.39	34	10	24	*	15	15	*	10	*8	11	*6
44.4	78	46	32	*	*9	21	48	13	18	28	18
44.42	*9	*5	*	*	*	*	*	*	*	*	*
44.43	63	39	24	*	*7	16	40	10	15	22	15
44.46	59	26	33	15	16	16	11	*9	*8	22	20
44.49	11	*	*7	*5	*	*	*	*	*	*	*
44.67	29	13	16	*6	*8	10	*	*5	*	11	9
44.69	*7	*	*	*	*	*	*	*	*	*	*5
44.9	15	*	11	—	*6	*9	*	*7	*	*7	*
44.95	13	*	9	—	*5	*7	*	*6	*	*6	*
45	2,220	981	1,239	39	300	645	1,236	462	503	793	461
45.0	*6	*	*	*	*	*	*	*	*	*	*
45.1	1,068	482	586	19	156	319	574	215	238	389	227
45.13	441	209	232	*6	66	124	244	93	106	147	95
45.14	*7	*	*	*	*	*	*	*	*	*	*
45.16	613	268	345	12	87	189	324	119	130	235	129
45.2	558	223	335	*7	77	145	329	122	126	196	114
45.23	287	116	171	*	30	77	179	61	64	101	61
45.24	46	19	27	*	*7	10	28	*8	13	18	*7
45.25	218	86	132	*5	39	54	119	51	47	74	45
45.3	11	*	*7	*	*	*	*7	*	*	*	*
45.4	133	70	63	*	*6	32	94	30	28	44	31
45.42	106	57	50	*	*	26	77	25	23	33	25
45.43	21	12	9	—	*	*6	14	*	*	*8	*5
45.6	84	35	49	*	15	22	42	18	18	33	15
45.62	78	32	46	*	13	21	41	18	16	30	15
45.7	266	123	143	*	32	89	142	52	66	95	52
45.72	14	*6	*8	*	*	*	*5	*	*	*7	*
45.73	98	43	55	*	10	29	58	18	26	36	18
45.74	14	*7	*7	*	*	*5	*8	*	*	*	*
45.75	29	14	16	*	*	10	15	*6	*8	10	*6
45.76	89	43	46	*	11	35	43	20	19	31	17
45.79	21	10	11	*	*	*	12	*	*6	*7	*5
45.8	12	*6	*7	*	*	*	*7	*	*	*	*
45.9	79	33	46	*	*8	29	39	16	19	28	15
45.91	22	*9	14	*	*	*6	11	*5	*6	*6	*
45.93	27	10	17	*	*	11	13	*	*8	10	*
45.94	22	10	12	*	*	*9	12	*	*	*9	*
46	227	110	117	12	45	70	101	44	48	82	53
46.0	23	12	11	*	*5	*8	9	*	*	*9	*6
46.01	12	*5	*7	*	*	*	*	*	*	*	*
46.03	10	*6	*	*	*	*	*6	*	*	*6	*
46.1	41	22	19	*	*6	13	20	10	*9	12	11
46.10	23	14	*9	*	*	*7	11	*6	*6	*6	*6
46.11	11	*	*6	*	*	*	*	*	*	*	*
46.13	*8	*	*	*	*	*	*	*	*	*	*
46.2	25	13	13	*	*7	*	12	*	*	12	*
46.20	13	*6	*6	*	*	*	*6	*	*	*7	*
46.23	*7	*	*	*	*	*	*5	*	*	*	*
46.3	22	13	9	*	*	*6	12	*	*	9	*6
46.39	18	10	*9	*	*	*5	10	*	*	*9	*
46.4	12	*5	*7	*	*	*	*7	*	*	*	*
46.5	40	18	22	*	9	16	12	*8	10	13	9
46.51	17	*6	11	*	*	*7	*	*	*	*6	*
46.52	23	12	11	*	*	*9	*8	*	*8	*7	*
46.7	43	18	25	*	10	16	15	*8	10	17	*8
46.73	22	*8	13	*	*5	*7	9	*	*	9	*
46.75	*5	*	*	*	*	*	*	*	*	*	*
46.79	*7	*	*	*	*	*	*	*	*	*	*
46.8	16	*8	*8	*	*	*	10	*	*	*	*
46.85	10	*	*5	*	*	*	*8	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
47	383	192	191	84	187	76	36	74	92	120	97
47.0	341	178	163	80	170	65	27	67	79	104	91
47.01	172	83	89	37	90	36	9	29	37	56	51
47.09	169	95	74	42	80	29	18	38	42	49	40
47.1	40	14	27	*	17	11	*9	*7	12	16	*5
47.11	*8	*	*5	*	*5	*	*	*	*	*	*
47.19	32	10	21	*	12	10	*7	*6	9	12	*
48	154	74	80	*8	31	41	74	34	32	59	29
48.2	62	31	32	*	16	13	29	16	15	21	11
48.23	22	12	10	*	*6	*7	*9	*6	*6	*7	*
48.24	38	18	20	*	10	*6	19	10	*9	12	*7
48.3	24	11	13	*	*	*6	16	*	*6	*8	*6
48.36	20	*9	11	—	*	*	13	*	*6	*6	*
48.5	*7	*	*	—	*	*	*	*	*	*	*
48.6	35	14	21	*	*	11	20	*8	*6	14	*7
48.63	23	*9	14	—	*	*8	14	*5	*	10	*
48.69	*8	*	*	—	*	*	*	*	*	*	*
48.7	*6	*	*	*	*	*	*	*	*	*	*
48.8	15	10	*5	*	*7	*6	*	*	*	*8	*
48.81	15	10	*5	*	*6	*6	*	*	*	*8	*
49	40	23	17	*	14	12	12	*8	12	12	*8
49.0	*9	*5	*	*	*	*	*	*	*	*	*
49.01	*7	*	*	*	*	*	*	*	*	*	*
49.2	*5	*	*	*	*	*	*	*	*	*	*
49.4	14	*9	*5	—	*	*5	*	*	*	*	*
49.46	11	*7	*	—	*	*	*	*	*	*	*
49.7	*5	*	*	*	*	*	*	*	*	*	*
50	118	59	59	*	21	52	41	27	22	47	23
50.1	80	35	45	*	14	33	29	19	18	28	15
50.11	50	23	27	*	*7	22	18	12	10	19	10
50.12	20	*9	11	*	*	*8	*9	*5	*6	*5	*
50.19	9	*	*6	—	*	*	*	*	*	*	*
50.2	14	*9	*	*	*	*7	*5	*	*	*6	*
50.22	*7	*	*	*	*	*	*	*	*	*	*
50.29	*7	*	*	*	*	*	*	*	*	*	*
50.6	*5	*	*	*	*	*	*	*	*	*	*
50.9	*8	*	*	—	*	*	*	*	*	*	*
50.91	*7	*	*	—	*	*	*	*	*	*	*
51	649	244	405	*	196	195	256	122	152	235	141
51.0	*7	*	*	—	*	*	*5	*	*	*	*
51.1	53	23	30	*	12	18	24	13	10	17	13
51.10	41	16	25	*	10	15	15	11	*7	13	9
51.14	*8	*5	*	—	*	*	*6	*	*	*	*
51.2	398	143	256	*	133	128	136	74	96	154	74
51.22	94	47	47	*	18	33	42	18	26	33	17
51.23	303	94	208	*	115	94	93	56	69	120	57
51.3	*7	*	*	*	*	*	*	*	*	*	*
51.36	*6	*	*	*	*	*	*	*	*	*	*
51.4	*7	*5	*	*	*	*	*	*	*	*	*
51.5	*7	*	*	—	*	*	*	*	*	*	*
51.8	158	58	99	*	46	36	75	24	37	51	45
51.84	*7	*	*	—	*	*	*	*	*	*	*
51.85	74	27	48	*	22	17	35	11	18	26	20
51.87	37	15	22	*	10	10	17	*8	*8	11	10
51.88	38	14	24	*	13	*7	19	*	10	13	11
51.9	9	*	*5	*	*	*	*6	*	*	*	*
51.98	*8	*	*	*	*	*	*6	*	*	*	*
52	30	16	13	*	*7	12	11	*6	*	11	*7
52.1	*7	*	*	—	*	*	*	*	*	*	*
52.9	9	*5	*	*	*	*	*	*	*	*	*
52.93	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
53	251	112	138	13	48	98	92	59	60	84	47
53.0	38	33	*5	*	*	10	19	9	*7	14	*8
53.00	*5	*	*	*	*	*	*	*	*	*	*
53.03	*9	*8	*	—	*	*	*5	*	*	*	*
53.04	9	*8	*	—	*	*	*6	*	*	*	*
53.05	*8	*6	*	—	*	*	*	*	*	*	*
53.1	*9	*7	*	*5	*	*	*	*	*	*	*
53.2	*6	*	*5	—	*	*	*	*	*	*	*
53.4	46	22	25	*	13	19	14	*8	12	13	13
53.41	13	*6	*6	—	*	*	*	*	*	*	*6
53.49	34	15	18	*	*9	14	10	*7	9	10	*7
53.5	45	12	33	*	*8	19	17	14	*8	17	*
53.51	27	*6	20	*	*5	12	10	10	*	10	*
53.59	18	*5	13	*	*	*7	*7	*	*	*7	*
53.6	91	34	58	*	18	42	32	18	27	31	15
53.61	71	26	45	*	13	32	26	15	22	23	12
53.69	20	*7	13	—	*	9	*6	*	*5	*8	*
53.7	*8	*	*6	*	*	*	*	*	*	*	*
53.9	*6	*	*	*	*	*	*	*	*	*	*
54	862	294	568	30	273	318	241	179	191	329	162
54.0	21	*7	14	*	*7	*7	*6	*	*5	*9	*
54.1	57	26	31	*	24	13	17	13	13	20	11
54.11	27	11	17	*	12	*6	*7	*6	*6	11	*
54.12	14	*8	*6	*	*6	*	*	*	*	*	*
54.19	16	*7	*8	*	*6	*	*	*	*5	*	*
54.2	109	33	76	*	38	35	31	24	25	40	20
54.21	66	23	43	*	29	20	15	13	17	24	12
54.23	16	*5	11	*	*	*6	*7	*	*	*	*
54.24	10	*	*8	—	*	*	*	*	*	*	*
54.25	14	*	13	*	*	*5	*	*	*	*7	*
54.3	13	*	10	*	*	*	*6	*	*	*	*
54.4	58	13	44	*	16	23	17	12	13	22	11
54.5	348	76	272	*6	133	114	95	66	82	141	60
54.51	66	12	53	*	26	25	14	12	22	22	9
54.59	283	64	219	*6	107	89	82	54	60	118	51
54.6	11	*6	*	*	*	*	*	*	*	*	*
54.7	12	*7	*6	*	*6	*	*	*	*	*	*
54.72	*5	*	*	*	*	*	*	*	*	*	*
54.9	232	123	109	11	42	116	63	52	44	87	49
54.91	168	94	74	*	24	93	48	41	33	60	35
54.92	*7	*	*	*	*	*	*	*	*	*	*
54.93	*7	*	*	*	*	*	*	*	*	*	*
54.95	17	*7	9	*6	*	*5	*	*	*	*6	*
54.98	31	13	18	*	10	12	*7	*	*6	14	*6
55–59	999	466	533	36	236	314	413	241	183	399	175
55	163	86	77	9	46	54	54	42	30	63	28
55.0	39	18	22	*	12	13	14	10	*7	15	*8
55.03	26	13	14	*	*8	*8	10	*7	*	11	*
55.04	*6	*	*	*	*	*	*	*	*	*	*
55.2	28	14	14	*	10	*9	*8	*8	*6	12	*
55.23	24	12	12	*	9	*8	*6	*6	*5	12	*
55.4	*7	*	*	*	*	*	*	*	*	*	*
55.5	56	31	25	*	12	21	22	11	11	21	14
55.51	54	30	25	*	12	19	22	10	10	20	13
55.6	11	*7	*	*	*6	*	*	*	*	*7	*
55.69	11	*7	*	*	*6	*	*	*	*	*7	*
55.8	9	*6	*	*	*	*	*	*	*	*	*
55.87	*8	*5	*	*	*	*	*	*	*	*	*
55.9	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
56	102	51	52	10	33	34	25	24	23	41	15
56.0	53	25	27	*	22	21	10	12	11	23	*6
56.3	18	11	*6	*	*6	*7	*	*	*	*8	*
56.31	15	10	*5	*	*6	*6	*	*	*	*7	*
56.5	11	*7	*	—	*	*	*8	*	*	*	*
56.51	10	*7	*	—	*	*	*7	*	*	*	*
56.7	15	*	11	*9	*	*	*	*	*	*	*
56.74	12	*	10	*8	*	*	*	*	*	*	*
57	390	214	176	12	60	91	226	110	62	146	72
57.0	24	20	*	*	*	*	18	*	*	*7	*8
57.1	25	16	9	*	*	*	15	*	*	13	*
57.17	10	*6	*	*	*	*	*6	*	*	*7	*
57.18	12	*7	*	*	*	*	*7	*	*	*	*
57.3	143	68	75	*	28	44	66	39	26	58	20
57.32	129	60	70	*	27	41	57	36	25	51	18
57.33	12	*8	*	—	*	*	*8	*	*	*6	*
57.4	46	31	14	*	*	*8	37	13	11	17	*
57.49	46	31	14	*	*	*8	37	13	11	17	*
57.7	*7	*5	*	—	—	*	*	*	*	*	*
57.71	*7	*5	*	—	—	*	*	*	*	*	*
57.8	23	10	13	*	*8	*5	*7	*5	*	*8	*
57.81	10	*	*7	*	*6	*	*	*	*	*	*
57.83	*5	*	*	—	*	*	*	*	*	*	*
57.9	114	60	54	*	16	19	77	40	*9	37	28
57.94	102	49	53	*	15	17	66	37	*7	32	26
58	45	33	12	*	*7	9	28	*8	*9	24	*
58.5	*6	*6	*	*	*	*	*	*	*	*	*
58.6	25	18	*7	*	*	*6	15	*	*	14	*
59	299	82	216	*	90	125	80	57	60	125	56
59.0	9	*	9	—	*	*	*	*	*	*	*
59.02	*6	*	*6	—	*	*	*	*	*	*	*
59.4	*8	*	*7	—	*	*	*	*	*	*	*
59.5	22	—	22	—	*7	12	*	*	*	11	*
59.7	81	*	81	—	20	37	24	11	16	31	23
59.79	80	*	80	—	20	36	24	10	16	31	23
59.8	168	77	92	*	55	64	46	39	36	72	22
59.9	*6	*	*	*	*	*	*	*	*	*	*
60–64	233	233	—	23	15	64	131	47	53	77	56
60	168	168	—	—	*	53	113	35	38	50	44
60.1	*6	*6	—	—	—	*	*	*	*	*	*
60.11	*5	*5	—	—	—	*	*	*	*	*	*
60.2	92	92	—	—	*	16	76	21	24	27	21
60.21	10	10	—	—	—	*	*8	*	*	*	*
60.29	82	82	—	—	*	14	68	19	21	25	17
60.5	56	56	—	—	*	34	20	9	11	16	19
60.9	*5	*5	—	—	—	*	*	*	*	*	*
61	12	12	—	*	*	*	*	*	*	*	*
61.0	*8	*8	—	*	*	*	*	*	*	*	*
62	10	10	—	*	*	*	*	*	*	*	*
63	*8	*8	—	*	*	*	*	*5	*	*	*
64	35	35	—	19	*	*	*6	*8	9	13	*
64.0	21	21	—	19	*	*	*	*	*8	*7	*
64.9	*8	*8	—	*	*	*	*	*	*	*	*
65–71	1,967	—	1,967	10	1,145	613	200	276	408	888	395
65	516	—	516	*	249	215	48	71	124	225	96
65.2	38	—	38	*	34	*	*	*7	*7	17	*7
65.25	*7	—	*7	*	*6	*	—	*	*	*	*
65.29	30	—	30	*	27	*	*	*5	*5	15	*5
65.3	16	—	16	*	12	*	*	*	*	*7	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
65.39	14	—	14	*	10	*	*	*	*	*7	*
65.4	88	—	88	*	54	28	*5	14	23	34	16
65.41	14	—	14	*	10	*	*	*	*	*	*
65.49	74	—	74	*	44	25	*	11	20	29	13
65.5	*8	—	*8	—	*	*	*	*	*	*	*
65.6	321	—	321	*	114	171	37	40	74	145	63
65.61	269	—	269	*	94	144	31	35	60	123	51
65.62	11	—	11	—	*6	*	*	*	*	*	*
65.63	40	—	40	—	12	24	*	*	*9	20	*8
65.8	26	—	26	*	20	*5	*	*	*6	10	*5
65.81	*6	—	*6	*	*5	*	*	*	*	*	*
65.89	20	—	20	*	14	*5	*	*	*	*8	*
65.9	*8	—	*8	*	*	*	—	*	*	*	*
65.91	*8	—	*8	*	*	*	—	*	*	*	*
66	416	—	416	*	406	*8	*	46	74	201	95
66.0	*5	—	*5	*	*	—	*	*	*	*	*
66.2	*6	—	*6	—	*6	*	—	*	*	*	*
66.3	340	—	340	—	338	*	—	35	59	169	77
66.32	249	—	249	—	247	*	—	24	39	127	59
66.39	90	—	90	—	90	*	—	11	20	42	17
66.4	*8	—	*8	*	*6	*	*	*	*	*	*
66.5	10	—	10	—	*7	*	*	*	*	*5	*
66.51	10	—	10	—	*6	*	*	*	*	*	*
66.6	38	—	38	*	37	*	*	*6	*7	16	*9
66.61	*8	—	*8	*	*7	*	—	*	*	*	*
66.62	20	—	20	—	20	—	—	*	*	11	*
66.69	*9	—	*9	*	*8	—	*	*	*	*	*
67	15	—	15	*	11	*	*	*	*	10	*
67.5	*5	—	*5	*	*	*	—	*	*	*	*
67.59	*5	—	*5	*	*	*	—	*	*	*	*
68	635	—	635	*	331	255	49	98	137	276	124
68.1	16	—	16	—	*9	*6	*	*	*	*8	*
68.12	10	—	10	—	*5	*	*	*	*	*6	*
68.2	42	—	42	—	33	*8	*	10	*	20	*7
68.29	39	—	39	—	31	*7	*	10	*	18	*6
68.3	43	—	43	—	23	18	*	10	*	16	12
68.31	13	—	13	—	*7	*6	*	*	*	*	*5
68.39	29	—	29	—	16	12	*	*8	*	12	*6
68.4	345	—	345	*	175	148	21	49	81	156	59
68.5	179	—	179	—	87	72	20	24	43	71	41
68.51	68	—	68	—	39	27	*	*8	17	33	10
68.59	111	—	111	—	48	45	18	15	26	38	32
68.6	*6	—	*6	—	*	*	*	*	*	*	*
69	101	—	101	*	80	14	*6	16	14	50	21
69.0	56	—	56	—	42	10	*	*8	*8	27	14
69.02	27	—	27	—	27	*	—	*	*	13	*8
69.09	29	—	29	—	15	10	*	*6	*	14	*6
69.2	*5	—	*5	—	*	*	*	*	*	*	*
69.5	25	—	25	*	25	*	*	*	*	*	15
69.52	19	—	19	—	19	—	—	*	*	11	*
69.9	*8	—	*8	*	*7	*	*	*	*	*	*
69.96	*6	—	*6	—	*6	—	—	*	*	*	*
70	249	—	249	*	52	111	84	36	51	108	53
70.1	*5	—	*5	*	*	*	*	*	*	*	*
70.2	*5	—	*5	*	*	*	*	*	*	*	*
70.3	10	—	10	*	*8	*	*	*	*	*6	*
70.32	*5	—	*5	—	*5	*	*	*	*	*	*
70.5	125	—	125	—	19	62	44	17	27	49	32
70.50	60	—	60	—	*7	29	24	*8	11	26	15
70.51	34	—	34	—	*6	16	12	*5	*9	11	9

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
70.52	31	—	31	—	*6	16	*8	*	*7	13	*7
70.7	53	—	53	*	*7	23	22	*7	13	23	*9
70.77	40	—	40	—	*	19	18	*5	10	19	*6
70.79	*8	—	*8	*	*	*	*	*	*	*	*
70.9	43	—	43	—	13	18	12	*7	*6	20	9
70.92	43	—	43	—	13	18	12	*7	*6	20	9
71	36	—	36	*	15	*8	10	*6	*8	18	*5
71.0	*6	—	*6	*	*	*	*	*	*	*	*
71.09	*6	—	*6	*	*	*	*	*	*	*	*
71.3	14	—	14	*	10	*	*	*	*	*8	*
71.7	11	—	11	*	*	*	*	*	*	*6	*
71.71	*5	—	*5	*	*	*	*	*	*	*	*
71.79	*6	—	*6	—	*	*	*	*	*	*	*
72–75	6,858	—	6,858	11	6,838	9	—	1,045	1,563	2,509	1,742
72	304	—	304	*	304	*	—	31	57	130	85
72.0	22	—	22	—	22	—	—	*	*6	11	*
72.1	29	—	29	*	29	—	—	*	*6	16	*
72.2	11	—	11	—	11	—	—	*	*	9	*
72.21	*7	—	*7	—	*7	—	—	*	*	*7	*
72.7	237	—	237	*	237	*	—	26	43	91	77
72.71	97	—	97	*	97	—	—	11	15	33	38
72.79	140	—	140	*	140	*	—	15	28	58	39
73	2,988	—	2,988	*5	2,979	*	—	426	713	1,098	751
73.0	928	—	928	*	926	*	—	120	275	317	216
73.01	149	—	149	*	149	—	—	14	47	57	31
73.09	779	—	779	*	777	*	—	106	228	260	185
73.1	12	—	12	—	12	*	—	*	*	*	*
73.4	616	—	616	*	614	*	—	88	144	225	158
73.5	1,022	—	1,022	*	1,019	*	—	145	211	386	280
73.51	*6	—	*6	—	*6	—	—	*	*	*	*
73.59	1,016	—	1,016	*	1,013	*	—	145	210	383	279
73.6	404	—	404	*	402	*	—	68	78	164	94
74	1,266	—	1,266	*	1,262	*	—	191	233	531	310
74.0	16	—	16	—	16	*	—	*	*	*6	*
74.1	1,237	—	1,237	*	1,233	*	—	187	227	519	304
74.9	*8	—	*8	—	*8	—	—	*	*	*	*
74.99	*7	—	*7	—	*7	—	—	*	*	*	*
75	2,300	—	2,300	*5	2,293	*	—	396	559	749	596
75.1	9	—	9	—	9	—	—	*	*	*	*
75.3	970	—	970	*	967	*	—	188	257	287	238
75.32	167	—	167	*	167	—	—	26	47	48	46
75.34	661	—	661	*	659	*	—	143	174	195	150
75.35	92	—	92	*	91	—	—	13	26	27	26
75.37	47	—	47	*	47	—	—	*6	10	16	15
75.4	31	—	31	—	31	*	—	*	10	*9	*7
75.5	*7	—	*7	—	*7	—	—	*	*	*	*
75.51	*6	—	*6	—	*6	—	—	*	*	*	*
75.6	1,251	—	1,251	*	1,247	*	—	192	282	432	345
75.61	40	—	40	*	40	—	—	*	*6	21	10
75.62	101	—	101	*	101	—	—	15	23	38	25
75.69	1,110	—	1,110	*	1,106	*	—	173	253	373	310
75.7	27	—	27	—	27	—	—	*8	*	13	*
76–84	4,139	1,972	2,167	168	893	1,444	1,634	782	971	1,472	914
76	78	47	31	*5	52	16	*5	12	11	37	18
76.6	25	10	15	*	20	*	*	*	*	12	*7
76.62	*6	*	*	*	*	*	—	*	*	*	*
76.65	*7	*	*5	*	*6	*	—	*	*	*	*
76.7	30	24	*6	*	21	*6	*	*	*7	13	*6
76.75	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
76.76	14	12	*	*	10	*	*	*	*	*7	*
76.9	13	*9	*	*	*7	*	*	*	*	*7	*
77	254	136	117	21	56	90	86	54	49	90	60
77.1	*6	*	*	*	*	*	*	*	*	*	*
77.3	14	*6	*8	*	*	*	*	*	*	*	*
77.4	47	24	22	*	*	13	27	11	*9	17	10
77.49	36	20	16	*	*	9	23	*8	*6	13	*9
77.5	*6	*	*	-	*	*	*	*	*	*	*
77.6	57	35	21	*	16	21	18	12	11	16	17
77.65	*9	*6	*	*	*	*	*	*	*	*	*
77.67	*9	*7	*	*	*	*	*	*	*	*	*
77.68	10	*6	*	*	*	*	*	*	*	*	*
77.69	16	*8	*8	*	*5	*6	*	*	*	*	*5
77.7	78	36	42	*6	22	34	15	15	16	33	14
77.79	72	34	38	*6	19	33	14	14	15	31	12
77.8	37	24	13	*	*6	15	16	*8	*6	15	*9
77.81	14	10	*	-	*	*5	*8	*	*	*7	*
77.88	*8	*5	*	*	*	*	*	*	*	*	*
77.89	*8	*	*	*	*	*	*	*	*	*	*
77.9	*6	*	*	-	*	*	*	*	*	*	*
78	226	109	117	24	56	71	74	41	52	89	44
78.0	28	13	15	*	9	9	*8	*6	*7	11	*
78.05	*8	*	*	*	*	*	*	*	*	*	*
78.07	*7	*	*	*	*	*	*	*	*	*	*
78.1	33	22	11	*	11	11	*7	*6	*7	14	*6
78.13	*9	*	*5	*	*	*	*	*	*	*	*
78.17	16	12	*	*	*6	*6	*	*	*	*6	*
78.4	10	*5	*5	*	*	*	*	*	*	*5	*
78.41	*5	*	*	*	*	*	*	*	*	*	*
78.5	53	23	30	*7	9	12	25	10	13	16	14
78.55	32	12	20	*	*	*	21	*6	10	*8	*8
78.57	*7	*	*	*	*	*	*	*	*	*	*
78.59	*6	*	*	*	*	*	*	*	*	*	*
78.6	94	41	52	*5	21	35	32	15	21	40	18
78.65	28	11	18	*	*	*6	17	*	*6	14	*
78.67	15	*9	*6	*	*	*6	*	*	*	*	*
78.69	36	14	23	*	9	19	*7	*5	*9	15	*8
79	688	330	358	50	203	159	277	136	159	252	142
79.0	65	30	35	10	18	15	23	14	17	21	13
79.01	*9	*	*7	*	*	*	*5	*	*	*	*
79.02	21	*7	13	*	*	*6	*9	*	*	*9	*
79.05	*7	*	*	*	*	*	*	*	*	*	*
79.06	25	13	12	*	*8	*6	*7	*5	*8	*5	*6
79.1	95	45	50	13	25	13	44	20	22	31	22
79.11	12	*6	*5	*8	*	*	*	*	*	*	*5
79.12	*8	*	*	*	*	*	*	*	*	*	*
79.15	54	22	32	*	*8	*6	37	12	15	16	10
79.16	14	*8	*6	*	*8	*	*	*	*	*	*
79.2	*6	*	*	*	*	*	*	*	*	*	*
79.3	424	190	235	17	121	102	184	85	98	157	85
79.31	37	17	19	*	*6	12	14	*7	10	13	*7
79.32	45	23	22	*	16	11	15	*9	11	16	10
79.33	*7	*6	*	*	*5	*	*	*	*	*	*
79.34	*8	*7	*	*	*6	*	*	*	*	*	*
79.35	160	47	113	*5	14	21	119	31	35	67	27
79.36	143	73	70	*	61	45	33	32	33	45	32
79.37	12	*7	*5	*	*8	*	*	*	*	*	*
79.39	12	*8	*	*	*6	*	*	*	*	*	*
79.6	54	38	16	*6	27	11	*9	*6	11	27	9
79.62	*8	*	*	*	*	*	*	*	*	*	*
79.66	25	17	*7	*	13	*	*	*	*	12	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
79.7	30	14	15	*	*6	11	13	*7	*6	*8	*8
79.71	*6	*	*	*	*	*	*	*	*	*	*
79.75	17	*8	9	*	*	*	10	*	*	*	*
79.8	9	*6	*	*	*	*	*	*	*	*	*
80	466	243	223	*8	139	214	105	81	100	181	104
80.0	15	*7	*8	-	*	*5	9	*	*	*	*
80.06	*8	*	*	-	*	*	*	*	*	*	*
80.1	34	18	16	*	10	10	10	*5	*7	14	*7
80.15	*6	*	*	*	*	*	*	*	*	*	*
80.16	14	*7	*7	*	*	*5	*	*	*	*6	*
80.2	27	11	15	*	*	11	10	*	*5	15	*
80.21	*8	*	*	-	*	*6	*	*	*	*	*
80.26	18	*8	10	*	*	*5	*8	*	*	11	*
80.3	*7	*5	*	*	*	*	*	*	*	*	*
80.4	15	10	*6	*	*	*5	*	*	*	*6	*
80.46	*8	*6	*	*	*	*	*	*	*	*	*
80.5	294	148	146	*	101	148	44	53	62	109	69
80.51	292	147	145	*	101	147	43	53	62	108	69
80.6	14	*9	*5	*	*	*8	*	*	*	*	*
80.7	20	11	9	*	*	*9	*7	*	*	*7	*
80.76	15	*9	*6	*	*	*6	*6	*	*	*6	*
80.8	36	22	14	*	*9	13	13	*6	10	16	*
80.81	*6	*	*	-	*	*	*	*	*	*	*
80.86	17	10	*7	*	*	*6	*6	*	*	*7	*
81	1,841	763	1,079	28	243	673	897	346	484	603	407
81.0	327	151	176	11	95	157	64	54	76	120	76
81.02	121	61	61	-	37	71	14	17	23	50	31
81.03	16	*9	*8	*	*	*8	*	*	*	*6	*
81.05	24	9	15	*8	9	*	*	*	*5	*9	*
81.06	22	10	13	*	11	10	*	*	*6	*8	*
81.07	11	*	*7	-	*	*	*	*	*	*5	*
81.08	123	56	68	*	30	58	34	19	29	44	31
81.1	13	*6	*7	*	*	*6	*	*	*	*	*
81.3	19	*8	11	*	*	9	*	*	*	*8	*
81.38	11	*	*7	-	*	*5	*	*	*	*	*
81.4	27	15	12	*	13	10	*	*6	*7	*7	*7
81.45	*5	*	*	*	*	*	*	*	*	*	*
81.47	13	*8	*5	*	*	*6	*	*	*	*	*
81.5	1,075	412	663	*	60	366	644	223	307	325	220
81.51	235	109	126	-	15	80	139	54	60	61	60
81.52	234	87	146	*	35	68	127	42	73	72	47
81.53	35	14	20	-	*	12	20	10	10	9	*5
81.54	534	184	349	*	*8	194	331	109	153	168	103
81.55	37	17	20	-	*	10	26	*7	10	14	*5
81.6	201	84	117	*6	46	80	70	22	37	87	55
81.62	127	60	66	*	39	62	25	11	20	55	41
81.63	19	*8	12	*	*	10	*6	*	*	11	*
81.64	*5	*	*	*	*	*	-	*	*	*	*
81.65	20	*	16	*	*	*	18	*	*7	*7	*
81.66	25	*8	17	-	*	*	21	*6	*	11	*
81.8	51	21	31	*	*	17	29	*7	13	18	13
81.80	15	*7	*8	-	*	*	10	*	*	*5	*
81.81	20	*	15	-	*	*5	14	*	*	*8	*
81.83	9	*5	*	-	*	*6	*	*	*	*	*
81.9	124	62	62	*	14	27	77	28	38	32	25
81.91	63	36	27	*	10	16	32	15	19	15	13
81.92	51	18	33	*	*	10	38	11	17	16	*7
81.97	*6	*5	*	*	*	*	*5	*	*	*	*
82	31	21	10	*	20	*8	*	*6	*	13	*8
82.0	*8	*5	*	*	*6	*	*	*	*	*	*
82.4	12	*8	*	*	9	*	*	*	*	*5	*
82.45	*5	*	*	*	*	*	*	*	*	*	*

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
83	261	149	112	26	63	88	85	53	48	92	67
83.0	33	23	10	*	10	11	*8	*	*6	11	12
83.02	*6	*5	*	*	*	*	*	*	*	*	*
83.09	20	13	*7	*	*6	*8	*	*	*	*7	*7
83.1	22	12	10	*	*6	*8	*	*5	*	*6	*6
83.14	11	*8	*	*	*	*5	*	*	*	*	*
83.2	21	13	*8	*6	*	*6	*6	*	*	11	*
83.21	21	13	*8	*6	*	*6	*6	*	*	11	*
83.3	26	12	14	*	*6	9	11	*7	*	11	*7
83.39	24	11	13	*	*5	*9	10	*6	*	10	*7
83.4	32	17	15	*	*8	10	11	*6	*9	11	*6
83.45	21	12	10	*	*	*6	*8	*	*6	*7	*
83.49	*5	*	*	*	*	*	*	*	*	*	*
83.5	*6	*	*	-	*	*	*	*	*	*	*
83.6	55	32	23	*	12	20	21	11	13	19	12
83.63	30	18	12	-	*	13	16	*	*6	12	*7
83.64	11	*6	*	*	*	*	*	*	*	*	*
83.65	11	*6	*	*	*5	*	*	*	*	*	*
83.7	*8	*	*	*	*	*	*	*	*	*	*
83.75	*5	*	*	*	*	*	*	*	*	*	*
83.8	42	25	17	*5	11	15	10	11	*7	14	10
83.82	12	*8	*	*	*	*6	*	*	*	*	*
83.85	14	*6	*8	*	*	*	*	*	*	*	*
83.88	11	*7	*	*	*	*	*	*	*	*	*
83.9	16	*7	9	*	*	*	10	*	*	*5	*
83.95	*5	*	*	*	*	*	*	*	*	*	*
84	294	174	119	*	62	126	102	54	63	114	63
84.0	10	*8	*	*	*	*	*	*	*	*	*
84.01	*7	*6	*	*	*	*	*	*	*	*	*
84.1	129	86	43	*	10	52	66	32	23	55	18
84.11	47	33	14	*	*	21	22	13	*8	19	*7
84.12	11	9	*	-	*	*6	*	*	*	*	*
84.15	36	25	11	-	*	17	16	*8	*8	15	*6
84.17	30	16	14	-	*	*6	23	*7	*	17	*
84.3	14	*9	*5	*	*	*	*8	*	*	*5	*
84.5	133	65	67	*	43	62	24	15	32	48	38
84.51	92	44	48	*	30	45	15	11	21	31	29
84.52	34	18	16	*	11	14	*8	*	10	14	*7
85–86	1,538	731	807	168	476	478	417	288	295	691	264
85	206	*	203	*	60	97	47	36	39	90	41
85.0	11	*	11	*	*	*	*	*	*	*	*
85.1	15	*	15	-	*	*	*7	*	*	*7	*
85.11	9	*	*9	-	*	*	*5	*	*	*	*
85.2	26	*	25	*	*7	12	*6	*5	*5	13	*
85.21	20	*	20	*	*6	10	*	*	*	11	*
85.3	12	*	12	-	*7	*	*	*	*	*6	*
85.32	*8	*	*8	-	*5	*	*	*	*	*	*
85.4	78	*	77	*	14	36	29	11	17	34	16
85.41	25	*	25	-	*	11	11	*	*5	12	*
85.42	*7	-	*7	-	*	*	*	*	*	*	*
85.43	44	*	42	-	*7	20	16	*7	11	18	*8
85.5	9	*	9	-	*	*6	*	*	*	*5	*
85.53	*5	*	*5	-	*	*	*	*	*	*	*
85.6	*6	*	*6	-	*	*	*	*	*	*	*
85.7	11	-	11	-	*	*6	*	*	*	*	*
85.8	10	-	10	-	*	*5	*	*	*	*	*
85.9	28	*	27	-	10	16	*	*	*	11	*9
85.94	*5	-	*5	-	*	*	*	*	*	*	*
85.95	17	*	17	-	*7	9	*	*	*	*7	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
86	1,332	728	604	167	415	381	369	252	255	602	223
86.0	396	212	184	47	114	136	99	83	78	167	69
86.01	15	*8	*7	*	*	*6	*	*	*	*5	*
86.04	208	118	90	24	76	66	41	41	45	90	31
86.05	56	27	29	10	12	20	13	13	*9	23	12
86.06	10	*7	*	*	*	*	*5	*	*	*	*
86.07	86	41	46	*8	16	32	31	18	15	38	14
86.09	17	10	*7	*	*	*8	*	*	*	*6	*
86.1	24	14	10	*	*6	*8	*8	*6	*	9	*
86.11	24	14	10	*	*6	*8	*8	*6	*	9	*
86.2	418	236	182	28	107	127	156	85	87	178	68
86.22	277	161	116	17	79	88	93	53	50	132	42
86.23	13	*8	*	*	*	*	*	*	*	*5	*
86.27	43	21	22	—	*	*8	32	16	13	11	*
86.28	81	44	37	*9	21	25	26	13	18	29	20
86.3	76	23	53	9	33	17	16	12	20	34	10
86.4	13	*6	*7	*	*	*	*	*	*	*6	*
86.5	158	92	66	10	69	29	51	27	41	48	41
86.59	158	92	66	10	69	29	51	27	41	48	41
86.6	107	76	31	32	35	24	17	12	10	76	9
86.66	14	10	*	*	*6	*	*	*	*	12	*
86.67	23	17	*6	*7	10	*	*	*	*	18	*
86.69	57	40	17	19	16	12	10	*9	*6	36	*6
86.7	58	36	22	17	17	12	12	*7	*5	36	10
86.74	44	27	18	15	13	9	*7	*5	*	29	*8
86.8	74	29	45	22	27	22	*	13	*6	45	10
86.83	27	*	24	*	15	12	*	*9	*	12	*
86.84	29	16	13	20	*8	*	*	*	*	28	*
86.89	12	*8	*	*	*	*6	*	*	*	*	*5
86.9	*7	*	*	*	*	*	*	*	*	*	*
87–99	13,243	6,366	6,877	950	2,497	3,727	6,069	3,795	2,761	4,120	2,567
87	902	418	483	58	202	257	384	297	177	239	188
87.0	315	148	167	19	57	85	155	151	48	62	54
87.03	308	144	164	17	55	83	153	150	46	59	53
87.2	15	*7	*9	*	*	*6	*	*	*	*7	*
87.21	*7	*	*	—	*	*	*	*	*	*	*
87.3	13	*6	*7	*	*	*5	*	*	*	*	11
87.39	11	*6	*6	*	*	*	*	*	*	—	11
87.4	240	119	121	12	42	76	110	85	61	38	57
87.41	148	73	75	*	27	48	70	65	26	25	32
87.44	79	39	40	9	13	24	33	14	31	12	22
87.49	12	*7	*6	—	*	*	*7	*6	*	*	*
87.5	130	47	83	*	45	39	46	17	32	53	27
87.53	119	42	77	*	44	35	40	15	30	50	24
87.54	*5	*	*	*	*	*	*	*	*	*	*
87.6	29	12	18	*7	*	*6	12	*9	*	9	9
87.61	*7	*	*	*	*	*	*	*	*	*	*
87.62	10	*	*6	*	*	*	*	*	*	*	*
87.63	*5	*	*	*	*	*	*	*	*	*	*
87.7	155	78	78	17	47	40	52	28	30	70	27
87.73	10	*5	*5	*	*	*	*	*	*	*5	*
87.74	104	48	57	*	35	30	36	21	24	47	13
87.76	15	9	*6	11	*	*	*	*	*	9	*
87.77	9	*6	*	*	*	*	*5	*	*	*	*
87.79	*7	*	*	*	*	*	*	*	*	*	*
88	3,616	1,879	1,736	111	517	1,260	1,728	1,029	798	1,123	666
88.0	255	108	148	10	78	70	97	122	51	33	49
88.01	253	106	146	10	77	69	96	121	50	33	49
88.1	17	10	*7	*	*6	*7	*	*	*6	9	*
88.11	*8	*6	*	—	*	*	*	*	—	*8	—
88.19	*8	*	*	*	*	*	*	*	*5	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
88.2	18	9	*8	*	*6	*	*6	*	*8	*	*
88.27	*5	*	*	*	*	*	*	*	*	*	*
88.3	104	53	51	10	30	29	35	50	28	16	10
88.38	100	52	47	*8	30	28	33	49	27	16	*8
88.4	501	265	236	*6	54	170	270	101	130	180	89
88.41	139	65	74	*	24	49	64	38	32	47	22
88.42	132	78	54	*	*8	44	78	17	41	53	21
88.43	20	11	10	*	*	*7	*7	*	*	*6	*7
88.44	21	9	12	*	*	*8	12	*7	*	*7	*
88.45	34	16	17	—	*	10	21	*6	*7	15	*7
88.47	30	15	15	—	*	9	19	*6	*7	11	*6
88.48	101	58	42	*	*8	35	58	20	31	31	19
88.49	24	12	12	*	*	*8	11	*	*5	10	*6
88.5	1,484	860	624	*7	126	624	728	259	358	574	294
88.51	20	10	10	*	*	*8	10	*5	*	*8	*
88.53	540	299	240	*	50	227	261	96	124	223	97
88.54	17	10	*7	*	*	*7	*8	*	*5	*	*
88.55	*9	*6	*	*	*	*	*	*	*	*	*
88.56	861	512	348	*	69	364	427	141	215	323	182
88.57	34	20	14	*	*	14	16	*7	*7	13	*7
88.6	30	16	14	*	*6	10	14	*7	9	10	*
88.67	15	*8	*6	*	*	*	*7	*	*	*6	*
88.7	884	411	473	44	150	249	441	318	150	245	170
88.71	43	18	25	*	*	*8	32	18	*	9	11
88.72	564	276	287	22	71	165	306	180	90	198	97
88.73	14	*5	*9	*	*	*5	*6	*	*	*	*6
88.74	14	*6	*8	*	*	*	*	*	*	*	*
88.75	29	14	15	*	*6	*6	14	17	*	*	*5
88.76	100	48	53	11	27	32	29	43	22	*8	27
88.77	79	34	46	*	13	24	42	37	21	13	9
88.78	17	—	17	—	17	*	*	*7	*	*	*5
88.79	23	10	13	*	*7	*	*9	10	*	*	*7
88.89	321	146	175	29	60	98	133	167	58	52	44
88.91	190	88	102	18	34	61	77	97	31	35	26
88.93	55	26	29	*	13	13	26	29	11	*7	*9
88.94	20	10	10	*	*	*	10	12	*	*	*
88.97	50	18	32	*	*8	18	19	26	10	9	*5
89	686	338	348	51	115	205	314	246	103	155	182
89.0	38	20	18	*	*7	13	15	*	—	*7	30
89.06	13	*8	*5	—	*	*5	*6	—	—	—	13
89.07	*8	*	*	—	*	*	*	—	—	—	*8
89.08	*6	*	*	*	*	*	*	*	—	*5	*
89.09	10	*5	*	—	*	*	*5	*	—	*	*8
89.1	86	43	44	24	21	18	23	34	9	15	28
89.14	54	26	27	12	*9	13	20	19	*7	*7	20
89.15	*7	*	*	*	*	*	*	*	*	*	*
89.19	24	11	14	11	9	*	*	14	*	*6	*
89.2	15	*5	10	*	*	*	9	*	*	*6	*
89.22	*6	*	*	—	*	*	*5	*	*	*	—
89.3	26	14	12	10	*	*6	*6	*8	*6	*5	*7
89.37	*7	*	*	*	*	*	*	*	*	*	*
89.38	*5	*	*	*	*	*	*	*	*	*	*
89.39	10	*6	*	*5	*	*	*	*	*	*	*
89.4	178	89	89	*	24	64	90	55	28	49	46
89.41	32	15	18	*	*6	16	10	*8	*	*7	14
89.44	102	49	53	—	13	36	53	33	13	28	28
89.45	18	*7	11	—	*	*	14	*6	*	*9	*
89.49	23	18	*6	—	*	9	12	*7	*8	*6	*
89.5	227	113	114	*9	42	66	110	122	33	35	36
89.50	*6	*	*	*	*	*	*	*	*	*	*
89.52	106	54	51	*7	21	32	45	43	24	17	21

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
89.54	101	50	51	*	17	30	53	69	*	15	13
89.59	14	*6	*8	—	*	*	*8	*	*5	*	*
89.6	116	55	60	*6	15	34	60	22	22	37	35
89.61	10	*6	*	*	*	*	*	*	*	*	*
89.62	*8	*	*	*	*	*	*	*	*	*	*
89.64	23	12	11	*	*	*8	13	*	*9	*7	*
89.65	56	23	33	*	*9	13	32	*9	*	21	22
89.68	15	*8	*7	*	*	*6	*6	*	*	*	*
90	34	13	21	*	11	*7	12	21	*7	*	*
90.5	27	10	17	*	10	*6	*9	20	*	*	*
90.52	*9	*	*5	*	*	*	*	*	*	*	*
90.59	17	*6	11	*	*7	*	*7	16	—	*	*
91	21	*7	14	*	*8	*	*8	13	*	*	*
91.3	17	*6	11	*	*6	*	*7	12	*	*	*
91.32	*8	*	*6	*	*	*	*	*	*	*	*
91.39	*6	*	*	—	*	*	*	*6	—	—	—
92	224	106	118	*8	29	78	110	78	23	63	60
92.0	103	46	57	*	14	36	51	32	*8	29	34
92.02	12	*5	*6	—	*	*	*	*6	*	*	*
92.04	12	*6	*6	*	*	*	*5	*	*	*	*
92.05	76	33	42	—	*7	29	40	20	*6	24	26
92.1	50	23	28	*	*6	16	26	24	*	*9	14
92.14	21	10	11	*	*	*5	11	10	*	*	*5
92.15	21	*8	13	—	*	*8	11	11	*	*	*6
92.2	66	36	31	*	*8	25	31	20	11	24	12
92.24	9	*6	*	*	*	*	*5	*	*	*	*
92.27	*7	*	*	*	*	*	*	*	*	*	*
92.29	44	24	20	*	*6	18	19	12	*8	17	*7
93	1,508	671	836	166	195	314	832	448	259	529	272
93.0	34	16	18	*	*6	9	17	19	*	*	*8
93.01	24	11	13	*	*	*7	14	17	*	*	*
93.08	*6	*	*	*	*	*	*	*	*	*	*
93.1	24	*9	15	*	*	*6	14	17	*	*	*
93.12	10	*	*6	*	*	*	*7	*7	*	*	—
93.2	34	13	21	*	*	12	18	23	*	*6	*
93.22	24	9	15	*	*	*8	14	18	*	*	*
93.3	349	145	204	*8	32	63	245	83	65	147	55
93.38	26	12	14	*	*	*6	17	*	*	*7	16
93.39	320	131	189	*8	28	56	228	79	63	138	39
93.4	11	*6	*5	*	*	*	*	*	*	*5	*
93.5	117	64	53	25	29	25	37	30	26	45	15
93.53	17	*7	10	*	*	*	*8	*5	*	*6	*
93.54	30	17	13	*	11	*7	10	*5	10	*8	*7
93.57	32	18	13	14	*8	*6	*	*8	*5	17	*
93.59	24	12	12	*	*	*7	13	*8	*	*9	*
93.7	36	17	19	*	*5	*7	22	*7	*7	13	*9
93.75	35	17	18	*	*5	*6	22	*7	*7	13	*9
93.8	430	165	265	9	49	84	288	105	54	195	75
93.81	56	22	34	*	16	13	25	*6	*5	34	11
93.83	208	84	123	*6	22	42	137	63	38	82	24
93.89	161	57	104	—	9	27	125	36	11	75	40
93.9	470	234	235	114	64	106	186	160	97	109	104
93.90	137	69	68	11	16	41	69	49	30	34	24
93.94	164	80	83	60	20	30	54	69	34	27	34
93.96	151	75	76	38	23	31	60	37	31	40	43
93.99	13	*7	*6	*	*	*	*	*	*	*6	*
94	672	395	277	41	379	203	49	309	171	150	41
94.0	12	*7	*5	*	*7	*	*	*	—	*8	—
94.08	*7	*	*	*	*	*	*	*	—	*	—
94.1	*9	*5	*	*	*5	*	*	*8	—	*	—

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
94.19	*6	*	*	*	*	*	*	*5	—	*	—
94.2	169	81	87	15	88	45	21	123	14	30	*
94.22	*7	*	*	*	*	*	*	*6	*	*	—
94.23	42	23	18	*7	26	*8	*	41	—	*	—
94.25	92	45	47	*7	49	24	11	67	9	15	*
94.27	28	9	19	—	*9	11	*8	*9	*	13	*
94.3	59	28	30	*9	33	12	*	19	26	13	*
94.38	14	*8	*6	*	10	*	*	14	—	*	—
94.39	43	20	23	*7	22	9	*	*	26	12	*
94.4	87	45	42	13	50	18	*6	25	30	32	—
94.42	*9	*	*	*	*	*	*	*	*	*	—
94.44	70	34	36	9	39	16	*6	14	26	29	—
94.6	337	228	109	*	197	124	16	130	101	66	39
94.61	*8	*5	*	*	*6	*	*	*	*	*	*
94.62	169	122	47	—	75	83	11	54	63	28	24
94.63	18	12	*6	*	10	*7	*	*	13	*	*
94.64	*7	*	*	*	*6	*	*	*	*	*	—
94.65	59	32	27	—	44	13	*	29	*9	14	*7
94.66	*8	*	*	—	*7	*	*	*	*	*	*
94.67	*8	*	*	*	*6	*	—	*	*	*	*
94.68	56	41	15	—	39	15	*	40	*	10	*
94.69	*5	*	*	*	*	*	—	—	*	*	*
95	12	*5	*7	10	*	*	*	*	*	*8	*
95.4	11	*	*6	9	*	*	*	*	*	*7	*
95.47	*8	*	*	*8	*	—	—	*	*	*6	*
96	1,613	798	815	159	328	390	737	376	337	537	364
96.0	618	320	299	43	100	169	306	135	125	213	145
96.04	506	268	238	40	81	136	249	108	102	181	115
96.05	*8	*	*	*	*	*	*	*	*	*	*
96.07	94	44	50	*	18	27	47	23	21	27	24
96.2	*6	*	*	*	*	*	*	*	*	*	*
96.3	18	*8	11	*	*7	*	*	*	*	*5	*6
96.33	*8	*	*	*	*	*	*	*	*	*	*
96.38	*6	*	*	*	*	*	*	*	*	*	*
96.4	93	*5	88	*	86	*	*5	9	26	30	27
96.49	89	*	86	*	85	*	*	*9	26	29	26
96.5	40	23	17	*6	12	*8	14	*6	*7	17	9
96.56	14	9	*	*	*	*	*8	*	*	*	*5
96.59	20	11	*8	*	*8	*6	*	*	*	10	*
96.6	146	67	80	30	12	26	79	72	31	25	18
96.7	691	372	319	75	109	181	325	146	139	244	161
96.70	10	*7	*	*	*	*	*	*	*	*6	*
96.71	439	236	203	47	77	113	202	91	94	155	99
96.72	242	130	112	27	29	67	119	53	45	83	61
97	107	51	57	10	22	33	43	25	19	37	26
97.0	31	13	18	*	*	*6	20	*8	*	10	9
97.02	22	*9	13	*	*	*	15	*6	*	*6	*7
97.05	*6	*	*	*	*	*	*	*	*	*	*
97.2	18	12	*6	*	*	9	*	*	*	*7	*5
97.23	14	10	*	*	*	*8	*	*	*	*	*
97.3	*9	*5	*	*	*	*	*	*	*	*	*
97.39	*6	*	*	*	*	*	*	*	*	*	*
97.4	11	*5	*6	*	*	*	*	*	*	*	*
97.49	*8	*	*	*	*	*	*	*	*	*	*
97.5	*8	*	*6	*	*	*	*	*	*	*	*
97.6	13	*5	*8	*	*	*	*	*	*	*	*
97.62	10	*	*7	*	*	*	*	*	*	*	*
97.8	14	*6	*7	*	*	*	*	*	*	*	*
98	31	16	15	*6	*8	*7	10	*	*7	13	*7
98.0	11	*7	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2005—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
98.02	*8	*5	*	*	*	*	*	*	*	*	*
98.1	*5	*	*	*	*	*	*	*	*	*	*
98.5	11	*	*8	—	*	*	*	*	*	*8	*
98.51	11	*	*8	—	*	*	*	*	*	*8	*
99	3,818	1,668	2,150	325	683	969	1,842	946	857	1,259	756
99.0	1,902	788	1,115	67	220	472	1,144	327	403	754	418
99.00	33	13	20	—	*	9	20	*7	*8	11	*7
99.02	56	21	35	—	*	26	27	*7	12	15	21
99.03	13	*	*8	—	*	*	*8	*	*	*6	*
99.04	1,537	620	917	43	185	362	947	268	329	627	313
99.05	83	37	46	13	11	23	36	15	14	33	21
99.07	166	84	82	—	15	46	100	26	31	58	51
99.1	415	165	250	45	133	102	135	125	90	99	102
99.10	38	21	16	—	*	14	19	*8	*8	13	*8
99.11	75	*	75	—	74	*	*	16	15	24	19
99.14	15	*8	*6	*7	*5	*	*	*	*	*5	*
99.15	134	62	72	30	19	31	54	35	32	38	30
99.17	35	18	17	—	*8	16	9	14	12	*	*
99.18	47	18	29	*6	*8	13	20	23	*	*	20
99.19	71	37	34	—	14	25	31	24	20	11	16
99.2	1,131	529	602	131	269	315	417	407	283	286	155
99.20	108	75	34	—	*7	49	52	22	24	32	31
99.21	342	145	197	44	96	78	124	142	95	63	42
99.22	*6	*	*	—	*	*	*	*	*	*	—
99.23	108	43	65	—	20	23	60	31	35	30	12
99.25	209	116	93	42	39	73	55	51	28	91	38
99.29	351	144	207	38	103	86	123	157	96	68	30
99.3	9	*7	*	—	*	*	*	*	*	*	*
99.4	20	*	20	—	20	*	*	*	*	*7	*7
99.47	11	—	11	—	11	*	—	*	*	*5	*
99.48	*8	—	*8	—	*8	—	—	*	*	*	*
99.5	77	34	43	*8	13	23	32	29	18	19	11
99.52	30	12	18	—	*5	10	13	14	*7	*7	*
99.55	43	19	23	*7	*6	12	18	15	10	10	*7
99.6	139	83	56	—	9	37	91	32	37	42	29
99.60	58	35	23	—	*	14	40	12	16	19	11
99.61	22	14	*8	—	*	*7	13	*6	*	*8	*
99.62	50	30	20	—	*	14	32	11	15	13	12
99.69	*8	*	*	—	*	*	*5	*	*	*	*
99.7	38	13	24	—	13	13	11	11	*	16	*6
99.71	10	*	*7	—	*	*	*	*	*	*	*
99.74	*5	*	*	—	*	*	*	*	*	*	*
99.77	16	*	12	—	*6	*	*	*	*	*8	*
99.8	70	41	29	66	—	*	*	*8	12	24	26
99.83	63	37	26	63	—	—	—	*7	11	21	24
99.9	17	*8	*8	—	*	*	*7	*	*	10	*
99.99	16	*8	*8	—	*	*	*7	*	*	10	*

* Figure does not meet standards of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see "Methods" section.

Appendix

Technical Notes

Abstract Form

Figure I presents the medical abstract form used for the 2005 NHDS. The hospital number, date of birth, and patient ZIP Code are confidential information and are not released to the public, although they can be used with privacy safeguards through the NCHS Research Data Center.

Population Estimates

Population estimates used to compute rates are shown in **Table I**. These estimates are for the civilian population of the United States, including institutionalized persons, as of July 1, 2005. The data are postcensal estimates based on the 2000 census and are from unpublished tabulations provided by the U.S. Census Bureau.

Reliability of Estimates

The statistics presented in this report are based on a sample survey and therefore may differ from the figures that would be obtained if a complete census had been taken. The standard error is primarily a measure of sampling variability that occurs by chance because only a sample rather than an entire universe is surveyed. Standard errors for the estimates from NHDS were calculated with SUDAAN software, which takes into account the complex sample design. A description of the software and the approach it uses has been published (23).

The chances are about 32 in 100 that an estimate from the sample would differ from a complete census by more than the standard error. The chances are 5 in 100 that the difference would be more than twice the standard error and about 1 in 100 that the difference would be more than 2.5 times the standard error. The relative standard error (RSE) expresses the standard error as a percentage of the estimate and can be multiplied by the estimate to obtain the standard error.

Table I. Civilian population by age, geographic region, and sex: United States, 2005

Age and geographic region	Both sexes	Male	Female
All ages		Number in thousands	
Total	295,194	144,959	150,235
Region:			
Northeast	54,583	26,480	28,103
Midwest	65,891	32,377	33,514
South	106,818	52,274	54,544
West	67,903	33,829	34,073
Under 15 years			
Total	60,700	31,057	29,644
Under 1 year	4,107	2,101	2,005
1–4 years	16,197	8,280	7,917
5–14 years	40,397	20,675	19,721
Region:			
Northeast	10,471	5,355	5,116
Midwest	13,259	6,786	6,473
South	22,159	11,329	10,829
West	14,812	7,586	7,226
15–44 years			
Total	124,910	63,006	61,904
15–24 years	41,585	21,233	20,352
25–34 years	39,705	20,047	19,658
35–44 years	43,620	21,726	21,894
Region:			
Northeast	22,618	11,314	11,303
Midwest	27,701	14,002	13,700
South	45,183	22,654	22,528
West	29,407	15,035	14,371
45–64 years			
Total	72,794	35,485	37,310
45–54 years	42,441	20,860	21,581
55–64 years	30,354	14,625	15,729
Region:			
Northeast	14,042	6,782	7,260
Midwest	16,488	8,095	8,393
South	26,160	12,674	13,486
West	16,104	7,932	8,171
65 years and over			
Total	36,790	15,413	21,377
65–74 years	18,640	8,529	10,110
75–84 years	13,054	5,279	7,775
85 years and over	5,096	1,604	3,492
Region:			
Northeast	7,453	3,028	4,425
Midwest	8,444	3,495	4,948
South	13,316	5,616	7,701
West	7,579	3,275	4,305

NOTE: Unpublished 2000-based postcensal estimates from the U.S. Census Bureau.

To derive approximate standard errors for the estimates in this report, first calculate the approximate RSE of the estimate using the parameters shown in **Table II**. Then multiply the approximate RSE by the estimate. The approximate RSE of an estimate X may be estimated from the formula:

$$\text{RSE}(X) = \sqrt{a + b/X}$$

where a and b are defined in **Table II**. When multiplied by 100, $\text{RSE}(X)$ is expressed as a percentage of X .

The approximate RSE for a percentage, $100p$, in which the numerator is a subclass of the

Notice – All information which would permit identification of an individual or an establishment will be held confidential, will be used only by persons engaged in and for the purposes of the survey, and will not be disclosed or released to other persons or used for any other purpose. Public reporting burden of this collection of information is estimated to average 4 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer; 1600 Clifton Road, MS D-74, Atlanta, GA 30333, ATTN: PRA (0920-0212).

FORM **HDS-1**
9-9-2005

U.S. DEPARTMENT OF COMMERCE
Economics and Statistics Administration
U.S. CENSUS BUREAU
ACTING AS COLLECTING AGENT FOR
DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION
NATIONAL CENTER FOR HEALTH STATISTICS

MEDICAL ABSTRACT – NATIONAL HOSPITAL DISCHARGE SURVEY**A. PATIENT IDENTIFICATION**

1. Hospital number	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	Month	Day	Year
2. HDS number	<input type="text"/> <input type="text"/> <input type="text"/>	<input type="text"/> - <input type="text"/>	<input type="text"/> - <input type="text"/>	<input type="text"/> <input type="text"/> <input type="text"/>
3. (Item deleted)				
4. Date of admission	<input type="text"/> <input type="text"/>	- <input type="text"/> <input type="text"/>	- <input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/> <input type="text"/>
5. Date of discharge	<input type="text"/> <input type="text"/>	- <input type="text"/> <input type="text"/>	- <input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/> <input type="text"/>
6. Residence ZIP Code	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>			

B. PATIENT CHARACTERISTICS

7. Date of birth	Month	Day	Year	11. Race – Mark all that apply		
	<input type="text"/> - <input type="text"/>	<input type="text"/> - <input type="text"/>	<input type="text"/> <input type="text"/> <input type="text"/>	1 <input type="checkbox"/> White		
				2 <input type="checkbox"/> Black or African American		
8. Age – Complete only if date of birth not given	Units			3 <input type="checkbox"/> American Indian or Alaska Native		
	<input type="text"/> <input type="text"/> <input type="text"/>	{ 1 <input type="checkbox"/> Years		4 <input type="checkbox"/> Asian		
		2 <input type="checkbox"/> Months		5 <input type="checkbox"/> Native Hawaiian or Other Pacific Islander		
		3 <input type="checkbox"/> Days		6 <input type="checkbox"/> Other – Specify <input checked="" type="checkbox"/>		
9. Sex – Mark (X) one				7 <input type="checkbox"/> Not stated		
10. Ethnicity – Mark (X) one	1 <input type="checkbox"/> Male	2 <input type="checkbox"/> Female	3 <input type="checkbox"/> Not stated	12. Marital status – Mark (X) one		
	1 <input type="checkbox"/> Hispanic	2 <input type="checkbox"/> Not Hispanic or Latino	3 <input type="checkbox"/> Not stated	1 <input type="checkbox"/> Married	3 <input type="checkbox"/> Widowed	5 <input type="checkbox"/> Separated
	2 <input type="checkbox"/> Urgent	4 <input type="checkbox"/> Newborn		2 <input type="checkbox"/> Single	4 <input type="checkbox"/> Divorced	6 <input type="checkbox"/> Not stated

C. ADMINISTRATIVE INFORMATION

13. Type of Admission – Mark (X) one	1 <input type="checkbox"/> Emergency	3 <input type="checkbox"/> Elective	5 <input type="checkbox"/> Items not available/unknown	16. Expected source(s) of payment	Principal	Other additional sources
	2 <input type="checkbox"/> Urgent	4 <input type="checkbox"/> Newborn			Mark one only	Mark all that apply
14. Source of Admission – Mark (X) one	1 <input type="checkbox"/> Physician referral	7 <input type="checkbox"/> Emergency room		1. Worker's compensation	<input type="checkbox"/>	<input type="checkbox"/>
	2 <input type="checkbox"/> Clinical referral	8 <input type="checkbox"/> Court/Law enforcement		2. Medicare	<input type="checkbox"/>	<input type="checkbox"/>
	3 <input type="checkbox"/> HMO referral	9 <input type="checkbox"/> Other – Specify <input checked="" type="checkbox"/>		3. Medicaid	<input type="checkbox"/>	<input type="checkbox"/>
	4 <input type="checkbox"/> Transfer from a hospital			4. Other government payments	<input type="checkbox"/>	<input type="checkbox"/>
	5 <input type="checkbox"/> Transfer from SNF			5. Blue Cross/Blue Shield	<input type="checkbox"/>	<input type="checkbox"/>
	6 <input type="checkbox"/> Transfer from other health facility	10 <input type="checkbox"/> Item not available		6. HMO/PPO	<input type="checkbox"/>	<input type="checkbox"/>
15. Status/Disposition of patient – Mark (X) appropriate box(es)	Status	Disposition		7. Other private or commercial insurance	<input type="checkbox"/>	<input type="checkbox"/>
	1 <input type="checkbox"/> Alive	a. <input type="checkbox"/> Routine discharge/discharged home		8. Self pay	<input type="checkbox"/>	<input type="checkbox"/>
		b. <input type="checkbox"/> Left against medical advice		9. No charge	<input type="checkbox"/>	<input type="checkbox"/>
		c. <input type="checkbox"/> Discharged, transferred to another short-term hospital		10. Other –Specify <input checked="" type="checkbox"/>		
		d. <input type="checkbox"/> Discharged, transferred to long-term care institution				
		e. <input type="checkbox"/> Other disposition/not stated				
	2 <input type="checkbox"/> Died					
	3 <input type="checkbox"/> Status not stated					
				No source of payment indicated	<input type="checkbox"/>	<input type="checkbox"/>

(Over)

Figure I. Medical abstract form for the National Hospital Discharge Survey, 2005

D. MEDICAL INFORMATION

17. Final Diagnoses (including E-code diagnoses) (Enter ICD-9-CM codes as well as narrative if available)

Principal: _____

Other/additional: _____

18. Surgical and Diagnostic Procedures (Enter ICD-9-CM codes as well as narrative if available)

Principal: _____

Other/additional: _____

Date of procedure(s)

Month Day Year

NONE

Completed by _____

Date _____

Table II. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics, by selected characteristics: 2005

Characteristic	Discharges or first-listed diagnosis		Days of care		All-listed diagnosis		All-listed procedures	
	a	b	a	b	a	b	a	b
Total	0.00207	346.164	0.00323	839.909	0.00253	357.843	0.00357	390.432
Sex								
Male	0.00222	334.327	0.00372	1,032.262	0.00241	312.599	0.00313	320.197
Female	0.00206	345.937	0.00328	1,088.173	0.00256	345.791	0.00326	299.400
Age								
Under 15 years	0.02222	211.185	0.03432	421.121	0.02380	259.063	0.03111	198.141
15–44 years	0.00250	283.693	0.00417	918.013	0.00239	335.039	0.00331	314.433
45–64 years	0.00183	366.971	0.00373	1,139.179	0.00256	354.531	0.00308	310.710
65 years and over	0.00266	277.413	0.00379	1,086.151	0.00291	310.255	0.00340	304.071
Region								
Northeast	0.01845	204.707	0.04005	271.786	0.01587	195.341	0.01490	185.349
Midwest	0.01409	215.440	0.01781	529.931	0.01567	252.282	0.02270	125.271
South	0.00305	379.475	0.00438	1,284.636	0.00326	325.059	0.00383	356.016
West	0.00552	384.878	0.01072	1,108.792	0.00929	346.631	0.00724	391.296
Principal expected source of payment								
Medicare	0.00259	296.049	0.00342	1,645.254	0.00261	340.962	0.00372	298.730
Medicaid	0.00629	275.580	0.01163	802.767	0.00570	296.800	0.00619	335.245
Worker's compensation and other government payment	0.00716	351.112	0.01578	1,187.014	0.00961	359.841	0.00739	378.745
HMO and PPO ¹	0.00600	217.785	0.00826	627.592	0.00627	284.264	0.00775	213.152
Blue Cross Blue Shield and other private insurance	0.00333	301.651	0.00569	819.794	0.00342	312.630	0.00480	262.275
Self pay	0.00386	258.913	0.00716	888.210	0.00371	308.890	0.00526	276.605
No charge and other	0.03576	175.844	0.04748	427.066	0.03111	255.961	0.03478	211.229

¹HMO is health maintenance organization, and PPO is preferred provider organization.

NOTE: The relative standard error (RSE) can be determined from the equation RSE(x)=SQRT(a+b/X).

denominator, may be calculated directly using the formula:

$$\text{RSE}(p) = \sqrt{b(1-p)/(pX)}$$

where 100p is the percentage of interest, X is the denominator of the percentage, and b is the parameter b in the formula for approximating the RSE(X). The values for b are given in Table II. When multiplied by 100, RSE(p) is expressed as a percentage of p>. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSEs of the numerator and denominator are both less than 10 percent (24,25).

The approximate RSE of a population-based rate is equivalent to the approximate RSE of the numerator (obtained by the method described above using figures from Table II) if the denominator is an estimate produced by the U.S. Census Bureau for the total population or for one or more of the age-sex groups of the total population.

The approximate RSE of a ratio (X/Y) in which the numerator (X) and the denominator (Y) are both estimated

from the same survey, but the numerator is not a subclass of the denominator, is calculated using the formula:

$$\text{RSE}(X/Y) = \sqrt{[\text{RSE}(X)]^2 + [\text{RSE}(Y)]^2}.$$

When multiplied by 100, the RSE(X/Y) is expressed as a percentage of the ratio X/Y>. This approximation is valid if the RSE of the denominator is less than 5 percent or if the RSEs of the numerator and denominator are both less than 10 percent (24,25).

Definitions of Selected Terms

Admission source—The source of admission variable was added to NHDS beginning in 2001 and was modeled after the variable in the National Uniform Bill (UB-92), the standard billing form used by Medicare for reimbursement of hospital inpatient care. The categories for admission source shown in this report include the following:

Physician referral—The patient was admitted to the hospital upon the

recommendation of his or her personal physician, the hospital's clinic physician, or a health maintenance organization physician.

Emergency department—The patient was admitted to the hospital upon the recommendation of the hospital's emergency department physician. The terms emergency department and emergency room are used interchangeably in this report.

Transfer from a hospital—The patient was admitted to the hospital as a transfer from an acute care hospital where he or she was an inpatient.

Transfer from other health facility—The patient was admitted to the hospital as a transfer from a skilled nursing facility or other health facility where he or she was an inpatient.

Other sources—The patient was admitted to the hospital upon the direction of a court of law, on the request of a law enforcement

agency representative, or for other reasons not categorized above.

Item not available—Information about the source of admission to the hospital was not available for the patient.

Admission type—Like admission source, the admission type variable was added to NHDS in 2001 and is modeled after the variable in the UB-92. Four admission types are shown in this report:

Emergency—The patient required immediate medical intervention as a result of severe life threatening or potentially disabling conditions.

Patients in this group are generally admitted through the emergency room.

Urgent—The patient required immediate care and treatment of a physical or mental disorder. Patients in this group are usually admitted to the first available and suitable accommodation.

Elective—The patient's condition permitted adequate time to schedule the availability of suitable accommodation.

Item not available—Information about the admission type for the patient was not available.

Age—The patient's age at the birthday prior to admission to the hospital.

Average length of stay—The mean length of stay for discharges, which is calculated by dividing the number of days of care by the number of discharges.

Barell injury diagnosis matrix—This matrix is a two-dimensional array that includes all the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM) codes describing injury (19). The code numbers from the matrix are shown in [Table III](#). This matrix was developed to provide a standard format for describing the site and type of injuries for use in epidemiological and clinical analyses of injury data.

Bed size of hospital—The number of beds, cribs, and pediatric bassinets

regularly maintained (set up and staffed for use) for patients, not including bassinets for newborn infants. In this report, the classification of hospitals by bed size is based on the number of beds listed in the Verispan files (16).

Days of care—The total number of days a patient spent in the hospital. A stay of less than 1 day (patient admitted as an inpatient and discharged on the same day) is counted as one day. For patients admitted and discharged on different days, the number of days of care is computed by counting all the days from (and including) the day of admission to (but not including) the day of discharge.

Death rate—The death rate is calculated by dividing the number of deaths for a category by the total number of discharges in that category and multiplying by 100. Discharges whose status is unknown (2.4 percent) are assumed to be discharged alive.

Diagnosis—A disease or injury (or other factor that influences a patient's health status or causes contact with health services) that is listed on the medical record of the patient at discharge. The principle diagnosis is the diagnosis established after the study to be chiefly responsible for the admission of the patient to the hospital. The *first-listed diagnosis* is the diagnosis specified as the principal diagnosis on the face sheet or discharge summary of the medical record, or, if the principal diagnosis is not specified, it is the first diagnosis listed on the face sheet or discharge summary of the medical record. The first-listed diagnosis is usually the main cause of the hospitalization. The number of first-listed diagnoses is the same as the number of discharges. *All-listed diagnoses* include first-listed diagnoses and secondary diagnoses and are included for users who want to know the total number of times conditions have been reported for hospitalized patients. A minimum of one and a maximum of seven diagnoses are coded in NHDS per discharge. The human immunodeficiency virus (HIV) data ([Table 23](#)) are for discharges with an HIV diagnosis listed in any position among the coded diagnoses.

Discharge—A completed inpatient hospitalization. A hospitalization may be completed by death or when the hospital releases the patient to the customary place of residence, a nursing home, another hospital, or another location.

Disposition—The placement or status of the patient when discharged from the hospital. Six disposition categories are shown in this report:

Routine or discharged home

Patients who were discharged to return to their homes, including those referred to home care services, discharged to a relative or foster home, or transferred to a corrections institute.

Transferred to other short-term hospital

Patients discharged and transferred to another short-term hospital for additional care.

Transferred to long-term care institution

Patients discharged and transferred or returned to a nursing home, skilled nursing facility, extended care facility, intermediate care facility, or custodial care facility.

Other discharged alive—Includes patients discharged alive whose disposition is unknown, patients who left the hospital against medical advice, patients transferred to another facility that could not be classified as short-term or long-term, and patients transferred to a state facility, psychiatric facility, terminal care or hospice facility, rehabilitation facility, or specialty hospital.

Died—Patients who died while hospitalized as inpatients.

Status not stated—No information was available on whether the patient was discharged alive or died in the hospital.

Hospital—Hospitals in this report are those with an average length of stay for all inpatients of less than 30 days (short stay) or those whose specialty is general (medical or surgical) or children's general. Federal hospitals, hospital units of institutions, and

Table III. Code numbers for Barell injury diagnosis matrix categories in Table 24[Code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification*]

Type of injury	All body sites	Traumatic brain injury	Other head, face, and neck	Spinal cord	Vertebral column	Torso	Upper extremities	Lower extremities	Other and unspecified
All types of injury . . .	800–908, 909.0–909.2, 909.4, 909.9, 910–994, 995.50–995.59, 995.80–995.85	800–801, 803, 804, 850–854, 950.1–950.3, 995.55	802, 807.5–807.6, 830, 848.0–848.2, 870–874, 900, 910, 918, 920–921, 925, 940–941, 947.0, 950.0, 950.9, 951, 953.0, 954.0, 957.0, 959.0	806, 952	805, 839.0–839.5, 847.0–847.4	807.0–807.4, 808–809, 839.6–839.7, 846, 847.9, 848.3–848.5, 860–868, 875–878, 879.0–879.7, 901, 902.0–902.5, 902.81–902.82, 911, 922, 926, 942, 947.3–947.4, 953.1–953.3, 953.5, 954.1–954.9, 959.1	810–818, 831–834, 840–842, 880–887, 903, 912–915, 923, 927, 943–944, 953.4, 955, 959.2–959.5	820–827, 835–838, 843–845, 890–897, 904.0–904.8, 916, 917, 924.0–924.5, 928, 945, 959.6–959.7	819, 828–829, 839.8–839.9, 848.8–848.9, 869, 879.8–879.9, 902.87, 902.89, 902.9, 904.9, 919, 924.8–924.9, 929, 946, 947.1–947.2, 947.8–947.9, 948, 949, 953.8–953.9, 956, 957.1–957.9, 959.8–959.9
Fracture	800–829	800–801, 803, 804	802, 807.5–807.6	806	805	807.0–807.4, 808, 809	810–818	820–827	819, 828–829
Dislocation	830–839	None	830	None	839.0–839.5	839.6–839.7	831–834	835–838	839.8–839.9
Sprains and strains . . .	840–848	None	848.0–848.2	None	847.0–847.4	846, 847.9, 848.3–848.5	840–842	843–845	848.8–848.9
Internal	850–854, 860–869, 952, 995.55	850–854, 995.55	None	952	None	860–868	None	None	869
Open wound	870–884, 890–894	None	870–874	None	None	875–878, 879.0–879.7	880–884	890–894	879.8–879.9
Amputation	885–887, 895–897	None	None	None	None	None	885–887	895–897	None
Blood vessels	900–904	None	900	None	None	901, 902.0–902.5, 902.81–902.82	903	904.0–904.8	902.87, 902.89, 902.9, 904.9
Contusions and superficial injuries . . .	910–924	None	910, 918, 920–921	None	None	911, 922	912–915, 923	916–917, 924.0–924.5	919, 924.8–924.9
Crush	925–929	None	925	None	None	926	927	928	929
Burns	940–949	None	940–941, 947.0	None	None	942, 947.3–947.4	943–944	945	946, 947.1–947.2, 947.8–947.9, 948–949
Nerves and spinal cord	950–951, 953–957	950.1–950.3	950.0, 950.9, 951, 953.0, 954.0, 957.0	None	None	953.1–953.3, 953.5, 954.1–954.9	953.4, 955	None	953.8–953.9, 956, 957.1–957.9
Unspecified	959	None	959.0	None	None	959.1	959.2–959.5	959.6–959.7	959.8–959.9
System-wide	905–908, 909.0–909.2, 909.4, 909.9, 930–939, 958, 960–994, 995.50–995.54, 995.59, 995.80–995.85								

SOURCE: CDC/NCHS, The Barrell injury diagnosis matrix, classification by body region and nature of the injury. Available from <http://www.cdc.gov/nchs/about/otheract/ice/barellmatrix.htm>.

hospitals with fewer than six beds staffed for patient use are excluded.

Newborn infant—An inpatient admitted to a hospital by birth. Sick newborn infants are those that had at least one of the ICD-9-CM (1) diagnostic codes 001–999.

Ownership of hospital—The type of hospital ownership is determined by the organization that controls and operates the hospital. Hospitals are grouped as follows:

Proprietary—Hospitals operated by individuals, partnerships, or corporations for profit.

Government—Hospitals operated by state or local governments.

Nonprofit—Hospitals operated by a nonprofit organization or church.

Principal expected source of payment—The primary expected method of payment for the hospitalization as reported on the patient's medical record. Types of payment include the following:

Medicare—The health insurance program for the aged and disabled administered by the Centers for Medicare and Medicaid Services (CMS). A person 65 years of age and over is included in this category only if Medicare is reported as the principal expected source of payment.

Medicaid—A jointly funded federal-state health insurance

program providing medical care to those unable to afford it. If Medicaid and health maintenance organization (HMO) or preferred provider organization (PPO) are reported, the discharge is included in the category listed first.

Worker's compensation—A state or municipal disability insurance or industrial accident insurance.

Other government—This includes payments under the Title V Program, the state-funded Maternal and Child Health Program that provides coverage for children up to 21 years of age with specific chronic medical conditions (e.g., the Crippled Children's Program), CHAMPUS (Civilian Health and Medical Program of the Uniformed Services) or TRICARE (the Defense Department's health care program for service families), casualty insurance paid by the state, vocational rehabilitation, federal or state research grant (medical research), or legal hold (prisoner in medical detention).

HMO or PPO—Any health maintenance organization (HMO) or preferred provider organization (PPO) sponsored by consumers, insurers, organizations, physicians, or hospitals. Medicare and Medicaid discharges are included in this category when HMO or PPO is

reported as their principal expected source of payment.

Blue Cross Blue Shield and other private—A private insurance plan not specified as an HMO or PPO. This includes Blue Cross Blue Shield plans; medical coverage provided by life insurance companies, private casualty insurance companies, and health insurance companies; and independent plans such as employer- and union-sponsored plans and self-funded plans (partial or total).

Self-pay—The majority of the costs for the hospitalization were expected to be paid by the patient, spouse, family, or next of kin.

No charge—Patients admitted with the understanding that the medical services would be free. This includes patients involved in hospital-sponsored special research or teaching.

Other—Includes any nonprofit source of payment, such as church welfare, United Way (United Appeal), or Shriner's Hospitals for Children.

Procedure—Procedures include surgical procedures (such as appendectomy), diagnostic procedures (such as spinal tap), and therapeutic treatments (such as injection or infusion

Table IV. Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey, by ICD-9-CM category, 2005

[Procedure categories and code numbers are based on the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM)]

Procedure category and ICD-9-CM code	Nonsurgical codes
Operations on the nervous system	01–05 01.18–01.19,03.31,03.39,04.19,05.19
Operations on the endocrine system	06–07 06.19,07.19
Operations on the eye	08–16 08.19,09.19,09.41–09.44,09.49,10.29,11.29,12.29,14.19,15.09,16.21,16.29
Operations on the ear	18–20 18.01,18.11,18.19,20.31,20.39
Operations on the nose, mouth, and pharynx	21–29 21.00–21.02,21.21,21.29,22.19,24.19,25.09,26.19,27.29,28.19,29.11,29.19
Operations on the respiratory system	30–34 31.41–31.42,31.48–31.49,33.21–33.23,33.29,34.21–34.22,34.28–34.29
Operations on the cardiovascular system	35–39 37.26–37.28,37.29,38.29,39.95
Operations on the hemic and lymphatic system	40–41 40.19,41.38–41.39
Operations on the digestive system	42–54 42.22–42.23,42.29,44.11–44.13,44.19,44.32,45.11–45.13,45.19,45.21–45.24,45.28–45.29,48.21–48.23,48.29,49.21,49.29,50.19,51.10–51.11,51.19,52.19,54.21,54.29
Operations on the urinary system	55–59 55.21–55.22,55.29,56.31,56.35,56.39,57.31–57.32,57.39,57.94–57.95,58.21–58.22,58.29,59.29
Operations on the male genital organs	60–64 60.18–60.19,61.19,62.19,63.09,64.19,64.94
Operations on the female genital organs	65–71 65.19,66.19,67.19,68.11,68.19,69.92,70.21–70.22,70.29,71.19
Obstetrical procedures	72–75 73.4–73.59,73.91–73.92,75.31–75.32,75.34–75.35,75.38,75.94
Operations on the musculoskeletal system	76–84 76.19,78.80–78.89,80.20–80.29,81.98,83.29,84.41–84.43,84.45–84.47
Operations on the integumentary system	85–86 85.19,86.19,86.92
Miscellaneous diagnostic and therapeutic procedures and new technologies	87–99,00 87.01–99.99,00.01–00.29

of cancer chemotherapeutic substance) reported on a patient's medical record. Data are for all-listed procedures, which include all occurrences of the procedure coded, regardless of their order in the medical record. Up to four procedures are coded per discharge in NHDS. The procedures classified as nonsurgical in this report are listed in [Table IV](#) by ICD-9-CM code numbers (1). All other procedures were considered surgical. This classification began in 1979 for the presentation of NHDS data and still remains an important distinction. However, the development of minimally invasive and noninvasive procedures has resulted in less difference between surgical and nonsurgical procedures with regard to operative or anesthetic risk and requirements for highly trained personnel or special equipment.

Region—Hospitals are classified by location in one of the four geographic regions of the United States that correspond to those used by the U.S. Census Bureau.

Region States included

Northeast	Maine, Vermont, New Hampshire, Massachusetts, Connecticut, Rhode Island, New York, New Jersey, and Pennsylvania
Midwest	Ohio, Illinois, Indiana, Michigan, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Kansas, and Nebraska
South	Delaware, Maryland, District of Columbia, West Virginia, Virginia, Kentucky, Tennessee, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, Oklahoma, Arkansas, and Texas
West	Washington, Oregon, California, Nevada, New Mexico, Arizona, Idaho, Utah, Colorado, Montana, Wyoming, Alaska, and Hawaii

Changes in ICD-9-CM Codes

Although the ICD-9-CM has been used for coding NHDS data since 1979, the classification system undergoes annual updating. Assignment of new diagnostic and procedure codes and fourth- and fifth-digit expansion of codes, as well as code deletions, are contained in addenda developed by the ICD-9-CM Coordination and Maintenance Committee and are approved by the Director of NCHS and the Administrator of CMS (formerly Health Care Financing Administration). Addenda to the ICD-9-CM have become effective October 1 of each year since 1986, except in 1999. No addendum was released in 1999 because of concerns about instituting coding changes before the millennium crossover.

The data in this report were coded using the addenda for October 1, 1986–2004. When hospitals reported data using code changes introduced in the October 1, 2005, addendum, the new codes were converted to their previous code assignments so that the annual estimates would be based on consistent coding.

The new ICD-9-CM codes from the 1986–2004 addenda, the date they were introduced, and the code to which the diagnosis or procedure had been assigned before development of the new code are shown in [Table V](#) (diagnoses codes) and [Table VI](#) (procedure codes). Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD-9-CM (1).

Table V. Changes in ICD-9-CM diagnosis codes, 1986–2004

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
005.81, 005.89.....	1995	005.8	204.91.....	1991	V10.61
007.4	1997	007.8	205.00.....	1991	205.0
007.5.....	2000	007.8	205.01.....	1991	V10.62
008.00–008.09.....	1992	008.0	205.10.....	1991	205.1
008.43–008.47.....	1992	008.49	205.11.....	1991	V10.62
008.6–008.69.....	1992	008.6	205.20.....	1991	205.2
031.2.....	1997	031.8	205.21.....	1991	V10.62
038.10–038.11, 038.19.....	1997	038.1	205.30.....	1991	205.3
040.82.....	2002	040.89	205.31.....	1991	V10.62
041.00–041.05, 041.09.....	1992	041.0	205.80.....	1991	205.8
041.04 (Code title restated).....	1997	041.04	205.81.....	1991	V10.62
041.10–041.19.....	1992	041.1	205.90.....	1991	205.9
041.81–041.85, 041.89.....	1992	041.8	205.91.....	1991	V10.62
041.86.....	1995	041.84	206.00.....	1991	206.0
042.....	1994	042.0–042.2, 042.9, 043.0–043.3, 043.9, 044.0, 044.9 (Codes deleted)	206.01.....	1991	V10.63
042.0–044.9	1986	279.19	206.10.....	1991	206.1
066.4.....	2002	066.3	206.11.....	1991	V10.63
066.40–066.42, 066.49.....	2004	066.4	206.20.....	1991	206.2
070.20–070.21.....	1991	070.2	206.21.....	1991	V10.63
070.22.....	1994	070.20	206.80.....	1991	206.8
070.23.....	1994	070.21	206.81.....	1991	V10.63
070.30–070.31.....	1991	070.3	206.90.....	1991	206.9
070.32.....	1994	070.30	206.91.....	1991	V10.63
070.33.....	1994	070.31	207.00.....	1991	207.0
070.41–070.43.....	1991	070.4	207.01.....	1991	V10.69
070.44.....	1994	070.41	207.10.....	1991	207.1
070.49.....	1991	070.4	207.11.....	1991	V10.69
070.51–070.53.....	1991	070.5	207.20.....	1991	207.2
070.54.....	1994	070.51	207.21.....	1991	V10.69
070.59.....	1991	070.5	207.80.....	1991	207.8
070.70.....	2004	070.51	207.81.....	1991	V10.69
070.71.....	2004	070.41	208.00.....	1991	208.0
077.98–077.99.....	1993	077.9	208.01.....	1991	V10.60
078.10–078.11, 078.19.....	1993	078.1	208.10.....	1991	208.1
078.88.....	1993	078.89	208.11.....	1991	V10.60
079.4.....	1993	079.8	208.20.....	1991	208.2
079.50–079.53, 079.59.....	1993	079.8	208.21.....	1991	V10.60
079.6.....	1996	079.89	208.80.....	1991	208.8
079.81.....	1995	079.89	208.81.....	1991	V10.60
079.82.....	2003	079.89	208.90.....	1991	208.9
079.88–079.89.....	1993	079.8	208.91.....	1991	V10.60
079.98–079.99.....	1993	079.9	237.70–237.72.....	1990	237.7
082.40–082.41, 082.49.....	2000	082.8	250.02.....	1993	250.90
088.81, 088.89.....	1989	088.8	250.03.....	1993	250.91
088.82.....	1993	088.89	250.12.....	1993	250.10
099.40–099.49.....	1992	099.4	250.13.....	1993	250.11
099.50–099.59.....	1992	078.89	250.22.....	1993	250.20
112.84–112.85.....	1992	112.89	250.23.....	1993	250.21
114.4–114.5.....	1993	114.3	250.32.....	1993	250.30
176.0–176.9.....	1991	173.0–173.9	250.33.....	1993	250.31
203.00.....	1991	203.0	250.42.....	1993	250.40
203.01.....	1991	V10.79	250.43.....	1993	250.41
203.10.....	1991	203.1	250.52.....	1993	250.50
203.11.....	1991	V10.79	250.53.....	1993	250.51
203.80.....	1991	203.8	250.62.....	1993	250.60
203.81.....	1991	V10.79	250.63.....	1993	250.61
204.00.....	1991	204.0	250.72.....	1993	250.70
204.01.....	1991	V10.61	250.73.....	1993	250.71
204.10.....	1991	204.1	250.82.....	1993	250.80
204.11.....	1991	V10.61	250.83.....	1993	250.81
204.20.....	1991	204.2	250.92.....	1993	250.90
204.21.....	1991	V10.61	250.93.....	1993	250.91
204.80.....	1991	204.8	252.00–252.02, 252.08.....	2004	252.0
204.81.....	1991	V10.61	255.10–255.14.....	2003	255.1
204.90.....	1991	204.9	256.31–256.39.....	2001	256.3
			273.4.....	2004	277.6
			275.40–275.42, 275.49.....	1997	275.4

Table V. Changes in ICD-9-CM diagnosis codes, 1986–2004—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
277.02–277.03, 277.09.....	2002	277.00	404.90–404.93.....	1989	404.9
277.7.....	2001	277.8	410.00–410.02.....	1989	410.0
277.81–277.84, 277.89.....	2003	277.8	410.10–410.12.....	1989	410.1
277.85–277.86.....	2004	277.89	410.20–410.22.....	1989	410.2
277.87.....	2004	277.89, 758.89	410.30–410.32.....	1989	410.3
278.00–278.01.....	1995	278.0	410.40–410.42.....	1989	410.4
282.41–282.42, 282.49.....	2003	282.4	410.50–410.52.....	1989	410.5
282.64.....	2003	282.63	410.60–410.62.....	1989	410.6
282.68.....	2003	282.69	410.70–410.72.....	1989	410.7
283.10–283.11, 283.19.....	1993	283.1	410.80–410.82.....	1989	410.8
285.21–285.22, 285.29.....	2000	285.8	410.90–410.92.....	1989	410.9
289.52.....	2003	289.59	411.81.....	1989	410.9
289.81–289.82, 289.89.....	2003	289.8	411.89.....	1989	411.8
291.81, 291.89.....	1996	291.8	414.00–414.01.....	1994	414.0
293.84.....	1996	293.89	414.02–414.03.....	1994	996.03
294.10–294.11.....	2000	294.1	414.04–414.05.....	1996	414.00
300.82.....	1996	300.81	414.06.....	2002	414.00
305.1.....	1994	305.10, 305.11, 305.12, 305.13 (Codes deleted)	414.07.....	2003	414.06
312.81–312.82, 312.89.....	1994	312.8	414.12.....	2002	414.11
315.32.....	1996	315.39	415.11.....	1995	997.3 + 415.1
320.81–320.89.....	1992	320.8	415.19.....	1995	415.1
331.11, 331.19.....	2003	331.1	428.20–428.23.....	2002	428.0
331.82.....	2003	331.89	428.30–428.33.....	2002	428.0
333.92–333.93.....	1994	333.99	428.40–428.43.....	2002	428.0
337.20–337.22, 337.29.....	1993	337.9	429.71, 429.79.....	1989	410.0–410.9
337.3.....	1998	337.9	433.00–433.01.....	1993	433.0
342.00–342.02.....	1994	342.0	433.10–433.11.....	1993	433.1
342.10–342.12.....	1994	342.1	433.20–433.21.....	1993	433.2
342.80–342.82.....	1994	342.9	433.30–433.31.....	1993	433.3
342.90–342.92.....	1994	342.9	433.80–433.81.....	1993	433.8
344.00–344.04, 344.09.....	1994	344.0	433.90–433.91.....	1993	433.9
344.30–344.32.....	1994	344.3	434.00–434.01.....	1993	434.0
344.40–344.42.....	1994	344.4	434.10–434.11.....	1993	434.1
344.81, 344.89.....	1993	344.8	434.90–434.91.....	1993	434.9
345.00–345.01.....	1989	345.0	435.3.....	1995	435.0 + 435.1
345.10–345.11.....	1989	345.1	437.7.....	1992	780.9
345.40–345.41.....	1989	345.4	438.0.....	1997	294.9 + 438
345.50–345.51.....	1989	345.5	438.10.....	1997	784.5 + 438
345.60–345.61.....	1989	345.6	438.11.....	1997	784.3 + 438
345.70–345.71.....	1989	345.7	438.12.....	1997	784.4 + 438
345.80–345.81.....	1989	345.8	438.19.....	1997	784.5 + 438
345.90–345.91.....	1989	345.9	438.20.....	1997	342.90 + 438
346.00–346.01.....	1992	346.0	438.21.....	1997	342.91 + 438
346.10–346.11.....	1992	346.1	438.22.....	1997	342.92 + 438
346.20–346.21.....	1992	346.2	438.30.....	1997	344.40 + 438
346.80–346.81.....	1992	346.8	438.31.....	1997	344.41 + 438
346.90–346.91.....	1992	346.9	438.32.....	1997	344.42 + 438
347.00–347.01.....	2004	347	438.40.....	1997	344.30 + 438
347.10–347.11.....	2004	347	438.41.....	1997	344.31 + 438
348.30–348.31, 348.39.....	2003	348.3	438.42.....	1997	344.32 + 438
355.71.....	1993	354.4	438.50–438.52.....	1997	344.89 + 438
355.79.....	1993	355.7	438.53.....	1998	438.50
357.81–357.82, 357.89.....	2002	357.8	438.6–438.7.....	2002	438.89
358.00–358.01.....	2003	358.0	438.81.....	1997	784.69 + 438
359.81, 359.89.....	2002	359.8	438.82.....	1997	787.2 + 438
365.83.....	2002	365.89	438.83–438.85.....	2002	438.89
371.82.....	1992	371.89	438.89.....	1997	438
372.81, 372.89.....	2000	372.8	438.9.....	1997	438
374.87.....	1990	374.89	440.20–440.22.....	1992	440.2
380.03.....	2004	733.99	440.23.....	1993	440.20 + (707.1 or 707.8 or 707.9)
403.00–403.01.....	1989	403.0	440.24.....	1993	440.20 + 785.4
403.10–403.11.....	1989	403.1	440.29.....	1993	440.2
403.90–403.91.....	1989	403.9	440.30–440.32.....	1994	996.1
404.00–404.03.....	1989	404.0	441.00–441.03.....	1994	441.0
404.10–404.13.....	1989	404.1	441.6.....	1993	441.1 + 441.3
			441.7.....	1993	441.2 + 441.4
			443.21.....	2002	442.81

Table V. Changes in ICD-9-CM diagnosis codes, 1986–2004—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
443.22	2002	442.2	530.12	2001	530.10
443.23	2002	442.1	530.20–530.21	2003	530.2
443.24, 443.29	2002	442.89	530.81	1993	530.1
445.01–445.02	2002	440.29	530.82–530.84, 530.89	1993	530.8
445.81	2002	440.1	530.85	2003	530.2
445.89	2002	440.8	530.86–530.87	2004	997.4
446.20–446.21, 446.29	1990	446.2	535.00–535.01	1991	535.0
451.82–451.84	1993	451.89	535.10–535.11	1991	535.1
453.40–453.42	2004	453.8	535.20–535.21	1991	535.2
454.8	2002	454.9	535.30–535.31	1991	535.3
458.2	1995	997.9 + 458.9	535.40–535.41	1991	535.4
458.21, 458.29	2003	458.2	535.50–535.51	1991	535.5
458.8	1997	458.9	535.60–535.61	1991	535.6
464.00–464.01	2001	464.0	536.3	1994	536.8
464.50–464.51	2001	464.0	536.40–536.42, 536.49	1998	997.4
466.11, 466.19	1996	466.1	537.82	1990	537.89
474.0 (Code title restated)	1997	474.0	537.83	1991	537.82
474.00–474.02	1997	474.0	537.84	2002	531.00
477.1	2000	477.8	556.0–556.6, 556.8–556.9	1994	556
477.2	2004	477.8	558.3	2000	558.9
480.3	2003	480.8	562.02	1991	562.00
482.30–482.39	1992	482.3	562.03	1991	562.01
482.40–482.41, 482.49	1998	482.4	562.12	1991	562.10
482.81–482.83, 482.89	1992	482.8	562.13	1991	562.11
482.84	1997	482.83	564.00–564.09	2001	564.0
483.0	1992	483	564.81, 564.89	1998	564.8
483.1	1996	078.88 + 484.8	569.60–569.61, 569.69	1995	569.6
483.8	1992	483	569.62	1998	569.69
491.20–491.21	1991	491.2	569.84	1990	557.1
491.22	2004	491.21	569.85	1991	569.84
493.02	2000	493.00	569.86	2002	569.82
493.12	2000	493.10	574.60	1996	574.00 + 574.30
493.20	1989	493.90	574.61	1996	574.01 + 574.31
493.21	1989	493.91	574.70	1996	574.10 + 574.40
493.22	2000	493.20	574.71	1996	574.11 + 574.41
493.81	2003	519.1	574.80	1996	574.00 + 574.10 + 574.30 + 574.40
493.82	2003	493.90–493.91	574.81	1996	574.01 + 574.11 + 574.31 + 574.41
493.92	2000	493.90	574.90	1996	574.20 + 574.50
494.0–494.1	2000	494	574.91	1996	574.21 + 574.51
512.1	1994	997.3	575.10–575.11	1996	575.1
517.3	2003	282.62	575.12	1996	575.0 + 575.1
518.6	1997	518.89	588.81, 588.89	2004	588.8
518.82–518.89	1987	518.8	593.70–593.73	1994	593.7
518.83–518.84	1998	518.81	596.51–596.53	1992	596.5
519.00–519.02, 519.09	1998	519.0	596.54	1992	344.61
521.00–521.05, 521.09	2001	521.0	596.55–596.59	1992	596.5
521.06–521.08	2004	521.09	599.81–599.89	1992	599.8
521.10–521.15	2004	521.1	600.0–600.3, 600.9	2000	600
521.20–521.25	2004	521.2	600.00–600.01	2003	600.0
521.30–521.35	2004	521.3	600.10–600.11	2003	600.1
521.40–521.45	2004	521.4	600.20–600.21	2003	600.2
523.20–523.25	2004	523.2	600.90–600.91	2003	600.9
524.00–524.06, 524.09	1992	524.0	602.3	2001	602.8
524.07	2004	524.09	607.85	2003	607.89
524.10–524.12, 524.19	1992	524.1	608.82	2001	608.83
524.20–524.29	2004	524.2	608.87	2001	608.89
524.30–524.37, 524.39	2004	524.3	618.00–618.05, 618.09	2004	618.0
524.50–524.57, 524.59	2004	524.5	618.81–618.83, 618.89	2004	618.8
524.60–524.63, 524.69	1991	524.6	621.30–621.33	2004	621.3
524.64	2004	524.69	622.10–622.12	2004	622.1
524.70–524.77, 524.79	1992	524.8	629.20–629.23	2004	629.8
524.75–524.76	2004	524.79	633.00–633.01	2002	633.0
524.81–524.82, 524.89	2004	524.8	633.10–633.11	2002	633.1
525.10–525.19	2001	525.1	633.20–633.21	2002	633.2
525.20–525.26	2004	525.2	633.80–633.81	2002	633.8
528.71–528.72, 528.79	2004	528.7			
530.10–530.11, 530.19	1993	530.1			

Table V. Changes in ICD-9-CM diagnosis codes, 1986–2004—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
633.90–633.91	2002	633.9	758.31–758.33, 758.39	2004	758.3
645.00–645.01, 645.03.	1991	645.0–645.1, 645.3	758.81	1996	758.8
645.10–645.11, 645.13.	2000	645.00–645.01, 645.03	758.89	1996	758.9
645.20–645.21, 645.23.	2000	645.00–645.01, 645.03	759.81–759.82	1989	759.8
651.30–651.31, 651.33.	1989	651.00–651.01, 651.03	759.83	1994	759.89
651.40–651.41, 651.43.	1989	651.10–651.11, 651.13	759.89	1989	759.8
651.50–651.51, 651.53.	1989	651.20–651.21, 651.23	760.75	1991	760.79
651.60–651.61, 651.63	1989	651.80–651.81, 651.83	760.76	1994	760.79
654.20–654.21, 654.23.	1990	654.2, 654.9	763.81–763.83, 763.89	1998	763.8
654.90–654.94	1990	654.2, 654.9	764.00–764.09	1988	764.0
655.70–655.71, 655.73.	1997	655.8	764.10–764.19	1988	764.1
657.00–657.01, 657.03.	1991	657.0–657.1, 657.3	764.20–764.29	1988	764.2
659.60–659.61, 659.63.	1992	659.80–659.81, 659.83	764.90–764.99	1988	764.9
659.70–659.71, 659.73.	1998	656.30–656.31, 656.33	765.00–765.09	1988	765.0
665.10–665.11	1992	665.10–665.12, 665.141 ¹	765.10–765.19	1988	765.1
670.00, 670.02, 670.04.	1991	670.0–670.1, 670.3	765.20–765.24	2002	765.00–765.09
672.00, 672.02, 672.04.	1991	672.0–672.1, 672.3	765.25–765.29	2002	765.10–765.19
674.50–674.54.	2003	674.80, 674.82, 674.84	766.21–766.22	2003	766.2
677.	1994	None	767.11–767.19	2003	767.1
686.00–686.01, 686.09.	1997	686.0	770.81–770.84, 770.89	2002	770.8
690.10, 690.18.	1995	690	771.81–771.83, 771.89	2002	771.8
690.11.	1995	691.8 + 704.8	772.10–772.14	2001	772.1
690.12.	1995	691.8	779.7	2001	772.1
690.8.	1995	690	779.81–779.82, 779.89	2002	779.8
692.72–692.74.	1992	692.79	779.83	2003	779.89
692.75.	2000	692.79	780.01–780.02, 780.09	1992	780.0
692.76–692.77.	2001	692.71	780.03	1993	780.01
692.82–692.83.	1992	692.89	780.31, 780.39	1997	780.3
692.84.	2004	692.89	780.57	1992	780.51, 780.53
702.0–702.8.	1991	702	780.58	2004	780.59
702.11, 702.19.	1994	702.1	780.71, 780.79	1998	780.7
704.02.	1993	704.09	780.91–780.92, 780.99	2002	780.9
705.21–705.22.	2004	780.8	780.93–780.94	2003	780.99
707.00–707.07, 707.09.	2004	707.0	781.8	1994	781.9
707.10–707.15, 707.19.	2000	707.1	781.91–781.92, 781.99	2000	781.9
709.00–709.01, 709.09.	1994	709.0	781.93	2002	723.5
710.5.	1992	288.3, 729.1	781.94	2003	781.99
718.70–718.79.	2001	718.80–718.89	783.21	2000	783.2
719.7.	2003	719.70, 719.75– 719.79 (Codes deleted)	783.22	2000	783.4
727.83.	2000	727.89	783.40–783.43	2000	783.4
728.86.	1995	729.4	783.7	2000	783.4
728.87.	2003	728.9	785.52	2003	785.59
728.88.	2003	728.89	786.03–786.07	1998	786.09
733.10–733.16, 733.19.	1993	733.1	787.01–787.03	1994	787.0
733.93.	2001	733.16	787.91	1995	558.9
733.94–733.95.	2001	733.19	787.99	1995	787.9
738.10–738.19.	1992	738.1	788.20–788.21, 788.29	1993	788.2
747.60–747.64, 747.69.	1993	747.6	788.30–788.39	1992	788.3
747.82.	1993	747.89	788.38	2004	788.39
747.83.	2002	747.89, 747.9	788.41–788.43	1993	788.4
752.51–752.52.	1996	752.5	788.61–788.62, 788.69	1993	788.6
752.61–752.63.	1996	752.6	788.63	2003	788.69
752.64–752.65, 752.69.	1996	752.8	789.00–789.07, 789.09	1994	789.0
752.81, 752.89.	2003	752.8	789.30–789.37, 789.39	1994	789.3
753.10–753.17, 753.19.	1990	753.1	789.40–789.47, 789.49	1994	789.4
753.20–753.23, 753.29.	1996	753.2	789.60–789.67, 789.69	1994	789.6
756.70–756.71, 756.79.	1997	756.7	790.01, 790.09	2000	790.0
			790.21–790.22, 790.29	2003	790.2
			790.91, 790.93, 790.99	1993	790.9
			790.92	1993	286.9
			790.94	1997	790.99
			790.95	2004	790.99
			792.5	2000	792.9
			793.80–793.81, 793.89	2001	793.8
			795.00–795.02, 795.09	2002	795.0
			795.03, 795.04	2004	622.1
			795.05, 795.08	2004	795.09

Table V. Changes ICD-9-CM in diagnosis codes, 1986–2004—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
795.31–795.39	2002	795.3	V08	1994	044.9, 795.8 (Codes deleted)
795.71	1994	795.8 (Code deleted)	V09.0–V09.91	1993	None
795.79	1994	795.7	V10.48	1998	V10.49
795.8	1986	795.7	V10.53	2001	V10.59
796.5	1997	796.9	V12.00–V12.03, V12.09	1994	V12.0
796.6	2004	796.9	V12.40–V12.41, V12.49	1997	V12.4
799.81, 799.89	2003	799.8	V12.50–V12.52, V12.59	1995	V12.5
813.45	2002	813.42	V12.70–V12.72, V12.79	1994	V12.7
823.40–823.42	2002	823.80–823.82	V13.00–V13.01, V13.09	1994	V13.0
840.7	2001	840.8	V13.21, V13.29	2002	V13.2
850.11–850.12	2003	850.1	V13.61, V13.69	1998	V13.6
864.05	1992	864.09	V15.01–V15.09	2000	V15.0
864.15	1992	864.19	V15.41–V15.42, V15.49	1996	V15.4
909.5	1994	909.9	V15.82	1994	305.13 (Code deleted)
922.31–922.33	1996	922.3	V15.84–V15.86	1995	V15.89
925.1–925.2	1993	925	V15.87	2003	V15.89
959.0 (Code title restated)	1997	959.0	V16.40–V16.43, V16.49	1997	V16.4
959.01	1997	854.00	V16.51, V16.59	1998	V16.5
959.09	1997	959.0	V18.61, V18.69	1998	V18.6
959.11–959.14, 959.19	2003	959.1	V21.30–V21.35	2000	None
965.61, 965.69	1998	965.6	V23.41, V23.49	2002	V23.4
989.81–989.84, 989.89	1995	989.8	V23.7	1989	V23.8
995.50–995.55, 995.59	1996	995.5	V23.81–V23.84, V23.89	1998	V23.8
995.59	1996	995.5	V25.03	2003	V25.01
995.60–995.69	1993	995.0	V25.43	1992	V25.49
995.7	2000	None	V25.5	1992	V25.8
995.80, 995.82–995.85	1996	995.81	V26.21–V26.22, V26.29	2000	V26.2
995.81 (Code title restated)	1996	995.81	V26.51, V26.52	1998	None
995.86	1998	995.89	V28.6	1997	V28.8
995.90–995.94	2002	038.0–038.9	V29.0–V29.1, V29.8	1992	V71.8
996.04	1994	996.09	V29.2	1994	V29.8
996.51–996.59	1987	996.5	V29.3	1998	V29.8
996.55	1998	996.52	V29.9	1992	V71.9
996.56	1998	996.59	V30.00–V30.01	1989	V30.0
996.57	2003	996.59	V31.00–V31.01	1989	V31.0
996.60–996.69	1989	996.6	V32.00–V32.01	1989	V32.0
996.68	1998	996.69	V33.00–V33.01	1989	V33.0
996.70–996.79	1989	996.7	V34.00–V34.01	1989	V34.0
996.80–996.84, 996.86, 996.89	1987	996.8	V35.00–V35.01	1989	V35.0
996.85	1990	996.89	V36.00–V36.01	1989	V36.0
996.87	2000	996.89	V37.00–V37.01	1989	V37.0
997.00–997.01, 997.09	1995	997.0	V39.00–V39.01	1989	V39.0
997.02	1995	997.9 + 430–434, 436	V42.81–V42.83, V42.89	1997	V42.8
997.71	2001	997.4	V42.84	2000	V42.89
997.72	2001	997.5	V43.21–V43.22	2003	V43.2
997.79	2001	997.2	V43.60–43.66, V43.69	1994	V43.6
997.91, 997.99	1995	997.9	V43.81–V43.82, V43.89	1995	V43.8
998.11–998.12	1996	998.1	V43.83	1998	V43.89
998.13	1996	998.89	V44.50–V44.52, V44.59	1998	V44.5
998.31–998.32	2002	998.3	V45.00, V45.02, V45.09	1994	V45.89
998.51, 998.59	1996	998.5	V45.01	1994	V45.0
998.81–998.82, 998.89	1994	998.8	V45.51	1994	V45.5
998.83	1996	998.89	V45.52, V45.59	1994	V45.89
V01.71, V01.79	2004	V01.7	V45.61, V45.69	1997	V45.6
V01.81, V01.89	2002	V01.8	V45.71	1997	611.8
V01.82	2003	V01.89	V45.72	1997	569.89
V01.83–V01.84	2004	V01.89	V45.73	1997	593.89
V02.51–V02.52, V02.59	1998	V02.5	V45.74	2000	593.89, 596.8
V02.60–V02.62, V02.69	1997	V02.6	V45.75	2000	V45.89
V03.81–V03.82, V03.89	1994	V03.8	V45.76	2000	518.89
V04.81–V04.82, V04.89	2003	V04.8	V45.77	2000	602.8, 607.89, 608.89, 620.8, 621.8, 622.8
V05.3–V05.4	1993	V05.8	V45.78	2000	360.89
V06.5–V06.6	1994	V06.8			
V07.31, V07.39	1994	V07.3			
V07.4	1992	V07.8			

Table V. Changes in ICD-9-CM diagnosis codes, 1986–2004—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
V45.79	2000	255.8, 289.59, 388.8, 569.49, 577.8 V45.89	V76.46–V76.47	2000	V76.49
V45.82	1994	V45.89	V76.50–V76.52	2000	V76.49
V45.83	1995	V45.89	V76.81, V76.89	2000	V76.8
V45.84	2001	None	V77.91, V77.99	2000	V77.9
V45.85	2003	V45.89	V82.81, V82.89	2000	V82.8
V46.11–V46.12	2004	V46.1	V83.01–V83.02	2001	None
V46.2	2002	V46.8	V83.81	2002	None
V49.60–V49.67	1994	V49.5	V83.89	2002	V19.8
V49.70–V49.77	1994	V49.5	V84.01–V84.04, V84.09	2004	None
V49.81	2000	None	V84.8	2004	None
V49.82	2001	None	E854.8	1995	E858.8
V49.83	2004	None	E869.4	1994	E869.8
V49.89	2000	V49.8	E880.1	1995	E884.9
V50.41–V50.42, V50.49	1994	V50.8	E884.3–E884.4	1995	E884.2
V53.01–V53.02, V53.09	1997	V53.0	E884.5–E884.6	1995	E884.9
V53.31	1994	V53.3	E885.0	2002	E885.9
V53.32, V53.39	1994	V53.9	E885.1–E885.4, E885.9	2000	E885
V53.90–V53.91, V53.99	2003	V53.9	E888.0	2001	E920
V54.01–V54.02, V54.09	2003	V54.0	E888.1, E888.8–E888.9	2001	E888
V54.10–V54.17, V54.19	2002	V54.8	E906.5	1995	E906.3
V54.20–V54.27, V54.29	2002	V54.8	E908.0–E908.4, E908.8–E908.9	1995	E908
V54.81, V54.89	2002	V54.8	E909.0–E909.4, E909.8–E909.9	1995	E909
V56.1	1995	V58.89	E917.3–E917.4	2001	E917.9
V56.1 (Code title restated)	1998	V56.1	E917.5	2001	E886.0
V56.2	1998	V56.1	E917.6	2001	E917.1
V56.31–V56.32	2000	V56.8	E917.7–E917.8	2001	E888
V57.21–V57.22	1994	V57.2	E920.5	1995	E920.4
V58.41, V58.49	1994	V58.4	E922.4	1997	E917.9
V58.42–V58.43	2002	V58.49	E922.5	2002	E922.8
V58.44	2004	V58.71–V58.78	E924.2	1995	E924.0
V58.61, V58.69	1995	V67.51	E928.3	2000	E928.8
V58.62	1998	V58.69	E928.4–E928.5	2003	E928.8
V58.63–V58.65	2003	V58.69	E955.6	1997	E955.9
V58.66–V58.67	2004	V58.69	E955.7	2002	E955.4
V58.71–V58.78	2002	V58.49	E967.2	1996	E967.0
V58.81, V58.89	1994	V58.8	E967.3	1996	None
V58.82	1995	V58.89	E967.4–E967.8	1996	E967.1
V58.83	2000	V58.89	E968.5	1995	E968.8
V59.01–V59.02, V59.09	1995	V59.0	E968.6	1997	E968.8
V59.6	1995	V59.8	E968.7	2000	E968.8
V61.10–V61.12	1996	V61.1	E985.6	1997	E985.4
V61.22	1996	V61.21	E985.7	2002	E985.4
V62.83	1996	V65.49	E979.0–E979.9	2002	E960.0–E966, E968.0–E968.9
V64.4	1997	None	E999.0	2002	E999
V64.41–V64.43	2003	V64.4	E999.1	2002	E969
V65.11, V65.19	2003	V65.1			
V65.40–V65.45, V65.49	1994	V65.4			
V65.46	2003	V65.1			
V66.7	1996	None			
V67.00–V67.01, V67.09	2000	V67.0			
V69.0–V69.3	1994	None			
V69.4	2004	V69.8			
V69.8–V69.9	1994	None			
V71.81, V71.89	2000	V71.8			
V71.82–V71.83	2002	V71.89			
V72.31–V72.32	2004	V72.3			
V72.40–V72.41	2004	V72.4			
V72.81 V72.85	1993	V72.8			
V73.88–V73.89	1993	V73.8			
V73.98–V73.99	1993	V73.9			
V76.10–V76.12, V76.19	1997	V76.1			
V76.44–V76.45	1998	V76.49			

¹The title for the subcategory 665.1 has been changed, making the fifth-digit subclassifications 665.12 and 665.14 invalid.

Table VI. Changes ICD-9-CM in procedure codes, 1986–2004

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
00.01.....	2002	99.99	37.67.....	1998	37.4
00.10.....	2002	99.25	37.68.....	2004	37.62
00.11.....	2002	99.19	37.70(Leads only).....	1987	37.70 (Leads/device)
00.12.....	2002	93.98	37.71–37.72 (Leads only).....	1987	37.74 (Leads/device)
00.13.....	2002	99.29	37.73 (Leads only).....	1987	37.73 (Leads/device)
00.14.....	2002	99.21	37.74 (Leads only).....	1987	37.76 (Leads/device)
00.15.....	2003	99.28	37.75 (Leads only).....	1987	37.89 (Leads/device)
00.16.....	2004	None	37.76 (Leads only).....	1987	37.81 (Leads/device)
00.17.....	2004	99.29	37.77 (Leads only).....	1987	37.83–37.84 (Leads/device)
00.21.....	2004	88.71	37.78.....	1987	37.71–37.72
00.22.....	2004	88.73	37.79.....	1987	86.09
00.23.....	2004	88.77	37.80–37.83 (Device only).....	1987	37.73–37.77 (Leads/Device)
00.24.....	2004	88.72	37.85–37.87.....	1987	37.85
00.25.....	2004	88.75	37.80–37.87.....	1992	89.49 (Code deleted; this procedure is included in the code for pacemaker insertion/replacement)
00.28–00.29.....	2004	88.79	37.89.....	1987	37.86 + 37.89
00.31–00.35, 00.39.....	2004	None	37.90.....	2004	37.99
00.50.....	2002	37.70–37.74, 37.76, 37.80–37.87	37.94–37.98.....	1986	37.99
00.51.....	2002	37.94	38.22.....	1986	38.29
00.52.....	2002	None	38.44 (Abdominal aorta only).....	1986	38.44 (Entire aorta)
00.53.....	2002	37.80–37.87	38.45 (Thoracic aorta added).....	1989	38.44–38.45
00.54.....	2002	37.96	38.95.....	1989	38.93
00.55.....	2002	39.90	39.28.....	1991	39.29
00.61–00.62.....	2004	39.50	39.50.....	1995	39.59
00.63–00.65.....	2004	00.55, 39.90	39.65.....	1988	39.61
00.91–00.93.....	2004	None	39.66.....	1990	39.65
02.96.....	1992	89.19	39.71, 39.79.....	2000	39.52
03.90.....	1987	03.99 (Insertion of catheter)	39.90.....	1996	39.50
05.25.....	1995	39.7	41.00–41.03.....	1988	41.0
11.75.....	1989	11.79	41.04.....	1994	99.79
11.76.....	1989	11.62	41.05–41.06.....	1997	None
20.96–20.98.....	1986	20.95	41.07.....	2000	41.04
22.12.....	1988	22.11	41.08.....	2000	41.05
26.12.....	1988	26.11	41.09.....	2000	41.01
27.64.....	2004	27.69	42.25.....	1988	42.24
29.31.....	1991	83.02	42.33.....	1989	42.32, 42.39
29.32–29.33, 29.39.....	1991	29.3	42.33.....	1990	42.91
31.45.....	1988	31.43–31.44	43.11.....	1989	43.1
31.95.....	1989	31.75	43.19.....	1989	43.1, 43.2
32.01, 32.09.....	1989	32.0	43.41.....	1989	43.41, 43.49
32.22.....	1995	32.29, 32.9	44.21, 44.29.....	1986	44.2
32.28.....	1989	32.29	44.22.....	1986	44.99
33.27.....	1987	33.22 + 33.27	44.32.....	2001	44.39
33.28.....	1987	33.27	44.38.....	2004	44.39
33.29.....	1987	33.28 33.29	44.43.....	1989	43.49, 45.32
33.50–33.52.....	1995	33.5	44.44.....	1989	38.86
33.6.....	1990	33.5 + 37.5	44.49.....	1989	43.0
34.05.....	1994	34.99	44.67.....	2004	44.66
35.84.....	1988	35.82	44.68.....	2004	44.69
35.96.....	1986	35.03	44.93–44.94.....	1986	44.99
36.00–36.03.....	1986	36.0	44.95–44.96.....	2004	44.69
36.04.....	1986	39.97	44.97–44.98.....	2004	44.99
36.05.....	1986	36.01, ¹ 36.02	45.16.....	1988	45.14 (45.15 before 1987)
36.05.....	1987	36.01	45.30.....	1989	45.31, 45.32
36.06.....	1995	36.01, 36.02, 36.03, 36.05	45.42.....	1988	45.41
36.07.....	2002	36.06	45.43.....	1989	45.49
36.09.....	1986	36.0	45.75 (Hartmann resection added).....	1988	48.66 (Code deleted)
36.09.....	1991	36.00 (Code deleted)	45.95.....	1987	45.93
36.17.....	1996	36.19	46.13.....	1992	46.12 (Code deleted)
36.31–36.32, 36.39.....	1998	36.3	46.32.....	1989	46.39
37.26–37.27.....	1988	37.29	46.85.....	1989	46.99
37.28.....	2001	88.72	46.97.....	2000	46.99
37.34.....	1988	37.33	47.01, 47.09.....	1996	47.0
37.35.....	1997	37.33	47.11, 47.19.....	1996	47.1
37.51.....	2003	37.5	48.36.....	1995	45.42
37.52.....	2003	37.62			
37.53–37.54.....	2003	37.63			
37.65–37.66.....	1995	37.62			

See footnotes at end of table.

Table VI. Changes ICD-9-CM in procedure codes, 1986–2004—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
49.31, 49.39	1989	49.3	77.56	1989	77.89, 78.49, 81.18
49.75–49.76.	2002	49.79	77.57	1989	77.89, 80.48, 81.18, 83.85
51.10.	1989	51.97	77.58	1989	77.59, 81.18
51.11.	1989	51.11, 51.97	78.10.	1991	78.40
51.14.	1989	51.12	78.11.	1991	78.41
51.15.	1989	51.97	78.12.	1991	78.42
51.21.	1996	51.22, 51.23	78.13.	1991	78.43
51.22.	1991	51.21 (Code deleted), 51.22	78.14.	1991	78.44
51.23.	1991	51.22	78.15.	1991	78.45
51.24.	1996	51.22, 51.23	78.16.	1991	78.46
51.64.	1989	51.69	78.17.	1991	78.47
51.84–51.88.	1989	51.97	78.18.	1991	78.48
51.97.	1986	52.91, 51.99, 51.82	78.19.	1991	78.49
51.98.	1986	51.99	78.20.	1991	78.10, 78.20, 78.30
52.13.	1989	51.97, 52.91	78.21.	1991	78.11, 78.31
52.14.	1989	52.11	78.22.	1991	78.12, 78.22, 78.32
52.21–52.22.	1989	52.2	78.23.	1991	78.13, 78.23, 78.33
52.84–52.86.	1996	99.29	78.24.	1991	78.14, 78.34
52.93.	1989	52.93 + 52.91	78.25.	1991	78.15, 78.25, 78.35
52.94.	1989	52.09	78.27.	1991	78.17, 78.27, 78.37
52.97–52.98.	1989	52.91	78.28.	1991	78.18, 78.38
52.99.	1989	52.93, 52.94, 52.99	78.29.	1991	78.11, 78.16, 78.19, 78.29, 78.39
54.24.	1987	54.23	78.39.	1991	78.31
54.25.	1993	54.98	78.90 ²	1987	78.40
54.51, 54.59.	1996	54.5	78.91 ²	1987	78.41
55.03–55.04.	1986	55.02	78.92 ²	1987	78.42
56.33–56.34.	1987	56.33	78.93 ²	1987	78.43
56.35.	1987	45.12	78.94 ²	1987	78.44
57.17–57.18.	1989	57.21	78.95 ²	1987	78.45
57.22.	1989	57.22, 57.82	78.96 ²	1987	78.46
58.31, 58.39.	1990	58.3	78.97 ²	1987	78.47
58.93.	1986	57.99	78.98 ²	1987	78.48
59.03.	1996	59.02	78.99 ²	1987	78.49
59.12.	1996	59.11	80.50–80.59.	1986	80.5
59.72.	1995	59.79	81.03.	1989	81.02
59.96.	1986	59.95	81.04–81.05.	1989	81.03, 81.04, 81.05
60.21, 60.29.	1995	60.2	81.06–81.07.	1989	81.06, 81.07
60.95.	1991	60.99	81.08.	1989	81.06, 81.07, 81.08
60.96–60.97.	2000	60.29	81.09.	1989	81.08
64.49.	1986	64.95	81.30–81.39.	2001	81.09
65.01, 65.09.	1996	65.0	81.40.	1989	81.69
65.13.	1996	65.12	81.51.	1989	81.51, 81.59
65.14.	1996	65.19	81.52.	1989	81.61–81.64
65.23.	1996	65.21	81.53.	1989	81.51, 81.59, 81.61–81.64
65.24.	1996	65.22	81.54–81.55.	1989	81.41
65.25.	1996	65.29	81.56.	1989	81.48
65.31, 65.39.	1996	65.3	81.57.	1989	81.31, 81.39
65.41, 65.49.	1996	65.4	81.59.	1989	81.39
65.53.	1996	65.51	81.61.	2002	81.00–81.08, 81.30–81.39
65.54.	1996	65.52	81.62–81.64.	2003	None
65.63.	1996	65.61	81.65–81.66.	2004	78.49
65.64.	1996	65.62	81.72.	1989	81.79
65.74.	1996	65.71	81.73–81.74.	1989	81.86 (Code deleted)
65.75.	1996	65.72	81.75.	1989	81.87 (Code deleted)
65.76.	1996	65.73	81.79.	1989	81.79, 81.87
65.81, 65.89.	1996	65.8	81.80.	1989	81.81
66.01.	1992	66.0	81.97.	1992	81.59
66.02.	1992	66.73	84.51–84.52.	2002	None
67.51, 67.59.	2001	67.5	84.53–84.54.	2004	78.30, 78.32–78.35, 78.37–78.39
68.15.	1987	68.14	84.55, 84.59.	2004	None
68.16.	1987	68.13	84.60–84.69.	2004	80.51
68.23.	1996	68.29	85.95–85.96.	1987	85.99
68.31, 68.39.	2003	68.3	86.06.	1987	86.09
68.51, 68.59.	1996	68.5	86.07.	1990	86.09
68.9.	1992	68.4	86.27.	1986	86.22–86.23
74.3.	1992	69.11 (Code deleted)	86.28.	1988	86.22
75.37.	1998	99.29	86.67.	1998	86.65
75.38.	2001	75.34			

Table VI. Changes in ICD-9-CM procedure codes, 1986–2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
86.93.....	1987	86.89	94.61–94.69	1989	94.25
86.94–86.96.....	2004	02.93, 03.93, 04.92	96.29.....	1998	96.39
88.90.....	1986	88.39	96.6.....	1986	96.35
88.91.....	1986	89.15	96.70–96.72.....	1991	93.92 (Code deleted)
88.92.....	1986	89.39	97.05.....	1989	51.97
88.93.....	1986	89.15	97.44.....	2001	37.64
88.94.....	1986	89.39	98.51–98.52, 98.59	1989	59.96 (Code deleted)
88.95.....	1986	89.29	99.00.....	1995	99.02
88.96.....	2002	88.91–88.97	99.10.....	1998	99.29
88.97.....	1989	88.99	99.15.....	1986	99.29
88.98.....	1989	88.90	99.20.....	1998	99.29
88.99.....	1986	89.39	99.28.....	1994	99.25
89.10, 89.19.....	1989	89.15	99.71–99.79 ³	1988	99.07
89.17–89.18.....	1988	89.15	99.75.....	2000	99.29
89.49.....	2004	89.59	99.76.....	2002	99.79
89.50.....	1991	89.54	99.77.....	2002	None
89.60.....	2002	89.65	99.78.....	2004	99.71
92.3.....	1995	01.59, 04.07, 07.63, 07.68	99.85.....	1987	93.35
92.30–92.33, 92.39.....	1998	92.3	99.86.....	1987	93.39
93.90.....	1988	93.92	99.88.....	1988	99.83

¹Before October 1986, contents of current code 36.05 would have been assigned to 36.0.²Codes 78.90–78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90–78.99 were reassigned to codes 78.40–78.49.³Codes 99.71–99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71–99.79 were reclassified as "Therapeutic apheresis".NOTE: ICD-9-CM is the *International Classification of Diseases, Ninth Revision, Clinical Modification*.

Vital and Health Statistics series descriptions

- SERIES 1. **Programs and Collection Procedures**—These reports describe the data collection programs of the National Center for Health Statistics. They include descriptions of the methods used to collect and process the data, definitions, and other material necessary for understanding the data.
- SERIES 2. **Data Evaluation and Methods Research**—These reports are studies of new statistical methods and include analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. These studies also include experimental tests of new survey methods and comparisons of U.S. methodology with those of other countries.
- SERIES 3. **Analytical and Epidemiological Studies**—These reports present analytical or interpretive studies based on vital and health statistics. These reports carry the analyses further than the expository types of reports in the other series.
- SERIES 4. **Documents and Committee Reports**—These are final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- SERIES 5. **International Vital and Health Statistics Reports**—These reports are analytical or descriptive reports that compare U.S. vital and health statistics with those of other countries or present other international data of relevance to the health statistics system of the United States.
- SERIES 6. **Cognition and Survey Measurement**—These reports are from the National Laboratory for Collaborative Research in Cognition and Survey Measurement. They use methods of cognitive science to design, evaluate, and test survey instruments.
- SERIES 10. **Data From the National Health Interview Survey**—These reports contain statistics on illness; unintentional injuries; disability; use of hospital, medical, and other health services; and a wide range of special current health topics covering many aspects of health behaviors, health status, and health care utilization. They are based on data collected in a continuing national household interview survey.
- SERIES 11. **Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey**—Data from direct examination, testing, and measurement on representative samples of the civilian noninstitutionalized population provide the basis for (1) medically defined total prevalence of specific diseases or conditions in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analyses of trends and relationships among various measurements and between survey periods.
- SERIES 12. **Data From the Institutionalized Population Surveys**—Discontinued in 1975. Reports from these surveys are included in Series 13.
- SERIES 13. **Data From the National Health Care Survey**—These reports contain statistics on health resources and the public's use of health care resources including ambulatory, hospital, and long-term care services based on data collected directly from health care providers and provider records.
- SERIES 14. **Data on Health Resources: Manpower and Facilities**—Discontinued in 1990. Reports on the numbers, geographic distribution, and characteristics of health resources are now included in Series 13.
- SERIES 15. **Data From Special Surveys**—These reports contain statistics on health and health-related topics collected in special surveys that are not part of the continuing data systems of the National Center for Health Statistics.
- SERIES 16. **Compilations of Advance Data From Vital and Health Statistics**—Advance Data Reports provide early release of information from the National Center for Health Statistics' health and demographic surveys. They are compiled in the order in which they are published. Some of these releases may be followed by detailed reports in Series 10–13.
- SERIES 20. **Data on Mortality**—These reports contain statistics on mortality that are not included in regular, annual, or monthly reports. Special analyses by cause of death, age, other demographic variables, and geographic and trend analyses are included.
- SERIES 21. **Data on Natality, Marriage, and Divorce**—These reports contain statistics on natality, marriage, and divorce that are not included in regular, annual, or monthly reports. Special analyses by health and demographic variables and geographic and trend analyses are included.
- SERIES 22. **Data From the National Mortality and Natality Surveys**—Discontinued in 1975. Reports from these sample surveys, based on vital records, are now published in Series 20 or 21.
- SERIES 23. **Data From the National Survey of Family Growth**—These reports contain statistics on factors that affect birth rates, including contraception, infertility, cohabitation, marriage, divorce, and remarriage; adoption; use of medical care for family planning and infertility; and related maternal and infant health topics. These statistics are based on national surveys of women of childbearing age.
- SERIES 24. **Compilations of Data on Natality, Mortality, Marriage, Divorce, and Induced Terminations of Pregnancy**—These include advance reports of births, deaths, marriages, and divorces based on final data from the National Vital Statistics System that were published as supplements to the *National Vital Statistics Report* (NVSR). These reports provide highlights and summaries of detailed data subsequently published in *Vital Statistics of the United States*. Other supplements to the NVSR published here provide selected findings based on final data from the National Vital Statistics System and may be followed by detailed reports in Series 20 or 21.

For answers to questions about this report or for a list of reports published in these series, contact:

Information Dissemination Staff
National Center for Health Statistics
Centers for Disease Control and Prevention
3311 Toledo Road, Room 5412
Hyattsville, MD 20782
1-866-441-NCHS (6247)
E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchs

**U.S. DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
3311 Toledo Road
Hyattsville, MD 20782

MEDIA MAIL
POSTAGE & FEES PAID
CDC/NCHS
PERMIT NO. G-284